



Published in final edited form as:

Can J Anaesth. 2012 August ; 59(8): 798–804. doi:10.1007/s12630-012-9737-1.

A brief review of practical preoperative cognitive screening tools

L. Stephen Long, M.D., William A. Shapiro, M.D., and Jacqueline M. Leung, M.D.

Department of Anesthesia & Perioperative Care, University of California, San Francisco 521 Parnassus Avenue, Room C-450 San Francisco, CA 94143-0648

Abstract

Purpose—Preoperative cognitive impairment is associated with the development of postoperative delirium, a common and consequential complication of major surgery in older patients. Screening for cognitive impairment should become a routine part of the preoperative evaluation of older patients. However, its implementation is hampered by limited clinical time and resources. The objective of this review was to identify cognitive screening tools that could be easily incorporated into the evaluation of older patients before major surgery.

Search Strategy—Using strict inclusion and exclusion criteria, we searched PubMed over a 15-year period for short and simple cognitive screening tools. In addition, studies examining the tools in a perioperative environment were reviewed.

Search Results—6 cognitive screening tools were identified. Each tool had an administration time of 2.5 minutes or less. Among the tools, sensitivity for cognitive impairment ranged from 79% to 99%, while specificity ranged from 70% to 98%. Of the 6 tools identified, only one (Mini-Cog) had been tested in a perioperative environment.

Conclusions—Incorporating a cognitive screening assessment into the preoperative evaluation of older patients is feasible. More research is needed to validate cognitive screening tools in the perioperative setting.

Introduction

Postoperative delirium is an acute and fluctuating confusional state (1, 2). Its incidence is high, affecting 40-50% of older patients after major surgery (3-6). The development of delirium impacts both short- and long-term outcomes, including a higher rate of complications after surgery, longer hospital stays, increased likelihood of discharge to long-term care facilities, and increased mortality (7-9).

Multiple preoperative risk factors for postoperative delirium have been identified. These include cognitive impairment, sensory impairment, older age, American Society for Anesthesiology (ASA) classification, low education level, psychotropic drug use, poor functional status, dehydration, medical comorbidities (especially cerebrovascular or other brain disease), electrolyte abnormalities, low albumin, and depression (3, 4, 10-25). Of these, pre-existing cognitive impairment [e.g., dementia or mild cognitive impairment (MCI)] is often cited as the strongest predictor of postoperative delirium (26) (see Table 1 for descriptions of different types of cognitive impairment).

Corresponding Author: L. Stephen Long, M.D. (415) 476-2131 longsl@anesthesia.ucsf.edu.

Conflict of Interest: None

Authors' Attestations: Drs. Long and Leung contributed to review design, data analysis, conduct of review, and manuscript preparation. Dr. Shapiro contributed to manuscript preparation.

It is important to recognize that “cognitive impairment” is a nonspecific term that can be used for a variety of cognitive problems. We believe screening for cognitive impairment before surgery should be thought of as screening for pre-existing dementia or MCI. Dementia and MCI are specific, well-described conditions used to classify chronic cognitive impairment, as detailed in Table 1. Accordingly, most outpatient cognitive screening tools are designed to detect one or both of these conditions.

Currently, routine preoperative evaluation does not include the evaluation of baseline cognitive functioning. Clinicians have considered “alert and oriented times three” to be “normal” cognitive status. In fact, this assumption has not been proven, and patients who do not have documented dementia may in fact have MCI. Accordingly, knowing the cognitive status of patients before surgery is critical for risk stratification to allow for subsequent prophylaxis, surveillance, and treatment. Indeed, a number of promising interventions and therapies have been proposed for postoperative delirium, including perioperative neuroleptic prophylaxis (27-32), reduction in sedative dosing (33), improved postoperative pain control (6, 10), and proactive postoperative nonpharmacologic management (5).

Previous investigators have examined the use of a variety of cognitive assessment tools to measure preoperative cognitive impairment. The most popular tool used for clinical research is the Mini-Mental State Examination (MMSE), developed by Folstein *et al.* (34). Despite its strengths and wide use, the MMSE may not be practical for preoperative cognitive screening due to its length of 7-10 minutes. Indeed, even a 5-minute test can add a substantial amount of work when implemented in a high-volume preoperative clinic. In addition, the MMSE has been criticized for copyright restrictions and age- and education-related biases (35).

Thus, an ideal preoperative cognitive screening tool should: (1) require a very short amount of time for administration and scoring; (2) detect cognitive impairment with moderately high sensitivity and specificity; (3) be validated in a preoperative geriatric sample.

The goal of this paper is to identify cognitive screening tools suitable for the preoperative setting.

Search Strategy

A literature search of PubMed was conducted. The database was searched over a 15-year period, from January 1, 1996 to January 1, 2011. A 15-year period was chosen to limit the search to contemporary tools only. The following search terms were used: *screen* or *screeener* or *test* or *tool* or *measure* or *instrument* or *assessment* or *battery* combined with *dementia* or *cognitive* combined with *quick* or *brief* or *short* combined with *Mini-Mental State Examination* or *MMSE*. Abstracts were limited to English language only. Inclusion criteria specified that the tool: (1) assesses at least two distinct domains of cognitive function (i.e., multidomain tools only); (2) has a mean administration time of 2.5 minutes or less in non-demented subjects; (3) has been developed in a preoperative, primary care, or community sample; (4) has been tested against or developed with the MMSE; and (5) has been developed on subjects 60 years or older. Studies were excluded for inadequate data (e.g., studies that did not report sensitivity or specificity). Tools requiring informant interviews or self-administration were also excluded.

After identifying cognitive screening tools using the aforementioned criteria, a second PubMed search was performed for each tool. All published English language abstracts were reviewed for each individual tool from January 1, 1996 to January 1, 2011. Studies examining the tools in perioperative settings were included.

Search Results

The preliminary search identified 513 abstracts. Most of the abstracts did not directly examine cognitive screening tools. Thirty-one abstracts were primary studies of multidomain cognitive screening tools. Twenty-five abstracts were excluded (19 due to length of administration, 3 for self-administration, 2 for inappropriate study samples, and 1 for informant interview). The remaining 6 abstracts described primary studies of cognitive screening tools (see Table 2 for a summary).

6-Item Screener (6-IS) (41)

The 6-IS consists of a 3-item recall (e.g., apple, table, penny) and a 3-item temporal orientation (day of the week, month, year). Each correct response earns 1 point for a total of 6 points. Administration time is 1 minute, not including the delay for the recall component.

The 6-IS was published in 2002 with the purpose to quickly screen for cognitive impairment with “an acceptable sensitivity and specificity” for dementia and mild cognitive impairment. The tool was developed in a community sample of 344 geriatric subjects, and then validated in a cohort of 651 referrals to an Alzheimer’s center. A geriatric psychiatrist or neurologist first evaluated subjects for MCI and dementia, then subjects were screened with both the 6-IS and MMSE. Using a 6-IS cutoff score of 3 points or less, the sensitivity and specificity for dementia was 88.7% and 88.0%, respectively, while the MMSE using a cutoff score of 23 produced values of 95.2% and 86.7%, respectively, in the community sample. At a cutoff score of 4 points or less, the 6-IS demonstrated a sensitivity and specificity for MCI of 74.2% and 80.2%, respectively, and at 5 points or less, 97.7% and 49.2%, respectively.

8-Item Screener (8-IS) (42)

The 8-IS employs a 3-item recall (e.g., bicycle, red, happiness) and an attention/calculation exercise, whereby subjects subtract 7 from 100 serially for 5 iterations (serial 7s). One point is awarded for each correct answer, totaling 8 points. It can be completed in 2 minutes or less.

The 8-IS was published in 2011 with a goal to rapidly screen for dementia in primary care clinics using 8 of the items that are included within the MMSE. The tool was developed in a cohort of 188 seniors from a geriatric clinic. Subjects were first screened with the complete MMSE (30 points). Those with scores less than 24 or 20, depending on education level, were referred to a neurologist for formal evaluation for dementia. The authors then calculated the sensitivity and specificity for dementia using only 8 points from the MMSE (recall plus attention/calculation). Scores of 6 or less produced a sensitivity and specificity for dementia of 94.9% and 59.1%, respectively, and these values changed to 85.9% and 78.2%, respectively, for scores of 5 or less. The authors advocate different 8-IS cutoff values for people with lower levels of education.

6-Item Cognitive Impairment Test (6-CIT) (43)

The 6-CIT involves a 3-item temporal orientation (year, month, time within 1 hour), a 5-item address recall (John, Brown, 42, West Street, Bedford), and two attention exercises (count backwards 20 to 1, say months in reverse order). Each incorrect response is given 1 point, and a formula is used to generate a weighted score. It can be completed in 1-2 minutes.

The 6-CIT was developed in 1999 for “usage as a screening tool” in primary care. The tool was tested against the MMSE in a sample of 287 geriatric subjects from England: 135 non-demented controls and 152 subjects who carried a previous diagnosis of dementia, selected

from both the community and outpatient settings. All subjects received the 6-CIT and MMSE. The 6-CIT was found to correlate strongly with the MMSE ($r^2=-0.911$). In addition, at a cutoff score of 6 points or higher, the 6-CIT produced a sensitivity and specificity for dementia of 92.1% and 95.6%, respectively, while the MMSE produced values of 78.6% and 100.0%, respectively, at a cutoff of 23 points or less.

The Sweet 16 (S-16) (44)

The S-16 includes 8 temporal/spatial orientation questions (i.e., orientation to time and place), 3 registration questions (i.e., immediate repetition of 3 items), 2 sustained attention questions (i.e., digit spans backward), and a 3-item recall, for a total of 16 points. The instrument and instruction manual are available at <http://hospitalelderlifeprogram.org>. In the pilot group study, the mean administration time was 2 minutes.

The S-16 was published in 2011 as an alternative to current cognitive screeners that are “underused, lack sensitivity, or may be restricted by copyright laws.” The tool was developed in 774 geriatric subjects who were recently hospitalized and then validated in 709 subjects who were randomly selected from a large national sample. An expert panel of clinicians assigned the diagnosis of dementia using DSM criteria. The performance of the S-16 was compared directly to that of the MMSE. The two instruments correlated well with $r^2=0.94$. At a cutoff score of 13 points or less, the S-16 demonstrated a sensitivity of 99% and specificity of 70% for dementia. The sensitivity and specificity for the MMSE at 23 points or less were 87% and 89%, respectively.

5-Item Recall and Fluency (5-IRF) (45)

The 5-IRF consists of a 5-item address recall (John, Brown, 42, Market Street, Chicago) and a 1-minute verbal fluency for animals (i.e., name as many different animals as possible in 1 minute). The tool is scored by counting the number recall errors and the number of animals named; 3 or more recall errors or 8 or fewer animals named correlates with dementia. It has an administration time of less than 2 minutes.

The tool was developed in 2005 to screen for dementia in patients with memory complaints. The authors retrospectively analyzed 2 geriatric cohorts. The first cohort consisted of 97 demented subjects (diagnosed using DSM criteria) matched with non-demented controls 1:1. The second cohort was comprised of 159 demented subjects (diagnosed using clinical criteria for dementia due to Alzheimer’s disease) matched 1:1 with non-demented controls. Subjects were screened for cognitive impairment. The cohorts were combined to allow for greater statistical power. The 5-IRF achieved a sensitivity of 79% and specificity of 98% for dementia using cutoff scores of 3 or more errors on the 5-item recall test and 8 animals or less on the verbal fluency test. At the same specificity (98%), the MMSE generated a sensitivity of 53% for dementia.

Mini-Cog (46)

The Mini-Cog is composed of a 3-item recall and a clock drawing task. One point is awarded for each correctly recalled word. The clock drawing is scored as normal if the clock has the correct time and is grossly normal. Recall scores of 0 irrespective of clock drawing score and recall scores of 1-2 with an abnormal clock drawing score correlate with dementia. In the pilot study, non-demented subjects required an average of 2.5 minutes for completion, whereas demented subjects took 3.7 minutes.

The Mini-Cog was published in 2000 for “discriminating demented from non-demented persons” in a diverse, geriatric community sample. The tool was developed on a multicultural, multilingual sample of 249 older adults, who were first classified as demented

or non-demented using formal diagnostic criteria. The subjects were then given the Mini-Cog and MMSE. The Mini-Cog's sensitivity (99%) and specificity (93%) for dementia were found to be higher than the MMSE (91% and 92%, respectively).

Two studies examining the Mini-Cog in a perioperative setting were identified. The first study sought to determine preoperative risk factors for the development of postoperative delirium in older patients scheduled for a major surgery. One hundred forty four subjects were studied, and 64 (44%) developed postoperative delirium. Subjects had received baseline cognitive and functional assessments preoperatively. The Mini-Cog was used to screen for pre-existing cognitive impairment, and the authors found that this factor was the most robust predictor of postoperative delirium (4).

The purpose of the second study was to identify preoperative variables associated with 6-month mortality after major surgery in older adults. One hundred ten subjects were studied, and the 6-month mortality was 15% (16 subjects). The Mini-Cog was used for preoperative cognitive assessment, and abnormal scores were shown to be significantly associated with 6-month postoperative mortality (47).

Discussion

The aim of this review was to identify practical screening tools that could be used to detect preoperative cognitive impairment in a clinical setting. During this review, we identified 6 screening tools which can be administered in 2.5 minutes or less. We believe any one of these tools could be used in a time-constrained preoperative environment.

Despite the similar lengths of administration, the tools differed in their ability to screen for cognitive impairment. The best screening tools were the S-16 and Mini-Cog, each with a sensitivity of 99% for dementia in their respective study populations. The Mini-Cog, which was also the only tool found to be tested in the perioperative environment, generated a higher specificity (93%) for dementia compared to the S-16 (70%). However, we wish to stress that direct comparison of sensitivities and specificities between tools is restricted by the diversity of methodology among the studies reviewed.

It should be noted that cognitive impairment detected by these tools is not diagnostic for dementia or MCI, but rather a screen only (these tools are also not designed to diagnose postoperative delirium). Accordingly, it would be prudent to discuss a positive screening with the patient and family to ensure referral to a primary care physician, neurologist, or psychiatrist for further evaluation before or after surgery. In addition, the possible significance of a positive screening should be discussed with the patient prior to the start of screening for cognitive impairment. Furthermore, if the screening is positive, proactive consultation with perioperative providers including surgeons, anesthesiologists, nurses, and pharmacists may be warranted to provide a strategy for delirium surveillance and possible care modification.

There are limitations to this review. First, none of the tools was designed specifically for surgical patients in a preoperative setting. Moreover, despite ample evidence linking preoperative cognitive impairment (usually detected by the MMSE) to postoperative delirium, only one of the tools (Mini-Cog) has been studied for preoperative risk stratification of postoperative delirium, and no tool has been studied extensively for this purpose. It is important to note that surgical patients may have unique characteristics affecting their cognition, including pain and anxiety, and medications to treat either or both. These factors may not be prevalent in non-surgical community dwellers.

Another potential limitation of the review is the strict inclusion and exclusion criteria. Our goal was to find cognitive screening tools that would be easy to incorporate into a time-constrained preoperative evaluation. Thus, we sought the simplest and shortest tools available. We acknowledge that screening tools with a longer administration time were excluded, and that some of these tools may be more comprehensive and therefore preferred by some clinicians.

A final limitation of this review is the inclusion of the MMSE criterion. This criterion was selected because of the MMSE's historical importance and widespread presence in published literature. However, the MMSE criterion likely reduced the number of tools identified in our search. We wish to emphasize that a tool need not be compared against the MMSE for it to be suitable for preoperative cognitive screening.

In summary, this review offers a starting point for preoperative cognitive screening, which, as pointed out by a recent editorial, should become a routine part of the evaluation of older patients before major surgery (48). Only after assessing cognitive function at baseline can we further understand how cognitive changes occur after anesthesia and surgery, and potentially intervene to mitigate these changes. Future studies are critically needed to prospectively validate the utility of cognitive screening tools as a means to identify patients at risk for postoperative delirium.

Acknowledgments

Financial Support: This project was supported in part by the National Institute of Aging, National Institutes of Health, Bethesda, MD, Grant # NIH 1R01AG031795-03 (Leung).

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th ed. American Psychiatric Association; Washington, DC: 2000.
2. Wong CL, Holroyd-Leduc J, Simel DL, Straus SE. Does This Patient Have Delirium? Value of Bedside Instruments. *JAMA*. 2010; 304(7):779–786. [PubMed: 20716741]
3. Leung JM, Sands LP, Mullen EA, Wang Y, Vaurio L. Are preoperative depressive symptoms associated with postoperative delirium in geriatric surgical patients? *J Gerontol*. 2005; 60A(12): 1563–1568.
4. Robinson TN, Baeburn CD, Tran ZV, Angles EM, Brenner LA, Moss M. Postoperative delirium in the elderly: Risk factors and outcomes. *Ann Surg*. 2009; 249(1):173–178. [PubMed: 19106695]
5. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc*. 2001; 49(5):516–522. [PubMed: 11380742]
6. Vaurio LE, Sands LP, Wang Y, Mullen EA, Leung JM. Postoperative delirium: the importance of pain and pain management. *Anesth Analg*. 2006; 102(4):1267–1273. [PubMed: 16551935]
7. Inouye SK. Delirium in older persons. *N Engl J Med*. 2006; 354(11):1157–1165. [PubMed: 16540616]
8. Marcantonio ER, Flacker JM, Michaels M, Resnick NM. Delirium is independently associated with poor functional recover after hip fracture. *J Am Geriatr Soc*. 2000; 48(6):618–624. [PubMed: 10855596]
9. Kiely DK, Marcantonio ER, Inouye SK, Shaffer ML, Bergmann MA, Yang FM, Fearing MA, Jones RN. Persistent delirium predicts greater mortality. *J Am Geriatr Soc*. 2009; 57(1):55–61. [PubMed: 19170790]
10. Morrison RS, Magaziner J, Gilbert M, Koval KJ, McLaughlin MA, Orosz G, Strauss E, Siu AL. Relationship between pain and opioid analgesics on the development of delirium following hip fracture. *J Gerontol*. 2003; 58(1):76–81.

11. Kalisvaart KJ, Vreeswijk R, de Jonghe JFM, van der Ploeg T, van Gool WA, Eikelenboom P. Risk factors and prediction of postoperative delirium in elderly hip-surgery patients: Implementation and validation of a medical risk factor model. *JAGS*. 2006; 54:817–822.
12. Brouquet A, Cudennec T, Benoist S, Moulias S, Beauchet A, Penna C, Teillet L, Nordlinger B. Impaired mobility, ASA status, and administration of tramadol are risk factors for postoperative delirium in patients aged 75 years or more after major abdominal surgery. *Ann Surg*. 2010; 251(4): 759–765. [PubMed: 20224380]
13. Marcantonio ER, Goldman L, Mangione CM, Ludwig LE, Muraca B, Haslauer CM, Donaldson MC, Whittlemore AD, Sugarbaker DJ, Poss R. A clinical prediction rule for delirium after elective noncardiac surgery. *JAMA*. 1994; 271(2):134–139. [PubMed: 8264068]
14. Litaker D, Locala J, Franco K, Bronson DL, Tannous Z. Preoperative risk factors for postoperative delirium. *General Hospital Psychiatry*. 2001; 23:84–89. [PubMed: 11313076]
15. Galanakis P, Bickel H, Gradinger R, Von Gumpengert S, Forstl H. Acute confusion state in the elderly following hip surgery: Incidence risk factors and complications. *Int J Geriatr Psychiatry*. 2001; 16(4):349–355. [PubMed: 11333420]
16. Dai YT, Lou MF, Yip PK, Huang GS. Risk factors and incidence of postoperative delirium in elderly Chinese patients. *Gerontology*. 2000; 46(1):28–35. [PubMed: 11111226]
17. Schneider F, Bohner H, Habel U, Salloum JG, Stierstorfer A, Hummel TC, Miller C, Friedrichs R, Muller EE, Sandmann W. Risk factors for postoperative delirium in vascular surgery. *Gen Hosp Psychiatry*. 2002; 24(1):28–34. [PubMed: 11814531]
18. Fisher BW, Flowerdew G. A simple model for predicting postoperative delirium in older patients undergoing elective orthopedic surgery. *J Am Geriatr Soc*. 1995; 43(2):175–178. [PubMed: 7836644]
19. Sasajima Y, Sasjima T, Uchida H, Kawai S, Haga M, Akasaka N, Kusakabe M, Inaba M, Goh K, Yamamoto H. Postoperative delirium in patients with chronic lower limb ischaemia: What are the specific markers? *Eur J Vasc Endovasc Surg*. 2000; 20:132–137. [PubMed: 10942684]
20. Schuurmans MJ, Duursma SA, Shortridge-Baggett LM, Clevers GJ, Pel-Littel R. Elderly patients with a hip fracture: the risk for delirium. *App Nurs Res*. 2003; 16(2):75–84.
21. Bohner H, Hummel TC, Habel U, Miller C, Reinbott S, Yang Q, Gabriel A, Friedrichs R, Muller EE, Ohmann C, Sandmann W, Schneider F. Predicting delirium after vascular surgery: a model based on pre- and intraoperative data. *Ann Surg*. 2003; 238(1):149–156. [PubMed: 12832977]
22. Dupplis GS, Wikblad K. Acute confusional states in patients undergoing hip surgery: a prospective observation study. *Gerontology*. 2000; 46(1):36–43. [PubMed: 11111227]
23. Kazmierski J, Kowman M, Banach M, Pawelczyk T, Okonski P, Iwaszkiewicz A, Zaslonka J, Sobow T, Kloszewska I. Preoperative predictors of delirium after cardiac surgery: a preliminary study. *Gen Hosp Psychiatry*. 2006; 28(6):536–538. [PubMed: 17088170]
24. Kazmierski J, Kowman M, Banach M, Pawelczyk T, Okonski P, Iwaszkiewicz A, Zaslonka J, Sobow T, Kloszewska I. Incidence and predictors of delirium after cardiac surgery: results from The IPDACS Study. *J Psychosom Res*. 2010; 69(2):179–185. [PubMed: 20624517]
25. Freter SH, Dunbar MJ, MacLeod H, Morrison M, MacKnight C, Rockwood K. Predicting postoperative delirium in elective orthopaedic patients: the Delirium Elderly At-Risk (DEAR) instrument. *Age and Ageing*. 2005; 34:169–171. [PubMed: 15713861]
26. Dasgupta M, Dumbrell AC. Preoperative risk assessment for delirium after noncardiac surgery: A systematic review. *JAGS*. 2006; 54:1578–1589.
27. Larsen KA, Kelly SE, Stern TA, Bode RH Jr, Price LL, Hunter DJ, Gulczynski D, Bierbaum BE, Sweeney GA, Hoikala KA, Cotter JJ, Potter AW. Administration of olanzapine to prevent postoperative delirium in elderly joint-replacement patients: a randomized, controlled trial. *Psychosomatics*. 2010; 51(5):409–418. [PubMed: 20833940]
28. Kaneko T, Jianhi C, Ishikura T, Kobayashi M, Naka T, Kaibara N. Prophylactic consecutive administration of haloperidol can reduce the occurrence of postoperative delirium in gastrointestinal surgery. *Yonaga Acta Medica*. 1999; 42:179–184.
29. Kalisvaart KJ, de Jonghe JFM, Bogaards MJ, Vreeswijk R, Egberts TC, Burger BJ, Eikelenboom P, van Gool WA. Haloperidol prophylaxis for elderly hip-surgery patients at risk for delirium: a

- randomized placebo-controlled study. *J Am Geriatr Soc.* 2005; 53:1658–1666. [PubMed: 16181163]
30. Schrader SL, Wellik KE, Demaerschalk BM, Caselli RJ, Woodruff BK, Wingerchuk DM. Adjunctive haloperidol prophylaxis reduces postoperative delirium severity and duration in at-risk elderly patients. *Neurologist.* 2008; 14:134–137. [PubMed: 18332845]
 31. Prakanrattana U, Prapaitrakool S. Efficacy of risperidone for prevention of postoperative delirium in cardiac surgery. *Anaesth Intensvie Care.* 2007; 35:714–719.
 32. Wang W, Li HL, Wang DX, Zhu X, Li SL, Yao GQ, Chen KS, Gu XE, Zhu SN. Haloperidol prophylaxis decreases delirium incidence in elderly patients after noncardiac surgery: A randomized controlled trial. *Crit Care Med.* 2012; 40(3):731–739. [PubMed: 22067628]
 33. Sieber FE, Zakriya KJ, Gottschalk A, Blute MR, Lee HB, Rosenberg PB, Mears SC. Sedation depth during spinal anesthesia and the development of postoperative delirium in elderly patients undergoing hip fracture repair. *May Clin Proc.* 2010; 85(1):18–26.
 34. Folstein MF, Folstein SE, McHugh PR. “Mini-Mental State”: a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975; 12:189–198. [PubMed: 1202204]
 35. Holsinger T, Deveau J, Boustani M, Williams JW. Does this patient have dementia? *JAMA.* 2007; 297(21):2391–2404. [PubMed: 17551132]
 36. Petersen RC, Stevens JC, Ganguli M, Tangalos EG, Cummings JL, DeKosky ST. Practice parameter: early detection of dementia: mild cognitive impairment (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology.* 2001; 56:1133–1142. [PubMed: 11342677]
 37. Hess TM. Memory and aging in context. *Psychol Bull.* 2005; 131:383–406. [PubMed: 15869334]
 38. Grady CL, Craik FI. Changes in memory processing with age. *Curr Opin Neurobiol.* 2000; 10:224–231. [PubMed: 10753795]
 39. Terrando N, Brzezinski M, Degos V, Eriksson LI, Kramer JH, Leung JM, Miller BL, Seeley WW, Vacas S, Weiner MW, Yaffe K, Young WL, Xie Z, Maze M. Perioperative cognitive decline in the aging population. *Mayo Clin Proc.* 2011; 86:885–893. [PubMed: 21878601]
 40. Tsai TL, Sands LP, Leung JL. An update on postoperative cognitive dysfunction. *Adv Anesth.* 2010; 28:269–284. [PubMed: 21151735]
 41. Callahan CM, Unverzagt FW, Hiu SL, Perkins AJ, Hendrie HC. Six-Item Screener to identify cognitive impairment among potential subjects for clinical research. *Medical Care.* 2002; 40(9): 771–781. [PubMed: 12218768]
 42. Chen CY, Leung KK, Chen CY. A quick dementia screening tool for primary care physicians. *Arch Gerontol Geriatr.* 2011; 53(1):100–103. Epub 2010 Jul 16.
 43. Brooke P, Bullock R. Validation of a 6 item cognitive impairment test with a view to primary care usage. *Int J Geriatr Psychiatry.* 1999; 14:936–940.
 44. Fong TG, Jones RN, Rudolph JL, Yang FM, Tommet D, Habtemariam D, Marcantonio ER, Langa KM, Inouye SK. Development and validation of a brief cognitive assessment tool. *Arch Intern Med.* 2011; 171(5):432–437. Epub 2010 Nov 8. [PubMed: 21059967]
 45. Kilada S, Gamaldo A, Grant EA, Moghekar A, Morris JC, O’Brien RJ. Brief screening tests for the diagnosis of dementia: Comparison with the Mini-Mental State Exam. *Alzheimer Dis Assoc Disord.* 2005; 9(1):8–16. [PubMed: 15764865]
 46. Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The Mini-Cog: A cognitive ‘vital signs’ measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry.* 2000; 15:1021–1027. [PubMed: 11113982]
 47. Robinson TN, Eiseman B, Wallace JI, Church SD, McFann KK, Pfister SM, Sharp TJ, Moss M. Redefining geriatric preoperative assessment using frailty, disability, and co-morbidity. *Ann Surg.* 2009; 250(3):449–453. [PubMed: 19730176]
 48. Crosby G, Culley DJ, Hyman BT. Preoperative cognitive assessment of the elderly surgical patient: A call for action. *Anesthesiology.* 2011; 114(6):1265–1268. [PubMed: 21490501]

Implication Statement

Postoperative delirium is a common complication after major surgery in older patients, and preoperative cognitive impairment is an important risk factor. This review describes brief cognitive screening tools that may be used to screen patients for cognitive impairment before surgery.

Table 1

Different Types of Cognitive Impairment

Types of Cognitive Impairment	Definitions
Delirium	<i>Delirium</i> is an acute and fluctuating confusional state characterized by impaired attention, perception, and cognition (1). It is usually triggered by an acute illness (e.g., urinary tract infection) or intervention (e.g., surgery, drug administration) and is reversible.
Dementia	<i>Dementia</i> is a combination of cognitive deficits that are chronic and nonfluctuating in nature. It always involves memory impairment (1). The most common causes are Alzheimer's disease and cerebrovascular disease. Dementia is viewed as a non-reversible disease.
Mild Cognitive Impairment (MCI)	<i>MCI</i> is the presence of memory impairment not explained by normal aging and not severe enough to meet criteria for dementia. MCI predicts progression to dementia, with an annual conversion rate reported as high as 25% (36).
Normal Aging	<i>Normal aging</i> results in cognitive changes not considered pathologic. Importantly, these individuals are able to retain learned information nearly as well as their younger counterparts and experience no functional deficits (37, 38).
Postoperative Cognitive Decline (POCD)	<i>POCD</i> is the precipitous worsening of cognitive function after surgery. The duration of POCD is on the order of months to years. At present, POCD is not coded as a disease despite garnering significant interest from researchers and clinicians (39, 40).

Table 2

Comparison of Cognitive Impairment Screening Tools

Tool	Setting	N	Average age (years)	Items	Admin. Time (min)	Scoring	Sn for dementia	Sp for dementia	Tested in a perioperative setting?
6-IS	Community and demented samples, USA	Cohort 1: 344 Cohort 2: 651	Cohort 1: 74.4 Cohort 2: 69.6	3 recall 3 orient	1	Simple adding	88.7%	88.0%	No
8-IS	Geriatric clinic sample, Taiwan	188	77.0	3 recall 5 serial 7s	<2	Simple adding	94.9% or 85.9%, depending on cutoff	59.1% or 78.2%, depending on cutoff	No
6-CIT	Community and 2 demented samples, UK	Cohort 1: 135 Cohort 2: 70 Cohort 3: 82	Cohort 1: 68.1 Cohort 2: 73.8 Cohort 3: 81.7	5 recall 3 orient	1-2	Rubric required	92.1%	95.6%	No
S-16	Post-acute hospitalization and national registry samples, USA	Cohort 1: 774 Cohort 2: 709	Cohort 1: 83.1 Cohort 2: 78.8	3 recall 8 orient 3 registers 2 digit spans	2	Simple adding	99%	70%	No
5-IRF	Community and demented samples, USA	Cohort 1: 194 Cohort 2: 318	Cohort 1: Demented: 84.2 Non-demented: 83.8 Cohort 2: Demented: 78.5 Non-demented: 78.0	5 recall Verbal fluency	<2	Simple adding	79%	98%	No
Mini-Cog	Community sample, USA	249	Demented: 77.9 Non-demented: 69.0	3 recall Clock drawing	2.5	Simple algorithm	99%	93%	Yes

Sn = sensitivity, Sp = specificity