

## *Cryptosporidium hominis* Infection Diagnosed by Real-Time PCR-RFLP

Hyeng-Il Cheun<sup>1,2</sup>, Kyungjin Kim<sup>3</sup>, Sejong Yoon<sup>3</sup>, Won-Ja Lee<sup>1</sup>, Woo-Yoon Park<sup>4</sup>, Seobo Sim<sup>3</sup> and Jae-Ran Yu<sup>3,\*</sup>

<sup>1</sup>Department of Malaria and Parasitic Diseases, National Institute of Health, Chungcheongbuk-do 363-951, Korea; <sup>2</sup>Division of Epidemic Intelligence Service, Korea Centers for Disease Control & Prevention, Chungcheongbuk-do 363-951, Korea; <sup>3</sup>Department of Environmental and Tropical Medicine, Konkuk University, School of Medicine, Seoul 143-701, Korea; <sup>4</sup>Department of Radiation Oncology, College of Medicine, Chungbuk National University, Cheongju 361-711, Korea

**Abstract:** There are approximately 20 known species of the genus *Cryptosporidium*, and among these, 8 infect immunocompetent or immunocompromised humans. *C. hominis* and *C. parvum* most commonly infect humans. Differentiating between them is important for evaluating potential sources of infection. We report here the development of a simple and accurate real-time PCR-based restriction fragment length polymorphism (RFLP) method to distinguish between *C. parvum* and *C. hominis*. Using the *CP2* gene as the target, we found that both *Cryptosporidium* species yielded 224 bp products. In the subsequent RFLP method using *TaqI*, 2 bands (99 and 125 bp) specific to *C. hominis* were detected. Using this method, we detected *C. hominis* infection in 1 of 21 patients with diarrhea, suggesting that this method could facilitate the detection of *C. hominis* infections.

**Key words:** *Cryptosporidium parvum*, *Cryptosporidium hominis*, real-time PCR, RFLP, *TaqI*

*Cryptosporidium parvum* is a parasitic protozoan that infects gastrointestinal epithelial cells of many vertebrates, including humans [1]. It causes watery diarrhea and can be fatal to immunocompromised individuals [1]. There are 20 known *Cryptosporidium* species and at least 44 genotypes, which differ significantly in their molecular signatures but have not been assigned species status [2]. Eight fully characterized *Cryptosporidium* species (*C. hominis*, *C. parvum*, *C. meleagridis*, *C. felis*, *C. canis*, *C. suis*, *C. muris*, and *C. andersoni*) and 5 partially characterized species (from the deer, monkey, skunk, rabbit, and chipmunk) infect humans [2-5], among which *C. hominis* and *C. parvum* are the most commonly detected [3]. A real-time PCR (qPCR) method using primers derived from the *CP2* gene is highly sensitive, specific, and accurate for the detection of cryptosporidiosis but cannot distinguish among species [6]. Therefore, we developed a qPCR-based restriction fragment length polymorphism (RFLP) method for the 2 *Cryptosporidium* spp. differentiation and we detected *C. hominis* in 1 of 21

patients with diarrhea.

DNA was prepared from *C. parvum* oocysts of (KKU isolates) collected from laboratory mice (C57BL6/J) that were infected with the parasite and maintained as described [6]. DNA was extracted from the oocysts and fecal materials by using a QIAquick Stool Mini Kit (QIAGEN Inc., Hamburg, Germany USA). The qPCR reactions were performed according to the method of Lee et al. [6] using a LightCycler® (Roche, Basel, Switzerland, USA). The results were analyzed using the LightCycler® software (version 4.05, Roche). DNase/RNase-free water was used in place of template DNA as a negative control. *CP2* sequences of *C. parvum* (AY471868) and *C. hominis* (XM\_661199) were aligned using Clone Manager Suite 7 (Sci-Ed Software, Cary, North Carolina, USA), and restriction enzyme cleavage sites were identified using NEBcutter V2.0 (<http://tools.neb.com/NEBcutter2/>). An aliquot (15 µl) of the qPCR product was digested with *TaqI* (Takara Bio Inc., Shiga, Japan) at 65°C for 2 hr, and DNA fragments were analyzed using 2.5% agarose gels. Stool samples from 21 patients with diarrhea in the Busan area of Korea were collected from June 1 to 30, 2011, by the Korea Centers for Disease Prevention and Control. Modified acid-fast staining was performed on stool samples that were positive for *C. hominis* by using qPCR-based RFLP.

The sequences of the *CP2* genes of *C. parvum* (GenBank AY-

•Received 19 February 2013, revised 22 April 2013, accepted 23 April 2013.

\*Corresponding author (maria205@kku.ac.kr)

© 2013, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

471868) and *C. hominis* (GenBank XM\_661199) are 94% identical (data not shown). Because analysis using NEBcutter V2.0 identified a *TaqI* site only in the sequence of *C. hominis* CP2 (Table 1), *TaqI* was chosen for restriction fragment length polymorphism (RFLP) analysis.

Using qPCR, we found that 1 of 21 patients with diarrhea tested was positive for CP2 (data not shown). The patient with positive results was a 6-year-old girl with watery diarrhea. No other information about the patient was available. To further identify the species of *Cryptosporidium* responsible for the infection, the sample was digested using *TaqI*, which generated 99 bp and 125 bp bands, indicating that the patient was infected by *C. hominis* (Fig. 1). *C. hominis* oocysts were also detected in the diarrheal stool sample by using the modified acid-fast staining (Fig. 2).

Morgan-Ryan et al. [7] proposed a new species of *Cryptosporidium*, *C. hominis*, to indicate its isolation from human feces. However, *C. hominis* and *C. parvum* oocysts are morphologi-

cally indistinguishable [7]. Species discrimination is important for molecular epidemiological purposes to evaluate potential sources of infections [8]. Real-time PCR increases the speed of sample analysis and decreases the risks of contamination with DNA present in the laboratory [8]. The present study showed that both major *Cryptosporidium* species can be detected simultaneously and distinguished from each other by using *TaqI* to digest the CP2 gene of *C. hominis*. Because the CP2 gene is highly specific, no genetic information is available for other *Cryptosporidium* species, except for *C. parvum* and *C. hominis* in GenBank ([www.ncbi.nlm.nih.gov/genbank/](http://www.ncbi.nlm.nih.gov/genbank/)).

The genotypes of *Cryptosporidium* in Korea have been reported [9-11]. Cheun et al. [11] studied *Cryptosporidium* sp. in 3 rural areas by using a PCR-RFLP method to detect 18S rDNA sequences and identified *C. parvum* in 12 patients with *Cryptosporidium* infection. Therefore, the case confirmed by the present study is very important, because it indicates the presence of *C. hominis* infection in Korea.


In the present study, we developed a simple and accurate qPCR-based RFLP method for differentiating *C. parvum* from *C. hominis*. This method could be helpful in facilitating the detection of *C. hominis* infection in Korea.

**Table 1.** DNA fragments produced using a qPCR-based *TaqI*-RFLP assay for the CP2 genes of *C. parvum* and *C. hominis*


	<i>C. hominis</i>	<i>C. parvum</i>
Size of PCR product	224 bp	224 bp
Enzyme cutting ( <i>TaqI</i> )	99 bp, 125 bp	224 bp

### ACKNOWLEDGMENTS

This work was supported by the Basic Science Research Program of the National Research Foundation of Korea (RF) fund-



**Fig. 1.** DNA fragments generated by *TaqI* digestion of real-time PCR-amplified CP2. The results are shown for fecal samples taken from 21 patients with diarrhea. *C. hominis* was detected in the stool samples. *C. parvum* isolated from laboratory mice was used as the control. (A) The qPCR product was confirmed using agarose gel electrophoresis. (B) *TaqI* digestion profile of the qPCR product.



**Fig. 2.** Modified acid-fast staining of a stool sample of a patient infected with *C. hominis*. Bar = 10  $\mu$ m.

ed by the Ministry of Education, Science, and Technology (2011-3333702) and by a grant from the National Institute of Health (NIH-2010E5401000), Ministry of Health and Welfare, the Republic of Korea. The authors declare no conflicts of interest.

## REFERENCES

- O'Donoghue PJ. *Cryptosporidium* and cryptosporidiosis in man and animals. *Int J Parasitol* 1995; 25: 139-195.
- Xiao L, Fayer R, Ryan U, Upton SJ. *Cryptosporidium* taxonomy: recent advances and implications for public health. *Clin Microbiol Rev* 2004; 17: 72-97.
- Caccio SM, Thompson RC, McLauchlin J, Smith HV. Unraveling *Cryptosporidium* and *Giardia* epidemiology. *Trends Parasitol* 2005; 21: 430-437.
- Nichols RA, Campbell BM, Smith HV. Molecular fingerprinting of *Cryptosporidium* oocysts isolated during water monitoring. *Appl Environ Microbiol* 2006; 72: 5428-5435.
- Feltus DC, Giddings CW, Schneck BL, Monson T, Warshauer D, McEvoy JM. Evidence supporting zoonotic transmission of *Cryptosporidium* spp. in Wisconsin. *J Clin Microbiol* 2006; 44: 4303-4308.
- Lee SU, Joung M, Ahn MH, Huh S, Song H, Park WY, Yu JR. CP2 gene as a useful viability marker for *Cryptosporidium parvum*. *Parasitol Res* 2008; 102: 381-387.
- Morgan-Ryan UM, Fall A, Ward LA, Hijjawi N, Sulaiman I, Fayer R, Thompson RC, Olson M, Lal A, Xiao L. *Cryptosporidium hominis* n. sp. (Apicomplexa: Cryptosporidiidae) from Homo sapiens. *J Eukaryot Microbiol* 2002; 49: 433-440.
- Jothikumar N, da Silva AJ, Moura I, Qvarnstrom Y, Hill VR. Detection and differentiation of *Cryptosporidium hominis* and *Cryptosporidium parvum* by dual TaqMan assays. *J Med Microbiol* 2008; 57: 1099-1105.
- Guk SM, Yong TS, Park SJ, Park JH, Chai JY. Genotype and animal infectivity of a human isolate of *Cryptosporidium parvum* in the Republic of Korea. *Korean J Parasitol* 2004; 42: 85-89.
- Park JH, Guk SM, Han ET, Shin EH, Kim JL, Chai JY. Genotype analysis of *Cryptosporidium* spp. prevalent in a rural village in Hwasun-gun, Republic of Korea. *Korean J Parasitol* 2006; 44: 27-33.
- Cheun HI, Choi TK, Chung GT, Cho SH, Lee YH, Kimata I, Kim TS. Genotypic characterization of *Cryptosporidium* oocysts isolated from healthy people in three different counties of Korea. *J Vet Med Sci* 2007; 69: 1099-1101.

