

# The Influence of Fasting, Diabetes, and Several Pharmacological Agents on the Pathways of Thyroxine Metabolism in Rat Liver

ALAN BALSAM and SIDNEY H. INGBAR with the technical assistance of FRANKLIN SEXTON, *Thorndike Laboratory, Harvard Medical School, and the Department of Medicine, Beth Israel Hospital, Boston, Massachusetts 02215*

**ABSTRACT** As judged from both paper and column chromatography, slices or homogenates of liver from rats fasted for 48 h displayed a lesser rate of generation of  $^{125}\text{I}$ -labeled 3,5,3'-triiodothyronine ( $\text{T}_3$ ) from  $^{125}\text{I}$ -labeled thyroxine ( $\text{T}_4$ ) added to incubation media than did preparations from normal chow-fed animals. A similar defect in the conversion of  $\text{T}_4$  to  $\text{T}_3$  in the livers of fasted animals was observed when preparations were incubated with substrate concentrations of  $\text{T}_4$  so that  $\text{T}_3$  generation could be assessed by radioimmunoassay. The effect of fasting could be prevented, wholly or in part, by administration of glucose in the drinking water to otherwise fasted animals, and the degree of prevention appeared to be proportional to the concentration of glucose employed. Diminished generation of  $\text{T}_3$  from  $\text{T}_4$  was similarly evident in the livers of animals with streptozotocin-induced diabetes mellitus, and this defect was overcome by the provision of insulin *in vivo*, but not *in vitro*. Decreased formation of  $\text{T}_3$  from  $\text{T}_4$  was also observed in preparations of liver from animals given dexamethasone, amiodarone, and propylthiouracil. In no case could these effects on the net formation of  $\text{T}_3$  from  $\text{T}_4$  be explained by effects of the experimental conditions on the degradation of the  $\text{T}_3$  generated, as judged from the rate of degradation of exogenous  $^{125}\text{I}$ - $\text{T}_3$  measured in parallel incubates.

An analysis of the rate of disappearance of  $^{125}\text{I}$ - $\text{T}_4$  from reaction mixtures in relation to the rate of appearance of  $^{125}\text{I}$ - $\text{T}_3$  and  $^{125}\text{I}$ -iodide was employed to estimate the activity of the 5-monodeiodinating pathway of  $\text{T}_4$  metabolism that leads to the formation of 3,3',5'-triiodothyronine (reverse  $\text{T}_3$ ). Such estimates indicated that reverse  $\text{T}_3$  formation was actively proceeding in the preparations studied, was slightly enhanced by

fasting, was unaffected by dexamethasone and amiodarone, and was markedly inhibited by propylthiouracil.

In view of the similarities between the effect of these experimental manipulations on the generation of  $\text{T}_3$  from  $\text{T}_4$  by rat liver *in vitro* to their effects on the production rates and serum concentrations of  $\text{T}_3$  in man, it is concluded that the rat liver system provides a suitable model for the study of factors that influence the conversion of  $\text{T}_4$  to  $\text{T}_3$  in man. In addition, the findings strongly indicate that this process, at least in the liver, is closely linked to the utilization of carbohydrate.

## INTRODUCTION

Recent studies have provided convincing evidence that 3,5,3'-triiodothyronine ( $\text{T}_3$ )<sup>1</sup> contributes a major portion to the overall metabolic action of the thyroid hormones within the peripheral tissues. It is evident, moreover, that much of the  $\text{T}_3$  found in the blood of normal man arises, not by direct secretion from the thyroid gland, but rather from the peripheral monodeiodination of thyroxine ( $\text{T}_4$ ) in its outer ring at the 5'-position. It is generally agreed that this reaction results in activation of the hormone molecule, as  $\text{T}_3$  is several times more potent than  $\text{T}_4$  in most all metabolic respects. An alternate pathway of peripheral  $\text{T}_4$  metabolism involves monodeiodination of  $\text{T}_4$  in its inner ring to yield 3,3',5'-triiodothyronine (reverse  $\text{T}_3$ ,  $\text{rT}_3$ ), a product that appears to be hormonally inactive, or nearly so. This pathway is, therefore, considered to be one leading to hormonal inactivation. Evidence in support of these conclusions has been extensively discussed in recent reviews (1, 2).

If the foregoing formulation is correct, then any situa-

Received for publication 9 November 1977 and in revised form 14 March 1978.

<sup>1</sup> Abbreviations used in this paper: PTU, propylthiouracil;  $\text{T}_3$ , 3,5,3'-triiodothyronine;  $\text{T}_4$ , thyroxine;  $\text{rT}_3$ , reverse  $\text{T}_3$ .

tion associated with an alteration in the relative activity of these two pathways should result in altered availability to the tissues of active thyroid hormone. As judged from concentrations of  $T_3$  and  $rT_3$  in serum, such shifts do indeed occur. A diversity of conditions has been shown to be associated with decreases in serum  $T_3$  concentration, among them starvation (3–6), acute or severe chronic illness (7, 8), operative stress (9, 10), anorexia nervosa (11–13), and the administration of glucocorticoids (14–16), the adrenergic blocking agents amiodarone (17), or propranolol (18), or propylthiouracil (PTU) (19, 20). In many of these situations there occurs an accompanying increase in serum  $rT_3$  concentration (4, 6, 8, 15, 17). In the case of patients with hepatic cirrhosis (21–23) or those undergoing starvation (24), decreased peripheral generation of  $T_3$  from  $T_4$  has been directly demonstrated. It would appear, therefore, that the pathway leading peripherally to the generation of  $T_3$  from  $T_4$  may be an important metabolic control point with respect to thyroid hormone action, one whose activity is subject to modification by factors as yet unknown.

In an effort to clarify the mechanisms that influence, and perhaps control, the processes by which  $T_3$  is generated from  $T_4$  in man, we have sought an *in vitro* model using animal tissue that would closely reflect the influence of factors known to or believed to affect the production of  $T_3$  from  $T_4$  in man. Preparations of rat liver have been shown in the present studies to fulfill this criterion, in that they displayed decreased generation of  $T_3$  from  $T_4$  when obtained from animals subjected to starvation, the diabetic state, or treatment with dexamethasone, amiodarone, or PTU. These and other influences on hepatic  $T_3$ -neogenesis form the subject of this report. A portion of these findings has been presented in abstract form (25).

## METHODS

*Isotopes, chemicals, and diet.* All hormones, whether isotopically labeled or unlabeled, as well as chemicals, drugs, and laboratory chow, were obtained from commercial sources<sup>2</sup>.

*Animals.* Experiments were performed with the use of

<sup>2</sup>  $^{125}\text{I}-T_4$  (sp act = 50–75  $\mu\text{Ci}/\mu\text{g}$ ) and  $^{125}\text{I}-T_3$  (sp act = 50–75  $\mu\text{Ci}/\mu\text{g}$ ) were purchased from Abbott Diagnostics, Diagnostic Products (North Chicago, Ill.)  $^{131}\text{I}-T_3$  (sp act = 85  $\mu\text{Ci}/\mu\text{g}$ ) was purchased from Industrial Nuclear Corporation (St. Louis, Mo.). Stable  $T_4$ ,  $T_3$ , and PTU were purchased from Sigma Chemical Co. (St. Louis, Mo.). NPH Iletin and regular insulin from Eli Lilly and Company (Indianapolis, Ind.). Streptozotocin was obtained from The Upjohn Company (Kalamazoo, Mich.), dexamethasone sodium phosphate (Decadron phosphate) from Merck, Sharp & Dohme (West Point, Pa.), and amiodarone [2-butyl-3-(4'-diethylaminoethoxy-3',5'-diiodobenzoyl) benzofurane] from La Bax, Brussels, Belgium. Pelleted laboratory chow, RMH 1000, was purchased from Agway-Country Foods, Agway Inc. (Syracuse, N. Y.).

tissues obtained from Sprague-Dawley rats of the CD strain (Charles River Breeding Laboratories, Wilmington, Mass.). Although the weights of the animals used in the various experiments varied between 150 and 250 g, the rats were closely matched for weight within a single experiment. Unless otherwise stated, animals were maintained on an *ad lib.* regimen of tap water and a standard pelleted laboratory chow, comprising 14% protein, 6% fat, and 54% carbohydrate for 1 wk before and during experiments.

*Fasting and glucose replacement.* In these experiments, a group of the animals was totally deprived of laboratory chow (fasted animals) for 48 h before sacrifice whereas fed controls were allowed continued access to food. In some experiments, some of the fasted animals were given solutions of glucose in place of drinking water, starting at the time that food was withdrawn.

*Experimental diabetes.* Diabetes mellitus was induced by the administration of a single intravenous dose of streptozotocin, 6.5 mg/100 g body wt. 1 day later, all animals so treated developed hyperglycemia and glycosuria without ketonuria, which persisted, in the absence of treatment, until the end of the experiment. Some of the diabetic animals were given a subcutaneous dose of protamine zinc insulin (3 U/100 g body wt) daily on the 2nd, 3rd, and 4th days of the experiment. This dose resulted in correction or amelioration of glycosuria and polyuria. Both untreated and treated diabetic animals, as well as controls, were allowed free access to laboratory chow until they were killed on the 5th day of the experiment.

*Laparotomy.* Under light ether anesthesia, wide vertical and horizontal incisions were made extending into the peritoneal cavity of the rat; muscle layers were then apposed with sutures and skin flaps were stapled together. After the procedure, which lasted approximately 20 min, animals were provided chow, fasted, or supplied glucose water only for the following 48 h.

*Drug treatments.* In some experiments, animals were treated with dexamethasone, amiodarone, or PTU according to treatment schedules described in the appropriate portion of the results section.

*Preparation and incubation of slices.* At the termination of each treatment regimen, animals were killed by cervical subluxation and the liver was quickly excised. With the aid of a Stadie-Riggs microtome, liver slices of uniform thickness, weighing ~200 mg, were prepared. After an initial weighing, slices from control and experimental animals were closely matched in weight by trimming. They were then suspended in vials containing 2 ml of Krebs-Ringer phosphate buffer, pH 7.4, enriched with either  $^{125}\text{I}-T_4$  (1  $\mu\text{Ci}/\text{ml}$ ; 0.020  $\mu\text{g}/\text{ml}$ ) or  $^{125}\text{I}-T_3$  (1.3  $\mu\text{Ci}/\text{ml}$ ; 0.025  $\mu\text{g}/\text{ml}$ ). In experiments in which  $T_3$  generation was to be assessed by radioimmunoassay, media were enriched with stable  $T_4$  (5  $\mu\text{g}/\text{ml}$ ). Vials were either incubated under room air or were closed by rubber stoppers and gassed continuously with purified  $\text{N}_2$ . Incubations were carried out at 37°C for 4 h. In most experiments, some vessels were incubated at 37°C, but without tissues, and others containing tissue slices were incubated at 0°C. No generation of  $^{125}\text{I}-T_3$  from  $^{125}\text{I}-T_4$  was observed in these controls, indicating the absence of artifactual conversion of  $T_4$  to  $T_3$  either during incubation or during chromatography.

At the end of incubation, vessels were plunged into cracked ice, and, when cold, the slices were homogenized in their own medium. Homogenates were then mixed with outdated blood bank plasma to inhibit any further metabolism of  $T_4$ , and the mixture was frozen and kept for 1 wk at most until paper chromatographic analysis was performed.

*Preparation and incubation of homogenates.* Pieces of liver were homogenized in Krebs-Ringer phosphate (1:3,

wt/vol). Homogenates were enriched with  $^{125}\text{I-T}_4$  ( $1 \mu\text{Ci/ml}$ ) or  $^{125}\text{I-T}_3$  ( $1.3 \mu\text{Ci/ml}$ ) and incubated under room air or  $\text{N}_2$  for 4 h, as described above for tissue slices, previous experiments having shown that generation of  $\text{T}_3$  from  $\text{T}_4$  increased in a linear manner during the first 6 h of incubation. In most experiments, control vessels containing homogenates were comparably incubated at  $0^\circ\text{C}$ . At the end of incubation, homogenates were mixed with outdated blood bank plasma (1:2) and the mixtures were frozen until chromatographic analysis was performed.

**Paper chromatography.** Homogenates were thawed and mixed vigorously, and  $10\text{-}\mu\text{l}$  aliquots were applied to Whatman 3 MM filter paper strips together with carrier iodide,  $\text{T}_4$ ,  $\text{T}_3$ , and, in some instances,  $r\text{T}_3$  or tetraiodothyroacetic acid. Chromatograms were developed in a descending hexane, tertiary amyl alcohol, 2 N ammonia (1:10:11) solvent system.<sup>3</sup> After development and drying of the chromatographic strips, iodothyronines were localized by fluorescent light, and iodide by staining with 0.1% palladium chloride. Zones corresponding to carriers, as well as to the origin, were excised and counted in a well-type scintillation counter. Preliminary experiments revealed that >98% of the  $^{125}\text{I}$  in the entire strip was present in these zones. Radioactivity in a given zone was calculated as a percent of the total  $^{125}\text{I}$  in these areas. Values for the percentage generation of the several products of  $\text{T}_4$  metabolism were always corrected for the percentage contamination by each in the substrates employed as assessed by paper chromatography. In the case of substrate  $^{125}\text{I-T}_4$ , 96% of the total radioactivity was  $\text{T}_4$ , between 0.2 and 0.5% was  $\text{T}_3$ , and ~2% was  $^{125}\text{I-iodide}$ . Substrate  $^{125}\text{I-T}_3$  was at least 95% pure and contained from 0.5 to 0.8%  $^{125}\text{I-iodide}$ .

**Sephadex chromatography.** In a number of instances, the generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  in liver slices incubated in air was assessed by chromatography on columns of Sephadex G-25 superfine in a modification (26) of the method described by Green (27). Incubation mixtures comprising slices homogenized in their media were diluted with human plasma (1:9, vol/vol) containing 8-anilino-1-naphthalene sulfonic acid (9 mg/ml) and marker  $^{131}\text{I-T}_3$ . The resulting mixture was applied to a  $25 \times 2\text{-cm}$  column containing preswollen Sephadex. The  $^{125}\text{I-T}_3$  peak was clearly separated from substrate  $^{125}\text{I-T}_4$ . The  $\text{T}_3$  generated was measured as the fractional contribution of the counts in the  $\text{T}_3$  peak to the total counts eluted.

**Stable  $\text{T}_3$  and  $\text{T}_4$  assays.** The concentrations of  $\text{T}_3$  and  $\text{T}_4$  in rat plasma were measured directly by radioimmunoassay and competitive binding techniques, respectively. In some experiments, the generation of stable  $\text{T}_3$  from  $\text{T}_4$  in liver slices was measured using a radioimmunoassay technique to assess the quantity of  $\text{T}_3$  formed. After incubation in air, slices were homogenized in their media and then extracted with 5 vol of 95% ethanol. In separate studies with  $^{125}\text{I-T}_3$ , this was shown to extract >95% of the  $\text{T}_3$  present. 10 or  $20 \mu\text{l}$  of extract was rapidly evaporated under nitrogen, the  $\text{T}_3$  was solubilized in rat serum freed of  $\text{T}_3$  by charcoal adsorption, and the  $\text{T}_3$  content was assayed. Preliminary studies indicated that 10–20  $\mu\text{l}$  of tissue extract incubated without substrate  $\text{T}_4$  could be introduced into the radioimmunoassay without influencing the configuration of the standard curve. Negligible generation of  $\text{T}_3$  from  $\text{T}_4$  was detected in simultaneously incubated tissue-free blanks.

**Statistical evaluation of data.** In experiments with two groups of animals, statistical analysis was performed using the

<sup>3</sup> The  $R_f$  values for carrier and(or) labeled compounds were:  $\text{I}^-$ , 0.11;  $r\text{T}_3$ , 0.23;  $\text{T}_4$ , 0.30; 3,3'-diiodothyronine (3,3'- $\text{T}_2$ ), 0.38; tetraiodothyroacetic acid, 0.43;  $\text{T}_3$ , 0.56.

Student's  $t$  test. When three or more treatment groups were studied simultaneously, a two-way analysis of variance was performed. Duncan's Multiple Range Test was then applied to identify significant differences among specific groups (28).

## RESULTS

The major purpose of these studies was to examine the effects of a variety of experimental manipulations on the generation of  $\text{T}_3$  from  $\text{T}_4$  in rat liver. In each experiment with  $^{125}\text{I-T}_4$ , however, the disappearance of labeled  $\text{T}_4$ , and generation of labeled iodide, origin material, tetraiodothyroacetic acid,  $r\text{T}_3$ , and conjugates of the iodothyronines was also examined. Detectable quantities of tetraiodothyroacetic acid,  $r\text{T}_3$ , and conjugates were not evident, only labeled  $\text{T}_3$ , iodide, and origin material being seen. The generation of labeled origin material ranged from 1 to 6% of added  $^{125}\text{I-T}_4$ , was not materially altered by any experimental manipulation, and will not be described further.<sup>4</sup> For purposes of brevity, results with respect to the generation of iodide will not be discussed, except when significant changes in the generation of this product were seen.

**Effects of fasting and glucose feeding.** Generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  was markedly reduced in slices of liver from fasted animals, whether incubated in air or in nitrogen (Fig. 1). This effect was completely prevented by providing 25% glucose in place of drinking water to animals whose food had been withdrawn. Indeed, in some experiments, the rate of  $\text{T}_3$ -neogenesis was greater in livers from glucose-fed than from chow-fed animals, but this difference was not statistically significant. In liver slices, generation of  $^{125}\text{I-T}_3$  was slightly, but not significantly, greater when incubations were conducted under nitrogen than when room air was employed.

Studies in slices from the same animals revealed that the changes in apparent  $^{125}\text{I-T}_3$  generation induced by fasting and by glucose replacement could not be explained by differences in the rate of deiodination of  $\text{T}_3$ , as judged from the disappearance of added  $^{125}\text{I-T}_3$ , since  $\text{T}_3$  disappearance was either the same or less rapid in slices from fasted animals than in those from fed controls (Fig. 1).

Entirely comparable results with respect to the effects of fasting and glucose replacement were observed in homogenates of rat liver, regardless of whether incubations were conducted in air or under nitrogen (Fig. 2). Here, however, as previous workers have found in rat kidney (29), a much greater generation of  $^{125}\text{I-T}_3$  under anaerobic rather than under aerobic conditions was observed. Again, differences

<sup>4</sup> For the experiments shown in Table I, the percentage generation of  $^{125}\text{I}$ -labeled origin material can be calculated from the formula  $[\text{A}-(\text{B} + \text{C})]$ .




FIGURE 1 Effect of fasting for 48 h or the provision of a 25% solution of glucose in the drinking water to food-deprived animals on the formation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  in rat liver slices.  $\text{T}_3$  metabolism was assessed in separate slices from the same animals incubated with  $^{125}\text{I-T}_3$  in air or nitrogen. In this and subsequent figures, bars and vertical brackets depict the mean and standard error of values obtained in the number of animals shown in parentheses, and absence of an indicated  $P$  value denotes lack of statistical significance between groups.

among the three experimental groups in respect to the rate of degradation of added  $^{125}\text{I-T}_3$  could not explain the differences that were observed in the generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$ .

In the case of homogenates incubated under nitrogen, fasting was associated with significant decreases

of about 50% both in the disappearance of  $^{125}\text{I-T}_4$  from the reaction mixture and in the generation of  $^{125}\text{I}$ -labeled iodide.

The ability of glucose replacement to prevent the reduction in  $\text{T}_3$  generation seen in livers from fasted animals was related to the concentration of glucose provided in the drinking water. In studies of liver slices incubated in air, in vivo administration of 1% glucose had no effect, whereas both 5 and 25% glucose significantly increased  $\text{T}_3$  generation to values higher than those seen in slices from fasted animals. Moreover, generation of  $\text{T}_3$  from  $\text{T}_4$  was greater when 25%, rather than 5%, glucose was administered (Fig. 3).

In contrast to the effects of glucose administration in vivo, enrichment of suspending media with glucose (5 mM), with or without insulin (0.001–1  $\mu\text{M}$ ), had no effect on the generation of  $\text{T}_3$  from  $\text{T}_4$  by liver slices from fasting animals (data not shown).

To validate the data obtained by paper chromatography of specimens incubated with  $^{125}\text{I-T}_4$ , two groups of experiments were performed. In the first,  $^{125}\text{I-T}_3$  generation by rat liver slices was assessed by chromatography of reaction mixtures on Sephadex columns, and the values obtained were compared to those obtained by paper chromatography. Close agreement between the two methods of analysis was noted, and fasting was again associated with decreased generation of  $\text{T}_3$  (Fig. 4).

In the second group of experiments, radioimmunoassay was employed in place of isotopic assay to assess the effects of fasting and glucose replacement on  $\text{T}_3$  generation by rat liver slices incubated in




FIGURE 2 Effect of fasting or provision of a 25% solution of glucose in the drinking water for 48 h on the formation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  in homogenates of rat liver incubated in air or nitrogen.  $\text{T}_3$  metabolism was assessed in parallel incubations with  $^{125}\text{I-T}_3$ .




FIGURE 3 Generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  by liver slices of fasted animals maintained on varying concentrations of glucose in the drinking water for 48 h. Incubations were performed in room air. Numbers below the bars indicate the concentration of glucose in the drinking water.

air (Fig. 5). The results obtained were completely confirmatory of those obtained by the isotopic technique.

**Effect of thyroid hormone replacement on the response to fasting.** Since it is known that thyroid function decreases in the fasted rat, studies were undertaken to determine what role, if any, hypothyroidism might play in the decreased generation of  $\text{T}_3$  from  $\text{T}_4$  seen in the livers of animals fasted for 48 h. In confirmation of previous findings, a 48-h fast was associated with a highly significant decrease ( $P < 0.001$ ), both in serum  $\text{T}_4$  concentration (control,  $4.3 \pm 0.2$ ; fasted,  $1.4 \pm 0.2 \mu\text{g/dl}$ ) (mean  $\pm$  SE) and in serum  $\text{T}_3$  concentration (control,  $39.6 \pm 1.4$ ; fasted,  $10.8 \pm 1.4 \text{ ng/dl}$ ).

Despite these changes, administration of two daily doses of either  $\text{T}_4$  ( $1.5 \mu\text{g}/100 \text{ g body wt}$ ) or  $\text{T}_3$  ( $0.5 \mu\text{g}/100 \text{ g body wt}$ ) during the 48-h period of fasting failed to influence the reduction in  $^{125}\text{I-T}_3$  generation from  $^{125}\text{I-T}_4$  seen in liver slices of matched animals not given exogenous hormone (data not shown).

**Effect of experimental diabetes.** Rats given streptozotocin and allowed free access to food became manifestly diabetic 2 days later and remained so during the ensuing 3 days. As compared to findings obtained in controls at that time, liver slices obtained from diabetic rats displayed greatly reduced generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  (Fig. 6). This effect was almost completely reversed by the administration of protamine zinc insulin ( $3 \text{ U}/100 \text{ g body wt}$ ) daily during the last 3 days of the experiment. However, no reversal

of the impairment in  $\text{T}_3$  generation was observed when liver slices from diabetic rats were incubated in media containing glucose ( $5 \text{ mM}$ ) and insulin ( $1 \mu\text{M}$ ).

The effects of diabetes and of insulin replacement on the apparent generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  could not be explained by an alteration in the rate of  $\text{T}_3$  degradation, as judged from the disappearance of added  $^{125}\text{I-T}_3$  in vessels containing matched slices (Fig. 6). Untreated diabetes was associated with a small, but not statistically significant, decrease in the disappearance of  $^{125}\text{I-T}_3$  from the reaction mixture. Moreover, when  $\text{T}_3$  degradation by liver slices from diabetic animals treated with insulin in vivo was compared to that of liver slices from diabetic animals incubated with insulin in vitro, a significantly ( $P < 0.05$ ) slower  $\text{T}_3$  disappearance was evident in the latter group.

**Effect of laparotomy.** Groups of five animals were fasted, provided laboratory chow, or supplied glucose in the drinking water for 48 h after laparotomy, whereas unoperated controls were maintained on parallel regimens. Mean values for the percent generation of  $\text{T}_3$  from  $\text{T}_4$  in liver slices from control and operated animals, respectively, were: chow-fed,  $4.3 \pm 0.3$  vs.  $4.8 \pm 0.7$ ; fasted,  $2.3 \pm 0.3$  vs.  $1.9 \pm 0.2$ ; glucose-fed,  $4.6 \pm 0.3$  vs.  $4.7 \pm 0.6$ . Analysis of variance confirmed to the characteristic effects of the dietary regimens, but demonstrated no effect of laparotomy.

**Effects of various drugs on  $\text{T}_3$  generation from  $\text{T}_4$**  (Table I). An approximate 50% inhibition of  $\text{T}_3$  generation from  $\text{T}_4$  was observed in liver slices from animals given  $1.0 \text{ mg}$  of dexamethasone subcutaneously daily for 3 days before sacrifice. However, no effect on  $^{125}\text{I-T}_4$  disappearance or generation of labeled iodide was observed.

Profoundly inhibited generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  was seen in liver slices from chow-fed animals given amiodarone as a  $0.1\%$  solution in  $25\%$  glucose for 3 days in place of the drinking water. Amiodarone also produced a slight, but statistically insignificant, decrease in  $^{125}\text{I-T}_4$  disappearance and a slight, but significant, decrease in the generation of labeled iodide.

PTU, when given in the drinking water as a  $0.05\%$  solution in  $25\%$  glucose for 3 days, almost completely abolished the generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  by liver slices. Concomitantly, disappearance of  $^{125}\text{I-T}_4$  and generation of  $^{125}\text{I}$ -labeled iodide were markedly inhibited.

**Effect of various drugs on the metabolism of  $\text{T}_3$**  (Table II). Animals treated with dexamethasone and PTU displayed slight, statistically insignificant reductions in the rate of disappearance of  $\text{T}_3$ , whereas those given amiodarone showed no change. Significantly decreased generation of  $^{125}\text{I}$ -iodide from added  $^{125}\text{I-T}_3$  was observed after treatment with all three agents.




FIGURE 4 Assessment of  $^{125}\text{I-T}_3$  generation from  $^{125}\text{I-T}_4$  by column chromatography of reaction mixtures on Sephadex G-25 superfine. Liver slices from rats fasted for 48 h or from fed controls were incubated with  $^{125}\text{I-T}_4$  and then homogenized in their media and homogenates were subjected to chromatography either on columns of Sephadex or on filter paper using a hexane-tertiary amyl alcohol-ammonia solvent system (HTA). Shown in the figure are elution patterns obtained during column chromatography of specimens from fed or fasted animals. Values for the percent generation of  $\text{T}_3$  from  $\text{T}_4$  as judged from paper chromatography (HTA- $\text{T}_3$ ) are compared to those obtained from column chromatography (Sephadex- $\text{T}_3$ ).

## DISCUSSION

In the present studies, we have sought to determine whether the rat liver constitutes a reasonable animal tissue model in which to study the factors that influence the peripheral conversion of  $\text{T}_4$  to  $\text{T}_3$  in man.




FIGURE 5 Effect of fasting or provision of a solution of 25% glucose in the drinking water on the generation of  $\text{T}_3$  from  $\text{T}_4$  in liver slices, as judged from radioimmunoassay. Incubations were performed in air with  $\text{T}_4$ ,  $5 \mu\text{g/ml}$ , added to the medium.

For this purpose we have determined the effect in this tissue of a variety of experimental manipulations that have been shown to inhibit the conversion of  $\text{T}_4$  to  $\text{T}_3$  in man, as judged from changes in serum  $\text{T}_3$  concentration,  $\text{T}_3$  production rate, or both. The data suggest that rat liver slices or homogenates are valid models, since excellent concordance was observed between changes in  $\text{T}_3$  generation from  $\text{T}_4$  by rat liver preparations and alterations in  $\text{T}_3$  metabolism in man in response to starvation or to the administration of dexamethasone, amiodarone, and PTU.

The apparent inhibitory effect of these experimental manipulations on  $\text{T}_3$  generation from  $\text{T}_4$  cannot be explained by an effect on the degradation of the  $\text{T}_3$  generated, at least as judged from the metabolism of exogenous  $\text{T}_3$  within these systems. Degradation of exogenous  $\text{T}_3$  was slow, and was retarded, rather than accelerated, by the factors that decreased  $\text{T}_3$  generation.

In previous studies of the generation of  $\text{T}_3$  from  $\text{T}_4$  by animal tissues in vitro, either paper chromatog-




FIGURE 6 The effect of streptozotocin-induced diabetes mellitus and of insulin replacement on the generation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$ . The formation of  $^{125}\text{I-T}_3$  from  $^{125}\text{I-T}_4$  and the metabolism of  $^{125}\text{I-T}_3$  were assessed in parallel incubations of liver slices in air. Diabetic animals either received injections of saline or protamine zinc insulin, 3 U/100 g for 3 days. Slices from diabetic animals were incubated in unsupplemented medium or in media containing  $1 \mu\text{M}$  insulin, and  $5 \text{ mM}$  glucose.

raphic analysis of preparations incubated with radioiodine-labeled  $\text{T}_4$  or radioimmunoassay of preparations enriched with unlabeled  $\text{T}_4$  has been used as a means of assessing  $\text{T}_3$  generation (29–34). In the

present studies, we have mainly employed the isotopic-paper chromatographic technique, and have validated the results obtained both by column chromatography of isotopically labeled preparations and by radioimmunoassay.

Few specific conclusions can be drawn as to the mechanisms by which the various experimental manipulations inhibit  $\text{T}_3$  generation from  $\text{T}_4$ . The similar effects of starvation and streptozotocin diabetes are not surprising, perhaps, in view of the variety of metabolic alterations common to both, including insulinopenia, and diminished activity of a variety of enzymes associated with glucose metabolism (35, 36). Similarly, the ability of glucose administration to prevent the effect of fasting, and of insulin in vivo to reverse the effect of experimental diabetes, suggests that, in the liver at least, the metabolism of glucose is somehow related to the process of  $\text{T}_3$  formation. A relationship between glucose metabolism and the generation of a cofactor necessary for  $\text{T}_3$  formation is suggested by our finding that the  $\text{T}_3$ -generating activity of broken cell systems from the livers of fasted rats is restored to normal by enrichment with an NADPH-generating system (isocitrate; isocitrate dehydrogenase; NADP,  $0.1 \text{ mM}$ ).<sup>5</sup>

The question arises as to whether  $\text{T}_3$ -neogenesis is specifically linked to carbohydrate utilization or whether it is more generally related to the caloric adequacy of the diet. Evidence that it is not solely a function of caloric intake is provided by studies conducted in our laboratory (data not reported) which

<sup>5</sup> Balsam, A., and S. H. Ingbar. Unpublished observations.

TABLE I  
The Metabolism of  $^{125}\text{I-T}_4$  in Slices of Rat Liver\*

Treatment	Treatment group	Number of animals	Measured values			Derived values			
			(A) $^{125}\text{I-T}_4$ degradation (% added $\text{T}_4$ )	(B) $^{125}\text{I-T}_3$ generation (% added $\text{T}_4$ )	(C) $^{125}\text{I}$ -Iodide generation (% added $\text{T}_4$ )	Excess $^{125}\text{I}$ - $\text{T}_4$ (% added $\text{T}_4$ ) (C - B)	Excess $\text{I}^-/\text{T}_4$ degradation (C - B)/A	$\text{T}_3$ generation/ $\text{T}_4$ degradation (2B/A)	Excess $\text{I}^-/\text{T}_3$ generation (C - B/B)
Fasting	Control	18	18.4 ± 1.3	3.0 ± 0.3	10.3 ± 0.9	7.3 ± 0.7	0.40 ± 0.02	0.32 ± 0.02	2.8 ± 0.3
	Exp.	18	18.0 ± 1.2	1.6 ± 0.2 <sup>†</sup>	10.1 ± 0.8	8.5 ± 0.7 <sup>†</sup>	0.47 ± 0.02 <sup>†</sup>	0.17 ± 0.01 <sup>†</sup>	5.7 ± 0.4 <sup>†</sup>
Diabetes	Control	4	30.8 ± 2.0	6.5 ± 0.5	23.8 ± 2.1	17.3 ± 2.2	0.56 ± 0.05	0.43 ± 0.04	2.7 ± 0.5
	Exp.	4	24.5 ± 1.6	3.7 ± 0.4 <sup>§</sup>	19.2 ± 2.1	15.5 ± 2.4	0.62 ± 0.05	0.31 ± 0.04	4.6 ± 1.3
Dexamethasone	Control	6	36.5 ± 3.4	5.1 ± 0.5	25.0 ± 2.3	19.9 ± 2.6	0.54 ± 0.05	0.30 ± 0.05	4.2 ± 0.8
	Exp.	6	36.4 ± 3.4	2.7 ± 0.4 <sup>†</sup>	24.1 ± 2.7	21.4 ± 2.5	0.62 ± 0.03 <sup>†</sup>	0.16 ± 0.02 <sup>†</sup>	8.4 ± 1.0 <sup>†</sup>
Amiodarone	Control	5	34.9 ± 2.8	6.3 ± 0.8	26.9 ± 1.9	20.6 ± 1.4	0.60 ± 0.03	0.36 ± 0.03	3.5 ± 0.5
	Exp.	5	26.1 ± 1.6	0.5 ± 0.02 <sup>§</sup>	21.8 ± 1.2 <sup>†</sup>	21.2 ± 1.3	0.82 ± 0.01 <sup>§</sup>	0.04 ± 0.004 <sup>†</sup>	42.2 ± 3.8 <sup>†</sup>
PTU	Control	5	29.7 ± 1.4	7.0 ± 0.08	17.8 ± 0.9	10.8 ± 0.5	0.37 ± 0.02	0.47 ± 0.04	1.7 ± 0.3
	Exp.	5	14.0 ± 0.6 <sup>†</sup>	0.6 ± 0.04	6.6 ± 0.7 <sup>†</sup>	6.0 ± 0.7 <sup>§</sup>	0.42 ± 0.03	0.09 ± 0.002 <sup>†</sup>	9.8 ± 0.6 <sup>†</sup>

\* Data shown represent mean ± SE. See text for precise description of experimental manipulations.

<sup>†</sup>  $P < 0.05$ .

<sup>§</sup>  $P < 0.01$ .

<sup>†</sup>  $P < 0.001$ .

TABLE II  
The Metabolism of  $^{125}\text{I-T}_3$  in Slices of Rat Liver\*

Treatment	Treatment group	No. of animals	$\text{T}_3$ degradation (% added $\text{T}_3$ )	Iodide generation (% added $\text{T}_3$ )
Fasting	Control	6	20.6±0.8	16.9±1.0
	Exp.	6	19.4±1.8	13.5±1.5
Diabetes	Control	7	27.3±1.0	20.1±0.6
	Exp.	7	23.7±1.3†	16.2±1.2§
Dexamethasone	Control	4	19.8±1.7	17.2±1.8
	Exp.	5	16.5±2.8	11.0±1.9†
Amiodarone	Control	6	20.4±2.6	13.7±1.5
	Exp.	6	19.5±1.9	7.2±0.9§
PTU	Control	6	20.4±2.6	13.7±1.5
	Exp.	6	16.3±0.4	5.5±0.5 <sup>¶</sup>

\* Data shown represent mean±SE. See text for precise description of experimental manipulations.

†  $P < 0.05$ .

§  $P < 0.01$ .

¶  $P < 0.001$ .

demonstrate that provision of fat in the form of an emulsion of peanut oil in the drinking water (5% vol/vol), which the animals readily consumed, failed to influence the defective formation of  $\text{T}_3$  observed in livers of animals deprived of chow. In fasting man, moreover, carbohydrate diets are more effective than equicaloric quantities of protein in restoring depressed serum  $\text{T}_3$  concentrations to normal (6), and the lesser effect of protein can itself perhaps be ascribed to enhancement of gluconeogenesis.

In man, a decrease in serum  $\text{T}_3$  concentration occurs in association with a variety of "stressor" stimuli, such as surgery (9, 10), and acute and chronic illnesses of wide variety of types (7, 8). Nonspecific stress does not appear to be a factor in the present in vitro system, however, since wide laparotomy failed to influence the hepatic generation of  $\text{T}_3$  from  $\text{T}_4$ . This finding also makes it unlikely that the effect of starvation or diabetes can be ascribed to adrenocortical activation, a possibility suggested by the inhibition of hepatic  $\text{T}_3$  generation induced by dexamethasone, albeit in very large doses, that we and others (32) have found.

Considerable evidence indicates that the peripheral metabolism of  $\text{T}_4$  in man proceeds mainly by a sequence of monodeiodinations occurring randomly in locus, though not in rate, at the inner and outer rings of the thyronine nucleus (37). Although it has obviously been possible to assess the rate of  $\text{T}_3$  generation from  $\text{T}_4$ , as for example in the present and previous studies (29–34), it has generally not been possible to demonstrate generation of  $\text{rT}_3$  from  $\text{T}_4$  in vitro. Such was certainly the case in the present studies in which

$^{125}\text{I-rT}_3$  was looked for, but never detected, in chromatographic analyses of the reaction mixtures containing labeled  $\text{T}_4$ . This is probably the result of the extreme rapidity with which  $\text{rT}_3$  is itself degraded in tissue systems (38). This being the case, we have attempted to utilize the ancillary data that we have obtained to assess the relative activities of the  $\text{T}_3$ - and  $\text{rT}_3$ -generating mechanisms and the manner in which they were affected by the experimental manipulations that were studied. In so doing, we have made the limiting assumption that deiodination of labeled  $\text{T}_3$  generated from labeled  $\text{T}_4$  is negligible and hence does not influence significantly either apparent  $\text{T}_3$  generation or the formation of labeled iodide. If that assumption is made, then estimates of the activity of 5-mono-deiodinating pathway, both absolute and relative to that of the 5'-monodeiodinating pathway, can be made.

Since neither conjugates nor deaminated derivatives of  $\text{T}_4$  were observed, then the fractional disappearance of  $^{125}\text{I-T}_4$  from the reaction mixtures must have represented the total proportion of added  $\text{T}_4$  converted to  $\text{T}_3$  and  $\text{rT}_3$ . Under the assumption that  $\text{T}_3$  formed undergoes no deiodination, all  $^{125}\text{I}$ -iodide generated in excess of that which could be accounted for by the generation of  $\text{T}_3$  must represent  $^{125}\text{I}$ -iodide that has passed from  $\text{T}_4$  through the  $\text{rT}_3$  pathway.<sup>6</sup> Hence, it was possible to obtain an estimate of the activity of this pathway. We have designated this as "excess iodide formation," and have calculated this function as percent  $^{125}\text{I}$ -iodide formed minus percent  $^{125}\text{I-T}_3$  generated. This function always comprised a substantial fraction of the  $\text{T}_4$  added and indeed always exceeded in magnitude by at least several-fold the measured fraction of  $\text{T}_3$  generated (Table I). Activity of the  $\text{T}_3$ -generating pathway as a fraction of overall  $\text{T}_4$  deiodination can be calculated as the ratio, percent  $\text{T}_3$  generation/percent  $\text{T}_4$  degradation; that of  $\text{rT}_3$  pathway as the ratio, percent excess iodide/percent  $\text{T}_4$  degradation; the activities of the two pathways relative to one another as ratio, percent excess  $^{125}\text{I}$ -iodide/percent  $^{125}\text{I-T}_3$  generation.

When the current data are examined in this way, several patterns of response to the various experimental manipulations employed emerge (Table I). Fasting decreased  $\text{T}_3$  generation, but did not alter significantly either  $\text{T}_4$  degradation or iodide generation. Hence, excess iodide generation increased. This indication of a significant, though small, increase in  $\text{rT}_3$

<sup>6</sup> It is theoretically possible that a portion of the excess iodide might have arisen from cleavage of the ether linkage of  $\text{T}_4$ , with deiodination of the outer-ring product of this reaction. Significant activity of any such pathway in rat liver has been excluded, however, by our failure to recover labeled diiodotyrosine when biosynthetically derived, randomly labeled  $\text{T}_4$  was incubated with liver slices in the presence of an inhibitor of iodotyrosine deiodinase, dinitrotyrosine (39), in concentrations sufficient to abolish deiodination of added  $^{125}\text{I}$ -diiodotyrosine.<sup>5</sup>



production in vitro as a result of starvation, as well as the clear evidence of an increase in the ratio of 5-monodeiodination/5'-monodeiodination, is in accord with the findings of recent studies on the effects of starvation on  $rT_3$  production rates in man (24, 40).

The apparent effect of diabetes differed slightly from that of starvation.  $T_4$  disappearance and total iodide generation were decreased slightly, though not significantly, and no evidence of an absolute increase in 5-monodeiodination was seen, as excess iodide formation was unchanged. Monodeiodination at the 5'-position ( $T_3$  generation) as a function of total  $T_4$  degradation was significantly decreased, whereas 5-monodeiodination, either in relation to total  $T_4$  degradation or to  $T_3$  generation, was increased, though not significantly so.

The effects of dexamethasone and amiodarone were qualitatively similar to those of diabetes in that, although  $T_3$  generation was inhibited, no evidence was obtained that  $rT_3$  formation was increased.

The effects of PTU were strikingly different from those of other experimental manipulations examined. As others have found in studies of  $T_3$  formation in the whole rat (41), PTU given in vivo greatly inhibited  $T_3$ -neogenesis in liver slices. In addition, PTU also appeared to inhibit 5-monodeiodination, as judged from estimates of excess iodide formation. The effect on 5-monodeiodination was less marked than the inhibition of  $T_3$  generation, however, since the ratio of excess iodide formation/ $T_3$  generation was greatly increased.

The finding that PTU inhibits the inner-ring pathway of  $T_4$  metabolism is in accord with data recently reported, showing that PTU inhibits the conversion of  $T_4$  to 3,3'- $T_2$ , a metabolite formed largely via the  $rT_3$  pathway in rat liver homogenate (42).

From the above, it emerges that all the manipulations studied resulted in decreased  $T_3$  generation, but that starvation may have enhanced  $rT_3$  formation; diabetes, dexamethasone, and amiodarone did not affect  $rT_3$  formation appreciably, while PTU inhibited  $rT_3$  formation, but less markedly than it inhibited the formation of  $T_3$ .

The calculations from which these conclusions are drawn are based upon the assumption that the  $^{125}I$ - $T_3$  generated from  $^{125}I$ - $T_4$  does not undergo significant deiodination. It is very unlikely that this is the case, however, since some deiodination of exogenous  $^{125}I$ - $T_3$  was always observed. Deiodination of endogenously generated  $T_3$  would, by the present methods of calculation, lead to overestimates of "excess iodide formation," a function of the activity of the  $rT_3$  pathway. However, as judged from the only source of information available, i.e., the rate of deiodination of exogenous  $^{125}I$ - $T_3$ , this factor could not have accounted for more than a small proportion of excess iodide formation, since, in relation to the magnitude of these derived

values, the proportionate conversion of  $T_4$  to  $T_3$  and the proportionate deiodination of added  $T_3$  were quite small.

For these reasons, the quantitative values of the derived functions must be regarded as approximations only. Nevertheless, we consider them to be, in the main, qualitatively reliable and productive of the only available evidence concerning the activity of the inner-ring monodeiodinating pathway of  $T_4$  metabolism in vitro and the influence thereon of a variety of factors that affect outer-ring monodeiodination.

The findings provide strong evidence that the inner- and outer-ring monodeiodinating pathways for  $T_4$  are biochemically distinct from one another, a conclusion consonant with studies which demonstrate divergent effects of both starvation and chronic liver disease on the production rates of  $T_3$  and  $rT_3$  in man (23, 24).

#### ACKNOWLEDGMENTS

Data analysis was performed, in part, on the PROPHET System, a national computer resource sponsored by the Chemical/Biological Information Handling Program, National Institutes of Health.

This work was supported in part by research grant AM-18416 from the National Institute of Arthritis, Metabolism and Digestive Diseases, and by grant RR-01032 from the General Clinical Research Centers Program of the Division of Research Resources, National Institutes of Health.

#### REFERENCES

1. Ingbar, S. H., and L. E. Braverman. 1975. Active form of the thyroid hormone. *Annu. Rev. Med.* **26**: 443-449.
2. Utiger, R. D. 1974. Serum triiodothyronine in man. *Annu. Rev. Med.* **25**: 289-302.
3. Portnay, G. I., J. T. O'Brian, J. Rush, A. G. Vagenakis, F. Azizi, R. A. Arky, S. H. Ingbar, and L. E. Braverman. 1974. The effect of starvation on the concentration and binding of thyroxine and triiodothyronine in serum and on the response to TRH. *J. Clin. Endocrinol. Metab.* **39**: 191-194.
4. Vagenakis, A. G., A. Burger, G. I. Portnay, M. Rudolph, J. T. O'Brian, F. Azizi, R. A. Arky, P. Nicod, S. H. Ingbar, and L. E. Braverman. 1975. Diversion of peripheral thyroxine metabolism from activating to inactivating pathways during complete fasting. *J. Clin. Endocrinol. Metab.* **41**: 191-194.
5. Chopra, I. J., and S. R. Smith. 1975. Circulating thyroid hormones in adult patients with protein-calorie malnutrition. *J. Clin. Endocrinol. Metab.* **40**: 221-227.
6. Spaulding, S. W., I. J. Chopra, R. S. Sherwin, and S. S. Lyall. 1976. Effect of caloric restriction and dietary composition on serum  $T_3$  and reverse  $T_3$  in man. *J. Clin. Endocrinol. Metab.* **42**: 197-200.
7. Bermudez, F., M. I. Surks, and J. H. Oppenheimer. 1975. High incidence of decreased serum triiodothyronine concentration in patients with nonthyroidal disease. *J. Clin. Endocrinol. Metab.* **41**: 27-39.
8. Chopra, I. J., U. Chopra, S. R. Smith, M. Reza, and D. H. Solomon. 1975. Reciprocal changes in serum concentrations of 3,3',5'-triiodothyronine (reverse  $T_3$ ) and 3,3',5'-triiodothyronine ( $T_3$ ) in systemic illnesses. *J. Clin. Endocrinol. Metab.* **41**: 1043-1049.
9. Burr, W. A., E. G. Glach, R. S. Griffiths, R. Hoffenberg, H. Meinhold, and K. W. Wenzel. 1975. Serum triiodothy-

- ronine concentrations after surgical operation. *Lancet*. II: 1277.
10. Brandt, M., H. Kehlet, J. M. Hansen, and L. Skovsted. 1976. Serum triiodothyronine and surgery. *Lancet*. I: 491.
  11. Miyai, K., T. Yamamoto, M. Azukizawa, K. Ishibashi, and Y. Kumahara. 1975. Serum thyroid hormones and thyrotropin in anorexia nervosa. *J. Clin. Endocrinol. Metab.* 40: 334-338.
  12. Moshang, T., J. S. Parks, L. Baker, V. Vaidya, R. D. Utiger, A. M. Bongiovanni, and P. J. Synder. 1975. Low serum triiodothyronine in patients with anorexia nervosa. *J. Clin. Endocrinol. Metab.* 40: 470-473.
  13. Croxson, M. S., and H. K. Ibbertson. 1977. Low serum triiodothyronine ( $T_3$ ) and hypothyroidism in anorexia nervosa. *J. Clin. Endocrinol. Metab.* 44: 167-174.
  14. Duick, D. S., D. W. Warren, J. T. Nicoloff, C. L. Otis, and M. S. Croxson. 1974. Effect of single dose dexamethasone on the concentration of serum triiodothyronine in man. *J. Clin. Endocrinol. Metab.* 39: 1151-1154.
  15. Chopra, I. J., D. E. Williams, J. Orgiazzi, and D. H. Solomon. 1975. Opposite effects of dexamethasone on serum concentrations of 3,3',5'-triiodothyronine (reverse  $T_3$ ) and 3,3',5-triiodothyronine ( $T_3$ ). *J. Clin. Endocrinol. Metab.* 41: 911-920.
  16. Degroot, L. J., and K. Hoyer. 1976. Dexamethasone suppression of serum  $T_3$  and  $T_4$ . *J. Clin. Endocrinol. Metab.* 42: 976-978.
  17. Burger, A., D. Dinichert, P. Nicod, M. Jenny, T. Le-Marchand-Beraud, and M. B. Valloton. 1976. Effect of amiodarone on serum triiodothyronine, reverse triiodothyronine, thyroxine, and thyrotropin: a drug influencing peripheral metabolism of thyroid hormones. *J. Clin. Invest.* 58: 255-259.
  18. Wiersinga, W. M., and J. L. Toubert. 1977. The influence of  $\beta$ -adrenoreceptor blocking agents on plasma thyroxine and triiodothyronine. *J. Clin. Endocrinol. Metab.* 45: 293-298.
  19. Saberi, M., F. H. Sterling, and R. D. Utiger. 1975. Reduction in extra-thyroidal triiodothyronine production by propylthiouracil in man. *J. Clin. Invest.* 55: 218-223.
  20. Geffner, D. L., M. Azukizawa, and J. Hershman. 1975. Propylthiouracil blocks extrathyroidal conversion of thyroxine to triiodothyronine and augments thyrotropin secretion in man. *J. Clin. Invest.* 55: 224-229.
  21. Chopra, I. J., D. H. Solomon, U. Chopra, R. T. Young, and G. N. Chua Teco. 1974. Alterations in circulating thyroid hormones and thyrotropin in hepatic cirrhosis: evidence for euthyroidism despite subnormal serum triiodothyronine. *J. Clin. Endocrinol. Metab.* 39: 501-511.
  22. Nomura, S., C. S. Pittman, J. B. Chambers, Jr., M. W. Buck, and T. Shimizu. 1975. Reduced peripheral conversion of thyroxine to triiodothyronine in patients with hepatic cirrhosis. *J. Clin. Invest.* 56: 643-652.
  23. Chopra, I. J. 1976. An assessment of daily production and significance of thyroidal secretion of 3,3',5'-triiodothyronine (reverse  $T_3$ ) in man. *J. Clin. Invest.* 58: 33-40.
  24. Eisenstein, Z., S. Hagg, A. G. Vagenakis, S. L. Fang, B. Ransil, A. Burger, A. Balsam, L. E. Braverman, and S. H. Ingbar. 1977. Effect of starvation on the production and peripheral metabolism of 3,3',5'-triiodothyronine in euthyroid obese subjects. *J. Clin. Endocrinol. Metab.* In press.
  25. Balsam, A., F. C. Sexton, and S. H. Ingbar. 1976. In vitro generation  $T_3$  from  $T_4$  by rat liver: reversible inhibition induced by starvation and diabetes mellitus. Abstract. 52nd Meeting of the American Thyroid Association. T-16.
  26. Rudolph, M., T. Sakurada, S. L. Fang, A. G. Vagenakis, L. E. Braverman, and S. H. Ingbar. 1978. Appearance of labeled metabolites in the serum of man following the administration of labeled thyroxine, triiodothyronine, and reverse  $T_3$ . *J. Clin. Endocrinol. Metab.* 46: 923-928.
  27. Green, W. L. 1972. Separation of iodocompounds in serum by chromatography on sephadex columns. *J. Chromatogr.* 72: 83-91.
  28. Dunnett, C. W. 1970. Multiple comparisons. In *Statistics in Endocrinology*. J. W. McArthur and T. Colton, editors. The M.I.T. Press, Cambridge, Mass. I: 86-87.
  29. Chirasevenupurand, P., U. Buerger, A. Goswami, and I. N. Rosenberg. 1975. Formation of triiodothyronine from L-thyroxine in rat kidney homogenate. In *Thyroid Research*. J. Robbins and L. E. Braverman, editors. American Elsevier Publishing Co., Inc., New York. 244.
  30. W. L. Green. 1975. Thyroxine metabolism by rat liver slices: evidence for a specific  $T_3$ -forming pathway. In *Thyroid Research*. J. Robbins and L. E. Braverman, editors. American Elsevier Publishing Co., Inc., New York. 239.
  31. Visser, T. J., I. Van der Does-Tobe, R. Docter, and G. Hennemann. 1975. Conversion of thyroxine into triiodothyronine by rat liver homogenate. *Biochem. J.* 150: 489-493.
  32. Hufner, M., and M. Knopfle. 1976. Pharmacological influences on  $T_4$  to  $T_3$  conversion in rat liver. *Clin. Chim. Acta.* 72: 337-341.
  33. Chopra, I. J. 1977. A study of extrathyroidal conversion of thyroxine ( $T_4$ ) to 3,3',5-triiodothyronine ( $T_3$ ) *in vitro*. *Endocrinology.* 101: 453-463.
  34. Kaplan, M. M., M. Schimmel, and R. D. Utiger. 1977. Changes in serum 3,3',5'-triiodothyronine (reverse  $T_3$ ) concentrations with altered thyroid hormone secretion and metabolism. *J. Clin. Endocrinol. Metab.* 45: 447-456.
  35. Cahill, G. F., Jr. 1970. Starvation in man. *N. Engl. J. Med.* 282: 668-675.
  36. Williams, R. H., and D. Porte, Jr. In *Textbook of Endocrinology*. R. H. Williams, editor. W. B. Saunders Company, Philadelphia. 5th edition. 542 pp.
  37. Sakurada, T., M. Rudolph, S. L. Fang, A. G. Vagenakis, L. E. Braverman, and S. H. Ingbar. 1978. Evidence that triiodothyronine and reverse triiodothyronine are sequentially deiodinated in man. *J. Clin. Endocrinol. Metab.* 46: 916-922.
  38. Cavaliere, R. R., L. A. Gavin, F. McMahon, and M. Hammond. 1977. Thyroxine ( $T_4$ ) deiodination in liver: subcellular localization of reverse- $T_3$  ( $RT_3$ ) forming and degrading systems. *Clin. Res.* 25:(3): 462A. (Abstr.)
  39. Green, W. L. 1968. Inhibition of thyroidal iodotyrosine deiodination by tyrosine analogues. *Endocrinology.* 83: 336-347.
  40. Suda, A., C. Pittman, and J. Chambers, Jr. 1977. Regulation of 3,5,3'-triiodothyronine ( $T_3$ ) production during fast and refeeding. Abstracts of the 53rd Meeting of the American Thyroid Association. T-14 (Abstr.)
  41. Oppenheimer, J. H., H. L. Schwartz, and M. I. Surks. 1972. Propylthiouracil inhibits the conversion of L-thyroxine to L-triiodothyronine: an explanation of the anti-thyroxine effect of propylthiouracil and evidence supporting the concept that triiodothyronine is the active thyroid hormone. *J. Clin. Invest.* 51: 2493-2497.
  42. Grussendorf, M., and M. Hufner. 1971. Deiodination of  $T_4$  in liver homogenate of normal and thyroid hormone substituted, thyroidectomized rats. Abstracts of the 8th Annual Meeting of the European Thyroid Association, Lyon. 32A. (Abstr.)