


Volume 13, 27 June 2013

Publisher: Igitur publishing

URL: <http://www.ijic.org>

Cite this as: Int J Integr Care 2013; Apr–Jun, URN:NBN:NL:UI:10-1-114599

Copyright: 

Submitted: 31 May 2012, revised 20 February 2013, accepted 21 February 2013

Integrated Care Case

The process flow and structure of an integrated stroke strategy

Emma F. van Bussel, MSc, Medical Doctor, BovenIJ Hospital, Amsterdam, The Netherlands

Thomas Jeerakathil, MD, Associate Professor, Neurology, University of Alberta, Canada; Chair, Evaluation and Quality Improvement Pillar, Alberta Provincial Stroke Strategy, Alberta, Canada

Augustinus J.P. Schrijvers, PhD, Professor of Public Health and Health Economist, Julius Center University Medical Center Utrecht, The Netherlands

Correspondence to: Augustinus Schrijvers, Julius Center University Medical Center Utrecht, Heidelberglaan 100, 3584 CX, Utrecht, The Netherlands, Phone: +3163007.4477, Fax: +3188 75 554 82, E-mail: mail@guusschrijvers.nl

Abstract

Introduction: In the Canadian province of Alberta access and quality of stroke care were suboptimal, especially in remote areas. The government introduced the Alberta Provincial Stroke Strategy (APSS) in 2005, an integrated strategy to improve access to stroke care, quality and efficiency which utilizes telehealth.

Research question: What is the process flow and the structure of the care pathways of the APSS?

Methodology: Information for this article was obtained using documentation, archival APSS records, interviews with experts, direct observation and participant observation.

Results: The process flow is described. The APSS integrated evidence-based practice, multidisciplinary communication, and telestroke services. It includes regular quality evaluation and improvement.

Conclusion: Access, efficiency and quality of care improved since the start of the APSS across many domains, through improvement of expertise and equipment in small hospitals, accessible consultation of stroke specialists using telestroke, enhancing preventive care, enhancing multidisciplinary collaboration, introducing uniform best practice protocols and bypass-protocols for the emergency medical services.

Discussion: The APSS overcame substantial obstacles to decrease discrepancies and to deliver integrated higher quality care. Telestroke has proven itself to be safe and feasible. The APSS works efficiently, which is in line to other projects worldwide, and is, based on limited results, cost effective. Further research on cost-effectiveness is necessary.

Keywords

stroke, neurology, telestroke, telemedicine, integrated care, Canada

Introduction

In the United States of America stroke is the third leading cause of death [1], and it is also a leading cause of functional impairment with 15–30% of survivors being permanently disabled [2]. The American Heart Association estimated that 780,000 strokes occur each year and annual costs are estimated to be 34 billion US dollar in the USA [1]. Stroke incidence increases with age. An aging population and unhealthy lifestyle choices threaten to cause a higher incidence of stroke and other chronic diseases in the years to come in the developed world [1, 3–5]. Risk factors are well known and guidelines for treatment and prevention are available [2, 6, 7]. Ischemic strokes account for approximately 87% of all strokes and hemorrhagic strokes for 10% [1].

A number of strategies are available to reduce the burden of stroke. Treatment of acute ischemic stroke with tPA (tissue plasminogen activator) increases neurological improvement. In eligible patients, the thrombolytic agent tPA has to be administered within 4.5 hours after onset of symptoms, the benefit is greater with earlier treatment [8, 9]. Of equal importance, timely secondary prevention is effective in reducing recurrent stroke in patients with stroke or TIA. Preventive measurements, such as blood pressure reduction, statin therapy and screening for additional risk factors, are proven effective [6].

Furthermore, coordinated stroke care results in improved outcomes and decreased length of hospital admission, as well as decreased costs. The implementation of telestroke as a component in a coordinated system of care increases the quality of care and the use of intravenous tPA at community hospitals [7]. Using the 'National Institutes of Health stroke scale', a reliable impression on the patients' state via video-conferencing can be derived [10–12]. Patients and healthcare providers report high levels of satisfaction with the telestroke intervention. International literature shows that the number of rural TIA and stroke patients that receive tPA treatment increases through the use of telestroke [12, 13].

The acute and preventive care for stroke patients is complex and requires the efforts and skills of all members of a multidisciplinary team [7]. In low population density countries like Canada delivery of standardized stroke care is even more challenging.

Systemic deficiencies of stroke care together with the increasing cardiovascular risk factor burden in the aging population led to the creation of a national initiative to improve stroke care in Canada. In 2005 the Canadian Stroke Strategy (CSS) was created. Following the CSS the province of Alberta developed the Alberta Provincial Stroke Strategy (APSS).

The APSS is a multidisciplinary and comprehensive approach to improve stroke care across the province throughout the continuum. Its goals are to improve access to specialized stroke care and improve the quality of acute care and follow up. Telestroke is an important component of the strategy.

This article relates the experience of a single province within Canada. The central question is: What is the process flow and the structure of the care pathway of the Alberta Provincial Stroke Strategy? The components the APSS introduced to achieve the present pathway, including telestroke, are explained.

Problem statement

Before the reformation of stroke care started in 2005, across Alberta, there was variable access to stroke care and little standardization of the approach to patients with a TIA. There was a gap between knowledge of best practice and actual practice for acute and non-acute stroke care, especially in rural hospitals. The situation before the creation of the APSS is reviewed here, to illustrate why a comprehensive intervention was adapted, and what elements were in the scope of the strategy.

In 2005 there were only four hospitals with neurologists, and only five hospitals that provided thrombolytic drugs to acute stroke patients, serving a population of 3.8 million Albertans, plus inhabitants of northern British Columbia (an adjoining province) and two northern Canadian territories which refer patients to Alberta. The province alone covers a 660,000 square kilometre area [14]. Other small hospitals in the province did not have neurologists, brain imaging possibilities or thrombolytic possibilities and rural physicians often lack training in neurological disorders. Often there was no timely communication with stroke neurologists.

People in northern Alberta and the adjoining territories had to travel up to 1500 kilometres to the capital city Edmonton to see a neurologist. In acute stroke situations their chance for timely thrombolysis was low only because of the travel time to the nearest hospital with the right expertise and equipment.

Besides, in many areas emergency medical services (EMS), lacked appropriate training in recognizing acute stroke symptoms and transported patients to the nearest hospital, irrespective of that facility's stroke expertise. This led both to patients not being in time for tPA treatment, and to many interhospital transports.

Furthermore, there was a deficiency of timely cross disciplinary communication in regards to stroke care. In a broader sense there was no multidisciplinary management of stroke at the provincial level to ensure

standards were being met and that access to appropriate care was available across the continuum.

There were no clear standards for the involvement of rehabilitation teams in the management of stroke inpatients, and transition of patients from inpatient or tertiary rehabilitation care to community living was not well developed.

The organization of preventive stroke care in Alberta initially was lacking. In 2005 there were three dedicated stroke prevention clinics providing information and care in the prevention of stroke, in the whole province.

Methods

This article describes the care pathway of the APSS and its structure using the case study methodology by Yin [15]. A care pathway can be evaluated by structure quality, process quality and outcome quality. The structure quality of the APSS is evaluated, using relevant subjects found in the literature [16].

Data collection is based on Yin's case study methodology. Yin [15] describes a case study as an empirical enquiry that investigates a contemporary phenomenon in depth and within its real-life context, especially when the boundaries between phenomenon and context are not clearly evident. The case is the Alberta Provincial Stroke Strategy and its telestroke program and the context is stroke care within Alberta. A characteristic of the case study is that it relies on multiple sources of evidence, with data to converge in a triangulation fashion. Yin distinguishes six sources of evidence: 1) documentation, 2) archival records, 3) interviews, 4) direct observations, 5) participant observations, and 6) physical artifacts or objects. In this case study physical artifacts were not used to support the findings. The five sources were used as follows.

Documentation: National and provincial policy papers [17, 18], local telestroke information [19, 20] protocols by the APSS [21–25] and an evaluation report published by the APSS [26] were collected. Reviews regarding telemedicine in stroke services worldwide were collected [12, 13].

Archival records: Organizational charts and a blueprint from the start of the APSS [27, 28] were used, as were statistics of the local and international population, its diseases and behavior [3–5, 14, 29]. International guidelines were used for evidence and evaluation of actions of the APSS [1, 2, 6–10].

Direct observations: The first author shadowed the stroke and telestroke service at the University of Alberta Hospital and the Walter C. Mackenzie Centre in Edmonton, during two weeks. She was involved with

daily practice and had many meetings with professionals, managers and patients.

Interviews: During the two weeks at the Comprehensive Stroke Centre the first author had extensive interviews with stroke specialists and paramedic personnel working with telestroke. For a broader view on the Albertan health system she interviewed twelve key players of health organisations in Alberta; Alberta Health Services and its subdivisions and Alberta Health and Wellness.

Participant observation: One of the three authors (TJ) works in the telestroke services of Edmonton and also serves as Chair of the Evaluation Pillar of the APSS. He brought to the research team his experience.

Triangulation of the data was done by the first two authors. Linkage and synthesis of collected data were accomplished. There was an extensive process of data cleaning and troubleshooting resulting in a final analysis with homogenous information. Where possible, data trends were validated through comparison to existing datasets, such as those of the Canadian Institute for Health Information.

Results

First the process flow of the integrated care pathway is described and outlined in flowcharts, with extra attention to telestroke. Then, the structure of the pathway of the APSS is described.

Care pathway of the APSS

The APSS adopted a broad approach to address all elements of stroke care, including health promotion, disease prevention, acute care, rehabilitation and community reintegration. **Box 1** presents a case to illustrate the process flow of stroke care in Alberta.

Flowcharts 1 and **2** give the general process flow of current stroke care, starting in the community, throughout the health system. The numbered hexagons indicate what factors were addressed and introduced by the APSS, and refer to the corresponding table (**Table 1**) explaining the improvements.

Characteristics of the care pathway

Organizational commitment

The Canadian Stroke Network and the Heart and Stroke foundation of Canada launched in 2005 the Canadian Stroke Strategy (CSS) to improve stroke care in Canada. Strategies were developed provincially to make best use of resources and customize local priorities. The vision and targets of the stroke strategy of Alberta, align with the visions and targets

Box 1. Patient case

Ft McMurray Alberta, Canada. The wife of Mr. S calls an ambulance because half an hour ago, her 65 year old healthy husband developed acute speech difficulties and was unable to move his right arm. Because of television commercials she was aware of the signs and symptoms of stroke and that she should call Emergency Medical Services (EMS). EMS personnel arrived within 6 minutes, recognized the symptoms of stroke through use of a stroke screening protocol, and brought Mr. S to the Provincial Stroke Center (PSC) in Fort McMurray.

At the PSC Mr. S is examined by a physician who activates the acute stroke protocol which involves standardized bloodwork, an urgent CT scan and activation of a critical care phone service (RAAPID) which alerts the telestroke neurologist from the CSC of Edmonton (450km south). After speaking with the physician via the RAAPID telephone service the telestroke neurologist coordinates the telestroke protocol. This involves reviewing the CT scan and the patient's lab results and past medical history on an electronic health record via secure internet connection. The telestroke neurologist then initiates a videoconference to conduct a very focused neurological examination of the patient centred around the National Institutes of Health Stroke Scale (NIHSS), finding him to have slurred speech, right facial and right arm weakness. The stroke specialist reviews the CT for Mr S. which shows normal cerebrum (no hemorrhage). Based on the examination, blood tests, and CT the specialist advises on further policy. Mr S received intravenous tPA and is admitted to hospital in Ft McMurray.

He receives a swallowing screen by a trained therapist and is allowed to eat after passing the screen. He is seen by rehabilitation therapists within 48 hours of his admission to work on physical and functional recovery. The physician in Ft McMurray, after consultation with the telestroke neurologist, orders appropriate investigations. It is determined by a multidisciplinary team that Mr. S will benefit from inpatient rehabilitation at a dedicated rehabilitation facility in Edmonton Alberta. Therefore, after a waiting period, Mr S. is transferred to Edmonton. After 4 weeks of rehabilitation he returns to his home in Ft McMurray functioning independently in the community.

Approximately 10 weeks after his stroke event he sees the Edmonton telestroke neurologist in followup via videoconference in Telestroke Stroke Prevention Clinic to ensure the workup has been complete and to optimize secondary prevention. The patient is accompanied in Ft McMurray by a local Stroke Navigator, in this case a nurse who coordinates care for stroke patients. She assists in ensuring that some followup blood tests are completed.

of the health department of the provincial government (Alberta Health and Wellness (AHW)) and Alberta Health Services (AHS) [18, 26]. Because of the aligning visions and the national movement to improve stroke care, these organisations have given the APSS financial and operational support. Telehealth is supported through AHW and AHS, and additionally through donor contributions, Health Canada First Nations and Inuit Health Branch, and Canada Health Infoway strategic investments.

Pathway project management and accountability

The APSS has a clearly defined organizational structure and chain of accountability [27]. The Alberta Stroke Council (ASC), formed to oversee the activities of the APSS is a collaboration of stakeholders in stroke care in Alberta: AHW, AHS, the Heart & Stroke Foundation of Alberta North West Territories & Nunavut and clinical experts [26]. The ASC reports to AHW and maintains accountability for all deliverables. Four pillars (1. health promotion and disease prevention, 2. emergency services and acute care, 3. rehabilitation and community reintegration, 4. evaluation and quality improvement) and other committees report to the ASC. The Pillars 1–3 manage the pathway and strongly collaborate to integrate the different aspects of the pathway. The ASC contained executive level membership from regional health authorities and later from AHS which allowed it to make binding decisions on allocation of funds. As a result the ASC could be quite flexible and responsive

to changing needs within the province and it could maintain its provincial perspective.

Format document


The APSS Blueprint is the initial overview document in which provincial capacities, gaps and proposed action plans are brought together. It provides the basis for the core business functions of the APSS and gives directions for moving forward to strengthen stroke services [28].

Professional content of pathway

At a national level Canadian best practice recommendations for stroke care [30] were developed and updated on a regular basis. These recommendations describe optimal care for stroke patients and best-practices for stroke prevention, medical care and recovery [17]. Most protocols of the APSS are based on these recommendations.

Multidisciplinary involvement

Stroke specialists in tertiary stroke centres initiated the APSS, and worked together with all stakeholders in stroke care in Alberta to make the strategy a success. They are all involved in the program and collaborate; care points are linked to one another in sequence resulting in a smoothed care process. This is evident in inpatient services between rehabilitation and medical teams, between primary care physicians and neurologists for TIA and acute stroke patients, and between administrators and all levels of health professionals.



Flowchart 1. Process flow of a patient with signs of stroke, from community to thrombolysis and admission.

*=this process flow is terminated; CSC: comprehensive stroke centre; CT: computed tomography; EMS: emergency medical services; GP: general practitioner; PSC: primary stroke centre; TIA: transient ischemic attack; tPA: tissue plasminogen activator.

Furthermore, provincial protocols were created by multidisciplinary teams.

Variance management

Professionals working with stroke patients are trained and educated and use their experience to be at variance from the process flow if needed. Each patient is evaluated several times by different specialist of the multidisciplinary team and decisions are based on the combination of protocols and the experts experience.

Patient involvement


Representatives of stroke survivors (the Stroke Recovery Association of Alberta) were involved in the original creation of the provincial stroke strategy and hold membership on Pillar 3 of the strategy. From the first

medical contact, patients are provided with disease information and the components of the care path. During follow up in the stroke prevention clinic and rehabilitation patients get tools and information to better lifestyle and set goals.

Telephone surveys of stroke survivors were conducted to evaluate and improve care. Quality of life, patient's experience while in hospital and on return to community were measured.

Implementation of pathway

In 2005 Alberta was successful in creating the APSS. Stroke advocates, consisting of the Heart and Stroke Foundation, stroke physicians, and other stroke clinicians submitted a business case to the government of



Flowchart 2. Process flow of a patient with stroke, from admission to community reintegration.

Alberta for a provincial stroke strategy. After the government of Alberta committed \$20 million CAD over two years in 2005 the APSS came into being.

One of the first actions of the APSS was to create the Pillar structure that would form the foundation for practice standards and for implementation of services (see project management). In addition, the nine health regions formed stroke steering committees to manage the APSS funds and to guide the implementation of stroke services in the health regions in the areas of prevention, emergency and acute care services, rehabilitation and community reintegration.

There remained challenges in the implementation of the strategy. At the present time there is one health care delivery organisation—AHS—that works with the Government of Alberta. This organization deals with current and future developments of the APSS. In 2005 AHS did not exist, and the Alberta Stroke Council had to collaborate with the 9 health regions that preceded AHS to make the strategy a success. At the start not all regions identified improvements in stroke care as a priority in the context of many other obligations. These regions needed extra effort to join APSS to improve stroke services at the provincial level.

Furthermore, not all practitioners conformed to best practise standards in stroke care and it was challenging to expose many to these standards. In many areas the problem was not willingness but rather the inability to hire, retain and maintain the skills of administrators and health care practitioners particularly in remote areas. The engagement of primary care physicians and generalist physicians in the process was also a challenge in some areas of the province. This was particularly so in the acute telestroke encounter which produced a new kind of interaction between emergency physicians, emergency room staff, and stroke neurologists.

The APSS includes one of Alberta's biggest telehealth initiatives: Telestroke. For its telestroke program, the APSS makes use of Alberta's pre-existing telehealth network.

Maintenance and development of the pathway

The Blueprint is not static but changes over time as events and new ideas evolve [28]. The evaluation pillar of APSS developed a de-identified data repository containing multiple datasets. The database uses administrative data and a variety of new datasets to track all stroke related hospital contacts, and admissions,

Table 1. Explanation to flowcharts items the APSS introduced to the stroke service in Alberta, and explanations

| | |
|----|--|
| 1 | To improve the awareness of common stroke symptoms and the knowledge of the importance of early treatment among the population, multimedia public awareness campaigns were conducted. |
| 2 | EMS were educated and trained to recognize stroke symptoms using screening tools [21, p.9]. They transport stroke patients directly to a primary stroke centre, bypassing small hospitals without thrombolytic facilities. Protocols [21] were developed in collaboration with EMS and moved into practice. Protocols indicate to transport patients with symptoms of acute stroke direct to a hospital with (access to) specialist care, and bypass hospitals that lack facilities for acute stroke care. |
| 3 | Stroke specialists are available in CSC 24/7 for stroke patients arriving at the CSC as well as patients in distant PSC. |
| 4 | Fifteen PSCs were introduced to allow for a better geographical coverage of stroke care. PSCs were developed by regional hospitals that were equipped with all necessary equipment: CT-scanner, swallowing assessments, physical therapy and occupational therapy. |
| 5 | Evidence based protocols for diagnostic testing [25, p.7] acute treatment [22], admission to a stroke unit [23, p.28], inpatient stroke care [23], rehabilitation [24], community reintegration [24] and secondary prevention [25] were developed, consistent with the Canadian Best Practice Recommendations for Stroke Care. [17] Best practice standards were created by multidisciplinary teams that involve physicians, administrators, nurses, occupational, physician and speech therapists, neurologists, internal medicine specialists and primary care physicians. |
| 6 | Rural professionals (physicians and nurses) of regional hospitals without stroke care facilities, had additional training and exposure to neurologic examination and stroke diagnosis and treatment. Protocols [21, 25] were developed in which specialist consultation is a standard step. |
| 7 | The critical care telephone service links PSC's 24/7 to a stroke specialist in a CSC using phone connection. The specialist advises on further management and investigations; in the case of a possible acute stroke CT-imaging and blood work. |
| 8 | To determine a definite diagnosis, the stroke-specialist has access to the blood work and brain imaging of the patient. CT scans are conducted at the PSC and can be retrieved in the CSC. |
| 9 | All PSC are connected to CSC by Telestroke . After the CT-scan is performed at the PSC the centres have renewed contact by Telestroke. Specialists in the CSC determine the definite diagnosis and advice on treatment in the PSC. The specialist assesses the imaging and the patients' status by doing a neurologic examination using videoconferencing. The specialist instructs the patient and with the assistance of the PSC physician and nurses, is able to determine the extent of the patient's deficits; the patient's score on the NIH stroke scale [10] is determined. The expert then decides, whether or not thrombolytic drugs should be administered, and the non-stroke specialist in the PSC administers the treatment [19]. Telestroke is also used to link urban rehabilitation professionals in physical therapy, occupational therapy, and especially speech therapy to patients in rural regions for treatment. |
| 10 | Early supported discharge teams were developed to allow for continued rehabilitation. The transition from hospital based care to community based care is smoothened. The inpatient rehabilitation teams refer inpatients to the early supported discharge teams which have been developed in several centres. Members of these community based teams will often meet the patient and family while they are in the hospital or, failing that, at least shortly after arrival back into their home. This provides the patient with multidisciplinary support (occupational therapy, physical therapy, speech therapy, social work) immediately upon their return to the community. |
| 11 | Patients can visit Stroke Prevention clinics , or interact with a neurologist via telehealth, using one of the 50 pre-existing hospital-based telehealth sites spread throughout the province. Usually a nurse or stroke care coordinator accompanies the patients at the telehealth site. Stroke standardized orders [24] now outline that occupational therapy, physical therapy and speech therapy need to be consulted within 48 hours of admission. Usage of protocols and telehealth sites, addresses access and referrals to these clinics for follow up and secondary prevention. |

outcome and costs. The repository is the basis for regular evaluation and improvements. In an interim evaluation report [26] the APSS stated goals for the next period, based on outcomes of the evaluation in terms of quality, new identified needs and disease outcomes.

Quality management

To control the quality of stroke care and allow for multi-stakeholder review the Alberta Stroke Council was introduced. Furthermore, APSS Pillar 4—Evaluation and Quality Improvement—was charged with both evaluating the effectiveness of the strategy and with creating a quality management framework. The

APSS governance recognized that to sustain the achieved gains there had to be continuing evaluation and measurement. The APSS identified fifty performance measures to evaluate progress made in delivery of stroke care [26]. For the near future the APSS still has goals to achieve, to make stroke care and prevention even better. In 2011 a quality management framework branded Alberta Stroke Improvement was created and funded through Pillar 4. This involved creation of a team of quality improvement consultants who conducted site visits of all primary and comprehensive stroke centres examining stroke services across the continuum.

Discussion

The APSS demonstrated that with the right effort, it is possible to successfully implement a comprehensive stroke strategy, with improvements in access and quality of care. In Canada the conditions were right for implementation when the APSS began. The Canadian Stroke Strategy had just been created and there was a pre-existing telehealth network although it needed upgrading and expansion to its current condition. Although the APSS benefited from favourable circumstances, much of the progress created was a result of the APSS itself therefore the APSS can serve as a model for other jurisdictions planning improvements in stroke care. Components that contributed to success which could be exported to other strategies are given here.

The APSS had province-wide influence on and control of care processes for stroke across the continuum. This influence was achieved by the comprehensive approach of the strategy. Though this approach professionals across almost all disciplines involved in the care of stroke patients in the province are involved in the program and collaborate. Care points are linked to one another in sequence improving the chances of a smooth care process. Transition points between community and inpatient services, medical team and rehabilitation team, inpatient teams and return to community services in particular have been enhanced. Better use of existing facilities, and the implementation of innovative technologies make the APSS a modern approach to improve care. The ability of the ASC to make final decisions on funding assisted its ability to control and guide the development of stroke care across the province.

Communication between different disciplines, urban and rural centres, the different therapies and the rehabilitation professionals also contributed to the success. Written protocols, the use of evidence-based interventions, such as stroke unit care, and abundant access to clinical stroke specialists are all present in the APSS and are associated with improved outcomes among patients treated for stroke [8]. Telestroke is also an important factor, it reduces travel time and gives experts the ability to cover a larger geographical area. Also important is the ongoing quality management and evaluation plan implemented in the APSS, increasing the chances of continued improvements.

Results on outcomes and quality of care, published in the APSS interim evaluation report [26] are promising. Outcome data were compared to the situation before the introduction of the APSS, and are available because of the thorough data repository of the APSS.

The proportion of patients who arrived within the treatable time window for tPA increased, possibly related

to the public awareness campaigns. Knowledge of symptoms decreased in the two years after the campaigns however, highlighting the need for continued public education. The proportion of patients receiving tPA increased due to a more streamlined care process, including the use of additional PSCs, and more patients were discharged back to their pre-admission place of residence. The number of patients referred to a stroke prevention clinic increased, and prescription of preventive medications on discharge increased reflective of better secondary preventive care. Protocols and training of EMS, together with the existence of PSC led to a reduction of the need to transfer stroke patients to comprehensive stroke centres. Patient satisfaction with care remained good.

The government invested 42.5 million Canadian dollars in the APSS over a 5-year period, averaging 8.5 million per year. A detailed economic analysis of the APSS is underway to determine the benefit for cost of the strategy. In the initial three years the actual number of stroke patients admitted to hospital decreased substantially. Whether this reduction can be sustained given the aging demographic of the population is unclear.

Studies and reviews on telestroke projects are difficult to compare, because projects were set up in areas with different needs. Nevertheless, all agree that the number of rural TIA and stroke patients that receive tPA appropriately, increases through telestroke [12, 13]. In Alberta about 23% of telestroke consultations resulted in thrombolytic treatment. In literature percentages range from 1.3% to 30%.

Strengths and limitations of this study

The availability of many documents, the explanations to them by many professionals and a fourteen days site visit in 2011 form the strength of this study. By using the theories and schemes of Yin we were able to give a systematic view on this innovative case of telestroke services. A limitation of the study is, that we do not offer controllable quantitative data on clinical outcome, utilization of resources, patient's experiences and costs. In this phase of sciences of telestroke services we can only say for sure: yes, telestroke service works and the results are promising but not yet evidence based.

Conclusion

In 2005 there was a national initiative to improve stroke care in Canada. In the province of Alberta, this led to the Alberta Provincial Stroke Strategy (APSS). The APSS is a comprehensive approach to improve stroke

care across the province. Stroke specialists in tertiary stroke centers initiated the APSS, and worked together with all stakeholders in stroke care in Alberta to make the strategy a success.

To address its goals, the APSS equipped small hospitals for diagnosing and treating acute stroke, introduced telestroke, and enhanced cross disciplinary communication and access to preventive care. Care efficiency is improved by coordinating the care, and integrating uniform use of best practice guidelines, assuring province wide uniformity in good practice. Quality of care improved by better use of stroke prevention clinics and improved multidisciplinary collaboration of health professionals within all disciplines.

Telestroke plays a major role in the approach of the APSS. People living in remote areas now have access to specialist stroke care, for acute care as well as follow up care. Furthermore, education and case conferencing between rural and urban professionals is conducted through telestroke.

The APSS has a clearly defined organizational structure and chain of accountability, with different committees bearing accountability for preventive care, acute care,

rehabilitation and evaluation. Based on a blueprint of the strategy the provincial Government gave its operational and financial commitment. The professional content of the strategy is based on national evidence-based guidelines, that were adjusted to the provincial needs in multidisciplinary teams. Data repositories make ongoing outcome and process evaluation possible. An evaluation and quality improvement committee is responsible for ongoing improvements and quality control.

Reviewers

Robbert Huijsman, Professor, Doctor, MBA, Professor Management and Organisation Elderly Care, Institute of Health Policy and Management, Erasmus University Rotterdam, The Netherlands

Lalit Kalra, Professor, Department of Neurosciences, Academic Neurosciences Centre, King's College London

Ron Heijnen, MD, Nursing Home Physician and stroke care researcher, Maastricht University, Maastricht, The Netherlands.

References

1. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, et al. Heart disease and stroke statistics - 2012 update: a report from the American heart association. *Circulation* 2012;125:e2–e220.
2. Goldstein LB, Bushnell CD, Adams RJ, Appel LJ, Braun LT, Chaturvedi S, et al. Guidelines for the primary prevention of stroke: a guideline from the American heart association/American stroke association. *Stroke* 2011;42:517–84.
3. Statistics Canada. Dependency ratio. [webpage on the internet]. [cited 2012 April 25; last updated: 2010 January 11]. Available from: <http://www.statcan.gc.ca/pub/82-229-x/2009001/demo/dep-eng.htm#hg>.
4. Statistics Canada. Health behaviours. [webpage on the internet]. [cited 2012 April 25; last updated: 2010 January 11]. Available from: <http://www.statcan.gc.ca/pub/82-229-x/2009001/deter/int3-eng.htm>.
5. Statistics Canada. Body mass index, overweight or obese, self-reported, adult, by age group and sex. [webpage on the internet]. [cited 2013 April 19; last updated 2013 June 17]. Available from: <http://www.statcan.gc.ca/tables-tableaux/sum-som/I01/cst01/health81a-eng.htm>.
6. Furie KL, Kasner SE, Adams RJ, Albers GW, Bush RL, Fagan SC, et al. Guidelines for the prevention of stroke in patients with stroke or transient ischemic attack: a guideline from the American heart association/American stroke association. *Stroke* 2011;42:227–76.
7. Summers D, Leonard A, Wentworth D, Saver JL, Simpson J, Spilker JA, et al. Comprehensive overview of nursing and interdisciplinary care of the acute ischemic stroke patient: a scientific statement of the American heart association. *Stroke* 2009;40:2911–44.
8. Adams HP Jr, Zoppo del G, Alberts MJ, Bhatt DL, Brass L, Furlan A, et al. Guidelines for the early management of adults with ischemic stroke: a guideline from the American heart association/American stroke association stroke council, clinical cardiology council, cardiovascular radiology and intervention council, and the atherosclerotic peripheral vascular disease and quality of care outcomes in research interdisciplinary working groups. *Stroke* 2007;38:1655–711.
9. Zoppo del GJ, Saver JL, Jauch EC, Adams HP. Expansion of the time window for treatment of acute ischemic stroke with intravenous tissue plasminogen activator: a science advisory from the American heart association/American stroke association. *Stroke* 2009;40:2945–8.
10. National institutes of health. NIH stroke scale. 2003. [cited 2012, December 12]. Available from: http://www.ninds.nih.gov/doctors/NIH_Stroke_Scale.pdf.
11. Wang S, Lee SB, Pardue C, Ramsingh D, Waller J, Gross H, et al. Remote evaluation of acute ischemic stroke: reliability of National Institutes of Health stroke scale via telestroke. *Stroke* 2003 Oct;34(10):e188–91.

12. Schwamm LH, Holloway RG, Amarenco P, Audebert HJ, Bakas T, Chumbler NR, et al. A review of the evidence for the use of telemedicine within stroke systems of care: a scientific statement from the American Heart Association/American Stroke Association. *Stroke* 2009 Jul;40(7):2616–34.
13. Johansson T, Wild C. Telemedicine in acute stroke management: systematic review. *International Journal of Technology Assessment in Health Care* 2010;26(2):149–55.
14. Statistics Canada. Land and freshwater area, by province and territory. [webpage on the internet]. [cited 2012 April 25; last updated: 2005 February 01]. Available from: <http://www.statcan.gc.ca/tables-tableaux/sum-som/l01/cst01/phys01-eng.htm>.
15. Yin RK. Case study research, design and methods. Fourth Edition. *Applied Social Research Methods series Vol. 5*. Thousand Oaks: SAGE Ltd.; 2009. p. 46–52.
16. Vanhaecht K, Witte de K, Depreitere R, Sermeus W. Clinical pathway audit tools: a systematic review. *Journal of Nursing Management* 2006;14:529–37.
17. Canadian Stroke Network. The quality of stroke care in Canada. 2011 [cited 2012 April 25]. Available from: <http://www.canadianstrokenetwork.ca/wp-content/uploads/2011/06/QoS-EN1.pdf>.
18. Government of Alberta. *Becoming the Best: Alberta's 5-year health action plan 2010–2015*. Canada: Alberta Health Services; Nov 2010.
19. Alberta Provincial Stroke Strategy. Telestroke. [webpage on the internet]. [cited 2012 April 25; last updated: 2010]. Available from: <http://www.strokestrategy.ab.ca/telestroke.html>.
20. Alberta Health and Wellness. Alberta Telehealth. [webpage on the internet]. [cited 2012 April 25]. Available from: www.health.alberta.ca/initiatives/telehealth.html.
21. Alberta provincial stroke strategy. Pre-hospital care. Canada; 2009 November. [cited 2012, December 12]. Available from: <http://strokestrategy.ab.ca/Prehospital%20Nov2409%20Final.pdf>.
22. Alberta provincial stroke strategy. Emergency management of acute stroke. Canada; 2010 January. [cited 2012, December 12]. Available from: <http://strokestrategy.ab.ca/ER%20Jan1910F.pdf>.
23. Alberta provincial stroke strategy. Inpatient care for acute stroke admissions. Canada; 2009 November. [cited 2012, December 12]. Available from: <http://strokestrategy.ab.ca/IP%20Nov1909%20Final.pdf>.
24. Alberta provincial stroke strategy. Rehabilitation and community reintegration. Canada; 2007 January. [cited 2012, December 12]. Available from: http://strokestrategy.ab.ca/PDFs/APSS_Rehabilitation_and_Community_Reintegration.pdf.
25. Alberta provincial stroke strategy. Secondary stroke prevention: best practice and pillar recommendations. Canada; 2009 February. [cited 2012 December 12]. Available from: <http://strokestrategy.ab.ca/Secondary%20Stroke%20Prevention%20March0909%20Final.pdf>.
26. Jeerakathil T, Burrige D, Thompson G, Fang S, Hill MD. *The Alberta Provincial Stroke Strategy: improving stroke care across Alberta. Interim evaluation report*. Edmonton: Alberta Provincial Stroke Strategy; Dec 2010.
27. Alberta Provincial Stroke Strategy. Organizational chart. Canada. [cited 2012 December 14]. Available from: http://www.strokestrategy.ab.ca/PDFs/APSS_Org_Chart.pdf.
28. Alberta provincial stroke strategy. Stroke blueprint: report of phase 1. Canada; 2006 May. [cited 2012 November 8]. Available from: http://www.strokestrategy.ab.ca/PDFs/APSS_Blueprint.pdf.
29. World Health Organization. Ten leading causes of death in 2008, by income group. [webpage on the internet]. [cited 2012 December 12] Available from: http://gamapserver.who.int/gho/interactive_charts/mbd/cod_2008/graph.html.
30. Canadian Stroke Network. Canadian best practice recommendations for stroke care. Canada; 2012 September. [cited 2012, December 15; last updated 2012 September 20]. Available from: <http://www.strokebestpractices.ca/index.php/overview/>.