

Management of Hepatitis C Virus/HIV Coinfection Among People Who Use Drugs in the Era of Direct-Acting Antiviral–Based Therapy

Lynn E. Taylor,¹ Tracy Swan,² and Gail V. Matthews^{3,4}

¹Department of Medicine, Brown University, Providence, Rhode Island; ²Treatment Action Group, New York, New York; ³Viral Hepatitis Clinical Research Program, The Kirby Institute, The University of New South Wales, and ⁴Infectious Diseases Clinical Services Unit, St Vincent's Hospital, Sydney, Australia

Where active antiretroviral therapy (ART) is accessible, human immunodeficiency virus (HIV) is a survivable illness and effective ART can reduce HIV transmission. Chronic hepatitis C virus (HCV) has emerged as a threat to the survival of individuals harboring both HCV and HIV, due to high prevalence and aggressive disease course. The HCV/HIV coinfection epidemic has been driven by people who inject drugs (PWID), although incident HCV is rising among HIV-infected men who have sex with men in the absence of drug injection. Coinfected individuals warrant aggressive treatment of both viruses; although early ART initiation is recommended to reduce the rate of liver disease progression, the most effective way to decrease HCV-related morbidity and mortality in coinfection is to achieve HCV viral eradication. Direct-acting antiviral (DAA) agents will soon revolutionize HCV treatment. Clinical data are needed regarding the efficacy of DAAs in coinfecting PWID. Drug–drug interaction studies between ART, DAAs, and opiate substitution therapy must be expedited. Coinfected PWID should have equitable and universal access to HIV/AIDS, HCV, and addiction prevention, care, and treatment. Essential basic steps include improving screening for both infections and engaging coinfecting PWID in HIV and HCV care early after diagnoses. Developing strategies to expand access to HCV therapy for coinfecting PWID is imperative to stem the HCV epidemic and limit the morbidity and mortality of those at greatest risk for HCV disease progression. The ultimate goal must be the elimination of HCV from all coinfecting PWID.

Keywords. hepatitis C virus; HIV; HCV/HIV coinfection; people who inject drugs (PWID); direct-acting antivirals (DAAs).

EPIDEMIOLOGY OF HCV/HIV

The human immunodeficiency virus (HIV) epidemic has been linked to injection drug use since the first reports of AIDS-related illness, and drug use continues to fuel the global HIV pandemic. Although injection drug use accounts for 10% of HIV infections overall, this proportion rises to 30% outside of Africa, and HIV incidence continues to rise among people who inject drugs (PWID) [1]. Sharing HIV-contaminated needles is the

most efficient means of spreading HIV after blood transfusion and is compounded by environmental conditions including stigma, discrimination, mental illness, poverty, and violence [1]. Unprotected sex is a further important HIV risk factor among PWID, in particular among women, female sex workers, and men who have sex with men (MSM). HIV-infected PWID transmit HIV sexually to noninjectors, and through vertical transmission to infants [1]. Whereas overall, 10% of HIV-infected persons are coinfecting with HCV, among PWID, HCV coinfection rates range from 50% to >90% [2].

An Emerging Incident HCV Epidemic

While prevalent HCV/HIV is driven by injection drug use, increasingly, incident HCV among HIV-infected MSM is associated with non-injection drug use. HCV incidence is rising among HIV-infected MSM internationally [3, 4]. In the Swiss HIV Cohort Study, from

Correspondence: Lynn E. Taylor, MD, The Warren Alpert Medical School of Brown University, The Miriam Hospital, 164 Summit Ave, Center for AIDS Research Bldg, Rm 156, Providence, RI 02906 (ltaylor@lifespan.org).

Clinical Infectious Diseases 2013;57(S2):S118–24

© The Author 2013. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

DOI: 10.1093/cid/cit326

1998 to 2011, HCV incidence rose 18-fold among HIV-infected MSM to an alarming rate of 4.1 cases per 100 person-years, while falling by 85% among PWID [5]. Most HIV-infected MSM acquiring HCV do not report injection drug use [4, 6]. In fact, among all HIV-infected persons, the majority of new HCV cases are occurring among non-PWID [5, 7, 8].

The reasons for this emerging epidemic are unclear. A conceptual model identifies a combination of drug use, sexual behaviors, and biologic factors (HIV status, syphilis) that work together to increase HCV risk [9]. The disturbing HCV incidence in a population previously thought to be at low risk motivates intervention. The European AIDS Treatment Network recommends quarterly screening for incident HCV in HIV-infected MSM, with the goal of rapid identification and treatment during the acute period of infection when interferon-based therapy is more successful [10–13].

NATURAL HISTORY OF HCV/HIV

Dual infection with HCV and HIV alters natural history of both diseases. HCV coinfection increases the risk of all-cause, AIDS-related, and liver-related death among HIV-infected people, despite use of antiretroviral therapy (ART) [14, 15]. HIV can accelerate HCV disease course, leading to more rapid progression to cirrhosis, liver failure, hepatocellular carcinoma (HCC), and increased HCV-related mortality [16–18]. Due to hastened progression and high prevalence in HIV-infected populations, HCV is a leading non-AIDS cause of death in HIV-infected persons in regions where ART is available [16, 18, 19]. The biologic basis for this is incompletely understood but may be related to impaired T-cell responses to HCV, to HIV's effect on hepatic cells, and to amplified microbial translocation promoting hepatic fibrosis [20, 21].

MEDICAL MANAGEMENT OF HCV/HIV

Coinfected individuals warrant aggressive treatment of both viruses. It has long been recognized that sustained virologic response (SVR) rates are lower in coinfection than in HCV monoinfection. Some of this may be due to biological and treatment-specific factors, but much may be linked to other issues found frequently in coinfecting populations including unstable socioeconomic circumstances, stigma, racial barriers, and competing priorities for healthcare. This is particularly true for individuals who have acquired disease through injection drug use. Improvements in HIV and HCV treatment and delivery, provide new hope that simple, well-tolerated, and effective treatment for both viruses may become widely available.

Screening and Diagnosis

The first challenge for providing universal treatment for HIV and HCV in coinfection is to improve screening for both

infections. HIV testing rates are improving, yet many HCV infections in HIV-infected individuals remain undiagnosed. In a report of long-term outcomes in a cohort of >2000 HIV-infected individuals, a third of whom were HCV antibody positive, mortality was 50% higher in the coinfecting group. Of concern, 30% of HCV-coinfecting persons were unaware of their HCV status [22]. Screening guidelines in HIV-infected populations support HCV antibody testing at HIV diagnosis and yearly thereafter to aid in the detection of often asymptomatic HCV infection (particularly relevant given the recent increases in newly acquired HCV infections seen within certain subgroups; see Epidemiology section) [13, 23]. Guidelines also advocate HCV RNA testing in the setting of new or unexplained alanine aminotransferase elevations and/or symptoms and risk exposure. If infection is confirmed, testing for HCV genotype, evaluation for other causes of liver disease, healthy liver lifestyle counseling, and evaluation and care for cirrhosis including screening for HCC should be performed [9]. HCV screening should be accompanied by education about prevention, transmission, natural history, and evolving therapies.

Treating HIV

Therapy for HIV infection is the cornerstone of effective medical management in coinfection. ART transforms HIV from a lethal process to a chronic, manageable disease; reduces AIDS-related morbidity and mortality and all-cause mortality; and is cost-effective in underresourced and well-resourced locales [24]. Twenty-three antiretroviral agents (ARVs) now exist in 6 mechanistic classes including 3 one-tablet, once-a-day combinations. Furthermore, the groundbreaking findings of a 2011 HIV Prevention Trials Network study verified the concept of HIV treatment as prevention [25]. This international randomized trial provided the strongest evidence to date that early ART initiation reduces rates of heterosexual transmission of HIV type 1 and clinical events, indicating both personal and public health benefits.

Previously, concerns around drug-induced liver injury (DILI) often discouraged clinicians from prescribing ART in coinfection. Although some earlier ARVs (ritonavir, nevirapine, stavudine, and didanosine) are associated with hepatotoxicity, mitochondrial toxicity, and/or steatosis, development of less hepatotoxic ARVs has reduced this risk. Additionally, untreated HIV may lead to high levels of immune activation, infection of hepatic stellate cells, and generation of inflammatory cytokines, thus promoting hepatic fibrosis and supporting the rationale to treat HIV early, especially in coinfection [20, 21]. Several studies suggest that better HIV control and higher CD4⁺ cell counts may be associated with slower fibrosis progression [26]. It is now clear that although coinfection increases risk of ART-related hepatotoxicity, liver disease progression is slower in patients receiving ART, and benefits outweigh risk. Irrespective of CD4⁺ cell count, HIV RNA level, HIV-related symptoms, and

opportunistic infections, early ART introduction is recommended to reduce the rate of progression of hepatic disease. In most cases, ART should precede HCV treatment, to decrease HIV-related morbidity and mortality and diminish HIV transmission. Treatments for HIV and HCV should not be started simultaneously, so that patients can adjust to each regimen sequentially. The most recent Department of Health and Human Services (DHSS) guidelines now state, “For most HIV/HCV-coinfected patients, including those with cirrhosis, the benefits of ART outweigh concerns regarding DILI. Therefore, ART should be considered for HIV/HCV-coinfected patients, regardless of CD4⁺ count (BII)” [27]. Given the wider availability of “liver-friendly” ART, including all 4 first-line regimens recommended by DHSS guidelines, ART can be safely prescribed to most coinfecting individuals—but only if issues of access, cost, and adherence are overcome.

Access and adherence are particularly relevant to PWID, who are less likely to be offered ART, and more likely to initiate ART at lower CD4⁺ cell counts than other populations [28, 29]. This is due in part to the perception that PWID may be less compliant with treatment and more likely to develop HIV drug resistance than other individuals. However, when appropriate support is provided, there is little evidence to support this. In fact, a recent meta-analysis examining ARV drug resistance in HIV-infected PWID versus non-IDUs found no difference in observed rates of resistance [30].

Treating HCV

ART may lessen the accelerated liver disease progression in coinfection but does not fully correct for it. In a meta-analysis of studies examining fibrosis rates in coinfecting individuals in the ART era, ART partially, but not fully, corrected the adverse impact of HIV on HCV progression [31]. Similarly, the Veterans Cohort Study recently reported that long-term liver disease-related outcomes were higher in their coinfecting versus HCV-monoinfected population, despite the fact that all HIV-infected patients were on ART [32]. The most effective way to reduce morbidity and mortality from liver disease in coinfection is to achieve SVR. SVR reduces hepatic fibrosis progression, liver stiffness, serum inflammatory markers of fibrosis, rates of hepatotoxicity, and overall mortality, as well as subsequent liver-related events (liver-related death, decompensation, HCC, or transplant) [33–38].

Treatment with pegylated interferon and ribavirin (peg-IFN/RBV) has typically resulted in SVR rates that are 10%–15% lower in HIV-infected individuals. For genotype 2/3 infections treated for 48 weeks, SVR rates have ranged from 44% to 73%, although this can be enhanced with weight-based RBV dosing. Data exist to support shortening treatment duration to 24 weeks in those with rapid virologic response; however, this is recommended only for those with low baseline HCV loads and minimal fibrosis—a fraction of the coinfecting population.

Genotype 1 SVR rates have been dismal, from 14% to 38%. Low SVR rates, treatment toxicities, polypharmacy, and the need for liver biopsy deter coinfecting individuals from attempting treatment, while these factors plus comorbidities prevalent in HIV deter physicians from prescribing HCV therapy.

HCV TREATMENT AMONG HCV/HIV-COINFECTED PWID

The goal of HCV viral eradication in coinfection has been challenging. Although international guidelines endorse considering all coinfecting patients for HCV treatment [27, 39], implementation of these guidelines has been limited because most coinfecting persons are current or former PWID. Historically, HCV therapy has been underused in this population. Few data are available regarding chronic HCV treatment outcomes among coinfecting PWID (Supplementary Table 1). Most studies of chronic HCV treatment efficacy in PWID involve HCV monoinfection; monoinfection data do not demonstrate differences in SVR rates between PWID and non-IDUs. Across the 11 published studies involving coinfecting PWID [40–50], 9 reported HCV treatment initiation specifically by HIV status, but SVR was not delineated by HIV status and is therefore unknown in more than half (6) of these. Data are lacking to consider coinfecting PWID for HCV treatment differently than non-IDUs.

DAAs in HCV/HIV

HCV treatment is rapidly evolving; in 2011, the first HCV protease inhibitors (PIs), boceprevir and telaprevir, were approved. Boceprevir- and telaprevir-based regimens significantly increase treatment efficacy for genotype 1 patients, regardless of HIV status. Treatment outcomes in coinfection from phase 2b studies with telaprevir and boceprevir have been reported [51]. In the telaprevir study, 38 genotype 1 treatment-naive coinfecting participants were treated with telaprevir plus peg-IFN/RBV for 12 weeks, followed by an additional 36 weeks of peg-IFN/RBV. Outcomes were compared to 22 coinfecting controls treated with peg-IFN/RBV for 48 weeks. SVR at week 24 data (defined as an undetectable HCV RNA level 24 weeks after treatment discontinuation) demonstrated efficacy to be considerably greater in the triple- (74%) versus the dual-therapy arm (45%). Adverse events occurred more frequently with triple therapy, and the pill burden was sizeable, particularly for those on efavirenz who were required to take 3 telaprevir pills 3 times daily. However there was no difference according to ART use and no loss of HIV control.

Although formal drug–drug interaction studies were not performed between telaprevir and ARVs, drug levels of telaprevir and ARVs the participants were already taking were measured. These results provided reassurance that interactions with

Table 1. Antiretroviral Agents Suitable for Concomitant Treatment of HIV and Hepatitis C Virus in the Context of Opioid Substitution Therapy

HCV PI	Antiretroviral Agents, by Class			
	NRTI	NNRTI	PI	II
Boceprevir	Tenofovir ^a Abacavir Lamivudine Emtricitabine	Etravirine ^b	An ongoing study is evaluating drug–drug interactions between boceprevir and HIV PIs	Raltegravir
Telaprevir	Tenofovir ^a Abacavir Lamivudine Emtricitabine	Etravirine Ralpivirine Efavirenz (telaprevir dose must be increased from 750 mg/tid to 1125 mg/tid)	^a Atazanavir/ritonavir	Raltegravir

Abbreviations: HIV, human immunodeficiency virus; II, HIV integrase inhibitor; NNRTI, nonnucleoside reverse transcriptase inhibitor; NRTI, nucleoside reverse transcriptase inhibitor; PI, protease inhibitor; tid, 3 times a day.

^a Monitoring for toxicity is recommended.

^b In healthy volunteers, boceprevir decreased etravirine area under the curve, C_{max} , and C_{min} , but clinical significance is not clear; when etravirine is coadministered with methadone or buprenorphine, clinical monitoring is recommended and dose adjustments may be necessary.

Sources: De Kanter C, Blonk M, Colbers A, et al. The influence of the HCV protease inhibitor boceprevir on the pharmacokinetics of the HIV integrase inhibitor raltegravir. In: 19th Conference on Retroviruses and Opportunistic Infections, Seattle, WA, 5–8 March 2012. Abstract 772LB.

Hammond KP, Wolfe P, Burton JR Jr, et al. Pharmacokinetic interaction between boceprevir and etravirine in HIV/HCV seronegative volunteers. *J Acquir Immune Defic Syndr* 2013; 62:67–73.

US Food and Drug Administration. Intelence (etravirine) Prescribing information. August 2012. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/022187s010lbl.pdf Accessed 5 February 2013.

Van Heeswijk RP, Beumont M, Kauffman RS, Garg V. Review of drug interactions with telaprevir and antiretrovirals [published online ahead of print 23 January 2013]. *Antivir Ther* 2013. PMID: 23344266.

tenofovir, atazanavir, and efavirenz plus telaprevir were manageable. Other ARVs that may be coadministered with telaprevir-based treatment include raltegravir, etravirine, rilpivirine, and 3 additional nucleoside reverse transcriptase inhibitors (NRTIs) (Table 1). However, telaprevir dose increase is necessary with concomitant efavirenz, clinical and laboratory monitoring for hyperbilirubinemia is recommended with atazanavir, and increased monitoring is warranted with tenofovir.

In the boceprevir study, similar outcomes were observed [52]. Sixty-four coinfecting genotype 1 treatment-naïve participants were randomized to 4 weeks of peg-IFN/RBV, followed by 44 weeks of peg-IFN/RBV/boceprevir, compared to 34 participants randomized to peg-IFN/RBV for 48 weeks. Again the SVR at week 12 data (defined as an undetectable HCV RNA level 12 weeks after treatment discontinuation) showed a similar efficacy benefit (61% in triple vs 27% in dual arm), but adverse events were greater in the triple arm. Although HIV breakthrough was similar in both arms (2 on triple and 3 on dual therapy), a subsequent drug–drug interaction study on healthy volunteers demonstrated considerable interactions between boceprevir and the HIV PIs. Area under the curve concentrations for darunavir, atazanavir, and lopinavir were reduced by 34%–44% in the presence of boceprevir, leading to a subsequent Food and Drug Administration recommendation not to use these drugs with boceprevir. Additionally, fosamprenavir, ritonavir, and efavirenz are not recommended in combination with boceprevir. ARVs that

may be coadministered with boceprevir-based treatment include raltegravir, etravirine, and 4 NRTIs (Table 1). In summary, although these drugs offer efficacy benefit, restrictions on ARV use, additional pill burden, and toxicity impede their widespread use.

OST IN HCV/HIV COINFECTION

HCV, HIV, and opioid dependence are overlapping epidemics. HCV-monoinfected PWID can be successfully treated for HCV with and without opioid substitution therapy (OST) [53]. OST facilitates HCV treatment among PWID in diverse settings [44, 45], reduces HIV acquisition, and is associated with enhanced HIV treatment outcomes (including increased ARV initiation rates, adherence, and CD4⁺ cell counts). Fewer interactions exist between buprenorphine and ARVs than between methadone and ARVs [54–56].

Opioid Substitution Treatment With DAAs

PWID and individuals on OST were excluded from the first HCV PI studies in coinfection. Among all coinfection telaprevir studies to date, exclusion criteria regarding PWID and OST are heterogeneous, ranging from no mention of excluding PWID or use of OST, to excluding individuals with alcohol intake and/or substance abuse that may represent an obstacle for participation. Among all coinfection boceprevir studies to date, exclusion criteria regarding PWID and OST are similarly varied. Some do not

mention excluding PWID or use of OST, whereas others exclude individuals with active drug or alcohol use or dependence that, in the opinion of the investigator, would interfere with adherence to study requirements; those receiving OST but not enrolled in an OST maintenance program; those with current evidence of substance abuse within 3 years of screening; those with a history of a clinical diagnosis within the past 6 months of substance abuse prior to day 1; or those with any current evidence of substance abuse of alcohol or other drugs. Some HCV-monoinfection PI studies have allowed inclusion of persons on OST, although safety and treatment outcome data have not been presented in this subpopulation. Fortunately, pharmacokinetic studies suggest that OST can be coadministered with boceprevir and telaprevir; although dose adjustments may not be necessary, clinical monitoring for withdrawal symptoms is recommended.

Opiate Substitution Treatment With ART and DAAs

Selecting ART for patients on methadone involves consideration of drug–drug interactions. Among the single-tablet regimens, Complera (rilpivirine/tenofovir/emtricitabine) cannot be used with methadone due to an increased risk of QTc prolongation and torsade des pointes. Efavirenz, the backbone of Atripla (efavirenz/tenofovir/emtricitabine), lowers methadone plasma concentration; dose adjustments are typically needed to avert withdrawal symptoms. Stribild (elvitegravir/cobicistat/tenofovir/emtricitabine) can be coadministered with methadone and buprenorphine, but there are no available data on drug–drug interactions with boceprevir or telaprevir.

Complex drug–drug interactions may exist between ARVs, DAAs, and OST. A list of ARVs suitable for concomitant treatment of HIV and HCV in the context of OST is provided in Table 1. ARV options may be further limited by multidrug resistance or financial constraints. Cotreatment of HCV and HIV is further complicated by overlapping drug toxicities, increased pill burden, and food restrictions.

Failure to perform drug–drug interaction studies that facilitate inclusion of high-prevalence patient groups in clinical trials perpetuates a vicious cycle; after approval, treatment is often withheld due to lack of information on safety, efficacy, and tolerability. Clinicians treating coinfecting patients on stable OST with advanced liver disease—individuals most likely to benefit from successful HCV treatment—often do so without adequate data. To date, drug–drug interaction studies with ARVs and HCV PIs have been performed only in healthy volunteers. Drug levels may be different in HIV-infected and HCV/HIV-coinfecting patients, especially those with hepatic impairment.

INVESTIGATIONAL DAAs IN HCV/HIV

Developments in HCV therapeutics will likely revolutionize HCV treatment for coinfecting individuals in the near future. It is anticipated that over time, difficulties seen with the first-

generation HCV PIs, including those of toxicity and drug–drug interactions, are likely to disappear. Studies are already under way with several second-generation drugs including simeprevir, faldeprevir, and daclatasavir, all of which have improved tolerability profiles. One of the most exciting drugs in development is sofosbuvir—a potent, tolerable direct-acting nucleotide polymerase inhibitor with a high genetic barrier to resistance, currently in phase 3 studies in monoinfection and coinfection. A phase 1b study of 19 coinfecting individuals suggests that HCV declines observed with 7 days' sofosbuvir monotherapy are identical to those in HCV monoinfection [57]. Drug–drug interaction studies with ARVs have begun; in a phase 1, fixed-sequence, 4-cohort study involving healthy volunteers, pharmacokinetic interactions between sofosbuvir and common ARVs demonstrated no clinically significant interactions or change in levels of either HIV or HCV drug [58].

ENHANCING DELIVERY OF CARE

Given the barriers to treatment that often exist for coinfecting PWID (lack of engagement in healthcare, incarceration, mental health disorders, and fractured social networks), urgent attention is needed to improve mechanisms through which this disenfranchised group can be treated. Programs that enhance engagement, retention, and adherence including directly observed therapy, supervised dispensing, patient–provider support, and OST have been shown to be effective. There are a growing number of successful models of HCV care for coinfecting PWID in settings including HIV clinics, community health centers, hospitals, and prisons. Common facilitators include multidisciplinary care, integrated addiction services, peer-based support, psychiatric care, “one-stop shopping,” risk reduction, case management, and nursing assessment. These approaches require implementation on a larger scale than that currently achieved, with investment in infrastructure at the healthcare service level.

CONCLUSIONS

Coinfecting PWID are among the most likely to be affected by HCV but are the least likely to have access to HCV treatment. Benefits of therapeutic advances in HCV will be limited for coinfecting PWID until barriers are overcome and HCV care enhanced on a massive scale. Only a revolution in HCV treatment delivery, similar to that occurring in HCV treatment development, will avert decades of preventable morbidity and mortality. Coordinated approaches to prevention and treatment of HCV/HIV among PWID are imperative.

Coinfecting PWID should have equitable and universal access to HIV/AIDS, HCV, and addiction prevention, care, and treatment. Drug use itself is not a contraindication to HCV therapy and treatment should not necessarily be deferred until this is

“stabilized.” Coinfected PWID must be engaged in HCV care early after HIV and HCV diagnoses. It may take time to adapt to ART and gain control of HIV before HCV treatment may be initiated. During this time, patients should have access to HCV education and evaluation. The stage of HCV, status of HIV, and patient preference should drive the timing of HCV therapy while DAAs evolve. If treatment is deferred, reevaluation should occur biannually and relative contraindications should be addressed.

In summary, the ultimate goal must be the elimination of HCV from all coinfecting PWID. To achieve this, substantial issues of cost, access, and treatment delivery will need to be overcome. Ensuring that all coinfecting PWID receive comprehensive HIV care and consideration for HCV treatment is the important first step.

Supplementary Data

Supplementary materials are available at *Clinical Infectious Diseases* online (<http://cid.oxfordjournals.org/>). Supplementary materials consist of data provided by the author that are published to benefit the reader. The posted materials are not copyedited. The contents of all supplementary data are the sole responsibility of the authors. Questions or messages regarding errors should be addressed to the author.

Notes

Acknowledgments. We thank Vaughn Bryant for his assistance with the literature review and Brittany Rizzo for administrative assistance.

Disclaimer. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse, the National Institute of Allergy and Infectious Diseases, or the National Institutes of Health.

Financial support. This work was supported by the National Institute on Drug Abuse (K23DA020383 to L. E. T.).

Supplement sponsorship. This article was published as part of a supplement entitled “Prevention and Management of Hepatitis C Virus Among People Who Inject Drugs: Moving the Agenda Forward,” sponsored by an unrestricted grant from the International Network on Hepatitis in Substance Users (INHSU), The Kirby Institute (University of New South Wales), Abbvie, Gilead Sciences, Janssen-Cilag, and Merck.

Potential conflicts of interest. G. M. has consulted for Janssen and ViiV; received payment for lectures from Bristol-Myers Squibb, Gilead, Roche, and Janssen; and received travel/accommodations/meeting expenses support from Gilead, MSD, and Roche. All other authors report no potential conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Stockman JK, Strathdee SA. HIV among people who use drugs: a global perspective of populations at risk. *J Acquir Immune Defic Syndr* **2010**; 55:S17–22.
2. Walsh N, Maher L. HIV and viral hepatitis C coinfection in people who inject drugs: implications of new direct acting antivirals for hepatitis C virus treatment. *Curr Opin HIV AIDS* **2012**; 7:339–44.
3. Bottieau E, Apers L, Esbroeck MV, Vandenbruaene M, Florence E. Hepatitis C virus infection in HIV-infected men who have sex with men: sustained rising incidence in Antwerp, Belgium, 2001–2009. *Euro Surveill* **2010**; 15:19673.
4. van der Helm JJ, Prins M, del Amo J, et al. The hepatitis C epidemic among HIV-positive MSM: incidence estimates from 1990 to 2007. *AIDS* **2011**; 25:1083–91.
5. Wandeler G, Gsponer T, Bregenzer A, et al. Hepatitis C virus infections in the Swiss HIV Cohort Study: a rapidly evolving epidemic. *Clin Infect Dis* **2012**; 55:1408–16.
6. Garg S, Taylor LE, Grasso C, Mayer KH. Prevalent and incident hepatitis C virus infection among HIV-infected men who have sex with men engaged in primary care in a Boston community health center. *Clin Infect Dis* **2013**; 56:1480–7.
7. Jin F, Prestage GP, Matthews G, et al. Prevalence, incidence and risk factors for hepatitis C in homosexual men: data from two cohorts of HIV-negative and HIV-positive men in Sydney, Australia. *Sex Transm Infect* **2010**; 86:25–8.
8. Rauch A, Rickenbach M, Weber R, et al. Unsafe sex and increased incidence of hepatitis C virus infection among HIV-infected men who have sex with men: the Swiss HIV Cohort Study. *Clin Infect Dis* **2005**; 41:395–402.
9. Taylor LE, Swan T, Mayer KH. HIV coinfection with hepatitis C virus: evolving epidemiology and treatment paradigms. *Clin Infect Dis* **2012**; 55(suppl 1):S33–42.
10. Lambers FA, Brinkman K, Schinkel J, et al. Treatment of acute hepatitis C virus infection in HIV-infected MSM: the effect of treatment duration. *AIDS* **2011**; 25:1333–6.
11. McGovern BH, Nagami EH, Birch CE, et al. Rate of sustained virologic response in relation to baseline hepatitis C virus (HCV) RNA level and rapid virologic clearance in persons with acute HCV infection. *J Infect Dis* **2009**; 200:877–81.
12. Piroth L, Larsen C, Binquet C, et al. Treatment of acute hepatitis C in human immunodeficiency virus-infected patients: the HEPAIG study. *Hepatology* **2010**; 52:1915–21.
13. Acute hepatitis C in HIV-infected individuals: recommendations from the European AIDS Treatment Network (NEAT) consensus conference. *AIDS* **2011**; 25:399–409.
14. Hernando V, Perez-Cachafeiro S, Lewden C, et al. All-cause and liver-related mortality in HIV positive subjects compared to the general population: differences by HCV co-infection. *J Hepatol* **2012**; 57:743–51.
15. van der Helm J, Geskus R, Sabin C, et al. Effect of HCV infection on cause-specific mortality after HIV seroconversion, before and after 1997. *Gastroenterology* **2013**; 144:751–60.e2.
16. Holmberg SD, Ly KN, Xing J, Klevens M, Jiles R, Ward JW. The growing burden of mortality associated with viral hepatitis in the United States, 1999–2007. In: 62th Annual Meeting of the American Association for the Study of Liver Diseases, San Francisco, California, **2011**.
17. Graham CS, Baden LR, Yu E, et al. Influence of human immunodeficiency virus infection on the course of hepatitis C virus infection: a meta-analysis. *Clin Infect Dis* **2001**; 33:562–9.
18. de Ledinghen V, Barreiro P, Foucher J, et al. Liver fibrosis on account of chronic hepatitis C is more severe in HIV-positive than HIV-negative patients despite antiretroviral therapy. *J Viral Hepat* **2008**; 15:427–33.
19. Weber R, Sabin CA, Friis-Moller N, et al. Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch Intern Med* **2006**; 166:1632–41.
20. Kim AY, Chung RT. Coinfection with HIV-1 and HCV—a one-two punch. *Gastroenterology* **2009**; 137:795–814.
21. Tuyama AC, Hong F, Saiman Y, et al. Human immunodeficiency virus (HIV)-1 infects human hepatic stellate cells and promotes collagen I and monocyte chemoattractant protein-1 expression: implications for the pathogenesis of HIV/hepatitis C virus-induced liver fibrosis. *Hepatology* **2010**; 52:612–22.
22. Branch AD, Van Natta ML, Vachon ML, Dieterich DT, Meinert CL, Jabs DA. Mortality in hepatitis C virus-infected patients with a diagnosis of AIDS in the era of combination antiretroviral therapy. *Clin Infect Dis* **2012**; 55:137–44.

23. Rockstroh JK, Bhagani S, Benhamou Y, et al. European AIDS Clinical Society (EACS) guidelines for the clinical management and treatment of chronic hepatitis B and C coinfection in HIV-infected adults. *HIV Med* **2008**; 9:82–8.
24. Walensky RP, Freedberg KA, Weinstein MC, Paltiel AD. Cost-effectiveness of HIV testing and treatment in the United States. *Clin Infect Dis* **2007**; 45(suppl 4):S248–54.
25. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med* **2011**; 365:493–505.
26. Brau N, Salvatore M, Rios-Bedoya CF, et al. Slower fibrosis progression in HIV/HCV-coinfected patients with successful HIV suppression using antiretroviral therapy. *J Hepatol* **2006**; 44:47–55.
27. US Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Available at: <http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>. Accessed 13 February 2013.
28. McGowan CC, Weinstein DD, Samenow CP, et al. Drug use and receipt of highly active antiretroviral therapy among HIV-infected persons in two U.S. clinic cohorts. *PLoS One* **2011**; 6:e18462.
29. Mehta SH, Kirk GD, Astemborski J, Galai N, Celentano DD. Temporal trends in highly active antiretroviral therapy initiation among injection drug users in Baltimore, Maryland, 1996–2008. *Clin Infect Dis* **2010**; 50:1664–71.
30. Werb D, Mills EJ, Montaner JS, Wood E. Risk of resistance to highly active antiretroviral therapy among HIV-positive injecting drug users: a meta-analysis. *Lancet Infect Dis* **2010**; 10:464–9.
31. Thein HH, Yi Q, Dore GJ, Krahn MD. Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis. *AIDS* **2008**; 22:1979–91.
32. Re VL. Increased risk of hepatic decompensation and hepatocellular carcinoma in HIV/HCV-co-infected patients compared to HCV-mono-infected patients despite combination antiretroviral therapy. In: 19th International AIDS Conference, Washington, DC, **2012**.
33. Ingiliz P, Valantin MA, Preziosi P, et al. Influence of interferon-based therapy on liver fibrosis progression in HIV/HCV coinfecting patients: a retrospective repeated liver biopsy analysis. *J Hepatol* **2012**; 56:49–54.
34. Labarga P, Soriano V, Vispo ME, et al. Hepatotoxicity of antiretroviral drugs is reduced after successful treatment of chronic hepatitis C in HIV-infected patients. *J Infect Dis* **2007**; 196:670–6.
35. Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology* **2009**; 50:1056–63.
36. Macias J, del Valle J, Rivero A, et al. Changes in liver stiffness in patients with chronic hepatitis C with and without HIV co-infection treated with pegylated interferon plus ribavirin. *J Antimicrob Chemother* **2010**; 65:2204–11.
37. Mira JA, Rivero-Juarez A, Lopez-Cortes LF, et al. Benefits from sustained virological response to pegylated interferon plus ribavirin in HIV/HCV-coinfecting patients with compensated cirrhosis. *Clin Infect Dis* **2013**; 56:1646–53.
38. Berenguer J, Alvarez-Pellicer J, Martin PM, et al. Sustained virological response to interferon plus ribavirin reduces liver-related complications and mortality in patients coinfecting with human immunodeficiency virus and hepatitis C virus. *Hepatology* **2009**; 50:407–13.
39. European AIDS Clinical Society. Guidelines, version 6.1. Available at: <http://www.europeanclinicalsociety.org/images/stories/EACS-Pdf/EacsGuidelines-v6.1-2edition.pdf>. Accessed 8 February 2013.
40. Seidenberg A, Rosemann T, Senn O. Patients receiving opioid maintenance treatment in primary care: successful chronic hepatitis C care in a real world setting. *BMC Infect Dis* **2013**; 13:9.
41. Jafferbhoy H, Miller MH, Dunbar JK, Tait J, McLeod S, Dillon JF. Intravenous drug use: not a barrier to achieving a sustained virological response in HCV infection. *J Viral Hepat* **2012**; 19:112–9.
42. Neukam K, Mira JA, Gilibert I, et al. Efficacy of chronic hepatitis C therapy with pegylated interferon and ribavirin in patients on methadone maintenance treatment. *Eur J Clin Microbiol Infect Dis* **2012**; 31:1225–32.
43. Melin P, Chousterman M, Fontanges T, et al. Effectiveness of chronic hepatitis C treatment in drug users in routine clinical practice: results of a prospective cohort study. *Eur J Gastroenterol Hepatol* **2010**; 22:1050–7.
44. Litwin AH, Harris KA Jr, Nahvi S, et al. Successful treatment of chronic hepatitis C with pegylated interferon in combination with ribavirin in a methadone maintenance treatment program. *J Subst Abuse Treat* **2009**; 37:32–40.
45. Grebely J, Raffa JD, Meagher C, et al. Directly observed therapy for the treatment of hepatitis C virus infection in current and former injection drug users. *J Gastroenterol Hepatol* **2007**; 22:1519–25.
46. Seal KH, Currie SL, Shen H, et al. Hepatitis C treatment candidacy and outcomes among 4318 US veterans with chronic hepatitis C virus infection: does a history of injection drug use matter? *J Clin Gastroenterol* **2007**; 41:199–205.
47. Mehta SH, Lucas GM, Mirel LB, et al. Limited effectiveness of antiviral treatment for hepatitis C in an urban HIV clinic. *AIDS* **2006**; 20:2361–9.
48. Sylvestre DL, Litwin AH, Clements BJ, Gourevitch MN. The impact of barriers to hepatitis C virus treatment in recovering heroin users maintained on methadone. *J Subst Abuse Treat* **2005**; 29:159–65.
49. Matthews G, Kronborg IJ, Dore GJ. Treatment for hepatitis C virus infection among current injection drug users in Australia. *Clin Infect Dis* **2005**; 40(suppl 5):S325–9.
50. Van Thiel DH, Anantharaju A, Creech S. Response to treatment of hepatitis C in individuals with a recent history of intravenous drug abuse. *Am J Gastroenterol* **2003**; 98:2281–8.
51. Sulkowski MS, Sherman KE, Soriano V, et al. Telaprevir in combination with peginterferon alfa-2a/ribavirin in HCV/HIV co-infected patients: SVR24 final study results. In: 63rd Annual Meeting of the American Association for the Study of Liver Diseases, Boston, Massachusetts, 2012.
52. Mallolas J, Pol S, Rivero A, et al. Boceprevir plus peginterferon/ribavirin for the treatment of HCV/HIV co-infected patients: end of treatment (week 48) interim results [SVR12 data in oral presentation]. In: 47th Annual Meeting of the European Association for the Study of the Liver, Barcelona, Spain, 2012.
53. Hellard M, Sacks-Davis R, Gold J. Hepatitis C treatment for injection drug users: a review of the available evidence. *Clin Infect Dis* **2009**; 49:561–73.
54. Moatti JP, Carrieri MP, Spire B, Gastaut JA, Cassuto JP, Moreau J. Adherence to HAART in French HIV-infected injecting drug users: the contribution of buprenorphine drug maintenance treatment. The Manif 2000 study group. *AIDS* **2000**; 14:151–5.
55. Altice FL, Bruce RD, Lucas GM, et al. HIV treatment outcomes among HIV-infected, opioid-dependent patients receiving buprenorphine/naloxone treatment within HIV clinical care settings: results from a multisite study. *J Acquir Immune Defic Syndr* **2011**; 56(suppl 1):S22–32.
56. Palepu A, Tyndall MW, Joy R, et al. Antiretroviral adherence and HIV treatment outcomes among HIV/HCV co-infected injection drug users: the role of methadone maintenance therapy. *Drug Alcohol Depend* **2006**; 84:188–94.
57. Rodriguez-Torres M, Gonzalez M, Rossi S, et al. HIV/HCV coinfecting and HCV mono-infected patients have similar early HCV viral kinetics with the potent HCV nucleotide polymerase inhibitor sofosbuvir (SOF). In: 52nd Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, California, 9–12 September **2012**.
58. Kirby B, Mathias A, Rossi S, Moyer C, Shen G, Brian P. Kearney Gilead Sciences, Inc. GS-7977 & HIV ARTs PK—no clinically significant pharmacokinetic interactions between sofosbuvir (GS-7977) and HIV antiretrovirals atirapla, rilpivirine, darunavir/ritonavir, or raltegravir in healthy volunteers. In: 63rd Annual Meeting of the American Association for the Study of Liver Diseases, Boston, Massachusetts, 9–13 November **2012**.