

Published in final edited form as:

Bioorg Med Chem Lett. 2010 August 1; 20(15): 4468–4471. doi:10.1016/j.bmcl.2010.06.042.

Synthesis and *in vitro* evaluation of *N*-alkyl-3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-one analogs as potential anticancer agents

Narsimha Reddy Penthala, Thirupathi Reddy Yerramreddy, Nikhil Reddy Madadi, and Peter A. Crooks*

Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, KY 40536, USA

Abstract

A series of novel 3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-one analogs (**3**) have been synthesized under microwave irradiation and conventional heating methods. These analogs were evaluated for *in vitro* cytotoxicity against a panel of 57 human tumor cell lines. Compound **3o** had GI₅₀ values of 190 nM and 750 nM against A549/ATTC non-small cell lung cancer and LOX IMVI melanoma cell lines, respectively, and both **3n** and **3o** exhibited GI₅₀ values ranging from 2–5 μM against CCRF-CEM, HL-60(TB), K-562, MOLT-4, and RPMI-8226 leukemia cell lines. These results indicate that *N*-4-methoxybenzyl-3-hydroxy-(2-imino-3-methyl-5-oxo-4-yl)indolin-2-one analogs may be useful leads for anticancer drug development.

Keywords

N-alkyl isatins; creatinine; microwave irradiation; *in vitro* cytotoxicity

Many isatin analogs have been reported as potent and selective cytotoxic agents against cancer cells. Vine et al¹ reported that the substitution of bromo groups at the C-5 and C-7 positions of isatin afforded an analog (Fig. 1. structure **A**) that exhibited potent cytotoxic activity. These investigators also reported that introduction of *N*-benzyl, *N*-phenethyl, *N*-phenethan-1-one, and *N*-2-naphthylmethyl moieties into the 5,7-dibromoisatin molecule (Fig. 1. structures **B**, **C**, **D** and **E**, respectively) significantly increased hydrophobicity and increased cytotoxicity toward lymphoma cells. These isatin analogs were also potent against a wide range of human cancer cell lines; including MDA-MB-231 metastatic breast adenocarcinoma cells and U937 human monocyte-like histiocytic lymphoma cells.^{1–3} Hall et al.⁴ studied the MDR1-selective mechanism of action of several isatin-β-thiosemicarbazones (Fig. 1. structure **F**) against a parental HeLa-derived cervical cancer cell line (KB-3-1), while Perrow et al.⁵ have described the cytotoxicity of a number of isatin derivatives conjugated to a cell targeting moiety via a spacer group (Fig. 1. structure **G**). In view of the general anticancer properties of these isatin analogs, and as part of an

© 2010 Elsevier Ltd. All rights reserved.

*Corresponding author: Tel.: +1-859-257-1718; fax: +1-859-257-7585; pcrooks@email.uky.edu.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:....

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

investigation devoted to the development of new anticancer agents derived from structural modification of indoles,^{6–8} the design and synthesis of a novel series of 3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-one analogs (**3a-w**) was undertaken. The resulting analogs were then evaluated for their cytotoxic activity against a panel of 57 human tumor cell lines.

The simple and *N*-alkyl substituted isatins (**1a-j**) were all prepared utilizing literature methods.^{9–11} A series of novel substituted 3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-one derivatives (**3a-w**) were synthesized by condensation of the appropriate substituted *N*-alkyl isatin with creatinine, in the presence of sodium acetate and acetic acid via both microwave irradiation and conventional heating methodologies (Scheme 1). Of these two methods, microwave irradiation was found to be advantageous over conventional heating, since the product yields were 83–94% for the former method, but only 70–83% for the latter method. In addition, the time course of the reaction was very fast using microwave irradiation (20–40 sec) compared to 6–8 h for conventional heating (Table 1). All the synthesized compounds were characterized by ¹H-NMR and ¹³C-NMR spectrometry.¹⁶ The geometry of the hydroxyl position in the representative compounds **3a**, **3b** and **3t** was established as *trans* to the 4'-methyne hydrogen from X-ray crystallographic data.^{12–14}

From the X-ray diffraction and ¹H-NMR data, analogs **3a-3w** were mixtures of *RR* and *SS* isomers. This is consistent with the mechanism of the aldol condensation reaction of **1** with **2**, which proceeds via the formation of the *E*-enolate, as per the Zimmerman–Traxler model, which favors *anti* products, and is predicted to lead to the formation of equimolar *RR* and *SS* enantiomers. We also determined from the crystal structures of **3a**, **3b** and **3t** that the 3-hydroxy group was *trans* to the 4'-methyne hydrogen, which may explain the inability of these analogs to undergo facile dehydration.

The single dose *in vitro* cytotoxicity screening assays of the analogs were carried out in accordance with the procedures described in Rubinstein et al.¹⁵ The human tumor cell line panel includes leukemia, non-small cell lung, colon, CNS, melanoma, ovarian, renal, prostate, and breast cancer cell lines. From the preliminary results of the *in vitro* single dose screen, analogs containing an *N*-benzyl group in the isatin moiety exhibited good growth inhibition properties. The introduction of chloro group at C-5 of the isatin moiety also afforded analogs with good growth inhibition properties.

The two most active compounds (**3n** and **3o**) from the preliminary 57 cell screen were subsequently evaluated in five dose-response studies for their *in vitro* cytotoxic effects on growth parameters against each of the 57 human tumor cell lines. Dose-response curves were created by plotting cytotoxic effect against the log₁₀ of the drug concentration for each cell line. Cytotoxic effects of each compound were determined as GI₅₀, TGI and LC₅₀ values, which represent the molar drug concentration required to cause 50% growth inhibition, total growth inhibition, and the concentration that kills 50% of the cells, respectively. The results are presented in Table 2.

Compound **3o** exhibited growth inhibitory properties against 93% of all cancer cell lines in the panel, with GI₅₀ values in the range of 0.19–30 μM (Table 2). Compound **3o** exhibited good growth inhibitory activity against A549/ATCC non-small cell lung cancer cell lines (GI₅₀=193 nM; TGI=1.96 μM, LC₅₀=10.9 μM) and LOX IMVI melanoma cell lines (GI₅₀=750 nM; TGI= 11.8 μM; LC₅₀=60.0 μM), and showed moderate growth inhibitory activity against NCI-H522 lung cancer (GI₅₀= 1.86 μM), HT29 colon cancer (GI₅₀= 1.38 μM), UACC-257 melanoma (GI₅₀= 1.43 μM), and OVCAR-8 and NCI/ADR-RES ovarian cancer

(GI₅₀= 1.35 and 1.88 μM, respectively), and CAKI-1 renal cancer (GI₅₀= 1.98 μM) cell lines.

Compound **3n** exhibited growth inhibitory properties against 89% of all cancer cell lines in the panel, with GI₅₀ values in the range of 2–60 μM. (Table 2). Moderate growth inhibitory activity was observed against CCRF-CEM leukemia (GI₅₀=2.18 μM), HCT-116 colon (GI₅₀=2.40 μM), and OVCAR-3 and NCI/ADR-RES ovarian (GI₅₀=2.29 and 1.99 μM, respectively) cell lines.

Of particular significance was the observation that both **3o** and **3n** were effective in inhibiting the growth of all five cell lines in the sub-panel of leukemia cancer cells, with GI₅₀ values in the narrow range of 2–5 μM.

In conclusion, a series of 3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-one analogs have been synthesized and evaluated for anticancer activity against a panel of 57 human tumor cell lines. Compounds **3n** and **3o** were identified as molecules of interest from a single dose assay, and were then evaluated for dose-dependent growth inhibition and cytotoxicity in all 57 human cancer cell lines. Compound **3o** had GI₅₀ values of 190 nM and 750 nM against A549/ATTC non-small cell lung cancer and LOX IMVI melanoma cell lines, respectively, and both **3n** and **3o** exhibited GI₅₀ values ranging from 2–5 μM against CCRF-CEM, HL-60(TB), K-562, MOLT-4, and RPMI-8226 leukemia cell lines. These results indicate that *N*-4-methoxybenzyl-3-hydroxy-(2-imino-3-methyl-5-oxo-4-yl)indolin-2-one analogs may be useful leads for anticancer drug development.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the NCI/NIH for their financial support under grant number CA 140409 and to the NCI Developmental Therapeutic Program (DTP) for screening data.

References and notes

1. Vine KL, Locke JM, Ranson M, Benkendorff K, Pyne SG, Bremner JB. *Bioorg Med Chem.* 2007; 15:931. [PubMed: 17088067]
2. Vine KL, Locke JM, Ranson M, Pyne SG, Bremner JB. *J Med Chem.* 2007; 50:5109. [PubMed: 17887662]
3. Matesic L, Locke JM, Bremner JB, Pyne SG, Skropeta D, Ranson M, Vine KL. *Bioorg Med Chem.* 2008; 16:3118. [PubMed: 18182300]
4. Hall MD, Salam NK, Hellawell JL, Fales HM, Kensler CB, Ludwig JA, Szakacs G, Hibbs DE, Gottesman MM. *J Med Chem.* 2009; 52:3191. [PubMed: 19397322]
5. Perrow, KL.; Ranson, M.; Locke, M.; Bremner, JB.; Pyne, SG. WO 2008/074078 A1.
6. Sonar VN, Thirupathi Reddy Y, Sekhar KR, Sasi S, Freeman ML, Crooks PA. *Bioorg Med Chem Lett.* 2007; 17(24):6821. [PubMed: 17980582]
7. Narsimha Reddy P, Thirupathi Reddy Y, Crooks PA. *Bioorg Med Chem Lett.* 2010; 20(2):591. [PubMed: 20006496]
8. Thirupathi Reddy Y, Sekhar KR, Sasi N, Narsimha Reddy P, Freeman ML, Crooks PA. *Bioorg Med Chem Lett.* 2010; 20(2):600. [PubMed: 20005706]
9. Macpherson LJ, Dubin AE, Evans MJ, Marr F, Schultz PG, Cravatt BF, Patapoutian A. *Nature (London, United Kingdom).* 2007; 445(7127):541. [PubMed: 17237762]
10. Jacobs TL, Winstein S, Linden GB, Robson JH, Levy EF, Seymour D. *Organic Synthesis.* 1948; 28:70.

11. Shindikar AV, Khan F, Viswanathan CL. Euro J Med Chem. 2006; 41(6):786.
12. Narsimha Reddy P, Thirupathi Reddy Y, Parkin S, Crooks PA. Acta Crystallographica. 2009; E65(11):o2909.
13. Narsimha Reddy P, Thirupathi Reddy Y, Parkin S, Crooks PA. Acta Crystallographica. 2009; E65(10):o2439–o2440.
14. Narsimha Reddy P, Thirupathi Reddy Y, Parkin S, Crooks PA. Acta Crystallographica. 2009; E65(3):o552.
15. Rubinstein LV, Shoemaker RH, Paull KD, Simo RM, Tosini S, Skehan P, Scudiero PA, Monks A, Boyd MR. J Natl Cancer Inst. 1990; 82:1113.
16. Analytical data for compound **3n**. MF: C₂₀H₂₀N₄O₄, mp: 213–215 °C, ¹H NMR (DMSO-*d*₆): δ 3.14 (s, 3H, CH₃), 3.71 (s, 3H, OCH₃), 4.20 (s, 1H, CH), 4.65–4.84 (Abq, 2H, CH₂), 6.56 (s, 1H, OH), 6.65–6.68 (d, *J* = 8.1 Hz, 1H, C₇H), 6.85–6.94 (m, 4H, Ar-H), 7.10–7.16 (m, 2H, C₅H, C₆H), 7.33–7.40 (d, *J* = 8.4 Hz, 1H, C₄H), 7.57 (bs, 2H, NH₂) *ppm*; ¹³C NMR (DMSO-*d*₆): δ 32.64 (CH₃), 42.30 (CH₂), 54.97 (OCH₃), 69.46 (CH), 75.99 (C-OH), 109.05, 111.00, 113.62, 121.79, 123.69, 127.43, 127.81, 128.55, 129.30, 137.74, 143.15, 158.20, 171.98 (C=N), 174.31 (isatin C=O), 182.29 (creatinine C=O) *ppm*. Analytical data for compound **3o**. MF: C₂₀H₁₉ClN₄O₄, mp: 188–190 °C, ¹H NMR (DMSO-*d*₆): δ 3.18 (s, 3H, CH₃), 3.72 (s, 3H, OCH₃), 4.21 (s, 1H, CH), 4.66–4.85 (Abq, 2H, CH₂), 6.67 (s, 1H, OH), 6.70–6.72 (d, *J* = 5.1 Hz, 1H, C₇H), 6.85–6.88 (d, *J* = 8.7 Hz, 2H, Ar-H), 7.06–7.07 (d, *J* = 2.4 Hz, 1H, C₄H), 7.24–7.27 (dd, *J* = 8.1 Hz, 2.4 Hz, 1H, C₆H), 7.37–7.40 (d, *J* = 8.4 Hz, 2H, Ar-H), 7.57 (bs, 1H, NH), 7.82 (bs, 1H, NH) *ppm*; ¹³C NMR (DMSO-*d*₆): δ 32.94 (CH₃), 42.42 (CH₂), 55.03 (OCH₃), 69.62 (CH), 76.05 (C-OH), 110.63, 113.72 (2C) 123.73, 125.88, 127.43, 128.62 (2C), 129.16, 129.45, 142.10, 158.31 (C-Cl), 172.21 (C=N), 174.00 (isatin C=O), 182.08 (creatinine C=O) *ppm*.





Fig. 1.
Structures of potent cytotoxic isatin derivatives

**Scheme 1.**

Reagents and conditions: (a) Method A: sodium acetate in acetic acid, microwave irradiation, 20–40 seconds, 83–94% yield; (b) Method-B: sodium acetate in acetic acid, 115–120 °C, 6–8 hours, 70–83% yield.

Table 1

Reaction times and yields of 3-hydroxy-3-(2-imino-3-methyl-5-oxoimidazolidin-4-yl)indolin-2-ones (**3a-w**)

Compd	R ¹	R ²	R ³	Method-A		Method-B	
				Yield (%)	Time (sec)	Yield (%)	Time (hr)
3a	H	H	H	92	20	80	6
3b	F	H	H	94	30	81	6
3c	Cl	H	H	91	40	78	6
3d	Br	H	H	87	40	75	7
3e	Br	Br	H	89	40	79	8
3f	NO ₂	H	H	92	40	83	8
3g	H	H	-CH ₃	88	30	78	7
3h	F	H	-CH ₃	90	40	76	7
3i	Cl	H	-CH ₃	86	40	77	8
3j	Br	H	-CH ₃	85	40	80	8
3k	H	H	-Bz	87	30	72	7
3l	Cl	H	-Bz	91	40	76	7
3m	Br	H	-Bz	88	40	75	8
3n	H	H	4-OCH ₃ Bz	84	30	70	6
3o	Cl	H	4-OCH ₃ Bz	85	30	74	8
3p	Br	H	4-OCH ₃ Bz	83	40	79	8
3q	H	H	4-Cl Bz	92	40	77	7
3r	H	H	4-COOCH ₃ Bz	87	20	75	6
3s	H	H	4-CN Bz	86	40	79	8
3t	H	H	-C ₆ H ₅	83	30	80	7
3u	H	H	-COCH ₃	87	30	73	7
3v	Cl	H	-COCH ₃	85	40	75	8
3w	H	H	-SO ₂ -C ₆ H ₅	89	30	78	6

Table 2

Antitumor activity ($GI_{50}/\mu M$)^a and toxicity ($LC_{50}/\mu M$)^b data of compounds selected for 5 dose studies for the NCI 57-cell lines screen

Panel/cell line	Compound 3o		Compound 3n	
	GI_{50}	LC_{50}	GI_{50}	LC_{50}
Leukemia Cancer				
CCRF-CEM	2.06	>100	2.18	>100
HL-60(TB)	3.22	>100	4.91	>100
K-562	5.04	>100	3.73	>100
MOLT-4	2.94	>100	4.79	>100
RPMI-8226	2.62	>100	2.87	>100
Non-Small Cell Lung Cancer				
A549/ATCC	0.19	10.9	43.6	>100
EKVX	28.5	>100	71.7	>100
HOP-62	5.12	>100	4.31	>100
NCI-H226	26.8	>100	38.4	>100
NCI-H23	4.74	>100	13.9	>100
NCI-H322M	14.3	>100	41.1	>100
NCI-H460	26.8	>100	29.0	>100
NCI-H522	1.86	20.5	3.65	>100
Colon Cancer				
COLO 205	14.3	>100	19.9	>100
HCC-2998	18.7	73.9	12.8	63.9
HCT-116	3.00	34.5	2.40	>100
HCT-15	12.6	>100	11.8	>100
HT29	1.38	44.1	4.06	>100
KM12	2.05	30.0	2.87	80.4
SW-620	2.59	49.6	4.12	>100
CNS Cancer				
SF-268	3.31	64.9	5.00	>100
SF-295	3.71	>100	26.0	>100
SF-539	3.12	79.1	3.73	>100
SNB-19	17.2	>100	30.6	>100
SNB-75	3.39	>100	28.8	>100
Melanoma Cancer				
LOX IMVI	0.75	60.0	8.09	>100
MALME-3M	29.6	>100	27.4	>100
M14	3.59	61.4	4.93	>100
MDA-MB-435	6.12	93.1	10.6	>100
SK-MEL-2	10.4	96.0	95.2	>100
SK-MEL-28	50.3	>100	36.9	>100
SK-MEL-5	22.6	>100	37.3	>100

Panel/cell line	Compound 3o		Compound 3n	
	GI ₅₀	LC ₅₀	GI ₅₀	LC ₅₀
UACC-257	1.43	19.4	6.28	>100
UACC62	23.4	>100	27.7	>100
Ovarian Cancer				
IGR-OV1	2.59	>100	3.26	>100
OVCAR-3	2.08	11.5	2.29	30.1
OVCAR-4	5.52	>100	7.13	>100
OVCAR-5	34.9	>100	42.1	>100
OVCAR-8	1.35	17.2	3.52	>100
NCI/ADR-RES	1.88	36.5	1.99	>100
SK-OV-3	30.1	>100	45.5	>100
Renal Cancer				
786-0	3.44	43.0	23.0	>100
A498	34.5	>100	>100	>100
ACHN	20.4	>100	>100	>100
CAKI-1	1.98	>100	53.9	>100
RXF 393	6.29	>100	7.40	>100
SN12C	5.94	>100	>100	>100
TK-10	4.11	>100	>100	>100
UO-31	4.15	>100	83.2	>100
Prostate Cancer				
PC-3	4.27	>100	46.9	>100
DU-145	3.76	74.6	21.4	>100
Breast Cancer				
MCF7	3.22	>100	6.93	>100
MDA-MB-231/ATCC	8.55	>100	19.4	>100
HS 578T	2.04	>100	60.3	>100
BT-549	2.98	54.9	8.45	97.0
T-47D	13.5	>100	32.0	>100
MDA-MB-468	2.29	>100	11.6	>100

NA: Not analyzed

^aGI₅₀: 50% Growth inhibition, concentration of drug resulting in a 50% reduction in net protein increase compared with control cells.

^bLC₅₀: Lethal concentration, concentration of drug lethal to 50% of cells.