

Thrombin Generation and Secretion of Platelet Factor 4 during Blood Clotting

MARC A. SHUMAN, *Division of Hematology-Oncology, Department of Medicine and the Cancer Research Institute, University of California, San Francisco, California 94143*

SHIRLEY P. LEVINE, *Division of Hematology and the Department of Medicine, Veterans Administration Hospital, University of Texas, San Antonio, Texas 78284*

ABSTRACT We have studied the platelet release reaction and thrombin generation during the spontaneous clotting of whole blood *in vitro*. Both thrombin formation and secretion of platelet Factor 4 were detected at least 12 min before clotting (clotting time, 22–26 min). Initially, at low thrombin concentrations (2–5 ng/ml), there is a small increase in plasma platelet Factor 4 (<1% of the amount present in serum). This is followed by a gradual increase in both platelet Factor 4 and thrombin concentrations over a 12 to 20-min interval. Finally, 5 min before clotting, there is a rapid increase in both thrombin generation and platelet secretion.

Thus, we have shown that the release of platelet Factor 4 is a prolonged reaction and the extent to which it occurs parallels thrombin generation. It is only when thrombin concentrations are high (45–90 ng/ml)—during the period of clot formation—that the major part of platelet Factor 4 secretion occurs. Release of platelet Factor 4, like fibrin formation, occurs in the last step of *in vitro* coagulation.

INTRODUCTION

Platelets have been assigned a dual role in hemostasis. Initially, after vascular injury, a platelet aggregate or plug is formed on the subendothelial surface. During plug formation, platelets undergo changes that result in the exposure of a phospholipid surface on which Factors X and II are converted to their active forms (1). Several recent observations indicate that the interaction of platelets with clotting factors during thrombin formation is more complex than initially postulated. Osterud et al. (2) have shown that platelets contain an activated form of Factor V and have postulated

that its release may be important in prothrombin activation. Experiments by Miletich et al. (3) demonstrate that the rate of thrombin generation is 10 to 100-fold faster when platelets are used in place of phospholipid and Factor V in the activation of prothrombin by Factor Xa.

Once thrombin is formed, it stimulates platelet secretion and aggregation (4). Some of the initial steps in this interaction have been characterized in experiments using purified thrombin and washed platelets. Thrombin binds specifically and reversibly to high affinity receptors on the platelet surface (5–8). There is good evidence to suggest that binding must occur before secretion can take place (8, 9). Although secretion does not occur if the active site of thrombin is blocked (4), the lack of thrombin turnover and the requirement for a threshold concentration of thrombin before release can occur (10) suggest that the thrombin-induced release is not a simple enzymatic process.

Our experiments were designed to study the interaction of platelets with thrombin in whole blood. Because we had previously demonstrated that thrombin is present early in clotting (11), we were interested in seeing whether or not there was evidence for an effect of thrombin on platelets at a similar time period. We found that the time-course of secretion of a platelet protein, platelet Factor 4 (PF₄)¹ parallels closely the generation of thrombin during *in vitro* blood clotting.

METHODS

Chemicals and radioisotopes were purchased from commercial sources as defined (11–14). Prostaglandin E₁ (PGE₁)

¹Received for publication 19 December 1977 and in revised form 23 January 1978.

¹Abbreviations used in this paper: PF₄, platelet Factor 4; dbcAMP, N⁶O²-dibutyryl adenosine 3',5' monophosphate cyclic AMP; PGE₁, prostaglandin E₁.

was generously provided by Dr. John Pike, Upjohn Co., Kalamazoo, Mich. Cohn fraction III paste and plasminogen-free fibrinogen used in these studies were provided by the American Red Cross National Fractionation Center with the partial support of the National Institutes of Health grant HL 13881. Outdated platelets for PF₄ purification were obtained from the American Red Cross, Tulsa, Okla. and Wichita, Kans., and the Wadley Institutes of Molecular Medicine, Dallas, Tex.

PF₄ secretion and thrombin generation in clotting blood. Venipuncture was performed on normal volunteers using 16 gauge needles. The first 5–10 ml was discarded, then 40 ml was collected rapidly into new polypropylene tubes and allowed to clot at room temperature (20°–24°C). Samples were removed at regular intervals, anticoagulated with 0.1 vol. of 3.8% trisodium citrate and assayed for either PF₄ or thrombin. To blood samples that were assayed for PF₄, 100 mM N⁶O²-dibutyl adenosine 3',5' monophosphate cyclic AMP (dbcAMP) (final concentration 1 mM) and 1 mg/ml PGE₁ in ethanol (final concentration 0.0028 mM) were added immediately. Aliquots assayed for thrombin were treated as described (11). All samples were kept on ice before centrifugation at 10,000 g for 10 min at 4°C. Plasma to be assayed for PF₄ was stored at –20°C, then shipped to San Antonio packed in dry ice. It had previously been determined that there was no difference in PF₄ concentration when measured in either fresh or frozen-thawed plasma.

To determine to what extent platelets might secrete PF₄, apart from that induced by clotting, 80 ml of blood was drawn and divided into two aliquots. One was immediately anticoagulated with either trisodium citrate (final concentration 0.38%), or 0.25 M EDTA (final concentration, 0.0058 M). The other was allowed to clot. Samples were removed at regular intervals, processed as described above, and assayed for PF₄ and thrombin.

Effect of dbcAMP and PGE₁ on thrombin-induced PF₄ secretion. Blood was drawn from normal volunteers using a 19-gauge needle. Samples were then immediately anticoagulated with 3.8% trisodium citrate (final concentration 0.38%), and divided into two equal portions. To one fraction, dbcAMP and PGE₁ were added in the concentrations as described for the clotting experiments. Samples of 2.0 ml were removed and thrombin, final concentrations of 0.4–5.0 U/ml was added. The other portion was treated identically except that neither dbcAMP nor PGE₁ were added. The samples were incubated at room temperature for 10 min, then centrifuged at 10,000 g for 10 min at 4°C, and the plasma removed for assay.

Thrombin radioimmunoassay. The techniques for human thrombin purification (12) and radioimmunoassay (11) have been published previously. The thrombin was assayed as described (13) and had a 3,200 U/mg sp act and was homogeneous based on sodium dodecyl sulfate polyacrylamide gel electrophoresis.

PF₄ radioimmunoassay. The methods for PF₄ purification (14) and radioimmunoassay (15) have been described. The interassay control was 7.3±1.8 ng. There was a 10–15% coefficient of variation when the same sample was run in the same assay.

RESULTS

Secretion of PF₄ in clotting blood. When freshly drawn blood was allowed to clot spontaneously in polypropylene tubes, the following observations were made: First, PF₄ secretion was initiated as early as 3 min after venipuncture and as long as 25 min before

visible clotting (Fig. 1A). The precise time of onset was variable among normal subjects. In some experiments, PF₄ secretion was not detected for as long as 10 min after venipuncture. In all instances there was at least a 12-min interval between initiation of PF₄ secretion and the time of clotting (clotting times in these experiments ranged from 22–26 min). The base-line PF₄ values tended to be higher than other published series, however, much larger volumes of blood were drawn and there was a longer interval before anticoagulant and platelet inhibitors were added. Second, once initiated, the secretory process occurred over a prolonged period (15–20 min) rather than as a rapid burst. Third, PF₄ secretion appeared to occur in at least two stages. Initially small amounts of secretion proceeded slowly for several minutes during which time there was at least one lag period with little or no increase in PF₄ concentration. This was of variable duration, lasting as long as 8 min. Then, in the second stage, there was a secretory burst over 4–6 min with at least a 20-fold increase in PF₄ concentration. This occurred just before the time of clotting.

Correlation of thrombin generation with PF₄ secretion. When prothrombin activation, as measured by thrombin radioimmunoassay (11), was compared with PF₄ secretion, certain similarities were seen (Fig. 1B). As described (11), small amounts of thrombin (2–5 ng/ml) appeared early after venipuncture (Fig. 1B). In most experiments, thrombin generation either preceded or occurred at the time of the first increase in PF₄. In some experiments, the initial secretion of PF₄ was observed before an increment in thrombin concentration could be detected. After the initial appearance of thrombin, a lag period in further prothrombin activation occurred (Fig. 1B) similar to that which took place in PF₄ secretion. A slow increase in thrombin concentration followed, again similar to that observed in PF₄ secretion. In all cases, if a delay in thrombin formation was observed, a delay was also seen in PF₄ secretion. Shortly before clotting, a marked increase in both thrombin and PF₄ concentrations was noted (Fig. 1A and B). At the time of clotting, when the thrombin concentration was 70–100 ng/ml, PF₄ in serum was 200 to 1,000-fold higher than base-line values.

To determine the extent to which PF₄ might be secreted from platelets apart from that induced by thrombin, PF₄ and thrombin levels were determined on blood samples which were obtained simultaneously and either anticoagulated immediately or allowed to clot. When the blood was anticoagulated immediately, the plasma PF₄ concentration did not rise until 10–14 min after venipuncture, irrespective of the anticoagulant (trisodium citrate or sodium EDTA, two experiments each). The PF₄ concentration then gradually increased with incubation over the next




FIGURE 1 PF₄ secretion and thrombin generation in clotting blood. Blood was collected with a 16-gauge needle and allowed to clot at room temperature in a polypropylene tube. Serial aliquots were removed and plasma assayed for (A) PF₄ (closed symbols), or (B) thrombin (open symbols). The results are based on blood obtained from three normal individuals. The arrows represent the time at which clotting occurred.

10–15 min with final levels of 260–360 ng/ml (<5% of the PF₄ present in serum). One possible explanation for this increase in PF₄ concentration is that incubation in anticoagulants is injurious to the platelets and results in secretion. Although we could detect no thrombin generation, a small amount may have occurred and induced secretion.

Net PF₄ secretion was determined by subtracting the PF₄ values for each time point measured in anticoagulated blood from values observed at the same time points in simultaneously obtained clotting blood. As shown in Fig. 2, ≈2 min after the first detectable thrombin generation and 10 min after venipuncture, there is a progressive rise in PF₄ concentrations. The initial PF₄ concentrations in these experiments—both clotting and anticoagulated specimens—were higher (39–125 ng/ml) than our normal base-line values. This appears to be related to the longer time necessary to collect the large volume of blood used in these experiments.

Inhibition of thrombin-induced secretion by dbcAMP and PGE₁. To study PF₄ secretion during clotting, it was necessary to demonstrate that platelet secretion subsequent to collection of samples at each

time point could be inhibited. Otherwise, the PF₄ concentration would represent additional secretion that might occur as the samples were prepared for radioimmunoassay. Although a small amount of secretion was seen when thrombin at concentrations of 0.8 U/ml (250 ng/ml) or less was added to whole blood containing dbcAMP and PGE₁, this represented <5% of the PF₄ secreted when these inhibitors were not present (Table I). At higher thrombin concentrations, i.e. 1.6 U/ml, the PF₄ concentration rose to 967 ng/ml. Still, this represents only 13% of the PF₄ secreted without inhibitors. At 3.2 U/ml and higher, inhibition is less effective with 25–50% of normal secretion.

DISCUSSION

We have studied platelet secretion in clotting whole blood that is unperturbed by anticoagulants or centrifugation before stimulation. Previously, we demonstrated that small amounts of thrombin are formed early in clotting, before a visible clot appears (11). We hypothesized that this thrombin could be a stimulus for the platelet release reaction. Our present observa-




FIGURE 2 PF₄ secretion and thrombin generation in anticoagulated and clotting blood. Blood was collected with a 16-gauge needle and distributed equally into two polypropylene tubes. One was anticoagulated immediately with 0.1 volume of 3.8% trisodium citrate and the other allowed to clot. (Clotting time 23 min.) Serial samples were removed and assayed for PF₄ or thrombin. Net PF₄ secretion (●) calculated by subtracting PF₄ values in the anticoagulated blood from values obtained at corresponding times in clotting blood; thrombin (○) in clotting blood.

tion that secretion of PF₄ also begins at least 12 min before clotting and is blocked in anticoagulated blood supports this hypothesis. Over 90% of the release of PF₄, however, occurs during the last 4–5 min

TABLE I
Effect of dbcAMP and PGE₁ on Thrombin-Induced Platelet Secretion

Thrombin final concentration		PF ₄	
U/ml	μg/ml	A	B
0		35	61
0.4	0.125	124	3,260
0.8	0.25	197	4,170
1.6	0.5	967	7,533
3.2	1.0	3,870	8,150
5	1.56	1,818	6,917

Blood was collected into 0.1 vol of 3.8% trisodium citrate and divided into two portions, "A" and "B." To fraction A, dbcAMP (final concentration 1 mM) and PGE₁ (final concentration 0.0028 mM) were added. Then, thrombin was added to 2.0-ml aliquots from each fraction and the samples placed on ice. After 10 min, the samples were centrifuged and the plasma removed for PF₄ assay.

of clotting. Platelet secretion, on the other hand, does not appear necessary for the initial appearance of thrombin. Low levels of PF₄ were always present at the earliest time points. Whether this is due to platelet activation during blood collection and handling has not been determined. There are several possible explanations for the observation that in some experiments PF₄ secretion occurred before thrombin generation could be detected. The radioimmunoassay may not have detected a small change in thrombin concentration which may have occurred at this earliest stage of blood clotting or thrombin bound to platelets may not have been detected in the radioimmunoassay. Alternatively, there may have been a stimulus for this initial PF₄ secretion other than thrombin. A third possibility is that centrifugation of blood samples alone resulted in the release of small amounts of PF₄.

The long duration of secretion can be explained by the previous observation that the extent of platelet secretion is dependent on the thrombin concentration (10). One possible explanation for the time-course observed, therefore, is that early on, the concentration of thrombin is just high enough to induce partial secretion, but only much later are concentrations sufficiently high to induce complete secretion. Alternatively, there may be different populations of platelets with different sensitivities to thrombin: a small population which is very sensitive and secretes at low thrombin concentrations, and larger populations that secrete later and at higher thrombin concentrations. Our data do not allow us to distinguish between these possibilities at this time. Nor do we know the effects of the secretion of adenosine diphosphate or the prostaglandin metabolites on this process. It may be that once a critical extracellular concentration of either of these substances is reached, secretion is greatly accelerated.

It has been reported that when thrombin is added to citrated platelet-rich plasma or whole blood, the concentration necessary to induce platelet secretion is 100-fold higher than that required to cleave fibrinopeptide A from fibrinogen (16). This observation may be misleading in that it suggests secretion might occur much later than fibrin formation or clotting. These experiments, however, may not correspond to what happens physiologically. When platelet secretion is stimulated by endogenous thrombin, the concentration on the platelet surface is probably much higher than that of plasma because that is where prothrombin activation presumably occurs (3). Furthermore, there is evidence to suggest that activated clotting factors may be protected from inhibition by antithrombin III when bound to a phospholipid surface (17, 18).

We do not know whether the secretory pattern that we have observed for PF₄ under these conditions

is different from that of other platelet intracellular constituents. It has been suggested that antiheparin activity is located in the alpha granules of platelets (19, 20). Holmsen et al. (21) have reported that higher concentrations of thrombin are needed for alpha granule release than for dense body secretion. Also, it has been reported that release of antiheparin activity is slower than that of serotonin when washed platelets are stimulated to secrete (22). If this is the case, one would expect secretion to occur earlier or be completed more rapidly than we observed for PF₄. However, the differences observed in these experiments may be related to the types of assays used to detect release rather than to different secretory time courses. Thus, a radioassay was used to measure serotonin release and functional assays to measure alpha granule release (22). It may be that the latter are not as sensitive as the former and do not detect low levels present in the early stages of secretion.

The precise role of platelet secretion in prothrombin activation is unclear. Miletich et al. (3) have shown that platelets have specific receptors for Factor Xa which appear on their surface only after secretion takes place. The receptor has characteristics similar to Factor V. The rapid formation of thrombin shortly before clotting, therefore, may be the result of increased binding of Factor Xa to Factor V secreted from platelets at this time.

We have developed a method for studying the interaction of platelets and coagulation factors which will allow further characterization of this process in normal and pathologic blood clotting. In the latter case, this may improve our understanding of the specific consequences of clotting factor deficiencies or platelet defects on coagulation.

ACKNOWLEDGMENTS

We wish to thank Linda S. Krentz for her expert technical assistance.

This work was supported by grants from the National Institutes of Health (HL 21403) and the Medical Research Service, Veterans Administration.

REFERENCES

- Marcus, A. J. 1972. Recent advances in platelet lipid metabolism research. *Ann. N. Y. Acad. Sci.* **201**: 102-108.
- Osterud, B., S. I. Rapaport, and K. Lavine. 1977. Factor V activity of platelets: evidence for an activated factor V molecule and for a platelet activator. *Blood.* **49**: 819-834.
- Miletich, J. P., C. M. Jackson, and P. W. Majerus. 1977. The interaction of coagulation factor Xa with human platelets. *Proc. Natl. Acad. Sci.* **74**: 4033-4036.
- Davey, M. G., and E. F. Luscher. 1967. Actions of thrombin and other coagulant and proteolytic enzymes on blood platelets. *Nature (Lond.)* **216**: 857-858.
- Tollefsen, D. M., J. R. Feagler, and P. W. Majerus. 1974. The binding of thrombin to the surface of human platelets. *J. Biol. Chem.* **249**: 2646-2651.
- Ganguly, P. 1974. Binding of thrombin to human platelets. *Nature (Lond.)* **247**: 306-307.
- Shuman, M. A., D. M. Tollefsen, and P. W. Majerus. 1976. The binding of human and bovine thrombin to platelets. *Blood.* **47**: 43-54.
- Martin, B. M., W. W. Wasiewski, J. W. Fenton II, and T. C. Detwiler. 1976. Equilibrium binding of thrombin to platelets. *Biochemistry.* **15**: 4886-4893.
- Shuman, M. A., and P. W. Majerus. 1975. The perturbation of thrombin binding to human platelets by anions. *J. Clin. Invest.* **56**: 945-950.
- Detwiler, T. C., and R. D. Feinman. 1973. Kinetics of the thrombin-induced release of calcium (II) by platelets. *Biochemistry.* **12**: 282-289.
- Shuman, M. A., and P. W. Majerus. 1976. The measurement of thrombin in clotting blood by radioimmunoassay. *J. Clin. Invest.* **58**: 1249-1258.
- Fenton, J. W. II, M. J. Fasco, A. B. Stackrow, D. L. Aronson, A. M. Young, and J. S. Finlayson. 1977. Human thrombins. Production, evaluation, and properties of α -thrombin. *J. Biol. Chem.* **252**: 3587-3598.
- Baughman, D. J. 1970. Thrombin assay. *Methods Enzymol.* **19**: 145-157.
- Levine, S. P., and H. Wohl. 1976. Human platelet factor 4: purification and characterization by affinity chromatography. *J. Biol. Chem.* **251**: 324-328.
- Levine, S. P. and L. S. Krentz. 1977. Development of a radioimmunoassay for human platelet factor 4. *Thromb. Res.* **11**: 673-686.
- Kaplan, K. L., and H. L. Nossel. 1977. Simultaneous measurement of platelet factor 4 and β -thromboglobulin release and fibrinopeptide A cleavage. *Clin. Res.* **25**: 341A. (Abstr.)
- Marciniak, E. 1973. Factor-Xa inactivation by antithrombin III. Evidence for biologic stabilization of factor Xa by factor V-phospholipid complex. *Br. J. Haematol.* **24**: 391-400.
- Yin, E. T. 1974. Effect of heparin on the neutralization of factor Xa and thrombin by the plasma alpha-2-globulin inhibitor. *Thromb. Diath. Haemorrh.* **33**: 43-50.
- DaPrada, M., M. Jakabova, E. F. Luscher, A. Pletscher, and J. G. Richards. 1976. Subcellular localization of the heparin-neutralizing factor in blood platelets. *J. Physiol.* **257**: 495-502.
- Broekman, M. J., R. I. Handin, and P. Cohen. 1975. Distribution of fibrinogen, and platelet factor 4 and XIII in subcellular fractions of human platelets. *Br. J. Haematol.* **31**: 51-55.
- Holmsen, H., C. A. Setkowsky, B. Lages, H. J. Day, H. J. Weiss, and M. C. Scrutton. 1975. Content and thrombin-induced release of acid hydrolases in gel-filtered platelets from patients with storage pool disease. *Blood.* **46**: 131-142.
- Walsh, P. N., and G. Gagnatelli. 1974. Platelet antiheparin activity: storage site and release mechanism. *Blood.* **44**: 157-168.