

Published in final edited form as:

Supramol Chem. 2013 June 1; 25(6): 315–322. doi:10.1080/10610278.2013.776170.

Effect of Bridging Anions on the Structure and Stability of Phenoxide Bridged Zinc Dipicolylamine Coordination Complexes

Edward J. O'Neil^a, Hua Jiang^{a,b}, and Bradley D. Smith^{a,*}

^aDepartment of Chemistry and Biochemistry, University of Notre Dame, 2365 Nieuwland Science Hall, Notre Dame, IN 46556

^bBeijing National Laboratory for Molecular Sciences, CAS Key Laboratory of Photochemistry, Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100190, China

Abstract

A series of four related phenol derivatives, with 2,2'-dipicolylamine substituents at the ortho positions, were prepared and their Zn²⁺ coordination complexes studied by spectroscopic methods. X-ray crystal diffraction analysis of a dinuclear zinc complex with two bridging acetate anions showed a ternary structure with highly charged interior and lipophilic exterior, which helps explain why this class of water-soluble complexes can effectively diffuse through cell membranes. The stability of the dinuclear zinc complexes in aqueous solution was found to be strongly anion dependent; that is, bridging oxyanions, such as acetate and pyrophosphate, lock the two Zn²⁺ cations to the surrounding ligand and greatly enhance ligand/zinc affinity. Overall, the results provide new insight into the structural and mechanistic factors that control the recognition and chemosensing performance of phenoxide bridged dipicolylamine molecular probes.

Keywords

Anion recognition; zinc coordination; nuclear magnetic resonance; X-ray crystallography; titration; spectroscopic analysis

Introduction

Phenol derivatives with 2,2'-dipicolylamine (DPA) substituents at the ortho positions are known to readily form coordination complexes with Zn²⁺, Cu²⁺, Ni²⁺, Co³⁺, Fe²⁺, and Fe³⁺ cations, and quite a few X-ray crystal structures of dinuclear complexes have been described.¹ This report focuses on the Zn²⁺ complexes with generic molecular structure, **BDPA•Zn₂** (Scheme 1). These compounds are part of a larger family of Zn²⁺-DPA complexes that have been developed for various supramolecular applications such as sensing of anionic biomolecules in aqueous solution, protein labeling, DNA targeting, RNA hydrolysis, biomembrane targeting, and cell penetration.^{2,3,4} The long term goal of our research is to invent new classes of molecular probes for biological imaging, and we have discovered that Zn²⁺-DPA complexes have a remarkable ability to recognize anionic cell

*smith.115@nd.edu, ph: +1 (574) 6318632, fax: +1 (574) 6318632.

Supporting Information Available NMR assignments and CIF file for X-ray structure can be obtained via <http://www.tandfonline.com>. The crystallographic data is also provided in CCDC 901795, which can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB21EZ, UK; fax (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).

membrane surfaces selectively in complex biological environments such as cell culture and living animals.⁵ During these studies we observed an unusual phase transfer feature with water-soluble **BDPA•Zn₂** complexes, namely, an ability to penetrate cell membranes.⁶ We exploited this finding by developing oligomers of **MDPA•Zn** as effective cell permeating peptides.⁷ However, non-selective cell penetration is not a desirable attribute for most bioimaging applications. In order to rationally improve the performance of **BDPA•Zn₂** complexes as bioimaging probes and chemical sensors, a better understanding is needed of the fundamental coordination chemistry that controls the molecular recognition.

Here, we report a study of the solid-state structure and solution dynamics of some **MDPA•Zn** and **BDPA•Zn₂** complexes using a combination of spectrometric methods. Specifically, we have examined the Zn²⁺ complexation properties of four ligands, **L1–L4** (Scheme 1). We find that Zn²⁺ affinity of these ligands in water is strongly anion dependent, that is, Zn²⁺ affinity is greatly enhanced by the presence of bridging oxyanions, such as acetate (OAc⁻) and pyrophosphate (PPi⁴⁻). Furthermore, oxyanion bridging of the two Zn²⁺ cations in **BDPA•Zn₂** produces a lipophilic ternary complex, which helps explain why water-soluble **BDPA•Zn₂** complexes can effectively diffuse through cell membranes. The article concludes with a discussion of how this study can be used to improve the performance of DPA coordination complexes as molecular imaging agents and optical chemosensors for Zn²⁺ and oxyanions.

Results and Discussion

Synthesis and Solid-State Structures

The ligands, **L1–L4**, were prepared in straightforward fashion using established procedures that produced a Mannich reaction of the parent phenol with one or two molar equivalents of paraformaldehyde and DPA.^{3c} An X-ray crystal structure was obtained for the zinc acetate complex of **L1**. Single crystals were obtained by recrystallizing a methanol solution containing **L1** mixed with two molar equivalents each of Zn(OAc)₂, triethylamine, and NaBF₄.^{1a} X-ray diffraction analysis revealed a molecular formula of Zn₂**L1**(OAc)₂(BF₄)•2MeOH. The C₂ symmetric structure (Figure 1) contains two, six-coordinate Zn²⁺ centers with distorted octahedral geometries and an N₃O₃ donor set. Each Zn²⁺ is coordinated by a DPA tertiary amine and two pyridyl nitrogen atoms, as well as the central phenoxy anion and two bridging OAc⁻ anions. The average Zn–N distances are 2.23 Å for the Zn–tertiary amine interactions and 2.17 Å for the Zn–pyridyl nitrogen interactions. The average Zn–O distances are 2.04 Å for the Zn–phenoxy bonds and 2.05 Å for the Zn–acetoxy bonds. The *trans*-axial angles through the distorted octahedral Zn atoms are all in the range of 162–171°, and the Zn1···Zn2 distance is 3.36 Å. These bond lengths and angles are consistent with similar Zn²⁺-DPA structures found in the literature.¹ The molecular structure has a highly charged interior and lipophilic exterior, and suggests a structural process that enables partitioning of water-soluble **BDPA•Zn₂** complexes into vesicle and cell membranes.⁷ The process involves reversible association of **BDPA•Zn₂** complexes with one or two fatty acids or phospholipids at the membrane surface to produce membrane-soluble coordination complexes that are analogous to the structure in Figure 1.

Solution-State UV Titrations

The two phenolic ligands, **L3** and **L4**, have appended naphthalene chromophores and they are known to exhibit ratiometric spectral changes when they form zinc complexes.^{3c} Thus, **L3** and **L4** were chosen as sensing ligands for a study of the effects of bridging oxyanions on relative zinc complex stoichiometry and stability. Shown in Figure 2 are the results of an absorbance titration experiment that added Zn(NO₃)₂ to control mono-DPA ligand **L4** in methanol:water (4:1). There is a large bathochromic shift in absorption maxima as the ligand

forms the expected $\mathbf{L4:Zn}^{2+}$ salt structure (see $\mathbf{MDPA}\cdot\mathbf{Zn}$ in Scheme 1) with 1:1 stoichiometry. The next set of absorbance titration experiments added $\text{Zn}(\text{NO}_3)_2$ to bis-DPA ligand $\mathbf{L3}$ in methanol:water (4:1) and yielded intriguing results. As shown in Figure 3 the stoichiometry at signal saturation depended on the identity of the counter anion. Addition of $\text{Zn}(\text{NO}_3)_2$ to a solution of $\mathbf{L3}$ reached saturation when the $\mathbf{L3:Zn}^{2+}$ stoichiometry was 1:1 (Figure 3a). In contrast, the saturated $\mathbf{L3:Zn}^{2+}$ stoichiometry was clearly 1:2 when the titration was conducted in the presence of Na_4PPi (Figure 3b) or NaOAc (Figure 3c). The data is rationalized by considering the two stepwise, zinc association equilibria in Scheme 2. When the free $\mathbf{L3}$ ligand (represented as generic ligand \mathbf{BDPA} in Scheme 2) is titrated with $\text{Zn}(\text{NO}_3)_2$, the first apparent Zn^{2+} association constant, K_1 , is larger than the second stepwise constant, K_2 .⁸ Thus, the absorbance titration isotherm in Figure 3a reflects the first Zn^{2+} association step in Scheme 2; that is, the conversion of free \mathbf{BDPA} ligand into mononuclear zinc complex $\mathbf{BDPA}\cdot\mathbf{Zn}$.⁹ When high-affinity, bridging oxyanions, such as PPi^{4-} or OAc^- , are present in the solution, the second Zn^{2+} association constant, K_2 , is greatly enhanced such that it is larger than K_1 (Scheme 2). Thus, the absorbance changes in Figure 3b and 3c reflect the conversion of generic ligand \mathbf{BDPA} directly into dinuclear zinc complex $\mathbf{BDPA}\cdot\mathbf{Zn}_2$ with no measurable accumulation of the intermediate mononuclear zinc complex $\mathbf{BDPA}\cdot\mathbf{Zn}$.

A meaningful quantitative analysis of the above absorption titration curves was not possible, primarily for two reasons — the aqueous methanol solvent was not buffered (K_1 is pH dependent) and the ligand/ Zn^{2+} association was too strong for the absorption titration method. However, semi-quantitative confirmation of the ligand/ Zn^{2+} stabilization provided by the bridging oxyanions was gained by conducting competition experiments using ethylenediaminetetraacetic acid (EDTA) as a competitive Zn^{2+} binder. The top of Figure 4 shows the change in absorption spectra when a solution of $\mathbf{L3:2}\cdot\mathbf{Zn}(\text{NO}_3)_2$ in water/methanol (4:1) was titrated with two molar equivalents of EDTA. The spectra clearly show that the EDTA stripped the two zinc cations and produced the free $\mathbf{L3}$ ligand. In contrast, the set of spectra at the bottom of Figure 4 show that addition of two molar equivalents of Na_4PPi to the solution of $\mathbf{L3:2}\cdot\mathbf{Zn}(\text{NO}_3)_2$ greatly stabilized the dinuclear $\mathbf{L3}\cdot\mathbf{Zn}_2$ structure and prevented EDTA stripping of the zinc cations. Even the addition of one hundred molar equivalents of EDTA was unable to remove any measurable amount of Zn^{2+} . The data show clearly that PPi^{4-} , a strongly binding, bridging oxyanion, can stabilize the dinuclear $\mathbf{L3}\cdot\mathbf{Zn}_2$ complex by several orders of magnitude compared to a weakly binding anion such as NO_3^- .

Solution-State NMR Titrations

Additional structural evidence for the bridging anion stabilization effect was obtained by monitoring analogous $\text{Zn}(\text{NO}_3)_2$ titration experiments using ^1H NMR spectroscopy. Shown in Figure 5 are partial ^1H NMR spectra from the titration of mono-DPA ligand $\mathbf{L2}$ with $\text{Zn}(\text{NO}_3)_2$ in $\text{CD}_3\text{OD:D}_2\text{O}$ (4:1). As expected, there was a smooth transition from free $\mathbf{L2}$ to $\mathbf{L2:Zn}$ complex with very strong affinity and 1:1 ligand/zinc stoichiometry. The two pyridyl rings in the DPA unit exhibit an equivalent set of four proton chemical shifts. Figures 6–8 show ^1H NMR spectra obtained during the titration of bis-DPA ligand $\mathbf{L1}$ with $\text{Zn}(\text{NO}_3)_2$ in $\text{CD}_3\text{OD:D}_2\text{O}$ (4:1) under three different conditions respectively: (i) no additional salt present, (ii) presence of NaOAc , and (iii) presence of Na_4PPi . The peak assignments on the spectra refer to the atom labels in Scheme 2 as elucidated by analyzing COSY spectra. The partial NMR spectra in Figure 6 show the titration of $\mathbf{L1}$ with $\text{Zn}(\text{NO}_3)_2$ with no other salt present. The first species to appear is the asymmetric $\mathbf{L1:Zn}$ complex with 1:1 stoichiometry. The most diagnostic peaks are the two aryl peaks on the central phenoxy ring which are non-equivalent at 6.98 and 7.25 ppm (assigned as f and f' on the generic mononuclear $\mathbf{BDPA}\cdot\mathbf{Zn}$ structure in Scheme 2). These peaks disappear as more $\text{Zn}(\text{NO}_3)_2$ is added, and they are replaced by the symmetric dinuclear $\mathbf{L1:2}\cdot\mathbf{Zn}$ complex. However, the

pyridyl proton peaks are very broad, indicating a fluxional structure. The only sharp NMR peak is at 6.86 ppm and corresponds to the two equivalent protons on the central phenoxy ring (labeled as *k* on the generic **BDPA**·**Zn**₂ structure in Scheme 2). The sodium salts of several anions were investigated to see if they influenced the zinc association equilibria. The presence of NaCl had no effect, with the titration spectra clearly showing stepwise production of the mononuclear species followed by the dinuclear complex (data not shown). In contrast, the presence of NaOAc promoted formation of the dinuclear zinc species with no evidence for any intermediate mononuclear structure. The spectra in Figure 7 show direct conversion of free **L1** into the dinuclear **L1**:2·**Zn** complex with eight, broadened inequivalent pyridyl proton signals. This spectral feature is consistent with the X-ray structure in Figure 1, which is a *C*₂ symmetric bis-acetoxy bridged complex with one zinc-coordinated pyridyl ring in pseudo axial position and the partner pyridyl ring in pseudo equatorial position. The broadness of the pyridyl peaks indicates that the pyridyl rings undergo exchange of pseudo axial/pseudo equatorial Zn²⁺-coordination positions. In Figure 8 is the titration of **L1** with Zn(NO₃)₂ in the presence of Na₄PPi. Again, the spectra show direct conversion of free **L1** into the dinuclear **L1**:2·**Zn** complex. In this case, the eight inequivalent pyridyl peaks of the dinuclear **L1**:2·**Zn** complex are quite sharp indicating that the PPI strongly bridges the two zinc cations and produces a rigid, *C*₂ symmetric structure. Indeed, it is extremely likely that the structure is analogous to that shown in Figure 1 with a single tetradentate PPI⁴⁻ replacing the two bidentate OAc⁻ anions.^{3b,3d} The increased rigidity of the PPI⁴⁻ bridged structure is due to the higher anionic charge and stronger tetradentate zinc-coordination by the PPI⁴⁻ which prevents the zinc-oxygen bond dissociation events that allow the pyridyl rings to exchange between pseudo axial and pseudo equatorial coordination positions.

Conclusions

The crystallographic and spectroscopic results demonstrate that strongly bridging oxyanions such as OAc⁻ and PPI⁴⁻ associate with water soluble, dinuclear **BDPA**·**Zn**₂ complexes in aqueous solution and form relatively rigid, *C*₂ symmetric coordination structures with octahedral geometries around each Zn²⁺ center (Figure 1).^{1,3b,3d} It is reasonable to conclude that if the bridging anions were fatty acids or phospholipids then the resulting coordination complexes would be lipophilic enough to partition into a biomembrane. The relatively high lipophilicity of DPA ligands has been reported by nuclear imaging researchers to produce poor pharmacokinetics and they have mitigated this problem by developing DPA analogues with increased hydrophilicity.¹⁰ It should be possible to refine these hydrophilic molecular designs and produce next-generation, membrane impermeable dinuclear zinc bis-DPA complexes with improved cell recognition and bioimaging performance.

The solution-state absorption and NMR titration data show clearly that bridging oxyanions lock the two Zn²⁺ cations to the surrounding ligand and greatly enhance ligand/zinc affinity (Scheme 2). Thus, at sufficiently low zinc concentrations the molecular recognition process can be viewed as a three-component assembly of mononuclear zinc-ligand complex, second Zn²⁺ cation, and bridging oxyanion. This stepwise process can be employed to create optical anion sensor designs that use the anion-mediated association of the second zinc cation to induce a change in ligand optical properties.^{2c,5b} At higher zinc concentrations the bis-DPA ligand is saturated with both Zn²⁺ cations (*i.e.*, it exists entirely as a dinuclear **BDPA**·**Zn**₂ complex) but the two sets of zinc-coordinated pyridyl rings are in a fluxional state. Association of a bridging oxyanion rigidifies the zinc-coordinated pyridyl rings, a process that may decrease non-radiative decay of excited state energy and be the basis of a “turn on” fluorescent sensor design.^{3c,11} In addition to anion sensing, many DPA-containing receptors have been investigated as Zn²⁺ cation sensors and while it has been reported that the response of a bis-DPA Zn²⁺ sensor can be anion dependent,¹² the impact of this anion

dependence on analytical performance for this class of zinc sensors has not been widely discussed. Finally, it is worth noting that the three-component molecular assembly describe here mimics a common biological strategy for targeting the surface of biomembranes.¹³ Metal cations often act as bridging cofactors that enable membrane association of peripheral membrane proteins (*e.g.*, membrane targeting by annexin V mediated by Ca²⁺ ions).¹⁴

Experimental

Synthesis

The known ligands **L3** and **L4** were prepared by a literature method,^{3c} (an improved procedure is described in reference 11) and ligands **L1** and **L2** were prepared by the following related procedure.

L1: A solution of 4-bromophenol (0.88 g), 1 M HCl (0.5 mL), paraformaldehyde (0.35 g) and 2,2'-dipicolylamine (2.1 g) in ethanol (20 mL) and water (40 mL) was refluxed for 36 hours, then cooled to room temperature and neutralized with Na₂CO₃. The neutral suspension was extracted into chloroform, dried with sodium sulfate, and the solvent was evaporated to yield a yellow oil that was purified on a silica gel column using an eluent of 97:3 CHCl₃: MeOH give to give **L1** as a pale yellow oil (2.5 g), 83% yield. ¹H NMR (300 MHz, CDCl₃, ppm): δ 8.53 (m, 4H), 7.61 (td, 4H, J₁=7.8 Hz, J₂=1.8 Hz), 7.44 (d, 4H, J= 8.1 Hz), 7.33 (s, 2H), 7.14 (m, 4H), 3.87 (s, 8H), 3.77 (s, 4H). ¹³C NMR (75 MHz, CDCl₃, ppm): δ 158.9, 155.1, 148.8, 136.6, 131.4, 126.5, 122.8, 122.0, 110.2, 59.7, 54.2. MS (FAB⁺): HRMS C₃₂H₃₂BrN₆O calcd 595.1821, found 595.1832.

L2: The above procedure was used with the following stoichiometry: 4-bromophenol (0.88 g), 1 M HCl (0.5 mL), paraformaldehyde (0.35 g), 2,2'-dipicolylamine (1.0 g). The same work up and purification gave **L2** as pale yellow oil (1.7 g), 87 % yield. ¹H NMR (300 MHz, CDCl₃, ppm): δ 8.55 (m, 2H), 7.60 (td, 2H, J₁=7.8 Hz, J₂=1.8 Hz), 7.31 (d, 2H, J=7.8 Hz), 7.22 (dd, 1H, J₁=8.7 Hz, J₂=2.4 Hz), 7.14 (m, 3H), 6.80 (d, 1H, t=8.7 Hz), 3.87 (s, 4H), 3.74 (s, 2H). ¹³C NMR (75 MHz, CDCl₃, ppm): δ 157.8, 156.7, 148.6, 136.7, 132.4, 131.5, 125.0, 122.9, 122.1, 118.3, 110.1, 58.6, 56.1. MS (FAB⁺): HRMS C₁₉H₁₉BrN₃O calcd 384.0711, found 384.0699.

X-ray Crystallography

L1 Dinuclear Zinc Acetate Complex—A solution of **L1** ligand and two molar equivalents of triethylamine, Zn(OAc)₂, and NaBF₄ in methanol was refluxed for 30 minutes and then allowed to cool to room temperature.^{4a} Colorless crystals were formed after 24 h and they were found to be suitable for X-ray diffraction analysis. C₃₈H₄₄BBBrF₄N₆O₇Zn₂, M_r=994.25, triclinic, unit cell dimensions *a*=10.2479(5), *b*=12.5046(6), *c*=17.2421(10) Å, α=69.363(3), β=81.066(3), γ=86.542(3)°, V=2042.60(18) Å³, T=100(2) K, p $\bar{1}$, Z=2. The structure was refined on *F*₂ to *wR*₂=0.0809, conventional *R*₁=0.0310 [11493 reflections with *I*>2σ(*I*)], and a goodness of fit=1.055 for 10254 refined parameters. The asymmetric unit contains the Zn₂L1 complex coordinated by two OAc anions, one BF₄ anion, and two molecules of CH₃OH. One of the CH₃OH molecules is disordered over two sites, and site occupancy for the principal component is 0.872(70).

Absorption Titrations

Appropriate aliquots of Zn(NO₃)₂ solution (20 mM in 4:1 CH₃OH:H₂O volume ratio) were added to 3.0 mL solutions of ligand **L3** or **L4** in the presence or absence of NaCl, NaOAc, or Na₄PPI in the same solvent and the absorption spectrum acquired at 25 °C.

¹H NMR Titrations

Appropriate aliquots of Zn(NO₃)₂ solution (100 mM in 4:1 CH₃OD:D₂O volume ratio) were added to 0.75 mL solutions of ligand **L1** or **L2** in the presence or absence of NaCl, NaOAc, or Na₄PPI in the same solvent and the ¹H NMR spectrum acquired at 25 °C.

Acknowledgments

This work was supported by the University of Notre Dame and the NIH (USA). We are grateful to B. Noll and A. Oliver for acquiring and analyzing the X-ray data.

Notes and References

- (a) Adams H, Bradshaw D, Fenton DE. *Inorg. Chim. Acta.* 2002; 332:195–200.(b) Matsufuji K, Shiraiishi H, Miyasato Y, Shiga T, Ohba M, Yokoyama T, Okawa H. *Bull. Chem. Soc. Jpn.* 2005; 78:851–858.(c) Dalgaard P, Hazell A, McKenzie CJ, Moubaraki B, Murray KS. *Polyhedron.* 2000; 19:1909–1915.(d) Torelli S, Belle C, Gautier-Luneau I, Pierre JL. *Inorg. Chem.* 2000; 39:3526–3536. [PubMed: 11196811] (e) Belle C, Beguin C, Gautier-Luneau I, Hamman S, Philouze C, Pierre JL, Thomas F, Torelli S. *Inorg. Chem.* 2002; 41:479–491. [PubMed: 11825074] (f) Suzuki M, Kanatomi H, Murase I. *Chem. Lett.* 1981:1745–1748.(g) Seo JS, Sung N-D, Hynes RC, Chin J. *Inorg. Chem.* 1996; 35:7472–7473.(h) Maeda Y, Ishida A, Ohba M, Sugihara S, Hayami S. *Bull. Chem. Soc. Jpn.* 2002; 75:2441–2448.(i) Albedyhl S, Averbuch-Pouchot MT, Belle C, Krebs B, Pierre J-L, Saint-Aman E, Torelli S. *Eur. J. Inorg. Chem.* 2001:1457–1464.(j) Torelli S, Belle C, Gautier-Luneau I, Hamman S, Pierre J-L. *Inorg. Chim. Acta.* 2002; 333:144–147.(k) Durot S, Hossain LH, Hamman S, Jamet H, Orio M, Gautier-Luneau I, Luneau D, Philouze C, Pierre J-L, Belle C. *Inorg. Chem.* 2010; 49:7832–7840. [PubMed: 20704332]
- (a) Sakamoto T, Ojida A, Hamachi I. *Chem. Comm.* 2009; 2:141–152. [PubMed: 19099054] (b) Kruppa M, König B. *Chem. Rev.* 2006; 106:3520–3560. [PubMed: 16967915] (c) O'Neil EJ, Smith BD. *Coord. Chem. Rev.* 2006; 250:3068–3080.(d) Kim SK, Lee DH, Hong J, Yoon J. *Acc. Chem. Res.* 2009; 42:23–31. [PubMed: 18798656] (e) Zhou Y, Xu Z, Yoon J. *Chem. Soc. Rev.* 2011; 40:2222–2234. [PubMed: 21336366] (f) Ngo HT, Liu X, Jolliffe KA. *Chem. Soc. Rev.* 2012; 41:4928–4965. [PubMed: 22688834] For reviews on molecular recognition using Zn²⁺-DPA complexes, see:references therein
- (a) Han MS, Kim DH. *Angew. Chem. Int. Ed.* 2002; 41:3809–3811.(b) Lee DH, Im JH, Son SK, Chung YK, Hong J-I. *J. Am. Chem. Soc.* 2003; 125:7752–7753. [PubMed: 12822964] (c) Lee DH, Kim SY, Son SK, Chung YK, Hong J-I. *Angew. Chem. Int. Ed.* 2004; 43:4777–4780.(d) Lee JH, Park J, Lah MS, Chin J, Hong J-I. *Org. Lett.* 2007; 9:3729–3731. [PubMed: 17705498] (e) Bae SW, Cho MS, Jeong AR, Choi BR, Kim D-E, Yeo W-S, Hong J-I. *Small.* 2010; 6:1499–1503. [PubMed: 20623738] (f) Oh DJ, Kim KM, Ahn KH. *Chem. Asian J.* 2011; 6:2033–2038.(g) Liu G, Choi KY, Bhirde A, Swierczewska M, Yin J, Lee SW, Park JH, Hong J-I, Xie J, Niu G, Kiesewetter DO, Lee S, Chen X. *Angew. Chem. Int. Ed.* 2012; 51:445–449.(h) Honda K, Fujishima S-H, Ojida A, Hamachi I. *ChemBioChem.* 2007; 8:1370–1372. [PubMed: 17590878] (i) Ojida A, Honda K, Shimi D, Kiyonaka S, Moria Y, Hamachi I. *J. Am. Chem. Soc.* 2006; 128:10452–10459. [PubMed: 16895410] (j) Honda K, Ojida A, Hamachi I. *Chem. Comm.* 2006:4024–4026. [PubMed: 17003887] (k) Drewry JA, Burger S, Mazouchi A, Duodu E, Ayers P, Gradinaru CC, Gunning PT. *Med. Chem. Commun.* 2012; 3:763–770.For papers on molecular recognition using dinuclear **BDPA•Zn₂** complexes, see:
- (a) Morrow JR, Iranzo O. *Curr. Opin. Chem. Biol.* 2004; 8:192–200. [PubMed: 15062781] (b) Ganesh V, Bodewits K, Bartholdson SJ, Natale D, Campopiano DJ, Marque-Rivas JC. *Angew. Chem. Int. Ed.* 2009; 48:356–370.(c) Kinoshita E, Takahashi M, Takeda H, Shiro M, Koike T. *Dalton Trans.* 2004:1189–1194. [PubMed: 15252659] (d) Ciavatta L, Mareque JC, Natale D, Salvatore F. *Ann. Chim-Rome.* 2006; 96:317–325.For papers on molecular recognition and catalysis using structurally related hydroxyl bridged dinuclear Zn²⁺-DPA complexes, see:
- (a) Koulov AV, Stucker K, Lakshmi C, Robinson JP, Smith BD. *Cell Death Diff.* 2003; 10:1357–1359.(b) Koulov AV, Hanshaw RG, Stucker KA, Lakshmi C, Smith BD. *Isr. J. Chem.* 2005; 45:373–379.(c) Hanshaw RG, Lakshmi C, Lambert TN, Johnson JR, Smith BD. *Chem. Bio. Chem.* 2005; 6:2214–2220.(d) Leevy WM, Gammon ST, Johnson JR, Lampkins AJ, Jiang H, Marquez M,

- Piwnica-Worms D, Smith BD. *Bioconjugate Chem.* 2008; 19:686–692.(e) Smith BA, Akers WJ, Leevy WM, Lampkins AJ, Xiao S, Wolter W, Suckow MA, Achilefu S, Smith BD. *J. Am. Chem. Soc.* 2010; 132:67–69. [PubMed: 20014845] (f) Smith BA, Xiao S, Wolter W, Wheeler J, Suckow MA, Smith BD. *Apoptosis.* 2011; 16:722–731. [PubMed: 21499791] (g) Smith BA, Xie B-W, van Beek ER, Que I, Blankevoort V, Xiao S, Cole EL, Hoehn. M.; Kaijzel EL, Löwik CWGM, Smith BD. *ACS Chem. Neurosci.* 2012; 3:530–537. [PubMed: 22860222]
6. (a) DiVittorio KM, Leevy WM, O'Neil EJ, Johnson JR, Vakulenko S, Morris JD, Rosek KD, Serazin N, Hilkert S, Hurley S, Marquez M, Smith BD. *ChemBioChem.* 2008; 9:286–293. [PubMed: 18076009] (b) Jiang H, O'Neil EJ, DiVittorio KM, Smith BD. *Org. Lett.* 2005; 7:3013–3016. [PubMed: 15987193]
 7. Johnson JR, Jiang H, Smith BD. *Bioconjugate Chem.* 2008; 19:1033–1039.
 8. Reference 4d indicates that a structurally related hydroxyl bridged dinuclear Zn²⁺-DPA complex exhibits a similar set of stepwise equilibria for binding two Zn²⁺ cations.
 9. There is no doubt that the second zinc association step subsequently occurs, however, it does not produce a detectable change in absorbance.
 10. Maresca KP, Marquis JC, Hillier SM, Lu G, Femia FJ, Zimmerman CN, Eckelman WC, Joyal JL, Babich JW. *Bioconjugate Chem.* 2010; 21:1032–1042.
 11. Pathberiya LG, Barlow N, Nguyen T, Graham B, Tuck KL. *Tetrahedron.* 2012; 68:9435–9439. For a discussion of other mechanisms that may control the fluorescence signal, see:
 12. Jang YJ, Jun EJ, Lee YJ, Kim YS, Kim JS, Yoon J. *J. Org. Chem.* 2005; 70:9603–9606. [PubMed: 16268641]
 13. Hanshaw RG, Stahelin RV, Smith BD. *Chem. – Eur. J.* 2008; 14:1690–1697. [PubMed: 18085538]
 14. Lambert TN, Smith BD. *Coord. Chem. Rev.* 2003; 240:129–141.




Figure 1. X-ray crystal structure of $Zn_2L1(OAc)_2(BF_4) \cdot 2MeOH$. The structure omits the BF_4^- anion and two MeOH molecules for clarity. Thermal ellipsoids are indicated for all non-hydrogen atoms at the 50% probability level.




Figure 2. (left) Absorbance spectra of **L4** (20 μM) upon titration of Zn(NO₃)₂ in methanol/water (4:1 volume ratio); (right) absorbance at 306 nm.




Figure 3.

(a) Absorbance spectra of **L3** (20 μM) upon titration of Zn(NO₃)₂ in methanol/water (4:1 volume ratio); (b) same titration repeated in the presence of Na₄PPi (40 μM); (c) same titration repeated in the presence of NaOAc (40 μM). In each case, the graph on the right shows absorbance intensity change for the spectral wavelength maxima band.




Figure 4.

(top) Absorption spectra of $\mathbf{L3:2\cdot Zn(NO_3)_2}$ complex ($20\ \mu\text{M}$) showing complete conversion to $\mathbf{L3}$ upon addition of two molar equivalents of EDTA in water/methanol (4:1 volume ratio). (a) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ with no EDTA, (b) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ plus 0.5 equiv. EDTA, (c) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ plus 1.0 equiv. EDTA, (d) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ plus 2.0 equiv. EDTA, (e) $\mathbf{L3}$. (bottom) Absorption spectra of $\mathbf{L3:2\cdot Zn(NO_3)_2}$ complex ($20\ \mu\text{M}$) in the presence of Na_4PPi ($40\ \mu\text{M}$) is unchanged by the addition of one hundred molar equivalents of EDTA in water/methanol (4:1 volume ratio). (a) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ plus Na_4PPi , (b) $\mathbf{L3:2\cdot Zn(NO_3)_2}$ plus Na_4PPi and 100 equiv. EDTA, (c) $\mathbf{L3}$.




Figure 5. Partial ¹H NMR spectra of **L2** (10 mM) with increasing amounts of Zn(NO₃)₂ in CD₃OD/D₂O (4:1 volume ratio). The asterisk indicates a solvent peak.




Figure 6. Partial ^1H NMR spectra of **L1** (10 mM) with increasing amounts $\text{Zn}(\text{NO}_3)_2$ in $\text{CD}_3\text{OD}/\text{D}_2\text{O}$ (4:1 volume ratio).




Figure 7. Partial ^1H NMR spectra of **L1** (10 mM) in the presence of NaOAc (20 mM) with increasing amounts $\text{Zn}(\text{NO}_3)_2$ in $\text{CD}_3\text{OD}/\text{D}_2\text{O}$ (4:1 volume ratio).






Figure 8. Partial ^1H NMR spectra of **L1** (10 mM) in the presence of Na_4PPI (20 mM) with increasing amounts $\text{Zn}(\text{NO}_3)_2$ in $\text{CD}_3\text{OD}/\text{D}_2\text{O}$ (4:1 volume ratio).



Scheme 1.

**Scheme 2.**

Stepwise association of **BDPA** with Zn^{2+} to form **BDPA•Zn₂** is pushed to the right by the presence of bridging oxyanions, $\text{A}^- = \text{OAc}^-$ or PPi^{4-} .