

Sleep quality among residents of an old folk's home in Malaysia

Abdul Rashid, Eng Keat Ong, Eleanor Shu Yi Wong

ABSTRACT

Background: Sleep is an essential part of life. Lack of sleep has been linked with increased morbidity and mortality. The objective of this study was to determine the sleep quality of older adults residing in a private elderly care institution in Malaysia.

Materials and Methods: This cross sectional study was conducted among consenting residents of a 200-bed non-governmental charity old folks home in Penang, Malaysia. The sleep quality of the respondents was measured using the Pittsburgh Sleep Quality Index (PSQI). Quality of life (WHOQOL-BREF), their attitude to ageing (AAQ), Barthel index (Activities of Daily Living) and body mass index were also measured. Data was analysed using PASW.

Results: The PSQI score ranged from 0 to 16 with a mean score of 7.1 (SD 3.4) and 76.8% (116) had scores ≥ 5 . The differences in the mean score for chronic illness ($t = 0.14/P = 0.04$), the people that could be counted on for help ($t = 4.09/P = 0.02$) and the feasibility of getting practical help from fellow residents ($t = 4.41/P = 0.01$) were statistically significant. There was a negative correlation between the PSQI score and the WHOQOL-BREF score (-0.318/0.00) and AAQ score (-0.332/0.00).

Conclusions: Staff working in an elderly care institution should understand the important relationship of illnesses, social support and sleep hygiene in the wellbeing of the residents.

Key words: Aged, home for the aged, Malaysia, sleep

INTRODUCTION

Most countries across the continents are experiencing an ageing population phenomenon. It is projected that the population of older adults in the world will quadruple and that the elderly population in the developing countries will increase to almost two billion.^[1] The population of Malaysia is 25 million and is made up of several ethnic groups, comprising mostly of Malays and other indigenous groups followed by the Chinese, Indian and non-Malaysian citizens who mainly comprise of immigrant labourers. Malaysia's elderly population was 2.3% of the total population in 1997, in the year 2000 this proportion increased to 6.2% and it is projected that the elderly population of Malaysia will increase to 9.5% of the population in the year 2020.^[2] At present the population of the elderly in Malaysia is approximately two million. This increase can be attributed to improved health, longer life expectancy, low mortality as well as declining fertility.^[3] There has been a change in the family structure from the

traditional extended families into nuclear families. Due to loss of spouse and inability to care for self, there has been an increase in the admissions of the elderly into institutional care facilities. This pattern of living arrangement has caused social implications for the older adults.

Sleep is an essential part of life. There are two types of sleep; non rapid eye movement (NREM) and rapid eye movement (REM).^[3] The restoration and repair of body tissue and the normal functioning of endocrine and immune system is influenced by sleep. Sleep quality of an individual encompasses sleep duration, sleep latency, depth of sleep and restfulness.^[4] A good quality of sleep which is usually associated with fewer sleep disturbances helps to maintain mood, memory and cognitive performance.^[5] Lack of sleep has been linked with irritability and emotional distress, compromised endocrine and immune function, cognitive impairment, hypertension, impaired healing and obesity.^[5-7]

Older adults generally require only 5 to 6 hours of sleep per night and due to the change in the circadian rhythm they tend to awaken early.^[3] Studies have shown that older adults take longer time to sleep, may awaken more frequently during the night and have greater difficulty getting back to sleep and sleep is shorter and periods of rapid eye movement are also infrequent and apnoeic episodes are common.^[3,8]

Department of Public Health Medicine, Penang Medical College,
4 Jalan Sepoy Lines, 10450 Georgetown, Pulau Pinang, Malaysia

Address for Correspondence: Assoc. Prof. Dr. Abdul Rashid,
Department of Public Health Medicine, Penang Medical College,
4 Jalan Sepoy Lines, 10450 Georgetown, Pulau Pinang, Malaysia.
E-mail: drrashid10@gmail.com

Insomnia affects up to 30% of adult population affecting more women than men.^[9,10] In the United States of America (USA) the economic effects of insomnia has been estimated to range from US\$ 92 to 107 billion.^[11] Factors affecting sleep include inactive lifestyles, poor sleep practices, anxiety, depression, pain, discomfort, urgency, frequency and nocturia, restless legs, cramps, nocturnal cough or breathlessness, certain drugs and drug withdrawal.^[8]

Majority of older adults with sleep disturbances do not seek help probably due to lack of knowledge about insomnia and due to under recognition and under diagnosis by health care personnel.^[12]

Early detection and intervention of sleep disturbances among older adults can help reduce the morbidity and mortality associated with this and helps increase their quality of life. The objective of this study was to determine the sleep quality of older adults residing in a private elderly care residential institution in Penang Malaysia.

MATERIALS AND METHODS

This cross sectional study was conducted in a 200-bed non-governmental charity elderly care residential institution in Penang, Malaysia from May to November 2011. The eligibility of being admitted into this institution includes those aged 60 and above, has no person to care for them, homeless or lacking funds for self-care. Participants were taken from among the residents of this institution. A purposive convenience sampling method was employed. Only residents who consented and were not debilitated with an illness which rendered them unable to communicate effectively were recruited into this study. The data was collected by the researchers using a questionnaire especially designed for this study. Besides the baseline demographic information the sleep quality of the respondents was measured using the Pittsburgh Sleep Quality Index (PSQI). PSQI is a reliable tool consisting of seven components including (a) subjective sleep quality (1 item) – overall sleep quality (b) sleep latency (2 items) – time spent trying to fall asleep each night (c) sleep duration (1 item) – hours of actual sleep each night (d) habitual sleep efficiency (2 items) – number of hours slept divided by the number of hours spent in bed multiplied by 100 (e) sleep disturbances (9 items) – frequency of trouble sleeping caused by certain events (f) use of sleep medication (1 item) – frequency of taking medicine to help sleeping and (g) day time dysfunction (2 items) – difficulties to stay awake while doing daily activities. Each component is rated on a likert scale ranging from '0-3'. '0' represents the absence of any disorder and 3 represent maximum disorder. The sum of the scores range from 0-21, a higher

score suggests poorer sleep quality. A score of 5 and above on the PSQI total score indicate worse sleep quality. PSQI has a sensitivity of 89.6% and specificity of 86.5%.^[4] In addition to PSQI, the quality of life of the respondents was measured using WHOQOL-BREF. This scale has four domains including physical, psychological, social and environment. Higher scores suggest higher quality of life.^[13] The participant's attitude to ageing was measured using Attitude to Ageing Questionnaire (AAQ). The scale has three domains which include psychological growth, psychosocial loss and physical change. Higher total scores for the psychosocial loss component indicate a negative attitude to ageing whereas higher total scores for the physical change and psychological growth components indicate a positive attitude to ageing. The scores for the psychosocial loss domain were reversed in order to be in line with the other domains where a higher score reflects a more positive attitude to ageing. The total score on the AAQ was used to give an indication of attitude to ageing. Higher total scores of the three domains indicate a positive attitude to ageing.^[14] Barthel index which is a well-established and commonly used nursing tool was used to assess the functional independence in the activities of daily living (ADL) of the participants. Body mass index which is used to accurately determine a person's nutritional status was also calculated. Data was tabulated, cross tabulated and analysed using PASW version 18. Inferential analysis was done using t tests. Regression analysis was attempted to determine the predictive risk factors. A probability value of $P < 0.05$ was considered to be statistically significant. All respondents were asked to give an informed written consent before starting the interview. The anonymity of the respondents is assured.

RESULTS

Out of the total 200 residents, 151 responded to the survey. As shown in Table 1 most of the respondents were between the age group of 70 and 79, female, Chinese, Taoist and single. Most of them either had non-formal education or the highest level of education was up to primary school. Majority of them were working as labourers previously. Most perceived that they did not have anyone that they could count on for help. However, most perceived that people do take interest in them and that it was possible to get help from fellow residents and their main source of emotional support was their relatives. Majority perceived their health to be very good and their activities of daily living were not limited due to ill health, however majority had some form of chronic illness and suffered from musculoskeletal pain in the past 12 months. The mean Barthel, BMI, AAQ and WHOQOL BREF score was 16.8 (SD 4.5), 21.5 (SD 4.5), 64.5 (SD 10.1) and 69.4 (SD 11.6) respectively.

Table 1: Baseline profile of the respondents

Variables	Frequency	Percentage
Age		
60-69	27	17.9
70-79	72	47.7
≥80	52	34.4
Sex		
Female	82	54.3
Male	69	45.7
Race		
Chinese	149	98.7
Others	2	1.3
Religion		
Taoist	124	82.1
Buddhist	16	17.9
Others	11	57.0
Marital status		
Single	86	56.9
Married	38	25.2
Widow/divorce	27	17.9
Education		
Illiterate	56	37.1
Non formal/primary school	61	40.4
Secondary school/tertiary	34	22.5
Previous occupation		
Labourer	59	39.1
Skilled	42	27.8
Non skilled	33	21.9
Housewife	17	11.3
Next of Kin		
Family	73	48.3
Alone	50	33.1
Others	28	18.5
People that can be counted on for help		
None	97	64.2
1 or 2	33	21.9
≥3	21	13.9
Interest taken by others		
People take interest	65	43.0
Uncertain	68	45.0
Little or None	18	12.0
Feasibility of getting practical help from fellow residents		
Easy	37	24.5
Possible	106	70.2
Difficult	8	5.3
Main source of emotional support		

Contd...

Table 1: Contd...

Variables	Frequency	Percentage
Relatives	54	35.8
No one	45	29.8
Friends	43	28.5
Spouse	5	3.3
Staff at institution	4	2.6
Daily activity limited due to ill health		
Yes	52	34.4
No	99	65.6
Self-perception of health		
Excellent	17	11.3
Very good	54	35.8
Good	47	31.1
Fair	28	18.5
Poor	5	3.3
Chronic illness		
Yes	94	37.7
No	57	62.3
Musculoskeletal pain past 12 months		
Yes	100	33.8
No	51	66.2
Activities of daily living		
Mean score (SD)	16.8 (4.5)	
Independent	114	75.5
Needs minimal help	22	14.6
Dependent	15	9.9
BMI		
Total Score – Mean (SD)	21.5 (4.5)	
Under nutrition	43	28.5
Normal	80	53.0
Overweight	21	13.9
Obese	7	4.6
Attitudes on ageing		
Mean (SD)	64.5 (10.1)	
Quality of life		
WHO QOL BREF Mean (SD)	69.4 (11.6)	

The PSQI score ranged from 0 to 16 with a mean score of 7.1 (SD 3.4). Majority (76.8%, $n = 116$) of the residents had high sleep quality index (≥ 5) indicating poor sleep quality as shown in Table 2.

As shown in Table 3, the mean PSQI score of those with chronic illness was higher than those without chronic illness. This difference was statistically significant. The difference in the mean scores of people that could be counted on for help and the feasibility of getting practical help from fellow residents were also statistically significant. Post hoc test

Table 2: Base line Pittsburgh Sleep Quality Index score

PSQI	Score
Total PSQI score	
Minimum	0
Maximum	16
Mean (SD)	7.1 (3.4)
PSQI score < 5-n (%)	35 (23.2)
PSQI score ≥ 5-n (%)	116 (76.8)
Component 1: Subjective sleep quality	
Minimum	0
Maximum	3
Mean (SD)	1.1 (0.8)
Component 2: Sleep Latency	
Minimum	0
Maximum	3
Mean (SD)	1.9 (1.0)
Component 3: Sleep duration	
Minimum	0
Maximum	3
Mean (SD)	1.4 (1.2)
Component 4: Habitual sleep efficiency	
Minimum	0
Maximum	3
Mean (SD)	1.1 (1.2)
Component 5: Sleep disturbances	
Minimum	0
Maximum	2
Mean (SD)	1.1 (0.4)
Component 6: Use of sleeping medication	
Minimum	0
Maximum	1
Mean (SD)	0.02 (0.1)
Component 7: Day time dysfunction	
Minimum	0
Maximum	2
Mean (SD)	0.5 (0.6)

showed that the mean score of respondents who perceived that they had no one to count on for help was higher than the mean score of those who perceived they have 1 or 2 persons they could count on help. And those who perceived it was difficult to get practical help from fellow residents had higher PSQI scores than those who perceived it was easier.

Table 4 shows the mean differences for the variables; people that could be counted for help, feasibility of getting help from fellow residents and chronic illness status to the seven components of the PSQI score. It can be seen here that the mean score for sleep duration and habitual sleep efficiency

Table 3: Comparison of mean Pittsburgh Sleep Quality Index scores of different variables

Variables	Global PSQI score mean (SD)	t test or ANOVA (F)/P value	Post hoc
Age			
60-69	6.69 (3.89)	0.76/0.47	
70-79	6.95 (3.12)		
≥80	7.61 (3.61)		
Sex			
Female	7.05 (3.27)	0.68/0.95	
Male	7.09 (3.66)		
Race			
Chinese	7.03 (3.41)	-1.213/0.23	
Others	10.0 (7.07)		
Religion			
Taoist	7.18 (3.23)	0.68/0.51	
Buddhist	7.00 (4.02)		
Others	5.91 (3.45)		
Marital status			
Single	6.83 (3.42)	2.39/0.09	
Married	8.08 (3.61)		
Widow/divorce	6.41 (3.10)		
Education			
Illiterate	7.70 (3.04)	1.502/0.23	
Non formal/primary school	6.72 (3.28)		
Secondary school/tertiary	6.65 (4.25)		
Previous occupation			
Labourer	7.12 (3.07)	2.61/0.054	
Skilled	7.95 (3.83)		
Non skilled	5.76 (3.25)		
Housewife	7.24 (3.48)		
Next of Kin			
Family	7.58 (3.84)	1.55/0.22	
Alone	6.60 (2.74)		
Others	6.57 (3.43)		
People that can be counted on for help			
None	7.63 (3.47)	4.09/0.019	None>1 or 2
1 or 2	5.76 (3.15)		
≥3	6.52 (3.29)		
Feasibility of getting practical help from fellow residents			
Easy	6.03 (2.97)	4.41/0.014	Difficult>Easy
Possible	7.23 (3.47)		
Difficult	9.75 (3.85)		
Main source of emotional support			

Contd...

Table 3: Contd...

Variables	Global PSQI score mean (SD)	t test or ANOVA (F)/P value	Post hoc
Relatives	7.37 (3.37)	1.38/0.24	
No one	7.36 (3.45)		
Friends	6.21 (3.43)		
Spouse	6.60 (1.95)		
Staff at institution	9.50 (5.26)		
Daily activity limited due to ill health			
Yes	7.13 (3.08)	-0.22/0.83	
No	7.01 (3.73)		
Self-perception of health	6.00 (3.10)	0.58/0.68	
Excellent			
Very good	6.94 (3.18)		
Good	7.38 (3.56)		
Fair	7.39 (4.12)		
Poor	7.20 (2.39)		
Chronic Illness			
Yes	7.51 (3.53)	-2.06/0.04	
No	6.33 (3.22)		
Activities of daily living			
Independent	7.05 (3.49)	0.138/0.87	
Needs minimal help	6.86 (4.04)		
Dependent	7.47 (3.45)		
BMI			
Under nutrition	7.05 (3.68)	0.39/0.76	
Normal	6.90 (3.24)		
Overweight	7.81 (3.60)		
Obese	6.86 (4.38)		
Musculoskeletal pain past 12 months			
Yes	7.29 (3.28)	-1.12/0.27	
No	6.63 (3.76)		

of those who perceived that no one could be counted for help was higher than that of those who perceived 1 or 2 persons could be counted for help. The mean score for habitual sleep efficiency of those who perceived it was difficult and those who perceived it was possible to get help from fellow residents was higher than those who perceived it was easy.

There was a negative correlation between the PSQI score and WHOQOL-BREF score (-0.318/0.00) and the AAQ score (-0.332/0.00). A linear regression was attempted to determine the significant predictive variables which were associated with the PSQI scores. The model in Table 5 shows that 19.7% (R^2 0.197) variability in the PSQI score

was explained by the variables in the model. It can be seen that WHOQOL-BREF score and AAQ score are inversely associated to the PSQI score meaning that the quality of sleep is better as the attitude towards ageing and the quality of life of the respondents are positive.

DISCUSSION

The prevalence of poor sleep quality among these residents was high and the factors associated with it was poor social relationship reflected by the complaints of difficulty in getting help from fellow residents and fewer people that could be counted on for help. Presence of chronic illness was another factor associated with poor sleep quality index.

Decreased quality of sleep may cause a decreased sense of well-being, increased risk of accidents and increased morbidity and mortality rates.^[5-7] Population study has shown that approximately 10% of the general population have insomnia and this prevalence of insomnia rises with age.^[15] It has been reported that 50% of elderly suffer sleep disorder and women ages 80-89 have the highest prevalence rates.^[16,17] In a survey conducted by the National Sleep Foundation on older adults aged 55 to 84 found that about two thirds of the participants had complained of one or more symptoms of sleep problems at least a few nights a week.^[18] Study among elderly in community dwelling in Australia found 28% had difficulty falling asleep and 42% had difficulty falling asleep and maintaining sleep.^[19] In residential care centres the prevalence of sleep disorders have been reported from 44.4% in Iran,^[20] 45% in USA^[21] to 69.3% in Taiwan^[22] which are lower than the 76.8% reported in the present study.

Poor social relationship as evident by the number of people that could be counted for help and the feasibility of getting help was another factor which was associated with poor sleep quality in this study. This is not surprising considering social relationships have been linked with sleep quality.^[23] Most elderly Asian expect family centred care, however due to the change in family structure, loss of spouse and inability to care for self has led some elders to be institutionalized.^[24] The new experience of living in an institution may lead to the feeling of isolation compounded by the possible grief may potentiate insomnia in the elderly.^[25] In Taiwan many older adults in institutional care feel that their families have abandoned them.^[26] To make matters worse people with sleep problem seldom share their experiences with others because they believe that sleep problems are personal and a private matter making them feel lonely and helpless.

The poor quality of sleep affects one's health and psychosocial well-being. Because social relationships affect

Table 4: Comparison of the mean of the different components of the Pittsburgh Sleep Quality Index score with significant variables

PSQI components	People that can be counted on for help mean (SD)			Feasibility of getting help from fellow residents mean (SD)			Chronic illness mean (SD)		Post hoc
	None	1 or 2	≥3	Easy	Possible	Difficult	No	Yes	
Subjective sleep quality	1.09 (0.81)	0.88 (0.65)	0.81 (0.60)	0.84 (0.55)	1.03 (0.79)	1.50 (0.93)	0.84 (0.75) [#]	1.11 (0.74) [#]	
Sleep Latency	1.98 (0.98)	1.70 (1.10)	1.95 (1.02)	1.84 (1.07)	1.88 (1.00)	2.75 (0.46)	1.93 (0.98)	1.90 (1.03)	
Sleep duration	1.61 (1.19) [*]	0.94 (1.12) [*]	1.29 (1.10) [*]	1.16 (1.07)	1.48 (1.19)	1.75 (1.49)	1.19 (1.17)	1.55 (1.18)	None>1 or 2
Habitual sleep efficiency	1.34 (1.18) ^{**}	0.64 (0.96) ^{**}	0.90 (1.26) ^{**}	0.62 (0.98) ^{***}	1.25 (1.17) ^{***}	1.88 (1.55) ^{***}	0.86 (1.16) ^{###}	1.29 (1.18) ^{###}	None>1 or 2
									Possible>easy Difficult>easy
Sleep disturbances	1.14 (0.41)	1.12 (0.33)	0.95 (0.38)	1.05 (0.33)	1.14 (0.40)	1.00 (0.54)	1.05 (0.35)	1.15 (0.41)	
Use of sleeping medication	0.02 (0.14)	0.00 (0.00)	0.05 (0.22)	0.03 (0.16)	0.01 (0.09)	0.13 (0.354)	0.02 (0.13)	0.02 (0.15)	
Day time dysfunction	0.44 (0.63)	0.51 (0.09)	0.57 (0.68)	0.49 (0.61)	0.44 (0.62)	0.75 (0.45)	0.44 (0.57)	0.49 (0.64)	

*F=4.25 / P=0.02 **F=5.02 / P=0.01 ***F=5.84 / P=0.004 *t=-2.12 / P=0.04 **t=-2.18 / P=0.03

Table 5: Linear regression

	B	t	Sig	95% CI
*QOL score	-0.063	-2.503	0.013	-0.113; -0.013
*Attitude to ageing	-0.053	-2.653	0.009	-0.092; -0.013
ADL score	0.91	1.478	0.142	-0.031; 0.213
BMI score	0.015	0.253	0.801	-0.100; 0.130
People who are close that can be counted on	-0.117	-0.282	0.779	-0.941; 0.706
Feasibility to get help from fellow residents	0.878	1.519	0.131	-0.265; 2.020
Chronic illness	0.929	1.697	0.092	-0.153; 2.011

*significant

the quality of life^[27] it was expected to find a correlation between quality of life and quality of sleep. Numerous studies have shown that poor sleep quality adversely affect the quality of life.^[28-31] Environment is an important component of the WHOQOL-BREF. Environment is strongly associated with quality of life in respect to social interaction, activity involvement, independence and psychological wellbeing.^[32,33] Studies among the Thai and Myanmar older adults have shown that poor social relationship, chronic illness and sleep difficulty to be associated with poor quality of life.^[34,35]

Attitude, perception and belief of the elderly towards themselves affect the sleep quality of the elderly. The AAQ scale also has psychosocial component which take into account close relationships with members of the community and involvement in activities. In a study conducted in Japan to determine the relationship between sleep disturbance and morale in elderly people found that sleep disturbance to be related to negative attitude towards own aging.^[36]

Studies have shown that sleep efficiency is affected by age related medical and psychiatric illnesses.^[19,37] Many older adults suffer from one or more chronic medical conditions that can contribute to sleep difficulties.^[38] In one estimate it was reported that 80% of older adults suffer at least a chronic disorder which makes them susceptible to sleep disorders.^[39] In a study conducted by the National Institute of Health in Malaysia, 81.4% of the older adults had at least one chronic illness and 12.7% had at least three or more chronic illnesses.^[40] In a study conducted to examine the interplay of social engagement, sleep quality and plasma levels of interleukin-6, regression analysis showed that the association of age and sleep efficiency was mediated by the presence of chronic health conditions.^[23] In another study conducted in Taiwan among elderly Taiwanese aboriginal women found that chronic illness as a significant factor associated with poor sleep quality.^[41]

CONCLUSION

The prevalence of poor quality of sleep in this study population was high. Poor social relationships and negative attitude towards ageing as evident by the inverse relationship found between the QOL and the attitude towards ageing scores were significantly associated with the PSQI scores. Administrators of the institutions of care for the aged should be made aware the importance of social interactions.

ACKNOWLEDGMENTS

The authors would like to thank the staff and the management of the Silver Jubilee home for the elderly for all their support.

REFERENCES

1. Venne R. Mainstreaming the concerns of older persons into the social development agenda. United Nations Secretariat. Division for Social Policy and Development; 2005.
2. Zaimi AR. Social Welfare Policies And Services For The Elderly: A Country Report (Malaysia): For The 5th ASEAN and Japan High Level Officials Meeting on Caring Societies: Collaboration of Social Welfare and Health Services, and Development of Human Resources and Community -Community Services for the Elderly. 27-90uyjh6730 August 2007, Tokyo, Japan. Available from: [http://www.jicwels.or.jp/about_jicwels/ASEAN and JapanHighLevelOfficialsMeeting/5th20Mtg.%20Country%20Report%202007%20-Community%20Services%20for%20the%20Elderly-Malaysia%20Welfare.pdf](http://www.jicwels.or.jp/about_jicwels/ASEAN%20and%20JapanHighLevelOfficialsMeeting/5th%20Mtg.%20Country%20Report%202007%20-Community%20Services%20for%20the%20Elderly-Malaysia%20Welfare.pdf). [Last accessed on 2012 Jan 05].
3. O'Neill PA. Caring for the older adult: A health promotion perspective. Philadelphia: W.B. Saunders Company; 2002.
4. Buysse DJ, Reynolds CF, Monk TH, Berman SR. Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.
5. National Sleep Foundation. Sleep-awake cycle: Its physiology and impact on health. Washington DC, 2006. Available from: http://www.sleepfoundation.org/atf/cf/%7BF6BF2668-A1B4-4FEB-8D1A-A5D39340D9CB%7D/Sleep-Wake_Cycle.pdf. [Last accessed on 2011 Jan 12].
6. Knutson KL. Role of sleep duration and quality in the risk and severity of type 2 diabetes Mellitus. *Arch Intern Med* 2006;166:1768-74.
7. Flaherty JH. Insomnia among hospitalized older adults. *Clin Geriatr Med* 2008;24:51-67.
8. Nicholl C, Wilson KJ, Webster S. Elderly care medicine. 7th ed. Oxford UK: Blackwell Publishing; 2008.
9. Krystal AD. Insomnia in women. *Clin Cornerstone* 2003;5:41-50.
10. Zhang B, Wing YK. Sex difference in insomnia: A meta-analysis. *Sleep* 2006;29:85-93.
11. Stoller MK. Economic effects of insomnia. *Clin Ther* 1994;16:873-97.
12. Hellstrom A, Hagell P, Fagerstrom C, Willman A. Measurement properties of the Minimal Insomnia Symptom Scale (MISS) in an elderly population in Sweden. *BMC Geriatr* 2010;10:84
13. World Health Organization. WHOQOL – BREF introduction, administration, scoring and generic version of the assessment. Geneva: Programme on mental Health; 1996.
14. Laidlaw K, Power MJ, Schimdt S. The WHOQOL-OLD group. The attitudes to ageing questionnaire (AAQ): Development and psychometric properties. *Int J Geriatr Psychiatry* 2007;22:367-79.
15. Ford DE, Kamerow DB. Epidemiologic study of sleep disturbances and psychiatric disorders. An opportunity for prevention? *JAMA* 1989;262:1479-84.
16. Shives LR. Basic Concepts of Psychiatric-Mental Health Nursing. 6th ed. Philadelphia: Lippincott Williams and Wilkins; 2004.
17. Lichstein KL. Insomnia in the elderly. *Sleep Med Clin* 2006;1:221-9.
18. National Sleep Foundation. Sleep in America poll. Washington DC. Available from: <http://www.kintera.org/arf/cf/%7BF6BF2668-A1B4-4FEB-8D1A-A5D39340D9CB%7D/2003SleepPollExecSummary.pdf>. [Last accessed on 2011 Jan 12].
19. Foley DJ, Monjan A, Brown SL, Simonsick E, Wallace R, Blazer D. Sleep complaints among elderly persons: An epidemiological study of three communities. *Sleep* 1995;18:425-32.
20. Hosseini H, Esfirizi MF, Marandi SM, Rezaei A. The effect of Tai Chi exercise on the sleep quality of the elderly residents in Isfahan, Sadeghieh elderly home. *Iran J Nurs Midwifery Res* 2011;16:55-60.
21. Cohen D, Eisdorfer C, Prinz P, Breen A, Davis M, Gadsby A. Sleep disturbances in institutionalized age. *J Am Geriatr Soc* 1983;32:79-82.
22. Lin CL, Su TP, Chang M. Quality of sleep its associated factors in the institutionalized elderly. *Formosan J Med* 2003;7:174-84.
23. Friedman EM, Hayney MS, Love GD, Urry HL, Rosenkranz MA, Davidson RJ, et al. Social relationships, sleep quality and interleukin-6 in aging women. *Proc Natl Acad Sci U S A* 2005;102:18757-62.
24. Liu YW, Chuang HM. Investigation of the decision making procession of how disable elders' families choose long term care welfare services. *Soochow J Soc Work* 2006;14:91-123.
25. Ancoli-Israel S. All I want is a Good Night's. Chicago, IL: Mosby –Year Book Inc.; 1996.
26. Chen KM, Chen MH, Lin MH, Fan JT, Lin HS, Li CH. Effects of Yoga sleep quality and depression in elders in assisted living facilities. *J Nurs Res* 2010;18:53-61.
27. Mudey A, Ambekar S, Goyal RC, Agarekar S, Wagh VV. Assessment of Quality of Life among rural and urban elderly population of Wardha district, Maharashtra, India. *Ethno Med* 2011;5:89-93.
28. Newman AB, Nieto FJ, Guidry U, Lind BK, Redline S, Pickering TG, et al. Relation of sleep disordered breathing to cardiovascular disease risk factors: The sleep heart health study. *Am J Epidemiol* 2001;154:50-9.
29. Shahar E, Whitney CW, Redline S, Lee T, Newman AB, Javier Nieto F, et al. Sleep disordered breathing and cardiovascular disease: A cross sectional results of the sleep heart health study. *Am J Respir Crit Care Med* 2001;163:19-25.
30. Baldwin CM, Ervin A-M, Mays MZ, Robbins J, Shafazand S, Walsleben J, et al. Sleep disturbances, quality of life and ethnicity: The sleep heart health study. *J Clin Sleep Med* 2010;6:176-83.
31. Faubel R, Lopez-Garcia E, Guallar-Castillon P, Balboa-Castillo T, Gutierrez-Fisac J, Banegas JR, et al. Sleep duration and health related quality of life among older adults: A population based cohort in Spain. *Sleep* 2009;32:1059-68.
32. Hwang HF, Liang WM, Chiu YN, Lin MR. Suitability of the WHOQOL-BREF for the community-dwelling older people in Taiwan. *Age Ageing* 2003;32:593-600.
33. Lawton MP. The physical environment of the person with Alzheimer's disease. *Aging Ment Health* 2001;5:556-64.
34. Assantachai P, Maranetra N. Nationwide survey of the health status and quality of life of elderly Thais attending clubs for the elderly. *J Med Assoc Thai* 2003;86:938-46.
35. Naing MM, Nanthamongkolchai S, Munsawaengsub C. Quality of life of the elderly people in Einne Township Irrawaddy Division, Myanmar. *ASIAPH* 2010;1:4-10.
36. Ichimiya A, Iqata R, Ogomori K, Iqata T. The relationship between sleep disturbance and morale in Japanese elderly people. *Int Psychogeriatr* 2005;17:443-9.
37. Kryger M, Monjan A, Bliwise D, Ancoli-Israel S. Sleep, health and ageing. Bridging the gap between science and clinical practice. *Geriatrics* 2004;59:24-30.
38. Ancoli-Israel S, Cooke JR. Prevalence and comorbidity of insomnia and effect on functioning in elderly population. *J Am Geriatr Soc* 2005;53:1299-304.

39. Mauk KL. Gerontological Nursing: Competencies of Care. Massachusetts: Jones and Bartlett Publishers; 2006.
40. Chua SL. 2006. Statistics Given in speech by Minister of Health Malaysia, at the Asia Pacific Geriatric Conference. 7th April 2006. Available from: <http://www.moh.gov.my/Mohportal/DownloadServlet?id=701> and type 2. [Last accessed on 2006 Jan 22].
41. Chueh KH, Yang MS, Chen CS, Chiou SM. Poor sleep quality and alcohol use problems among elderly Taiwanese aboriginal women. *Int Psychogeriatr* 2009;21:593-9.

How to cite this article: Rashid A, Ong EK, Wong ESY. Sleep quality among residents of an old folk's home in Malaysia. *Iranian J Nursing Midwifery Res* 2012;17:512-9.

Source of Support: None, **Conflict of Interest:** None.