



Research

Cite this article: Warnecke L, Turner JM, Bollinger TK, Misra V, Cryan PM, Blehert DS, Wibbelt G, Willis CKR. 2013 Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing damage to mortality. *Biol Lett* 9: 20130177.
<http://dx.doi.org/10.1098/rsbl.2013.0177>

Received: 21 February 2013
Accepted: 8 May 2013

Subject Areas:

health and disease and epidemiology,
environmental science

Keywords:

dehydration, plasma electrolytes, *Geomyces destructans*, hibernation, hypovolemia

Author for correspondence:

Craig K. R. Willis
e-mail: c.willis@uwinnipeg.ca

[†]Present address: Department of Animal Ecology and Conservation, University Hamburg, 20146 Hamburg, Germany.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2013.0177> or via <http://rsbl.royalsocietypublishing.org>.

Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing damage to mortality

Lisa Warnecke^{1,†}, James M. Turner^{1,†}, Trent K. Bollinger², Vikram Misra³, Paul M. Cryan⁴, David S. Blehert⁵, Gudrun Wibbelt⁶ and Craig K. R. Willis¹

¹Department of Biology, University of Winnipeg, Winnipeg, Manitoba, Canada R3B 2E9

²Department of Veterinary Pathology, and ³Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, Saskatchewan, Canada S7N 5B4

⁴US Geological Survey, Fort Collins Science Center, Fort Collins, CO 80526, USA

⁵US Geological Survey, National Wildlife Health Center, Madison, WI 53711, USA

⁶Leibniz Institute for Zoo and Wildlife Research, 10315 Berlin, Germany

White-nose syndrome is devastating North American bat populations but we lack basic information on disease mechanisms. Altered blood physiology owing to epidermal invasion by the fungal pathogen *Geomyces destructans* (*Gd*) has been hypothesized as a cause of disrupted torpor patterns of affected hibernating bats, leading to mortality. Here, we present data on blood electrolyte concentration, haematology and acid–base balance of hibernating little brown bats, *Myotis lucifugus*, following experimental inoculation with *Gd*. Compared with controls, infected bats showed electrolyte depletion (i.e. lower plasma sodium), changes in haematology (i.e. increased haematocrit and decreased glucose) and disrupted acid–base balance (i.e. lower CO₂ partial pressure and bicarbonate). These findings indicate hypotonic dehydration, hypovolaemia and metabolic acidosis. We propose a mechanistic model linking tissue damage to altered homeostasis and morbidity/mortality.

1. Introduction

Infectious fungal diseases are causing unprecedented wildlife die-offs [1]. Amphibian chytridiomycosis has caused the most severe disease-related loss of biodiversity ever observed [2], and bat white-nose syndrome (WNS) in North America has caused the fastest decline of wild mammals in history, threatening common species with extinction [3]. Both diseases are caused by fungal skin infections. In amphibians, disrupted cutaneous function appears to cause a cascade of pathophysiological changes leading to mortality [4]. In bats, the WNS pathogen, *Geomyces destructans* (*Gd*), affects the skin of the nose, muzzle and ears of bats but most severely damages the wings [5,6], which play a crucial role in thermoregulation, water economy, gas exchange and immune function [7,8]. Altered water and electrolyte balances due to wing lesions have been hypothesized as important in WNS pathogenicity [9–11], while mortality from WNS is preceded by a progressive increase in the frequency of periodic arousals from torpor during hibernation and premature fat depletion [12]. Therefore, interactions among cutaneous infection, hibernation physiology and mortality are likely, but mechanisms linking these processes are not understood.

Skin lesions that occur with *Gd* infection superficially resemble burn injuries and could lead to similar complications, including fluid loss [9,10,13]. This has led to the ‘hypotonic dehydration hypothesis’ that affected bats exhibit increased water and solute loss across wing lesions, replenish body water by drinking inside hibernacula but eventually suffer electrolyte depletion because they have no access to food [10]. Measurements of sodium and chloride

from wild and experimentally infected, WNS-positive little brown bats (*Myotis lucifugus*) are consistent with this hypothesis [10]. However, pronounced fluid loss and hypotonic dehydration predict other changes in blood physiology such as hypovolaemia and metabolic acidosis [14]. In addition to shedding light on the pathophysiology of WNS, understanding how these factors influence morbidity and mortality could be valuable for rehabilitation of infected bats and potentially disease mitigation.

We studied blood physiology of *M. lucifugus* experimentally inoculated with *Gd* by Warnecke *et al.* [12] to test the hypotonic dehydration hypothesis [10], identify other consequences of *Gd* infection and help understand WNS mortality. Specifically, we tested for changes in electrolytes, haematology and acid–base balance. We predicted that infected bats would exhibit: (i) altered extracellular electrolyte concentrations (e.g. reduced sodium) owing to fluid loss over the epidermis from wing lesions [10]; (ii) haematological evidence of dehydration and starvation (e.g. elevated haematocrit owing to fluid loss [9] and decreased glucose because of depleted fat reserves [12]); and (iii) evidence of disrupted acid–base balance associated with hypovolaemia (e.g. reduced pH with reduction of bicarbonate and respiratory compensation resulting in decreased carbon dioxide). We also predicted that the magnitude of these changes would be associated with severity of wing tissue necrosis.

2. Material and methods

We collected 54 *M. lucifugus* from a WNS-negative cave and inoculated them with either a North American isolate of *Gd*, a European isolate or sham-inoculate ($n = 18$ each) as described in Warnecke *et al.* [12] (see the electronic supplementary material). All inoculated bats, but no controls, contracted WNS based on histopathology [12]. Bats were housed at 7°C and greater than 97% relative humidity with ad libitum water. After four months, we increased enclosure temperature to 25°C for approximately 1 h to assist rewarming from torpor, anaesthetized bats and collected blood (see the electronic supplementary material). Sampling from infected bats was complicated by high blood viscosity, but we obtained samples from eight inoculated individuals and all controls. Pathogenicity of both *Gd* isolates was similar [12] so we pooled results for inoculated bats. Using a blood analyser (i-STAT1, CG8+ cartridge, Abaxis, Union City, CA) we measured, (i) electrolytes: concentrations of sodium ($[Na^+]$, $mmol\ l^{-1}$) and potassium ($[K^+]$, $mmol\ l^{-1}$); (ii) haematology: haematocrit (Hct, % packed cell volume) and glucose concentration ($[Glu]$, $mg\ dl^{-1}$); (iii) acid–base: carbon dioxide partial pressure (pCO_2 , mmHg) and pH. The analyser calculated bicarbonate concentration ($[HCO_3^-]$, $mmol\ l^{-1}$) based on pH and pCO_2 .

To quantify severity of infection, we processed, using standard histological techniques and periodic acid-Schiff stain, a total of 15–20 cm of 5 μ m-thick sections of skin from the entire left wing of each bat and evaluated them using light microscopy [6]. We scored necrosis based on the percentage of skin surface with fungal hyphae present that was necrotic (i.e. loss of epidermis and disruption of underlying connective tissue): 0, no necrotic tissue; 1, less than 1 per cent; 2, 1–10%; 3, 10–30%; 4, 30–50%; 5, greater than 50 per cent. We used *t*-tests to assess the effect of infection on blood parameters, and regression to test for relationships between necrosis and blood parameters using Statistix v. 1.9 (see the electronic supplementary material). We adjusted alpha levels ($\alpha = 0.1$ due to our small sample size) using the false discovery rate procedure suggested by Narum [15]: *t*-tests: $\alpha = 0.07$, regressions: $\alpha = 0.04$ (see the electronic supplementary material for details).

Table 1. Statistics for the effects of *Geomyces destructans* infection on blood parameters, and regression analyses describing the relationship between necrosis score and blood parameters.

category	variable	<i>t</i> -test (infected versus control)			regression (effect of necrosis score)			slope coefficient	$\pm 95\%$ CI	
		<i>t</i>	d.f.	<i>p</i>	<i>F</i>	d.f.	<i>p</i>			intercept
electrolytes	$[Na^+]$	5.06	24	<0.001 ^a	10.28	1,6	0.018 ^a	141.9	–4.714	–8.311/–1.117
	$[K^+]$	–0.52	24	0.608 ^b	3.67	1,6	0.104	3.5	0.697	–0.194/1.588
haematology	Hct	–2.44	8.44	0.040 ^a	2.41	1,6	0.172	42.3	5.120	–2.956/13.196
	$[Glu]$	2.60	24	0.016 ^a	0.13	1,6	0.730	78.7	–3.593	–27.938/20.752
acid–base	pCO_2	3.03	24	0.006 ^a	8.18	1,6	0.029 ^a	38.1	–4.272	–7.928/–0.617
	pH	1.32	24	0.200	2.71	1,6	0.150	7.6	–0.049	–0.121/0.024
	$[HCO_3^-]$	3.65	8.24	0.006 ^a	15.93	1,6	0.007 ^a	36.1	–5.577	–8.997/–2.158

^aRepresents $p < 0.07$ (*t*-tests) and $p < 0.04$ (regression).

^bData log-transformed.




Figure 1. (a) Blood parameters of bats infected with *Geomyces destructans* (inf., black bars, $n = 8$) versus controls (con., white bars, $n = 18$) illustrating electrolyte concentrations, haematology and acid–base balance. Error bars indicate standard error, and asterisk represents $p < 0.07$ (table 1). (b) Relationship between wing necrosis score and sodium concentration and carbon dioxide partial pressure.

3. Results

Blood parameters differed between infected and control bats (table 1). In terms of electrolytes, sodium concentration was lower for infected bats than controls but potassium did not differ (figure 1a and table 1). Haematology of infected bats differed from controls, with increased haematocrit and decreased glucose concentration (figure 1a and table 1). With regards to acid–base status, we found reduced $p\text{CO}_2$ and bicarbonate concentrations, but no change in pH (figure 1a and table 1).

Mean necrosis score of infected bats was 2.2 ± 0.39 (range 1.0–4.0). Sodium concentration, as well as $p\text{CO}_2$ and bicarbonate concentrations, were negatively related to the severity of necrosis but other variables were not (figure 1b and table 1).

4. Discussion

Bats inoculated with *Gd* showed pronounced changes in blood physiology. Specifically, we observed (i) extracellular electrolyte depletion (reduced sodium), (ii) changes in haematology indicative of dehydration (increased haematocrit) and possibly starvation (decreased glucose) and (iii) evidence consistent with disrupted acid–base balance (reduced $p\text{CO}_2$ and bicarbonate). Sodium, $p\text{CO}_2$ and bicarbonate were negatively associated with wing necrosis score suggesting a mechanistic link between wing damage and pathophysiology. This supports the hypothesis that hypotonic dehydration associated

with fungal infection contributes to WNS mortality, and implies that water and electrolyte imbalance, reduced blood supply to tissues and altered acid–base balance play a role in disease-related morbidity.

The decreased sodium concentrations we observed are consistent with severe fluid loss. Sodium concentrations measured previously for naturally and experimentally infected *M. lucifugus*, as well as healthy controls [10], were nearly identical to the values we observed. Cryan *et al.* [10] also found reduced chloride (not measured in our study) but ruled out renal failure based on normal urine specific gravity values [10], and histological examination showed no evidence of kidney damage in their study, or ours [12]. Hypotonic dehydration can be caused by severe burns [16], and burn victims may suffer hypovolaemia and increased evaporative water loss (EWL; [13]). In addition to sodium depletion, we found elevated haematocrit, which could be explained by erythrocyte swelling in hypotonic plasma but is also consistent with hypovolaemia and dehydration [14] and, given the critical water economy of hibernating bats in general [17], a dehydrated state is highly plausible. Taken together, these results support the hypothesis that electrolyte imbalance resulted from fluid loss across the compromised epidermis.

Based on these results, and the emaciated state of infected bats [12], we propose a mechanistic model connecting wing lesions caused by *Gd* with a cascade of pathophysiological responses, disrupted homeostasis and morbidity/mortality (figure 2). Wing damage in our model has two direct




Figure 2. Theoretical model connecting damage to bat wings caused by *Geomyces destructans* with disruption of physiological processes which maintain homeostasis during winter hibernation, thus ultimately leading to mortality. Dashed line indicates alternative possible relationship. EWL, evaporative water loss; RR, respiration rate.

consequences: damage to the epidermis enables fluid loss and depletion of sodium (and presumably chloride [10]), while damage to underlying connective tissue increases vascular permeability, further accelerating fluid loss [9,10]. Both contribute to reduced plasma volume (i.e. hypovolaemia), causing elevated haematocrit and increased blood viscosity [14]. Hypovolaemia reduces blood pressure and inhibits capillary refill, causing local hypoxia. Hypovolaemia stimulates thirst [18] and could trigger bats to arouse from torpor to drink. EWL also influences torpor bout duration in hibernators even at high relative humidity [17]. Thus, fluid loss over the compromised epidermis could explain the increased frequency of arousals from torpor and premature fat depletion we reported previously [12]. Results here suggest that increased arousals reflected a response to elevated EWL, hypovolaemia and thirst.

Our model explains the disruption in acid–base balance suggested by our results. Dehydration and hypovolaemia can cause metabolic acidosis due to anaerobic lactic acid production in tissues with reduced blood flow. Plasma pH was not significantly reduced with infection probably owing to buffering by bicarbonate and the quick response by peripheral chemoreceptors to acidosis triggering an increase in respiration rate to off-load CO_2 [19]. This is consistent with the decreased $p\text{CO}_2$ we observed. Increased respiration

would further increase EWL and therefore arousal frequency [17]. Thus, the model accounts for altered torpor patterns as a combined result of hypotonic dehydration and respiratory compensation for metabolic acidosis. Starvation, hypovolaemia, metabolic acidosis or some combination of the three could all be the final cause of death.

Our model is hypothetical due to lack of data on the normal physiology of healthy, hibernating bats and on other pathophysiological effects of *Gd*. There are alternative explanations for the changes we observed. Hibernating ground squirrels (*Ictidomys tridecemlineatus*) preferentially rely on fatty acids and fat-derived ketones to fuel metabolism and, although healthy individuals are not acidotic, ketosis can cause metabolic acidosis in starvation [20]. Therefore, evidence for metabolic acidosis, combined with the reduced glucose we observed, could reflect increased energy expenditure with keto-acidosis and/or hypoglycaemia as potential causes of death. Owing to small blood volumes collected, we could not measure all relevant parameters. We suggest that future studies quantify serum/plasma protein and albumin concentrations to confirm hypovolaemia; plasma chloride, sodium, potassium and bicarbonate to calculate anion gap and test for metabolic acidosis; and lactate and ketone concentrations to differentiate lactic from keto-acidosis. Our model provides testable hypotheses that can be targeted by studies of particular pathways.

Epidermal invasion by *Gd* and subsequent pathophysiological changes are reminiscent of amphibian chytridiomycosis [9]. Both pathogens directly affect only skin but cause fatal disruptions of homeostasis. Although underlying mechanisms appear to differ (i.e. in amphibians, electrolyte imbalance leads to cardiac arrest [4]), WNS is superficially similar. Given the scale of wildlife population declines caused by fungal pathogens [1], understanding the pathophysiology underlying these diseases is important for developing strategies to hopefully mitigate mortality of affected species.

Methods were approved by the University Committee on Animal Care and Supply of the University of Saskatchewan.

We thank ACU and CCWHC staff in Saskatoon. K. Castle, C. Cooper, L. McGuire, J. Voyles, N.R. Willis, P. Withers, M. Wojciechowski and two reviewers gave outstanding feedback. Financially supported by US Fish and Wildlife Service, Natural Sciences and Engineering Research Council (Canada), Canada Foundation for Innovation, Manitoba Research and Innovation Fund, Government of Canada Post-doctoral Fellowship and German Academic Exchange Service (DAAD). Any use of trade products or firm names is for description only and does not imply endorsement by the US Government.

References

- Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012 Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**, 186–194. (doi:10.1038/nature10947)
- Wake DB, Vredenburg VT. 2008 Are we in the midst of the sixth mass extinction? A view from the world of amphibians. *Proc. Natl Acad. Sci. USA* **105**, 11 466–11 473. (doi:10.1073/pnas.0801921105)
- Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, Butchkoski CM, Kunz TH. 2010 An emerging disease causes regional population collapse of a common North American bat species. *Science* **329**, 679–682. (doi:10.1126/science.1188594)
- Voyles J *et al.* 2009 Pathogenesis of chytridiomycosis, a cause of catastrophic amphibian declines. *Science* **326**, 582–585. (doi:10.1126/science.1176765)
- Bleher DS, Hicks AC, Behr M. 2009 Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**, 227. (doi:10.1126/science.1163874)
- Meteyer CU, Buckles EL, Bleher DS, Hicks AC, Green DE, Shearn-Bochsler V, Thomas NJ, Gargas A, Behr MJ. 2009 Histopathologic criteria to confirm white-nose syndrome in bats. *J. Vet. Diagn. Invest.* **21**, 411–414. (doi:10.1177/104063870902100401)
- Herreid CF, Bretz WL, Schmidt-Nielsen K. 1968 Cutaneous gas exchange in bats. *Am. J. Physiol.* **215**, 506–508.
- Dongaonkar RM, Stewart RH, Laine GA, Davis MJ, Zawieja DC, Quick CM. 2009 Venomotion modulates lymphatic pumping in the bat wing. *Am. J. Physiol.*

- Heart Circul. Physiol.* **296**, H2015–H2021. (doi:10.1152/ajpheart.00418.2008)
9. Cryan PM, Meteyer CU, Boyles JG, Blehert DS. 2010 Wing pathology of white-nose syndrome in bats suggests life-threatening disruption of physiology. *BMC Biol.* **8**, 135. (doi:10.1186/1741-7007-8-135)
 10. Cryan PM *et al.* 2013 Electrolyte depletion in white-nose syndrome bats. *J. Wildl. Dis.* **49**, 398–402. (doi:10.7589/2012-04-121)
 11. Willis CKR, Menzies AK, Boyles JG, Wojciechowski MS. 2011 Evaporative water loss is a plausible explanation for mortality of bats from white-nose syndrome. *Integr. Comp. Biol.* **51**, 364–373. (doi:10.1093/icb/icr076)
 12. Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, Cryan PM, Wibbelt G, Blehert DS, Willis CKR. 2012 Inoculation of bats with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. *Proc. Nat. Acad. Sci. USA* **109**, 6999–7003. (doi:10.1073/pnas.1200374109)
 13. Keck M, Herndon DH, Kamolz LP, Frey M, Jeschke MG. 2009 Pathophysiology of burns. *Wien. Med. Wochenschr.* **159**, 327–336. (doi:10.1007/s10354-009-0651-2)
 14. Beck N. 2009 *Blood hematology*. London, UK: Springer.
 15. Narum SR. 2006 Beyond Bonferroni: less conservative analyses for conservation genetics. *Conserv. Genet.* **7**, 783–787. (doi:10.1007/s10592-005-9056-y)
 16. Gangopadhyay H. 2008 Clinical hyponatremia and hypernatremia. In *Renal failure and replacement therapies* (ed. S Blakeley), pp. 77–80. London, UK: Springer.
 17. Thomas DW, Geiser F. 1997 Periodic arousals in hibernating mammals: is evaporative water loss involved? *Funct. Ecol.* **11**, 585–591. (doi:10.1046/j.1365-2435.1997.00129.x)
 18. Stricker EM. 1968 Some physiological and motivational properties of the hypovolemic stimulus for thirst. *Physiol. Behav.* **3**, 379–385. (doi:10.1016/0031-9384(68)90066-8)
 19. Nestler JR. 1990 Relationships between respiratory quotient and metabolic rate during entry to and arousal from daily torpor in deer mice (*Peromyscus maniculatus*). *Physiol. Zool.* **63**, 504–515.
 20. Andrews MT, Russeth KP, Drewes LR, Henry P-G. 2009 Adaptive mechanisms regulate preferred utilization of ketones in the heart and brain of a hibernating mammal during arousal from torpor. *Am. J. Physiol.* **296**, R383–R393. (doi:10.1152/ajpregu.90795.2008)