

Use of cumulative mortality data in patients with acute myocardial infarction for early detection of variation in clinical practice: observational study

Richard A Lawrance, Micha F Dorsch, Robert J Sapsford, Alan F Mackintosh, Darren C Greenwood, Beryl M Jackson, Christine Morrell, Michael B Robinson, Alistair S Hall for the EMMACE (Evaluation of Methods and Management of Acute Coronary Events) Study Group

BHF Heart Research Centre, Leeds General Infirmary Leeds LS2 9JT
Richard A Lawrance
British Heart Foundation research fellow

Micha F Dorsch
British Heart Foundation research fellow

Robert J Sapsford
specialist registrar in cardiology

Beryl M Jackson
research nurse

Christine Morrell
research nurse

Alistair S Hall
professor of clinical cardiology

St James's University Hospital, Leeds LS9 7TF

Alan F Mackintosh
consultant cardiologist

Nuffield Institute for Health, University of Leeds, Leeds LS2 9PL

Darren C Greenwood
medical statistician

Michael B Robinson
consultant in public health and epidemiology

Correspondence to: A Hall
a.s.hall@leeds.ac.uk

Members of the EMMACE Study Group are listed at the end of the paper

BMJ 2001;323:324-7

Abstract

Objectives Use of cumulative mortality adjusted for case mix in patients with acute myocardial infarction for early detection of variation in clinical practice.

Design Observational study.

Setting 20 hospitals across the former Yorkshire region.

Participants All 2153 consecutive patients with confirmed acute myocardial infarction identified during three months.

Main outcome measures Variable life-adjusted displays showing cumulative differences between observed and expected mortality of patients; expected mortality calculated from risk model based on admission characteristics of age, heart rate, and systolic blood pressure.

Results The performance of two individual hospitals over three months was examined as an example. One, the smallest district hospital in the region, had a series of 30 consecutive patients but had five more deaths than predicted. The variable life-adjusted display showed minimal variation from that predicted for the first 15 patients followed by a run of unexpectedly high mortality. The second example was the main tertiary referral centre for the region, which admitted 188 consecutive patients. The display showed a period of apparently poor performance followed by substantial improvement, where the plot rose steadily from a cumulative net lives saved of -4 to 7. These variations in patient outcome are unlikely to have been revealed during conventional audit practice.

Conclusions Variable life-adjusted display has been integrated into surgical care as a graphical display of risk-adjusted survival for individual surgeons or centres. In combination with a simple risk model, it may have a role in monitoring performance and outcome in patients with acute myocardial infarction.

Introduction

The UK Department of Health has recently published a national service framework for coronary artery disease, which aims to coordinate efforts to improve medical care of patients by the implementation of

minimal standards of care and audit.¹ Many risk models have been derived in an attempt to reliably compare care at different centres. It is self evident that if a doctor or hospital is responsible for a high proportion of sick patients then simply assessing crude mortality will result in an unfavourable comparison of care. Most of the risk-adjusted models that have been derived suffer from overcomplexity, exclude substantial proportions of patients with higher risk profiles, use administrative data without clinical details, or are applicable only to selected subgroups.^{2,3} We have developed a model based on three objective admission clinical characteristics—age, heart rate, and systolic blood pressure. We have validated our model with subgroups of our patients with acute myocardial infarction, and it seems to accurately reflect patient outcome.⁴⁻⁶

Scoring systems, such as the Parsonnet score,⁷ have been used to assess individual surgeons' performance by comparing actual and expected death rates. Lovegrove et al have described the variable life-adjusted display,⁸ a graphical technique based on the cumulative sum (cusum) method^{9,10} that incorporates information of estimated risks for each individual case. Cusum charts record successive cases horizontally, with plots ascending by single units for each outcome event (such as death) reached. The variable life-adjusted display incorporates outcome information, simultaneously accounting for prior estimated risk in each case and plotting the difference in cumulative expected and observed mortality. Sherlaw-Johnson et al recently described a method for estimating variability for such displays in order to assess the probability that observed deviations occurred as a result of chance rather than because of a difference in quality of care.¹¹

In this study we extend the use of variable life-adjusted displays in combination with a simple, accurate, and validated risk model for assessing care of patients with acute myocardial infarction and suggest indications for more detailed review of patient care.

Subjects and methods

Patient population

Between 1 September and 30 November 1995, 3684 potential recruits to the evaluation of methods and

management of acute coronary events (EMMACE) study were identified in the 20 adjacent hospitals that comprised all units in the former Yorkshire region that admitted patients with acute myocardial infarction. We identified potential recruits from coronary care registers, clinical coding, and biochemistry records of requests for cardiac enzyme assays. We evaluated the patients' medical records and confirmed 2153 consecutive cases of acute myocardial infarction according to either the World Health Organization's criteria or to the clinical diagnosis reached by the attending physician. We included these 2153 patients, of whom 1643 were discharged from hospital alive after a first myocardial event, in our study. For data abstraction, we applied a hierarchical system of preferred data sources to maintain consistency among our research assistants: for example, we used the first recorded blood pressure and heart rate obtained from either the accident and emergency department or from medical or nursing notes.

We completed a 250 item record of demographic, clinical, and treatment variables for each patient according to a standardised operations manual and entered the data on a computer database. We included all consecutive patients with acute myocardial infarction regardless of age or place of care within a hospital. We considered only patients' first presentation with acute myocardial infarction (during the recruitment window), and patients transferred to a tertiary centre were counted only once for their index admission. We recorded patients' clinical characteristics on admission from the following sources in order of preference: (a) emergency department medical notes, (b) the admitting medical team's first clerking, or (c) nursing notes.

Calculation of expected mortality

We developed and externally validated a risk model with a demonstrable clinical utility in stratifying patients after myocardial infarction.^{4,5} The model calculates an individual's probability of death at 30 days (P_{30}) on the basis of three robust admission characteristics (age, heart rate, and systolic blood pressure). The calculation is

$$P_{30} = 1 / (1 + \exp(-L_{30})),$$

where

$$L_{30} = -5.624 + (0.085 \times \text{age}) + (0.014 \times \text{heart rate}) - (0.022 \times \text{systolic blood pressure}).$$

In our patient cohort its accuracy was demonstrated by the area under the receiver-operating characteristics (ROC) curve of 0.78. For the patients studied, the mean expected 30 day mortality was 23.5% and the observed mortality was 24.4%, giving a standardised mortality ratio of 1.04.

Variable life-adjusted display

Using the variable life-adjusted display, we can illustrate the cumulative mortality for a hospital or an individual doctor and the variation either side of that predicted. By calculating a patient's expected probability of death after myocardial infarction, we can also estimate the probability of survival. The line plotted in this graphical display varies about zero, ascending, for a patient who survives, by an amount equivalent to the estimated probability of death. For those patients who die by 30 days, the plot descends by an amount equal to the estimated probability of survival. It can be shown




Fig 1 Variable life-adjusted display of a small cottage hospital for a series of 30 consecutive patients with acute myocardial infarction. Predicted mortality was calculated by a validated risk score based on admission characteristics of age, systolic blood pressure, and heart rate

that there is a proportionally greater shift in the plot for patients with unexpected outcomes.

We examined two hospitals for variations in risk-adjusted 30 day survival over the three month period of study—hospital A, the smallest of the district hospitals studied, and hospital B, the supraregional tertiary referral centre for the area.

Results

The variable life-adjusted displays (figs 1 and 2) show the number of cases on the x axis and the cumulative difference between the predicted and observed deaths on the y axis. Figure 1 shows the results for hospital A, the centre with the highest standardised mortality ratio. This small hospital had a series of only 30 patients with acute myocardial infarction. The observed and expected mortalities were 44.2% and 27.5% respectively (standardised mortality ratio 1.61). The risk model would predict eight deaths, whereas the actual




Fig 2 Variable life-adjusted display of a large teaching hospital and regional cardiac centre for a series of 188 consecutive patients with acute myocardial infarction. Predicted mortality was calculated by a simple risk score based on admission characteristics of age, systolic blood pressure, and heart rate

number was 13. The display reveals minimal variation from that predicted for the first 15 patients followed by a run of unexpectedly high mortality.

Figure 2 shows the for hospital B, which admitted 188 patients with acute myocardial infarction during the study period. The plot terminates with 3.5 extra "lives" above expected (observed number of deaths 42, expected 45.5; standardised mortality ratio 0.92). Between case numbers 35 and 135 the plot ascends from a cumulative (expected minus observed) net lives gained of -4 to 7 .

Discussion

Using just patients' age, blood pressure, and heart rate to adjust for case mix, we derived a continuous plot to compare observed and expected outcome for patients with acute myocardial infarction.

The main purpose of audit is to provide high quality data analyses with which to monitor the required standards and to allow clinicians to evaluate the performance of their units. Block audit, however, is retrospective and does not allow immediate identification, and thus rectification, of potential problems. With application of a simple risk-adjusted model to populations of patients with acute myocardial infarction, we have produced variable life-adjusted displays. Such plots have become increasingly integrated into surgical care, allowing immediate monitoring of performance for individual surgeons or units over time, but they are not yet used in routine audit of medical patients.

Recently, Sherlaw-Johnson et al showed that prediction intervals can be derived for plots of standardised mortality ratio, with the preoperative case mix taken into account.¹¹ We did not calculate prediction intervals for our populations because we consider that the usefulness of the graphical displays is enhanced by being qualitative rather than quantitative. We do not consider that achievement of a P value less than 0.05 is helpful in the context of continuous, real time monitoring, nor should it provide evidence that is not in keeping with other qualitative indicators. For example, if hospital B (fig 2) had a major shortage of beds in its coronary care unit beds that coincided with the period of increased deaths, actions might be taken even without a significant P value. Although an apparent run of poor results may not represent a real change, a failure to act on the qualitative trend observed until a significant P value had been achieved could result in the needless loss of life.

Our display method of monitoring performance or outcome over time can be used as an early warning system to allow more detailed audit to be performed.¹² For example, the sharp descent of the plot in the second half of figure 1 highlights the possibility of poor performance. Conversely, in figure 2 the graph ascends from a cumulative net lives saved of -4 to 7 in the space of 100 consecutive cases, suggesting that any possible problem in the hospital was identified and corrected. Such periods of poor or above average performance are liable to be missed with block audit, especially when the overall figures are not significantly different from those forecast. As stated by de Mol when commenting on the original description of the variable life-adjusted display,¹³ differences in care measured by patient outcome are common. A judgment of whether

What is already known on this topic

The national service framework for coronary artery disease requires minimal standards of care and audit of patients with acute myocardial infarction but does not integrate clinical status into the audit tool

Predictive models using only a few factors to adjust for case mix are easy to use and may be as accurate as more complicated methods

Early identification of variations in patient outcome is not revealed by block audit, and, instead, a continuous monitoring process is required

What this study adds

Using just patients' age, blood pressure, and heart rate to adjust for case mix, a continuous plot can be derived to compare observed and expected outcome for patients with acute myocardial infarction

This method of monitoring outcome over time can be used as an early warning system to allow more detailed audit to be performed

this performance is above or below average could be based on predefined limits in order to adequately identify important outliers as well as permitting more cost effective and logical use of audit time. However, we would further suggest that rigorous application of a P value (which is itself subject to errors in the context of multiple observations) may mean that early qualitative trends towards harm remain unheeded. Solely relying on quantification of variability in performance rather than a qualitative assessment is therefore laden with shortcomings.¹⁴ A hypothetical example of this limitation is that 16 of 160 patients would have to die after routine cardiac surgery (10% mortality) before a statistically significant doubling of deaths could be detected with 90% chance.^{15 16} Common sense suggests that such a trend should be identified and corrected long before the problem was so far progressed, especially if other strong qualitative indicators were present.

Any risk model can become less accurate over time because of progressive improvements in care. The Parsonnet scoring system, derived in 1989, has consistently been shown to predict risk strata that correlate with the observed mortality in adult acquired heart disease. However, it equally consistently overestimates the risk of contemporary coronary surgery by up to 100%.^{7 17 18} In order to avoid this problem if our model were to be in general use, we would recommend that patients' heart rate and blood pressure on admission should be added to the core dataset of the national service framework for coronary artery disease. If hospitals were then to provide the data required for the risk model to a centralised database the model could be continually updated and made available on the internet. This would reduce the likelihood of overestimating risk in patients with acute myocardial infarction.

As stated in the national service framework documentation, good quality data can provide a powerful engine for change and improvement. The method we describe constitutes an improvement over simple clinical intuition and has a potentially important role in indicating the need for more detailed audit. In the interests of patient safety, we suggest that there is a need for continuous, qualitative observations of deviations from expected outcomes.

We thank all the staff in the biochemistry departments, coronary care units, audit and coding departments, pharmacies and medical records departments, and all non-cardiology consultants of the 20 acute hospitals in the former Yorkshire region, as well as M Allen at the Office for National Statistics. This paper is dedicated to the memory of Julia Oldham.

Members of the EMMACE study group: Alistair S Hall, Michael B Robinson (principal investigators); Robert J Sapsford (clinical coordinator); Beryl M Jackson, Christine Morrell (research assistants); Darren C Greenwood (statistical advisor); J Oldham (database manager); R J I Bain, S G Ball, P D Batin, K E Berkin, R M Boyle, J L Caplin, R S Clark, J C Cowan, J Dhawan, D Garg, G Kaye, S Khan, H Larkin, R V Lewis, A F Mackintosh, J McLenachan, M A Memon, L C A Morley, G W Morrison, M S Norrell, E J Perrins, M M Pye, G Reynolds, N P Silvertown, J H Smyllie, U Somasundram, R N Stevenson, J B Stoker, A P Tandon, L B Tan, C J P Welsh, C Weston, G J Williams, P T Wilmshurst, J I Wilson, A V Zezulka (key investigators).

Contributors: AFM conceived the original idea after the publication by Lovegrove et al.⁸ RAL and MFD performed the data analysis and wrote the original and revised drafts. RJS, CM, and BMJ performed the main data collection, with JO responsible for database design and management. DCG advised on statistical methods. All authors contributed to the conception, design, and interpretation of the data and approved the final version of the article. MBR and ASH are guarantors for the study.

Funding: This project was commissioned and funded by the NHS Research and Development Programme.

Competing interests: None declared.

- 1 National service framework for coronary heart disease: modern standards and service models. London: Department of Health, 2000.
- 2 Iezzoni LI, Ash AS, Coffman GA, Moskowitz MA. Predicting in hospital mortality. A comparison of severity measurement approaches. *Med Care* 1992;30:347-59.
- 3 Lee KL, Woodlief LH, Topol EJ, Weaver WD, Betriu A, Col J, et al for the GUSTO-I Investigators. Predictors of 30-day mortality in the era of reperfusion for acute myocardial infarction. Results from an international trial of 41,021 patients. *Circulation* 1995;91:1659-68.
- 4 Dorsch MF, Lawrance RA, Sapsford RJ, Oldham J, Greenwood DC, Jackson BM, et al on behalf of the EMMACE (Evaluation of Methods and Management of Acute Coronary Events) Study Group. A simple benchmark for evaluating quality of care of patients following acute myocardial infarction. *Heart* (in press).
- 5 Lawrance RA, Dorsch MF, Sapsford RJ, Hall AS. National service framework—Core data set. *eHeart* [online journal] 2000;84(2). <http://heart.bmjournals.com/cgi/eletters/84/2/116#EL1>.
- 6 Birkhead J. Re: National service framework—core data set. *eHeart* [online journal] 2000;84(2). <http://heart.bmjournals.com/cgi/eletters/84/2/116#EL1>.
- 7 Parsonnet V, Dean D, Bernstein AD. A method of uniform stratification of risk for evaluating the results of surgery in acquired adult heart disease. *Circulation* 1989;79(6 Pt 2):13-12.
- 8 Lovegrove J, Valencia O, Treasure T, Sherlaw-Johnson C, Gallivan S. Monitoring the results of cardiac surgery by variable life-adjusted display. *Lancet* 1997;350:1128-30.
- 9 De Leval MR, Francois K, Bull C, Brawn W, Spiegelhalter D. Analysis of a cluster of surgical failures. Application to a series of neonatal arterial switch operations. *J Thorac Cardiovasc Surg* 1994;107:914-23, 923-4.
- 10 Williams SM, Parry BR, Schlup MM. Quality control: an application of the cusum. *BMJ* 1992;304:1359-61.
- 11 Sherlaw-Johnson C, Lovegrove J, Treasure T, Gallivan S. Likely variations in perioperative mortality associated with cardiac surgery: when does high mortality reflect bad practice? *Heart* 2000;84:79-82.
- 12 Nashef SA. Monitoring of cardiac surgical results. *Lancet* 1998;351:143-4.
- 13 De Mol BA. Monitoring of cardiac surgical results. *Lancet* 1998;351:144.
- 14 Sergeant P, Meyns B. La critique est aisee mais l'art est difficile. *Lancet* 1997;350:1114-5.
- 15 Ecochard R, De Gevigney G, Colin C. Fair comparison of mortality data following cardiac surgery. *Heart* 2000;84:5-6.
- 16 Poloniecki J, Valencia O, Littlejohns P. Cumulative risk adjusted mortality chart for detecting changes in death rate: observational study of heart surgery. *BMJ* 1998;316:1697-700.
- 17 Nashef SAM, Carey F, Silcock MM, Oommen PK, Levy RD, Jones MT. Risk stratification for open heart surgery: trial of the Parsonnet system in a British hospital. *BMJ* 1992;305:1066-7.
- 18 Anderson JR, Parker DJ, Unsworth-White MJ, Treasure T, Valencia O. Training surgeons and safeguarding patients. *Ann R Coll Surg Engl* 1996;78(3 suppl):116-8. (Accepted 29 June 2001)

Life hangs on chance and the generosity of others

Laura came to us from the paediatric cystic fibrosis centre when she was 16 years old. She had severe lung disease and been placed on the transplant list in January 1997. She had just completed her GCSEs and had got top marks. She had just started her A levels and wanted to study theatre and drama at university. Her disease progressed; we put her on to overnight ventilation and permanent intravenous antibiotics.

Our patients listed for transplantation continued to die on the waiting list because of the shortage of donors. Laura's activities became more limited at school, and she needed help with mobility and portable oxygen. Characteristically, despite her limitations and against advice, she went on theatre trips to London and Dublin. Last summer, she completed her A levels and came into hospital exhausted from her efforts. She picked up a cold in hospital and deteriorated. Her chest radiograph showed confluent shadowing associated with terminal disease.

Claire, a close friend with severe disease, came back from holiday with a severe chest infection. Laura and Claire were too breathless to visit each other, although they were only a few rooms apart. Claire failed to respond and became breathless despite high flow oxygen. She asked us to allow her to die comfortably. We withdrew the antibiotics, and she died comfortably surrounded by her family.

Laura had now been in hospital for seven weeks and was bedbound. I told her mother that she would probably die. We informed the transplant team that the situation was desperate. Laura had now been on the transplant list for three years. At the ward round, we decided that medical treatment had failed and agreed to discuss with Laura as to whether she would like us to stop treatment as she was suffering so much. I went to see her on

my own at the end of the ward round to talk to her about dying, but, just as I started, the sister came in to tell me the senior transplant surgeon was on the phone; I had talked to him that morning about potential donors and he wondered whether Laura was now too sick to transplant. Now he told me that they had found a suitable donor.

I went back to Laura and now spoke to her about living rather than the dying as I had intended a few moments before. That night Laura received her transplant. After a stormy postoperative period, she was extubated on the day the A level results came out; her university place was assured with two As and a B. Nearly a year later she continues to prosper, talking continuously, taking a gap year before university, and sending emails under the name of demolition babe (I thought at first it was a virus).

We don't have enough organ donors. Other countries do better than us. The current government needs to do more, but currently two out of three transplant units (Sheffield, Birmingham, and Manchester) are proposed to close. Although all three sites were inspected over a year ago, the government has still not released the decision.

How can organ donation be improved? It might help if it was known that the current minister of health was on the donor registry. Is the prime minister? How many members of the cabinet are on the donor registry? The number of people carrying donor cards might increase if there was top down government support.

Anthony Kevin Webb *clinical director, Manchester Adult Cystic Fibrosis Unit, Wythenshawe Hospital, South Manchester Trust*