

Specific inhibitors of HCV polymerase identified using an NS5B with lower affinity for template/primer substrate

Ginette McKercher, Pierre L. Beaulieu¹, Daniel Lamarre, Steven LaPlante¹, Sylvain Lefebvre, Charles Pellerin, Louise Thauvette and George Kukolj*

Department of Biological Sciences and ¹Department of Chemistry, Boehringer Ingelheim Canada Ltd R&D, 2100 Cunard Street, Laval, Québec H7S 2G5, Canada

Received September 25, 2003; Revised and Accepted November 10, 2003

ABSTRACT

The interaction of the hepatitis C virus (HCV) RNA-dependent RNA polymerase with RNA substrate is incompletely defined. We have characterized the activities of the HCV NS5B polymerase, modified by different deletions and affinity tags, with a routinely used homopolymeric substrate, and established apparent affinities of the various NS5B constructs both for the NTP and the template/primer substrates. We identified a uniquely tagged HCV NS5B RNA polymerase construct with a lower affinity (higher K_m) than mature HCV NS5B for template/primer substrate and highlighted the use of such a polymerase for the identification of inhibitors of NS5B activity, particularly inhibitors of productive RNA binding. The characterization of specific benzimidazole-5-carboxamide-based inhibitors, identified in a screening campaign, revealed that this class of compounds was non-competitive with regard to NTP incorporation and had no effect on processive elongation, but inhibited an initiation phase of the HCV polymerase activity. The potency of these compounds versus a panel of different NS5B polymerase constructs was inversely proportional to the enzymes' affinities for template/primer substrate. The benzimidazole-5-carboxamide compounds also inhibited the full-length, untagged NS5B *de novo* initiation reaction using HCV 3'-UTR substrate RNA and expand the diversifying pool of potential HCV replication inhibitors.

INTRODUCTION

Discovered more than a decade ago, hepatitis C virus (HCV) prevalence is now estimated at approximately 170 million people worldwide (1,2). The plus-strand HCV RNA genome is ~9600 nt in length and encodes at least one open-reading frame with approximately 3010 amino acids. In infected cells, this polyprotein is cleaved at multiple sites by cellular and

viral proteases to produce structural and non-structural (NS) proteins (3,4). One of the NS proteins, NS5B, is an RNA-dependent RNA polymerase (RdRp) that catalyzes the replication of HCV (5). The enzyme is a prime target in the search for inhibitors of HCV replication and a variety of *in vitro* assays for HCV NS5B polymerase activity have been developed. Though primer-independent *de novo* initiation of complementary strand RNA polymerization can be reconstituted *in vitro* with templates that represent the 3' end of either the plus- or minus-strand genome (6–12), many screening assays utilize synthetic homopolymeric templates/primers (5,7,13–22). Specific inhibitors of the HCV polymerase recently identified from such screening campaigns can be broadly classified as either non-nucleoside compounds that may affect an initiation step (23–26) or nucleoside analogs that inhibit polymerase elongation (27,28).

The recombinant HCV NS5B polymerase enzymes commonly used in *in vitro* assays are produced and isolated from either *Escherichia coli* or baculovirus-infected insect cells. Expression of the full-length HCV NS5B, either untagged or tagged (such as a hexa-histidine tag or GST tag), results in insoluble protein requiring extraction with detergents, salt and glycerol (5,7,14,15,18–20,22,29). The HCV NS5B protein has a highly conserved C-terminal hydrophobic segment and truncation of this portion in recombinant clones results in the expression of a soluble form of the enzyme that retains *in vitro* activity (16,17,19). In addition to their use in screening campaigns, these soluble forms of the enzyme have been particularly effective in crystallizing the NS5B to reveal an X-ray derived structure similar to other polymerases, but with an encircled active site (30–32). Recent X-ray derived structures of compounds bound to NS5B reveal a variety of potentially distinct inhibitor pockets, many of which localize to the thumb domain (26,33,34).

The activity of NS5B in *in vitro* polymerase reactions with homopolymeric RNA requires interaction with multiple substrates that include a template/primer and ribonucleotide triphosphate. Steady-state kinetic parameters, such as K_m , can be determined for both the ribonucleotide triphosphate (6,14,17,19,20) and the template/primer (17) substrates. The C-terminally truncated, soluble forms of recombinant HCV polymerases have high affinity (a low K_m value in the

*To whom correspondence should be addressed. Tel: +1 450 682 4640; Fax: +1 450 682 4642; Email: gkukolj@lav.boehringer-ingelheim.com

nanomolar range) for homopolymeric template/primer (17). In order to identify compounds that interfere with template/primer binding, compound concentration and/or affinity must surpass the K_m of the substrate in a competitive inhibition assay. In this report we describe a modified HCV polymerase enzyme, distinct from the commonly used truncated forms, that has lower affinity for primer/template substrate. The characteristics of this modified enzyme in an *in vitro* assay facilitated the identification of a class of benzimidazole-5-carboxamide-based compounds that specifically inhibit HCV NS5B productive RNA binding. The compounds' mode of inhibition is confirmed by steady state kinetics and order of addition experiments wherein we demonstrate that they interfere with the initiation process of RNA replication, rather than processive elongation. This distinct class of inhibitors would not only complement inhibitors of other HCV targets, but may also complement nucleoside analogs and other non-nucleoside NS5B inhibitors to expand the repository of potential HCV therapeutics.

MATERIALS AND METHODS

Production and purification of the different polymerase constructs

HT-NS5B. Briefly, the entire HCV NS5B region was amplified by PCR from a full-length HCV 1b genotype clone (HCV1b-40) and cloned into a pFastBacHTa vector (Invitrogen). The resulting vector, encoding the NS5B sequence with a hexahistidine N-terminal fusion under the control of the polyhedrin promoter, was used as a donor to introduce the NS5B into a recombinant baculovirus. *Sf21* insect cells were grown to a density of 1×10^6 cells/ml and infected with BacHTaNS5B16-3 recombinant baculovirus expressing the HCV polymerase. Cells expressing the HCV polymerase were harvested by centrifugation, washed with PBS and resuspended in lysis buffer (25 mM Tris-HCl pH 7.5, 2 mM 2-mercaptoethanol, 5 mM MgCl₂, 1 mM EDTA, 500 mM NaCl, 50% glycerol, 2% Triton X-100, protease inhibitor cocktail). Cells were homogenized and treated with bovine pancreas DNaseI. The homogenate was sonicated and centrifuged at 105 000 *g* for 45 min. Supernatants were pooled and subjected to metal affinity chromatography using a Qiagen Ni-NTA column. The polymerase was eluted with an 85–400 mM imidazole linear gradient. The material was then applied onto a DEAE-Sephacel column. The flow-through and washes were pooled for subsequent purification using heparin-Sephacel chromatography. Bound protein was eluted using a linear gradient of 200–1000 mM NaCl. In order to maintain solubility of the HT-NS5B, all of the chromatography buffers contained 0.05% Triton X-100 and 0.1% NP-40. Fractions enriched (>90% purity) in NS5B (according to Coomassie-stained SDS-PAGE) were pooled. The protein concentration of this pool was determined by the micro-Bradford method (Bio-Rad) using BSA as standard. This pool was aliquoted and stored at -80°C (in 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 800 mM NaCl, 20% glycerol, 0.05% Triton X-100 and 0.1% NP-40) without any significant loss of activity during 3 years of storage. The yield of purified protein was 1 mg/l of cultured *Sf21* cells and it was used in the standard HCV RdRp assay at a final concentration of 10 nM.

HT-NS5B Δ 21 and NS5B Δ 21-HT. The recombinant HCV NS5B polymerase can be produced in soluble form by expression of a variant that lacks the C-terminal 21 amino acids (16,17,19). We expressed this NS5B Δ 21 with an N-terminal hexahistidine (termed HT-NS5B Δ 21) and with a C-terminal hexahistidine tag (termed NS5B Δ 21-HT). Expression of these genes from pET vectors in *E.coli* strain JM109 (DE3) was induced with 0.4 mM IPTG for 3 h at 24°C . Cells were harvested and lysed in a microfluidizer. The lysate, after centrifugation, was purified according to the HT-NS5B protocol: Ni-NTA, DEAE-Sephacel and heparin-Sephacel chromatography, in buffers lacking detergent. The proteins were thereafter concentrated on a Resource S column, and applied to a Superdex 200 column where peak fractions containing highly pure (>98%) and monomeric histidine tag NS5B Δ 21 were pooled and stored at -80°C in (20 mM Tris pH 7.5, 10% glycerol, 5 mM DTT and 300 mM NaCl) for more than 3 years without significant loss in activity. The yield of both enzymes was ~ 2 mg/l of *E.coli* culture and they were used in the standard HCV RdRp assay at final concentration of 2–3 nM.

NS5B Δ 57-HT. This polymerase variant that lacks the C-terminal 57 amino acids was expressed with a C-terminal hexahistidine tag (termed NS5B Δ 57-HT). Expression of this NS5B gene construct from a pET vector in *E.coli* strain JM109 (DE3) was induced with 0.4 mM IPTG for 3 h at 24°C . Cells were harvested and lysed in a microfluidizer. The lysate, after centrifugation, was first purified according to the HT-NS5B Δ 21 protocol: Ni-NTA, DEAE-Sephacel and heparin-Sephacel chromatography and then loaded onto a Q-Sephacel column. The flow-through from this last step was collected and concentrated with centrifugal concentrators. Analytical gel filtration revealed highly pure (>95%), monomeric protein with an overall yield of 1 mg/l of culture. This enzyme was used at a concentration of 2 nM in the HCV RdRp assay.

NS5B. The mature, full-length NS5B polymerase was produced as a tagged precursor (HTaA5B) in *Sf21* insect cells infected with a recombinant baculovirus encoding the HCV NS5B, with an N-terminal hexahistidine tag linked by the C-terminal portion of NS5A (BacHTaA5B) to the NS5B segment. The N-terminal histidine tag was removed by processing with the HCV NS3 protease and a NS4A peptide cofactor, resulting in a mature, untagged form of the HCV RdRp (12). Briefly, the precursor histidine-tagged polymerase was purified according to the standard protocol in buffers containing 0.15% *n*-dodecyl- β -D-maltoside and then cleaved with the NS3/4A protease using a 1:50:1.25 molar ratio of NS3 protease:NS4A cofactor:HTaA5B. The reaction was performed at room temperature for 45 min followed by incubation at 4°C for 5 h. The mature NS5B RdRp was then isolated by removing the cleaved histidine-tag leader and any uncleaved HTaA5B protein with metal affinity resin. Unbound mature NS5B was subjected to heparin chromatography as described above in order to separate the NS3 protease from NS5B RdRp. The NS5B fraction obtained from heparin chromatography underwent a final step of gel filtration from which mature monomeric NS5B enzyme was recovered. The enzyme (final yield ~ 0.1 mg/l of *Sf21* cells) was stored at

–80°C for more than 3 years without loss in activity, and was used at 1–2 nM in the standard HCV RdRp assay.

Poliovirus RNA polymerase. The enzyme was produced in *E. coli* BL21(DE3) harboring a T7 expression plasmid encoding the poliovirus RNA polymerase (a kind gift from K. Kirkegaard, Stanford University). The protein was purified by precipitation with ammonium sulfate at 40% of saturation, followed by S-Sepharose and Q-Sepharose chromatographies. After a final gel-filtration step, the preparation was homogeneous as revealed by protein gel-electrophoresis. The enzyme was stored at –20°C in storage buffer [25 mM Tris–HCl pH 7.5, 300 mM NaCl, 5 mM DTT, 1 mM EDTA, 0.1% (v/v) Igepal CA-630 and 30% (v/v) glycerol].

Calf thymus RNA polymerase II. RNA polymerase II was purified from fresh calf thymus tissue extracts by polyethyleneimine and ammonium sulfate precipitation followed by standard chromatography techniques according to the method of Kim and Dahmus (35). The enzyme stock solution was stored in 50 mM Tris–HCl pH 7.9, 10 mM EDTA, 10 mM EGTA, 400 mM ammonium sulfate, 0.25 mM DTT and 20% glycerol (v/v).

HCV RNA-dependent RNA polymerase assay

The standard HCV RdRp assay was performed in 96-well plates using 2–10 nM of enzyme (as indicated), 0.5 μ Ci of [³H]UTP, 1 μ M UTP, 250 nM 5'-biotinylated oligo(rU₁₂), 10 μ g/ml poly(rA) in 20 mM Tris–HCl pH 7.5, 5 mM MgCl₂, 1 mM EDTA, 1 mM DTT, 0.2 U/ μ l of RNasin, 5% DMSO, 3% glycerol, 30 mM NaCl, 0.33% dodecyl- β -D-maltoside, 0.01% IGEPAL. The 60 μ l reaction was terminated after 90 min at 22°C by the addition of 20 μ l of stop solution (150 μ g/ml of tRNA in 0.5 M EDTA) and 30 μ l of streptavidin-coated beads [8 mg/ml in 20 mM Tris–HCl pH 7.5, 25 mM KCl, 0.025% (w/v) sodium azide] for a Scintillation Proximity Assay (Perkin Elmer-Amersham). After 30 min at room temperature, 75 μ l of 5 M cesium chloride were added to the wells and the plate was left at room temperature for 1 h before quantifying the radioactive UMP incorporated onto the biotinylated primer by counting for 60 s on a TopCount (Packard).

Poliovirus RNA-dependent RNA polymerase assay

The protocol was identical to the standard HCV RdRp assay, except that the poliovirus RNA polymerase enzyme was used at a final concentration of 20 nM.

Calf thymus RNA polymerase II assay

This assay measures the incorporation of ³³P-UTP during the transcription of the calf thymus DNA by calf thymus RNA polymerase II (35). The assay was performed in 96-well plates using 4 μ g/ml of calf thymus polymerase enzyme, 100 μ g/ml of calf thymus DNA, 250 μ M each of GTP, CTP and ATP, 1 μ M UTP, 2.5 μ Ci/ml of ³³P-UTP in 80 mM Tris–HCl pH 7.9, 80 mM (NH₄)₂SO₄, 1 mM MnSO₄, 2.25 mM MgCl₂, 0.035 mM EDTA, 1 mM DTT, 0.1 U/ μ l of RNasin, 5% DMSO, 6% glycerol. The 60 μ l reaction was terminated after 120 min at 37°C by TCA precipitation and filtration of the RNA transcripts. The retained RNA products were quantified using a TopCount instrument (Packard).

Enzymatic characterization of the different NS5B polymerase constructs

In order to determine the K_m for the template/primer [poly(rA)/oligo(U₁₂)], a saturating amount of UTP (25–50 μ M; 0.08–0.2 μ Ci/ μ l of ³³P-UTP) was used in the assay in the presence of increasing concentrations of poly(rA)/oligo(U₁₂) (with a template/primer ratio maintained at 10). Similarly, for the determination of the K_m for UTP, a saturating amount of template/primer was used (250–1000 nM, depending on the enzyme) in the presence of increasing concentrations of UTP (0.1–200 μ M; 0.03–0.07 μ Ci/ μ l). Conditions of other reagents in the reaction were identical to the biochemical RdRp assay. Depending on the specific activity of the different enzyme constructs, their concentration used in enzymatic characterization varied between 0.5 and 10 nM. Aliquots of 8 μ l were removed at specific times, spotted onto DE81 filter paper discs, washed in phosphate buffer to remove unincorporated UTP, rinsed and counted to quantify the amount of UMP incorporated into bound product RNA. Velocities of the reactions with each concentration of poly(rA)/oligo(U₁₂) (or UTP) were determined. The data were processed and analyzed with kinetics software (GraFit Erithacus Software) to obtain the K_m for both template/primer and UTP.

Determination of the K_i and mode of compound 1 inhibition

Two series of reactions were performed to determine the mode of inhibition of compound 1. In the first series of experiments, velocities of reactions were determined at different template/primer and inhibitor concentrations. The concentration of template/primer ranged from 25 to 1000 nM with a fixed concentration of UTP at 25 μ M (reactions contained up to 0.2 μ Ci/ μ l of ³³P-UTP). The concentration of enzyme used in the assay was 5 nM and the concentration of inhibitor ranged from 0.25- to 4-fold the IC₅₀ value. For each of these reactions, velocities were determined by withdrawing aliquots at defined times, transferring them on DE81 filter discs and measuring bound radioactivity on a DE81 filter as described above.

In the second series of experiments, velocities were monitored at different UTP and inhibitor concentrations. The template/primer concentration was fixed (at 250 nM), the UTP concentration ranged from 0.25 to 50 or 100 μ M (with 0.02–0.2 μ Ci/ μ l of ³³P-UTP), and the concentration of the inhibitor ranged from 0.25- to 4-fold the IC₅₀ value. The NS5B polymerase used in this series of experiments ranged in concentration from 10 to 25 nM. Kinetic results were then plotted according to the method of Cornish-Bowden (36), allowing for determination of the mode of inhibition as well as the constant(s) of inhibition (K_i as the competitive part of inhibition and K_i' as the uncompetitive part of inhibition).

Determination of IC₅₀ values with different NS5B polymerase constructs

The IC₅₀ values for a series of compounds were determined with the five different constructs of the NS5B polymerase. All of the RdRp assays were performed in SPA format precisely as described in the standard protocol with the indicated enzyme construct. From serial dilution of the test compound, the percentage inhibition was plotted against the compound

concentration and a non-linear curve was fitted (Hill model) to the percentage inhibition–concentration data. The calculated percentage inhibition values were then used to determine the median inhibitory concentration IC_{50} , slope factor (n) and maximum inhibition (I_{max}) by the non-linear regression procedure of SAS (Statistical Software System, SAS Institute Inc., Cary, NC, USA) using the following equation:

$$\% \text{ inhibition} = \frac{I_{max} \times [\text{inhibitor}]^n}{[\text{inhibitor}]^n + IC_{50}^n}$$

The specificity of compounds as inhibitors of HCV polymerase was evaluated through a poliovirus RNA polymerase counter screen in an SPA assay format that utilizes the identical substrates and conditions previously optimized for the standard HCV polymerase assay. The compounds were further profiled for inhibition of DNA-dependent RNA polymerase II (purified from calf thymus).

Modified order of addition single cycle HCV NS5B RdRp assay

A single-cycle of HCV RdRp initiation and elongation was assayed by modifying the order of addition, followed by the addition of heparin. The template/primer (250 nM) was first pre-incubated with the enzyme (10 nM) for 20 min at room temperature, after which heparin (final concentration of 1.2 $\mu\text{g/ml}$), the inhibitor and 0.5 μM [^3H]UTP (0.013 $\mu\text{Ci}/\mu\text{l}$) were successively added to the reaction. Following a period of 90 min incubation at room temperature, stop solution and streptavidin beads were added as in the standard HCV RdRp assay. Radioactivity in the wells was quantified on a TopCount and the inhibitory potential of a series of compounds was evaluated.

HCV NS5B *de novo* initiation reaction

The primer-independent *de novo* initiation activity of the mature full-length NS5B construct was reconstituted with substrates and conditions as previously described (12). Products were analyzed on denaturing PAGE gels or quantified on DE81 filters after washing and removal of unincorporated nucleotides.

Compound synthesis

Compounds 1, 1a–1e inclusively, and compound 2 were prepared using published procedures (37–39). All compounds had $^1\text{H-NMR}$ and mass spectral data consistent with their assigned structures. Compound formulas are as indicated: compound 1, $\text{C}_{37}\text{H}_{47}\text{N}_5\text{O}_5$; compound 1a, $\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_6$; compound 1b, $\text{C}_{27}\text{H}_{27}\text{N}_3\text{O}_5$; compound 1c, $\text{C}_{29}\text{H}_{29}\text{N}_3\text{O}_7$; compound 1d, $\text{C}_{29}\text{H}_{28}\text{N}_4\text{O}_5$; compound 1e, $\text{C}_{31}\text{H}_{30}\text{N}_4\text{O}_7$; and compound 2, $\text{C}_{16}\text{H}_{10}\text{Cl}_3\text{NO}_6\text{S}$.

RESULTS

Enzymatic characterization of the different NS5B polymerase constructs

We examined a panel of purified HCV NS5B polymerase constructs that had different terminal modifications and placements of affinity tags, as depicted in Figure 1. Kinetic parameters, such as the K_m , were determined for both the

template/primer and the UTP. Determination of a K_m for template/primer RNA with the HT-NS5B construct is exemplified in Figure 2, where the reaction velocity, as measured by UMP incorporation onto the extending RNA primer, is plotted against a range of template/primer concentrations and performed with saturating (50 μM) amounts of UTP. Similarly, the K_m for UTP was determined for this and other NS5B constructs by varying the UTP concentration in the presence of saturating amounts of template/primer RNA. The K_m for UTP among the different NS5B constructs ranged from 1.8 to 12 μM (Table 1). The K_m for the template/primer varied <3-fold among four of the five NS5B constructs (HT-NS5BA21, NS5BA57-HT, NS5BA21-HT and NS5B) and ranged from 0.025 to 0.058 μM (Table 1); however, the HT-NS5B (full-length NS5B with an N-terminal hexa-histidine tag) displayed a significantly lower affinity ($K_m = 0.21 \pm 0.02 \mu\text{M}$).

Identification and characterization of a specific HT-NS5B inhibitor

Screening of broad chemical compound collections for specific inhibitors of HCV polymerase activity was performed with the HT-NS5B construct. We postulated that the lower affinity of this construct for template/primer RNA may render it more susceptible to competitive inhibitors of template/primer binding. As a result, compound 1 (Fig. 3) a member from a benzimidazole-5-carboxamide library, was identified as a specific inhibitor of the HT-NS5B polymerase with an IC_{50} of 13.6 μM (Table 2). In order to determine its mode of inhibition, two series of kinetic experiments quantifying polymerase velocity were performed with HT-NS5B and compound 1. First, the concentrations of UTP and inhibitor were varied in the presence of a constant amount of template/primer (Fig. 4), and secondly, the concentrations of template/primer and inhibitor were varied in the presence of a constant amount of UTP (Fig. 5). Compound 1 demonstrated non-competitive inhibition of UTP incorporation (K_i competitive = K_i uncompetitive = 20 μM), as both Dixon and Cornish-Bowden plots intercept on the x -axis (Fig. 4). In contrast, compound 1 displayed a mixed mode of inhibition towards the template/primer, with a significant competitive component (K_i competitive = 26 μM and K_i uncompetitive = 105 μM); the mode of inhibition is characterized by the Dixon and Cornish-Bowden plots of the data in Figure 5, where the location of the intercepts on the two plots are above and below the x -axis, respectively.

Compound IC_{50} s with different HCV polymerase constructs

The IC_{50} values for a series of five compounds related to compound 1 (Fig. 3) were determined with the five different constructs of NS5B polymerase as illustrated in Table 2. HCV RdRp assays that were performed with the HT-NS5B enzyme used in our screening campaign and which displayed the lowest affinity for template/primer (Table 1), provided the lowest relative IC_{50} values for this class of compounds. Compounds (38) with modified amino acid derived R1 groups (1a–1e, Fig. 3) had improved potencies, such that compound 1e with a 5-indole-oxyacetic acid displayed an IC_{50} of 19 nM. The IC_{50} values determined under the same conditions with the other NS5B constructs were significantly higher for all six




Figure 1. Schematic representation of the five NS5B constructs cloned and produced from an HCV 1b genome. The full-length N-terminal hexa-histidine HT-NS5B fusion and the full-length untagged NS5B were produced and purified from recombinant baculovirus-infected insect cells. Three NS5B clones were expressed in *E.coli*: the HT-NS5B Δ 21, harboring an N-terminal hexa-histidine tag and C-terminal 21 amino acid deletion; the NS5B Δ 57-HT encoding a C-terminal 57 amino acid deletion fused to hexa-histidine tag; and the NS5B Δ 21-HT encoding a C-terminal 21 amino acid deletion fused to a hexa-histidine tag.




Figure 2. Determination of the K_m for the template/primer poly(rA)/oligo(U) with the HT-NS5B enzyme. Reaction velocity, measured as ^{33}P -UMP incorporated into product, was plotted versus increasing concentrations (ranging from 25 to 2000 nM) of poly(rA)/oligo(U) substrate. The curve was fit with GraFit Software (GraFit Erithacus Software) to obtain a K_m of 214 nM.

of the benzimidazole-5-carboxamide derivatives (Table 2), yet the ranking of the compound potencies among all of the constructs was identical; i.e. IC_{50} for compound 1e < compound 1d < compound 1c < compound 1b ~ compound 1a < compound 1. Hence, the structure-activity relationship for this class of HCV polymerase inhibitor could be followed with any one of these enzyme constructs.

Notably, the fold increase in IC_{50} values of the benzimidazole-5-carboxamide class of compounds in assays performed with the different HCV polymerase constructs was inversely proportional to the K_m of the respective NS5B constructs. For example, assays using the construct with the

highest K_m for template/primer (HT-NS5B) provided the lowest relative IC_{50} values, whereas constructs with a low K_m (NS5B) provided the highest relative IC_{50} values. This shift in IC_{50} values for a common inhibitor among the different HCV polymerase constructs was unique to the benzimidazole-5-carboxamide series, as compound 2, an unrelated pyrophosphate product-mimic inhibitor of HCV polymerase (39) displayed less than a 4-fold difference in potency in polymerase assays with the different NS5B enzyme constructs.

Benzimidazole-5-carboxamide compounds do not inhibit HCV polymerase initiation or elongation from pre-bound RNA

The standard HCV RdRp assay is a continuous polymerization reaction with multiple cycles of initiation and elongation events. The assay can be modified by pre-incubating the enzyme with the RNA substrate and delaying the addition of NTP and test compound after polyanionic heparin is added to quench any dissociated polymerase; such a modification quantifies only a single cycle of initiation and elongation from pre-bound template/primer substrate (19,23). The potential of compound 1 and its analogs to inhibit initiation or subsequent elongation from pre-formed HCV polymerase-RNA complexes was evaluated in this assay format. Sequential, ordered addition of heparin, inhibitor and UTP to the pre-incubated NS5B-RNA complex indicated that the benzimidazole-5-carboxamide class of compounds did not inhibit HCV polymerase initiation or elongation from pre-bound RNA complexes (Table 3). Furthermore, simply pre-incubating the NS5B with RNA substrate followed by the addition of compound and UTP (without heparin) also alleviated

Table 1. Kinetic analyses of different constructs of NS5B polymerases

	HT-NS5B	HT-NS5B Δ 21	NS5B Δ 57-HT	NS5B Δ 21-HT	NS5B
K_m (P/T) ^a (μM)	0.21 ± 0.02	0.058 ± 0.01	0.034 ± 0.01	0.025 ± 0.005	0.025 ± 0.004
K_m (UTP) ^b (μM)	6.2 ± 0.63	12 ± 2.3	1.8 ± 0.3	5.2 ± 1.3	3.3 ± 1.3

^aValues are the mean of two ($n = 2$) determinations except for HT-NS5B ($n = 3$).

^bValues are the mean of two ($n = 2$) determinations except for NS5B Δ 21-HT ($n = 3$).




Figure 3. Structure of the compounds used in the study. Compound 1 and its analogs (1a–1e, inclusive) are a series of benzimidazole-5-carboxamide derivatives identified as specific inhibitors of HCV polymerase. Compound 2 is an unrelated di-ketoacid inhibitor of the HCV polymerase (39,49).

Table 2. IC₅₀ values^a (μM) for a series of compounds with different polymerase constructs

Compound	HT-NS5B	HT-NS5BA21	NS5BA57-HT	NS5BA21-HT	NS5B	Poliovirus polymerase	CT RNA Pol II
1	13.6	29	>500	>500	>500	>500	>500
1a	0.59	2.7	13.2	20	20	>120	>200
1b	0.49	3.0	23	19	28	>100	>200
1c	0.14	1.2	4.5	7.3	11.2	100	>200
1d	0.054	0.44	2.2	3.0	5.7	79	>200
1e	0.019	0.16	0.73	1.1	1.6	100	>200
2	0.95	0.63	0.68	0.3	0.25	n.d.	n.d.

^aValues are the mean of three or four determinations for each compound with coefficient of variation below 25%.

inhibition by compound 1 (data not shown). In contrast, compound 2, the di-ketoacid derivative that specifically interferes with NTP incorporation by HCV polymerase (39), displayed similar potencies in both the standard HCV RdRp and single cycle initiation and elongation assays. The mechanism of action of the benzimidazole-5-carboxamide compounds inferred from these order of addition experiments, is consistent with the kinetic experiments that this class of compounds interferes with an early step in the polymerase reaction such as productive RNA binding.

Inhibition of HCV polymerase *de novo* initiation

Though these compounds inhibited reactions with primer-dependent homopolymeric substrates, the HCV polymerase is

proposed to commence complementary strand RNA synthesis on genomic template RNA in a primer independent *de novo* initiation reaction. Using a previously characterized HCV plus-strand 3'-UTR template RNA substrate (12), we examined the effect of compound 1e on product formation in a *de novo* initiation reaction catalyzed by the full-length, untagged NS5B construct. The major products of *de novo* initiation in this reaction are the 246 and 227 nt products (12,40,41) depicted in Figure 6A. Reactions reconstituted with increasing amounts of compound 1e inhibited formation of the product of *de novo* RNA synthesis with an IC₅₀ of 2.3 μM (Fig. 6B). The IC₅₀s of compound 1e in this assay and the standard HCV polymerase assay run with the mature NS5B (IC₅₀ = 1.6 μM, Table 2) are similar and reflect the similar K_m of the NS5B




Figure 4. K_i determination of compound 1 and mode of inhibition with regard to UTP substrate. Reaction velocities were measured at different UTP concentrations (0.25, 0.50, 1.0, 2.5, 5.0, 10, 25 and 50 μM , plotted as open diamonds, closed diamonds, open circles, closed circles, open squares, closed squares, open triangles and closed triangles, respectively) with the template/primer fixed at 0.25 μM . Dixon and Cornish-Bowden plots of the reciprocal velocity with compound 1 at concentrations of 0, 2.5, 5, 10, 20 and 40 μM . Both plots display intersects on the x -axis and reflect non-competitive inhibition with regard to UTP (K_i competitive = K_i uncompetitive = 20 μM).




Figure 5. K_i determination of compound 1 and mode of inhibition with regard to template/primer substrate. Reaction velocities were measured at different template/primer concentrations (0.05, 0.075, 0.15 and 0.2 μM plotted as closed diamonds, open diamonds, closed squares and open squares, respectively) with the UTP fixed at 25 μM . Dixon and Cornish-Bowden plots of the reciprocal velocity with compound 1 at concentrations of 0, 15, 30 and 60 μM . Compound 1 displayed a mixed mode of inhibition towards template/primer, with a major competitive component (K_i competitive = 26 μM ; intercept on the Dixon plot above the x -axis) and a minor uncompetitive component (K_i uncompetitive = 105 μM ; intercept extrapolated from the Cornish-Bowden plot below the x -axis).

Table 3. IC_{50} values^a (μM) for a series of compounds in a single-cycle HT-NS5B elongation assay

Compound	HT-NS5B (elongation)
1	>250
1a	>100
1b	>100
1c	>100
1d	>100
1e	>100
2	1.52 \pm 0.37

^aValues are the mean of two or three determinations.

construct both for the homopolymeric substrate RNA (Table 1) and the 3'-UTR template (data not shown).

DISCUSSION

The viral nucleic acid polymerase inhibitors constitute the majority of currently approved antiviral drugs (42) and portray

polymerases as prime therapeutic targets for emerging viral infections. Consequently, the NS5B enzyme has become a primary target in the search for novel inhibitors of HCV replication and a variety of *in vitro* assays for NS5B polymerase activity have been developed for such a purpose. In this study, we have characterized steady-state kinetic parameters of both nucleotide triphosphate and template/primer RNA substrates for NS5B constructs encoding a variety of common truncations and affinity tag fusions. The K_m for UTP varied from 1.8 to 12 μM , depending on the polymerase construct, which is a range consistent with previously published values using the full-length or truncated enzyme, produced in insect cells or in *E.coli*, with or without a histidine tag (14,19,20,43). The K_m for the template/primer with our panel of NS5B constructs ranged from 25 to 214 nM. Most of the NS5B constructs have low nanomolar affinity for the homopolymeric substrate, which is in agreement with a report by Ferrari *et al.* (17) of a K_m for a template/primer of 30 nM with the common C-terminal $\Delta 21$ truncation of the HCV polymerase using a poly(C)/oligo(G) substrate. However, the full-length N-terminally tagged polymerase, HT-NS5B, displayed a markedly lower affinity for the RNA substrate and offered a potential advantage in a screen for inhibitors.

A competitive inhibitor of enzyme activity must have affinities comparable with (or greater than) the substrate. For viral polymerases such as the NS5B, the triphosphate form of nucleoside analogs represent classic competitive inhibitors of NTP incorporation whose IC_{50} value is directly proportional to the substrate concentration and inversely proportional to the K_m value (44). The widely established *in vitro* NS5B assays that use low micromolar NTP concentration are therefore sensitive to a broad range of inhibition by nucleotide mimics (27,28). Towards this end, and before the development of cell-based HCV RNA replication assays (45), nucleoside analogs (reviewed in 46) were originally identified by screening their triphosphorylated derivatives in *in vitro* assays with NTP concentrations at or below the K_m value. The HCV RdRp assay that we used in a broad screening campaign, was optimized to detect competitive inhibitors of both NTP and template/primer RNA substrates, by using the HT-NS5B enzyme construct under conditions that reconstituted *in vitro* activity with the concentration of both substrates at or below their K_m value. Hence, the low affinity that the HT-NS5B construct had for the poly(rA)/oligo(U) homopolymeric substrate, supported the design of a screening assay with a relatively high RNA substrate concentration [i.e. 250 nM of oligo(U) primer and a 10-fold molar-nucleotide excess of poly(rA) template] that was within the K_m value. In contrast, the optimized standard assay that used the other NS5B polymerases constructs (Fig. 1 and Table 1) had template/primer concentrations that were 4–10-fold higher than the K_m values, and accounted (44) for the significantly higher IC_{50} (Table 2) values displayed by inhibitors of RNA binding.

Using the HT-NS5B-based assay, compound 1 was identified in a broad screen of the Boehringer Ingelheim proprietary compound collection, as a non-nucleoside inhibitor (IC_{50} = 13.6 μM) originating from a benzimidazole-5-carboxamide combinatorial chemistry library. Compound 1 and derivatives displayed specific and reversible inhibition of the HCV polymerase and failed to inhibit poliovirus RdRp and a




Figure 6. Compound 1e inhibits *de novo* initiation by HCV NS5B polymerase. (A) Lane 1 shows the major products of NS5B *de novo* initiation using a short 3'-UTR plus strand template as 246 and 227 nt RNAs (12). Lanes 2–8 display the products of NS5B *de novo* initiation in the presence of 1.6, 3.13, 6.25, 12.5, 25, 50 and 100 μM of compound 1e. (B) Dose–response curve displaying the IC_{50} of compound 1e in a NS5B *de novo* initiation assay. The points on the curve represent an average (with error bars) of three independent determinations.

DNA-dependent RNA polymerase II activity (Table 2). Moreover, inhibition by compound 1 was only observed with the N-terminally tagged NS5B and not with the other NS5B constructs (Table 2). In order to determine the mode of HT-NS5B inhibition by compound 1, reaction velocities were measured over a range of template/primer, NTP and inhibitor concentrations. Reciprocal plots of velocity (36) revealed a mixed mode of inhibition for compound 1 towards the template/primer with a significant competitive component, and non-competitive inhibition with UTP. HIV-1 RT non-nucleoside inhibitors, such as nevirapine, are non-competitive inhibitors with UTP, but in contrast to compound 1, are also non-competitive with template/primer substrate (47,48).

At least three other chemically distinct series of non-nucleoside inhibitors of the HCV polymerase have been recently reported: (i) a benzothiadiazine series (23); (ii) phenylalanine derivatives (26); and (iii) a cyclopentyl dihydropyran-2-one inhibitor (34). The latter two bind to a distinct allosteric site in the thumb domain distally removed from the central active site (26,34). Representative compounds of these three non-nucleoside inhibitors are all non-competitive inhibitors of NTP incorporation, and their mode of action with respect to template/primer are incompletely characterized. Though the benzothiadiazine series binding site has not been reported, an uncompetitive mode of inhibition with respect to template/primer RNA is suggested from the unproductive enzyme–substrate–inhibitor complex formed by the NS5B, template RNA substrate and benzothiadiazine (24). The benzimidazole-5-carboxamide compounds' mode of action clearly differed from the benzothiadiazine inhibitor. The latter series effectively inhibits NS5B initiation from pre-bound RNA (24), whereas the benzimidazole-5-carboxamides did not inhibit the activity of pre-bound NS5B–RNA complexes (Table 3). Besides the modified order of addition experiments, the steady-state kinetic data revealed that benzimidazole-5-carboxamides interfere with RNA binding. Combination experiments with a benzothiadiazine and a benzimidazole-5-carboxamide showed that the compounds were not mutually exclusive and probably bind to separate sites (McKercher,G., unpublished observations).

The distinct mode of inhibition by the benzimidazole-5-carboxamide class of compounds was also manifested by the dramatically different IC_{50} values obtained with the modified enzyme constructs. Though the potency ranking of the related derivatives was identical between the different constructs, increases in IC_{50} s ranged from ~6-fold with the HT-NS5B Δ 21 enzyme up to 72-fold with the untagged full-length NS5B. Notably, the magnitude of the increases in IC_{50} s was inversely proportional to the K_m of the respective enzyme constructs for template/primer RNA [K_m (P/T)] and is fully consistent with inhibitors that are predicted to interfere with RNA binding. As a result, compound screening with the higher affinity NS5B constructs would have greatly diminished the assay's sensitivity to inhibitors of template/primer binding and failed to detect weak to moderately potent compounds of this class. Conversely, compounds that function by an alternative mechanism, such as the di-ketoacid class (compound 2) that mimic pyrophosphate product (39,49), would be unaffected by an enzyme's K_m (P/T) property and thereby displayed similar potencies against this group of modified enzymes.

The position of the hexa-histidine tag on the NS5B affected the enzyme's affinity for RNA substrate and, in turn, the IC_{50} of the benzimidazole-5-carboxamides. Lower IC_{50} values were effectively obtained with the two polymerase constructs that had the hexa-histidine tag at the N-terminus. The tagged, full-length polymerase (HT-NS5B) had the lowest affinity for RNA substrate and we ensured that this was not an artifact of reduced enzyme solubility by extracting, purifying and maintaining the protein (without any significant loss in activity) in detergent. The truncated, soluble polymerase that can be purified without detergent also had slightly lower affinity for RNA substrate, relative to the untagged NS5B, only when the tag was positioned at the N-terminus (HT-NS5B Δ 21); notably, inhibition by compound 1 was also detected with this construct.

The optimized compound 1e inhibited the primer-independent *de novo* initiation activity and demonstrated efficacy and similar potencies in assays with homopolymeric or heteropolymeric RNA template using the untagged, full-length NS5B. Compound 1e was also tested for inhibition of

HCV RNA replication in a cell-based replicon assay (45,50) and demonstrated marginal potency (i.e. 63% inhibition at 50 μ M; data not shown). Current medicinal chemistry efforts are directed towards improving the cell permeability of these compounds (Beaulieu, P., manuscript in preparation).

A uniquely modified NS5B enzyme construct facilitated the identification of the benzimidazole-5-carboxamide series of compounds as a novel class of HCV RNA polymerase inhibitors with a mode of action distinct from any previously characterized viral polymerase inhibitors. Studies are ongoing to improve their potency and assess their benefit as a potential therapy in the treatment of chronic HCV infection.

ACKNOWLEDGEMENTS

We thank Samuel April, Dr Professor Volkhard Austel, Yves Bousquet, Michael Bös, James Gillard, Gulrez Fazal, Jean Gauthier, Sylvie Goulet and Marc-André Poupart for the medicinal chemistry contributions. We thank Carol Homon, Paul Kaplita, Gustavo Rodriguez and George Rogers for high-throughput screening. We are grateful for the discussions with Pierre Bonneau, Michael Cordingley and Peter White throughout the course of this work.

REFERENCES

- Choo, Q.L., Kuo, G., Weiner, A.J., Overby, L.R., Bradley, D.W. and Houghton, M. (1989) Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science*, **244**, 359–362.
- Wasley, A. and Alter, M.J. (2000) Epidemiology of hepatitis C: geographic differences and temporal trends. *Semin. Liver Dis.*, **20**, 1–16.
- Hijikata, M., Kato, N., Ootsuyama, Y., Nakagawa, M. and Shimotohno, K. (1991) Gene mapping of the putative structural region of the hepatitis C virus genome by *in vitro* processing analysis. *Proc. Natl Acad. Sci. USA*, **88**, 5547–5551.
- Grakoui, A., Wychowski, C., Lin, C., Feinstone, S.M. and Rice, C.M. (1993) Expression and identification of hepatitis C virus polyprotein cleavage products. *J. Virol.*, **67**, 1385–1395.
- Behrens, S.E., Tomei, L. and De Francesco, R. (1996) Identification and properties of the RNA-dependent RNA polymerase of hepatitis C virus. *EMBO J.*, **15**, 12–22.
- Luo, G., Hamatake, R.K., Mathis, D.M., Racela, J., Rigat, K.L., Lemm, J. and Colonna, R.J. (2000) *De novo* initiation of RNA synthesis by the RNA-dependent RNA polymerase (NS5B) of hepatitis C virus. *J. Virol.*, **74**, 851–863.
- Oh, J.W., Ito, T. and Lai, M.M. (1999) A recombinant hepatitis C virus RNA-dependent RNA polymerase capable of copying the full-length viral RNA. *J. Virol.*, **73**, 7694–7702.
- Zhong, W., Ferrari, E., Lesburg, C.A., Maag, D., Ghosh, S.K., Cameron, C.E., Lau, J.Y. and Hong, Z. (2000) Template/primer requirements and single nucleotide incorporation by hepatitis C virus nonstructural protein 5B polymerase. *J. Virol.*, **74**, 9134–9143.
- Zhong, W., Uss, A.S., Ferrari, E., Lau, J.Y. and Hong, Z. (2000) *De novo* initiation of RNA synthesis by hepatitis C virus nonstructural protein 5B polymerase. *J. Virol.*, **74**, 2017–2022.
- Kao, C.C., Yang, X., Kline, A., Wang, Q.M., Barket, D. and Heinz, B.A. (2000) Template requirements for RNA synthesis by a recombinant hepatitis C virus RNA-dependent RNA polymerase. *J. Virol.*, **74**, 11121–11128.
- Kao, C.C., Singh, P. and Ecker, D.J. (2001) *De novo* initiation of viral RNA-dependent RNA synthesis. *Virology*, **287**, 251–260.
- Pellerin, C., Lefebvre, S., Little, M.J., McKercher, G., Lamarre, D. and Kukolj, G. (2002) Internal initiation sites of *de novo* RNA synthesis within the hepatitis C virus polypyrimidine tract. *Biochem. Biophys. Res. Commun.*, **295**, 682–688.
- Yuan, Z.H., Kumar, U., Thomas, H.C., Wen, Y.M. and Monjardino, J. (1997) Expression, purification and partial characterization of HCV RNA polymerase. *Biochem. Biophys. Res. Commun.*, **232**, 231–235.
- Lohmann, V., Roos, A., Korner, F., Koch, J.O. and Bartenschlager, R. (1998) Biochemical and kinetic analyses of NS5B RNA-dependent RNA polymerase of the hepatitis C virus. *Virology*, **249**, 108–118.
- Lohmann, V., Korner, F., Herian, U. and Bartenschlager, R. (1997) Biochemical properties of hepatitis C virus NS5B RNA-dependent RNA polymerase and identification of amino acid sequence motifs essential for enzymatic activity. *J. Virol.*, **71**, 8416–8428.
- Yamashita, T., Kaneko, S., Shirota, Y., Qin, W., Nomura, T., Kobayashi, K. and Murakami, S. (1998) RNA-dependent RNA polymerase activity of the soluble recombinant hepatitis C virus NS5B protein truncated at the C-terminal region. *J. Biol. Chem.*, **273**, 15479–15486.
- Ferrari, E., Wright-Minogue, J., Fang, J.W., Baroudy, B.M., Lau, J.Y. and Hong, Z. (1999) Characterization of soluble hepatitis C virus RNA-dependent RNA polymerase expressed in *Escherichia coli*. *J. Virol.*, **73**, 1649–1654.
- Ishii, K., Tanaka, Y., Yap, C.C., Aizaki, H., Matsuura, Y. and Miyamura, T. (1999) Expression of hepatitis C virus NS5B protein: characterization of its RNA polymerase activity and RNA binding. *Hepatology*, **29**, 1227–1235.
- Tomei, L., Vitale, R.L., Incitti, I., Serafini, S., Altamura, S., Vitelli, A. and De Francesco, R. (2000) Biochemical characterization of a hepatitis C virus RNA-dependent RNA polymerase mutant lacking the C-terminal hydrophobic sequence. *J. Gen. Virol.*, **81**, 759–767.
- Johnson, R.B., Sun, X.L., Hockman, M.A., Villareal, E.C., Wakulchik, M. and Wang, Q.M. (2000) Specificity and mechanism analysis of hepatitis C virus RNA-dependent RNA polymerase. *Arch. Biochem. Biophys.*, **377**, 129–134.
- Qin, W., Luo, H., Nomura, T., Hayashi, N., Yamashita, T. and Murakami, S. (2002) Oligomeric interaction of hepatitis C virus NS5B is critical for catalytic activity of RNA-dependent RNA polymerase. *J. Biol. Chem.*, **277**, 2132–2137.
- Qin, W., Yamashita, T., Shirota, Y., Lin, Y., Wei, W. and Murakami, S. (2001) Mutational analysis of the structure and functions of hepatitis C virus RNA-dependent RNA polymerase. *Hepatology*, **33**, 728–737.
- Dhanak, D., Duffy, K.J., Johnston, V.K., Lin-Goerke, J., Darcy, M., Shaw, A.N., Gu, B., Silverman, C., Gates, A.T., Nonnemacher, M.R. *et al.* (2002) Identification and biological characterization of heterocyclic inhibitors of the hepatitis C virus RNA-dependent RNA polymerase. *J. Biol. Chem.*, **277**, 38322–38327.
- Gu, B., Johnston, V.K., Gutshall, L.L., Nguyen, T.T., Gontarek, R.R., Darcy, M.G., Tedesco, R., Dhanak, D., Duffy, K.J., Kao, C.C. *et al.* (2003) Arresting initiation of hepatitis C virus RNA synthesis using heterocyclic derivatives. *J. Biol. Chem.*, **278**, 16602–16607.
- Chan, L., Reddy, T.J., Proulx, M., Das, S.K., Pereira, O., Wang, W., Siddiqui, A., Yannopoulos, C.G., Poisson, C., Turcotte, N. *et al.* (2003) Identification of *N,N*-disubstituted phenylalanines as a novel class of inhibitors of hepatitis C NS5B polymerase. *J. Med. Chem.*, **46**, 1283–1285.
- Wang, M., Ng, K.K., Cherney, M.M., Chan, L., Yannopoulos, C.G., Bedard, J., Morin, N., Nguyen-Ba, N., Alaoui-Ismaili, M.H., Bethell, R.C. *et al.* (2003) Non-nucleoside analogue inhibitors bind to an allosteric site on HCV NS5B polymerase. Crystal structures and mechanism of inhibition. *J. Biol. Chem.*, **278**, 9489–9495.
- Carroll, S.S., Tomassini, J.E., Bosserman, M., Getty, K., Stahlhut, M.W., Eldrup, A.B., Bhat, B., Hall, D., Simcoe, A.L., LaFemina, R. *et al.* (2003) Inhibition of hepatitis C virus RNA replication by 2'-modified nucleoside analogs. *J. Biol. Chem.*, **278**, 11979–11984.
- Stuyver, L.J., Whitaker, T., McBrayer, T.R., Hernandez-Santiago, B.I., Lostia, S., Tharnish, P.M., Ramesh, M., Chu, C.K., Jordan, R., Shi, J. *et al.* (2003) Ribonucleoside analogue that blocks replication of bovine diarrhoea and hepatitis C viruses in culture. *Antimicrob. Agents Chemother.*, **47**, 244–254.
- Al, R.H., Xie, Y., Wang, Y. and Hagedorn, C.H. (1998) Expression of recombinant hepatitis C virus non-structural protein 5B in *Escherichia coli*. *Virus Res.*, **53**, 141–149.
- Ago, H., Adachi, T., Yoshida, A., Yamamoto, M., Habuka, N., Yatsunami, K. and Miyano, M. (1999) Crystal structure of the RNA-dependent RNA polymerase of hepatitis C virus. *Struct. Fold. Des.*, **7**, 1417–1426.
- Bressanelli, S., Tomei, L., Roussel, A., Incitti, I., Vitale, R.L., Mathieu, M., De Francesco, R. and Rey, F.A. (1999) Crystal structure of the RNA-dependent RNA polymerase of hepatitis C virus. *Proc. Natl Acad. Sci. USA*, **96**, 13034–13039.

32. Lesburg,C.A., Cable,M.B., Ferrari,E., Hong,Z., Mannarino,A.F. and Weber,P.C. (1999) Crystal structure of the RNA-dependent RNA polymerase from hepatitis C virus reveals a fully encircled active site. *Nature Struct. Biol.*, **6**, 937–943.
33. Bressanelli,S., Tomei,L., Rey,F.A. and De Francesco,R. (2002) Structural analysis of the hepatitis C virus RNA polymerase in complex with ribonucleotides. *J. Virol.*, **76**, 3482–3492.
34. Love,R.A., Parge,H.E., Yu,X., Hickey,M.J., Diehl,W., Gao,J., Wriggers,H., Ekker,A., Wang,L., Thomson,J.A. *et al.* (2003) Crystallographic identification of a noncompetitive inhibitor binding site on the hepatitis C virus NS5B RNA polymerase enzyme. *J. Virol.*, **77**, 7575–7581.
35. Kim,W.Y. and Dahmus,M.E. (1988) Purification of RNA polymerase IIO from calf thymus. *J. Biol. Chem.*, **263**, 18880–18885.
36. Cornish-Bowden,A. (1974) A simple graphical method for determining the inhibition constants of mixed, uncompetitive and non-competitive inhibitors. *Biochem. J.*, **137**, 143–144.
37. Beaulieu,P.L., Haché,B. and von Moos,E. (2003) A practical Oxone®-mediated, high-throughput, solution-phase synthesis of benzimidazoles from 1,2-phenylenediamines and aldehydes and its application to preparative scale synthesis. *Synthesis*, **2003**, 1683–1692.
38. Beaulieu,P.L., Fazal,G., Gillard,J., Kukulj,G. and Austel,V. (2002) *Viral Polymerase Inhibitors*. Patent US 6448281, US 6479508. PCT/WO 02/04425.
39. Altamura,S., Tomei,L., Koch,U., Neuner,P.J.S. and Summa,V. (2000) *Diketoacid-Derivatives as Inhibitors of Polymerases*. Patent US 6492423. PCT/WO 00/06529.
40. Oh,J.W., Sheu,G.T. and Lai,M.M. (2000) Template requirement and initiation site selection by hepatitis C virus polymerase on a minimal viral RNA template. *J. Biol. Chem.*, **275**, 17710–17717.
41. Reigadas,S., Ventura,M., Sarih-Cottin,L., Castroviejo,M., Litvak,S. and Astier-Gin,T. (2001) HCV RNA-dependent RNA polymerase replicates *in vitro* the 3' terminal region of the minus-strand viral RNA more efficiently than the 3' terminal region of the plus RNA. *Eur. J. Biochem.*, **268**, 5857–5867.
42. De Clercq,E. (2001) Antiviral drugs: current state of the art. *J. Clin. Virol.*, **22**, 73–89.
43. Carroll,S.S., Sardana,V., Yang,Z., Jacobs,A.R., Mizenko,C., Hall,D., Hill,L., Zugay-Murphy,J. and Kuo,L.C. (2000) Only a small fraction of purified hepatitis C RNA-dependent RNA polymerase is catalytically competent: implications for viral replication and *in vitro* assays. *Biochemistry*, **39**, 8243–8249.
44. Cheng,Y. and Prusoff,W.H. (1973) Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (IC₅₀) of an enzymatic reaction. *Biochem. Pharmacol.*, **22**, 3099–3108.
45. Lohmann,V., Korner,F., Koch,J., Herian,U., Theilmann,L. and Bartenschlager,R. (1999) Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science*, **285**, 110–113.
46. Beaulieu,P.L. and Llinàs-Brunet,M. (2002) Therapies for hepatitis C infection: targeting the non-structural proteins of HCV. *Curr. Med. Chem. Anti-Infect. Agents*, **1**, 163–176.
47. Merluzzi,V.J., Hargrave,K.D., Labadia,M., Grozinger,K., Skoog,M., Wu,J.C., Shih,C.K., Eckner,K., Hattox,S., Adams,J. *et al.* (1990) Inhibition of HIV-1 replication by a nonnucleoside reverse transcriptase inhibitor. *Science*, **250**, 1411–1413.
48. Kopp,E.B., Miglietta,J.J., Shrutkowski,A.G., Shih,C.K., Grob,P.M. and Skoog,M.T. (1991) Steady state kinetics and inhibition of HIV-1 reverse transcriptase by a non-nucleoside dipyrindodiazepinone, BI-RG-587, using a heteropolymeric template. *Nucleic Acids Res.*, **19**, 3035–3039.
49. De Francesco,R., Tomei,L., Altamura,S., Summa,V. and Migliaccio,G. (2003) Approaching a new era for hepatitis C virus therapy: inhibitors of the NS3-4A serine protease and the NS5B RNA-dependent RNA polymerase. *Antiviral Res.*, **58**, 1–16.
50. Pause,A., Kukulj,G., Bailey,M., Brault,M., Do,F., Halmos,T., Lagace,L., Maurice,R., Marquis,M., McKercher,G. *et al.* (2003) An NS3 serine protease inhibitor abrogates replication of subgenomic hepatitis C virus RNA. *J. Biol. Chem.*, **278**, 20374–20380.