

Increased Glyoxalase-1 Levels in *Fkbp5* Knockout Mice Caused by Glyoxalase-1 Gene Duplication

Lorenz K. Kollmannsberger, Nils C. Gassen, Andrea Bultmann, Jakob Hartmann, Peter Weber, Mathias V. Schmidt, and Theo Rein¹

Max Planck Institute of Psychiatry, 80804 Munich, Germany

ABSTRACT *Fkbp5* is genetically linked to stress-related diseases. *Fkbp5* knockout mice are available and widely used to explore the role of *Fkbp5* in health and disease. We found that these mice carry a gene duplication of glyoxalase-1, which explains why glyoxalase-1 levels are increased in the *Fkbp5* knockout mice.

KEYWORDS

FKBP51
flanking gene
problem
glyoxalase-1
knockout mice

In several genetic studies researchers linked FKBP5 binding protein 5 (*Fkbp5*) to stress-related diseases and phenotypes such as major depression, posttraumatic stress disorder, and recovery from psychosocial stress (Binder *et al.* 2004; Zimmermann *et al.* 2011; Klengel *et al.* 2013). In addition, *Fkbp5* is also linked to treatment response in depression (Binder *et al.* 2004; Lekman *et al.* 2008). To elucidate the role of FKBP5 in an animal model, a conventional knockout mouse has been constructed and made available to the scientific community (Tranguch *et al.* 2005; Touma *et al.* 2011). These *Fkbp5*-deficient mice show no overt phenotype unless they are older than 10 months of age (O'Leary *et al.* 2011) or exposed to stress (Touma *et al.* 2011; Hartmann *et al.* 2012).

To elucidate the effects of *Fkbp5*-deletion on molecular pathways, we compared the expression profile of *Fkbp5*^{+/+} and *Fkbp5*^{-/-} litter mates. A marked difference in glyoxalase-1 (*Glo1*) mRNA was observed with *Fkbp5*^{-/-} mice expressing greater levels (not shown). Consistent with this observation, about 2-fold more GLO1 protein was found in *Fkbp5*^{-/-} mice (Figure 1A). For more detailed molecular analyses, we sought to establish a cellular model. Therefore, we overexpressed FKBP5 by transient transfection in either primary rat astrocytes or HEK293 cells. However, overexpression of FKBP5 did not change *Glo1* mRNA (not shown) and also not alter protein levels of GLO1 (Figure 1B).

We noted that the genes *Fkbp5* and *Glo1* are only approximately 2 Mb apart from each other on chromosome 17 of the mouse (Figure 1C). In addition, gene duplication around *Glo1* was observed in several mouse strains (Egan *et al.* 2007; Williams *et al.* 2009). The *Fkbp5* deletion was constructed in 129SvJ ES cells, and the resulting mice were then crossed with C57BL/6 animals; 129SvJ mice carry the *Glo1* gene duplication but C57BL/6 mice do not (Williams *et al.* 2009).

Therefore, it appeared likely that through selection of *Fkbp5*^{+/+} and *Fkbp5*^{-/-} alleles in the subsequent crossings the *Glo1* gene duplication originating from 129SvJ mice was coselected with the *Fkbp5*^{-/-} allele, whereas the unduplicated *Glo1* cosegregated with the *Fkbp5*^{+/+} allele. To test this hypothesis, we used polymerase chain reaction (PCR) primers designed for monitoring the *Glo1* gene duplication (Williams *et al.* 2009). DNA samples from *Fkbp5*^{-/-}, *Fkbp5*^{+/+} and *Fkbp5*^{+/+} mice were probed. No *Glo1* gene duplication was detectable in *Fkbp5*^{+/+} mice, whereas the PCR signal in *Fkbp5*^{-/-} mice was clearly detectable and twice as high as in *Fkbp5*^{+/+} mice (Figure 1D). Therefore, the greater levels of mRNA and protein of GLO1 in *Fkbp5*^{-/-} mice compared with wild-type mice are likely due to the double *Glo1* gene dose in these mice. In general, this so-called “flanking allele” problem is a well-known and likely common phenomenon in gene knockout via homologous recombination (Gerlai 1996; Crusio *et al.* 2009). It could be avoided, for example, by genome editing with engineered nucleases or by using inducible gene knock out techniques (Sauer 1998; Carbery *et al.* 2010).

GLO1 is a ubiquitously expressed enzyme involved in the detoxification of methylglyoxal (Thornalley 2008). Methylglyoxal is a toxic byproduct of glycolysis that leads to protein modification and apoptosis (Thornalley 2008) and influences behavior when acting as GABA_A receptor agonist (Distler *et al.* 2012). GLO1 has been linked to diabetic complications, anxiety disorders, schizophrenia, seizure susceptibility, pain, cancer, and aging (Thornalley 2008; Distler and Palmer 2012). At least some of these diseases and phenotypes also

Copyright © 2013 Kollmannsberger *et al.*

doi: 10.1534/g3.113.006445

Manuscript received April 15, 2013; accepted for publication May 24, 2013

This is an open-access article distributed under the terms of the Creative Commons Attribution Unported License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

¹Correspondence: Max Planck Institute of Psychiatry, Kraepelinstr. 10, 80804 Munich, Germany. E-mail: theorein@mpipsykl.mpg.de




Figure 1 (A) Comparison of GLO1 protein expression in hippocampi from *Fkbp5*^{-/-} and *Fkbp5*^{+/+} mice. Hippocampi were prepared, and GLO1 expression was determined after protein extraction by Western blotting (polyclonal antibody; Santa Cruz Biotechnologies); signals were normalized to ACTIN (polyclonal antibody; Santa Cruz Biotechnologies). Expression difference was analyzed by Tukey's test ($n = 12$ per genotype; $P < 0.001$). (B) Overexpression of *Fkbp5* in HEK-293 cells by transient transfection did not affect GLO1 levels. Cells were transfected with *Fkbp5* expressing or control vector, and protein levels were determined in cell extracts by Western blotting 3 d later. Mean protein levels \pm SEM of GLO1 and FKBP5 (polyclonal antibody; Bethyl Laboratories) normalized to ACTIN are indicated ($n = 5$). (C) Scheme of genomic arrangement of *Fkbp5* (28.5–28.4 Mb) and *Glo1* (30.6–30.6 Mb) on chromosome 17 (28.3–30.7 Mb), without (upper) and with (lower) *Glo1* gene duplication. The wild-type *Fkbp5* allele (originating from C57BL/6 mice) is usually coinherit with a single copy of *Glo1*, whereas the knockout *Fkbp5* allele (originating from 129SvJ mice) is coinherit with two copies of *Glo1*. (D) Verification of coinheritance of the *Fkbp5* knockout allele with *Glo1* duplication. Genomic duplications of the *Glo1* spanning region were determined by quantitative reverse-transcription PCR (two independent PCRs per mouse) with primers against the duplication transition region [fw 5'-CTCTGCCCGAGAGAACAGTC and rv 5'-TGATAGAGGCCACACAGCAG (Williams *et al.* 2009)] and normalized to genomic levels of *Npsr1* (determined by quantitative reverse-transcription PCR with the following primers: fw 5'-CAGCTGCTGCCCGGCTAAC and rv 5'-GGTTGGCTGGCATGGCTCAGG).

have been associated with *Fkbp5*, making *Fkbp5*^{-/-} mice potentially very useful genetic model for further investigation. Our observation of *Glo1* gene duplication in *Fkbp5*^{-/-} mice suggests that the *Glo1* status should be taken into consideration when interpreting data. Studies on neuroendocrine and stress effects of *Fkbp5* gene deletion published so far are likely not biased by the *Glo1* gene duplication, in particular because no differences between *Fkbp5*^{+/+} and *Fkbp5*^{-/-} mice have been observed under basal conditions when neuroendocrine parameters or behavior, including anxiety-like behavior, is assessed (Touma *et al.* 2011; Hartmann *et al.* 2012).

LITERATURE CITED

- Binder, E. B., D. Salyakina, P. Lichtner, G. M. Wochnik, M. Ising *et al.*, 2004 Polymorphisms in FKBP5 are associated with increased recurrence of depressive episodes and rapid response to antidepressant treatment. *Nat. Genet.* 36: 1319–1325.
- Carbery, I. D., D. Ji, A. Harrington, V. Brown, E. J. Weinstein *et al.*, 2010 Targeted genome modification in mice using zinc-finger nucleases. *Genetics* 186: 451–459.
- Crusio, W. E., D. Goldowitz, A. Holmes, and D. Wolfer, 2009 Standards for the publication of mouse mutant studies. *Genes Brain Behav.* 8: 1–4.
- Distler, M. G., and A. A. Palmer, 2012 Role of Glyoxalase 1 (Glo1) and methylglyoxal (MG) in behavior: recent advances and mechanistic insights. *Front Genet.* 3: 250.
- Distler, M. G., L. D. Plant, G. Sokoloff, A. J. Hawk, I. Aneas *et al.*, 2012 Glyoxalase 1 increases anxiety by reducing GABAA receptor agonist methylglyoxal. *J. Clin. Invest.* 122: 2306–2315.
- Egan, C. M., S. Sridhar, M. Wigler, and I. M. Hall, 2007 Recurrent DNA copy number variation in the laboratory mouse. *Nat. Genet.* 39: 1384–1389.
- Gerlai, R., 1996 Gene-targeting studies of mammalian behavior: is it the mutation or the background genotype? *Trends Neurosci.* 19: 177–181.
- Hartmann, J., K. V. Wagner, C. Liebl, S. H. Scharf, X. D. Wang *et al.*, 2012 The involvement of FK506-binding protein 51 (FKBP5) in the behavioral and neuroendocrine effects of chronic social defeat stress. *Neuropharmacology* 62: 332–339.
- Klengel, T., D. Mehta, C. Anacker, M. Rex-Haffner, J. C. Pruessner *et al.*, 2013 Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat. Neurosci.* 16: 33–41.
- Lekman, M., G. Laje, D. Charney, A. J. Rush, A. F. Wilson *et al.*, 2008 The FKBP5-gene in depression and treatment response—an association study in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) Cohort. *Biol. Psychiatry* 63: 1103–1110.

- O'Leary, J. C., III, S. Dharia, L. J. Blair, S. Brady, A. G. Johnson *et al.*, 2011 A new anti-depressive strategy for the elderly: ablation of FKBP5/FKBP51. *PLoS ONE* 6: e24840.
- Sauer, B., 1998 Inducible gene targeting in mice using the Cre/lox system. *Methods* 14: 381–392.
- Thornalley, P. J., 2008 Protein and nucleotide damage by glyoxal and methylglyoxal in physiological systems—role in ageing and disease. *Drug Metabol. Drug Interact.* 23: 125–150.
- Touma, C., N. C. Gassen, L. Herrmann, J. Cheung-Flynn, D. R. Bull *et al.*, 2011 FK506 binding protein 5 shapes stress responsiveness: modulation of neuroendocrine reactivity and coping behavior. *Biol. Psychiatry* 70: 928–936.
- Tranguch, S., J. Cheung-Flynn, T. Daikoku, V. Prapapanich, M. B. Cox *et al.*, 2005 Cochaperone immunophilin FKBP52 is critical to uterine receptivity for embryo implantation. *Proc. Natl. Acad. Sci. USA* 102: 14326–14331.
- Williams, R., J. E. Lim, B. Harr, C. Wing, R. Walters *et al.*, 2009 A common and unstable copy number variant is associated with differences in *Glo1* expression and anxiety-like behavior. *PLoS ONE* 4: e4649.
- Zimmermann, P., T. Brückl, A. Nocon, H. Pfister, E. B. Binder *et al.*, 2011 Interaction of variants in the *FKBP5* gene and adverse life events in predicting the first depression onset: results from a ten-year prospective community study. *Am. J. Psychiatry* 168: 1107–1116.

Communicating editor: I. M. Hall