



Published in final edited form as:

J Trauma Stress. 2012 April ; 25(2): 171–178. doi:10.1002/jts.21683.

Identification of Trauma Exposure and PTSD in Adolescent Psychiatric Inpatients: An Exploratory Study

Jennifer F. Havens^{1,2}, Omar G. Gudiño^{1,2}, Emily A. Biggs^{1,2}, Ursula N. Diamond^{1,2}, J. Rebecca Weis^{1,2}, and Marylene Cloitre^{2,3}

¹Department of Child & Adolescent Psychiatry, Bellevue Hospital Center, New York, New York, USA

²Department of Child & Adolescent Psychiatry, New York University School of Medicine, New York, New York, USA

³National Center for PTSD, Palo Alto, California, USA

Abstract

Trauma exposure and posttraumatic stress disorder (PTSD), though prevalent among adolescent psychiatric inpatients, are under-identified in standard clinical practice. In a retrospective chart review of 140 adolescents admitted to a psychiatric inpatient unit, we examine associations between probable PTSD identified through the Child PTSD Symptom Scale and adolescents' service use and clinical characteristics. Results suggest a large discrepancy between rates of probable PTSD identified through standardized assessment and during the emergency room psychiatric evaluation (28.6% vs. 2.2%). Adolescents with probable PTSD had greater clinical severity and service utilization, an increased likelihood of being diagnosed with bipolar disorder (27.5% vs. 9.2%) and being prescribed antipsychotic medications (47.5% vs. 27.6%), and were prescribed more psychotropic medications. Upon discharge, those with probable PTSD were more likely to be assigned a diagnosis of PTSD (45% vs. 7.1%), a comorbid diagnoses of major depressive disorder (30% vs. 14.3%), to be prescribed an antidepressant medication (52.5% vs. 33.7%), and they continued to be prescribed more medications. The under-identification of trauma exposure and PTSD have important implications for the care of adolescents given that accurate diagnosis is a prerequisite for providing effective care. Improved methods for identifying trauma-related problems in standard clinical practice are needed.

Posttraumatic stress disorder (PTSD) is prevalent in youth utilizing inpatient and outpatient mental health services (Allwood, Dyl, Hunt, & Spirito, 2008; Mueser & Taub, 2008), but PTSD may be under-identified in standard clinical practice. In fact, there are great disparities between rates of PTSD identified through structured research assessments (43-66%) and rates documented in patient charts (2-6%; Craine, Henso, Collier, & MacLean, 1988; Miele & O'Brien, 2010; Mueser, Rosenberg, Goodman, & Trumbetta, 2002). In adolescent outpatients with severe emotional disorders, Mueser and Taub (2008) found that only 26% of youth meeting Diagnostic and Statistical Manual of Mental Disorders, 4th Edition Text Revision (DSM-IV-TR; American Psychiatric Association, 2000) criteria for PTSD, based on structured clinical interviews, had a chart diagnosis of PTSD.

Youth diagnosed with PTSD present with high levels of clinical severity and complexity. For example, youth diagnosed with PTSD are more likely to have (a) higher levels of anxiety and depression symptoms; (b) high degrees of comorbidity with other mental health disorders; (c) engage in high-risk (e.g., running away), self-harming, or delinquent behaviors; and (d) evidence poorer functioning relative to youth without PTSD (Ackerman, Newton, McPherson, Jones, & Dykman, 1998; Copeland, Keeler, Angold, & Costello, 2007; Giacon, Reinherz, Silverman, Pakiz, Frost, & Cohen, 1995; Lipschitz, Winegar, Hartnick, Foote, & Southwick, 1999; Mueser & Taub, 2008). Similarly, adolescents with PTSD in psychiatric inpatient settings have a greater number of co-occurring disorders, more suicidal ideation, more prior psychiatric hospitalizations, and longer hospitalizations (Allwood, Dyl, Hunt, & Spirito, 2008, Lipschitz et al., 1999).

Despite the prevalence and clinical importance of identifying trauma exposure and PTSD in mental health settings, currently there are few empirical data that shed light on the treatment implications of under-recognition of these issues. Mueser and Taub (2008) found that youth meeting criteria for PTSD were more likely to have a recorded diagnosis of depression (47% vs. 16%) and were more likely to be on multiple psychotropic medications (31% vs. 11%) when compared to youth without PTSD. Given the high rates of comorbidity between abuse history/PTSD and depression, recent evidence supports the need to understand that a trauma history is present to select the most appropriate treatment for depression (Asarnow et al., 2009; Lewis et al., 2010; Shamseddeen et al., 2011). Also, accurately identifying trauma and PTSD when present is a necessary step for providing the most appropriate and effective treatments for PTSD, especially since a considerable evidence base supports the use of cognitive behavioral therapy (Cohen, 2010; Silverman et al., 2008). Furthermore, if trauma-related clinical problems are not identified and addressed in adolescence, changing cognitive, emotional, and behavioral patterns becomes increasingly more difficult in later development (Ford, 2009).

The current study examined rates of trauma exposure and probable PTSD in a sample of adolescents admitted to a psychiatric inpatient unit. Utilizing structured assessments, we first compared youth with probable PTSD to those without on demographic characteristics (age, gender, race/ethnicity), service use (level of care, past hospitalizations, past outpatient care, and length of current hospitalization), and clinical severity (depressive symptoms, suicidal ideation, prior suicidal behavior). Based on previous research, we anticipated that trauma exposure and probably PTSD would be highly prevalent in this sample and that probable PTSD would be associated with greater service use and severity of clinical presentation. Secondly, we examined associations between a probable diagnosis of PTSD, the intake diagnoses given upon admission, and the medication regimen youth were prescribed prior to admission. Third, we examined associations between a probable diagnosis of PTSD and a patient's discharge diagnoses and medication regimen. Lastly, we examined the concordance between a diagnosis of probable PTSD based on a standardized assessment and mention of a diagnosis of PTSD prior to admission anywhere in the medical record, during the emergency room admission process, and at discharge. Consistent with prior research (e.g., Mueser & Taub, 2008), we predicted poor concordance between rates of probable PTSD identified through standardized assessment and rates identified in prior treatment or during the admission process. Given that treating clinicians in the inpatient unit, however, had access to the assessment results, we expected improved concordance between rates of probable PTSD and discharge diagnoses of PTSD.

Method

Participants and Procedure

Medical records for 140 adolescents ranging in age from 12-18 years ($M = 15.11$, $SD = 1.6$) admitted consecutively to a psychiatric inpatient unit at a major public hospital in New York City between September 2008 and November 2009 were reviewed. Approximately half of the participants (50.7%) were male and the racial/ethnic breakdown was as follows: 37.3% African American, 35.6% Hispanic, 16.1% Non-Hispanic White, 8.5% Asian American, and 2.1% other. The sample was of relatively low socioeconomic status, with 89% of youth qualifying for public health insurance. Patients with mental retardation, autism, or those who were actively psychotic were excluded from this sample.

Patients admitted to the psychiatric inpatient unit initially presented to the emergency department, where they received a psychiatric evaluation conducted by a child psychiatry resident in consultation with an attending child and adolescent psychiatrist. As part of standard clinical practice, all patients admitted to the adolescent psychiatric inpatient unit then completed a battery of measures assessing trauma exposure, PTSD, and depressive symptoms within the first week of their hospitalization. Psychology and psychiatry trainees administered the battery of assessment measures, with all questions being read aloud to patients. Results of this assessment were subsequently available to the entire clinical team working with a particular patient.

For the current study, trained research assistants reviewed medical records of patients and abstracted the relevant clinical information described below. The Institutional Review Boards of the hospital and medical school approved the use of this archival clinical data for research purposes. As data were initially collected as part of standard clinical practice, approval included a waiver of informed consent.

Measures

Service use characteristics—The medical record was reviewed to assess past psychiatric hospitalizations (“yes/no”), number of prior psychiatric hospitalizations, and whether the adolescent had been in outpatient treatment. Given an extreme range of prior hospitalizations (ranging from 0-14) and the fact that 93.5% of youth had 5 or fewer hospitalizations, the 9 patients with more than 5 hospitalizations were coded as having 5 hospitalizations. Where previous mental health treatment was documented anywhere in the medical record, we also coded whether a diagnosis of PTSD was given to the patient by a previous provider (based on any available information in the medical record). Length of the current hospital stay (in days) was also obtained. Length of stay also included extreme values (ranging from 6-222 days). Over 95% of youth had a stay of 81 days or less, so values for the 9 more extreme cases were recoded to 81.

Trauma history—The NYU Child and Adolescent Stressors Checklist (Cloitre, Morin, & Silva, 2002; Mullett-Hume, Anshel, Guevara, & Cloitre, 2008) is a 66-item checklist assessing various potentially traumatic events that was included in the chart. Youth reported whether they had experienced each event in the past (“yes/no”). For the current study, we focused specifically on the 29 items assessing exposure to events that satisfy PTSD Criterion A1 as per the DSM-IV-TR (experiencing, witnessing, or being confronted with events that involve actual or threatened death or serious injury or a threat to physical integrity). Items assessing exposure to stressors that do not meet criterion A1 (e.g., placement in foster care, parental divorce, etc.) were not included. Items were grouped to form the following dichotomous (yes/no) trauma exposure variables: Accidents, disasters, community violence victimization, witnessed community violence, physical abuse, exposure to domestic

violence, and sexual abuse. A sum score representing the total number of different types of traumatic events reported was also computed (possible range from 0-7). The obtained Cronbach's alpha in the current sample was .75.

Probable PTSD—The results of the Child PTSD Symptom Scale (CPSS; Foa, Johnson, Feeny, & Treadwell, 2001) in the chart were used to obtain a probable diagnosis of PTSD. The CPSS contains 17 items corresponding to the 17 symptoms of PTSD outlined in the DSM-IV-TR and is designed for use with children ages 8-18. Items were rated on a 4-point Likert scale ranging from *not at all* (0) to *almost always* (3). The CPSS has demonstrated good test-retest reliability as well as convergent and discriminant validity (Foa, Johnson, Feeny, & Treadwell, 2001). In the current sample, coefficient $\alpha = .90$. The CPSS can be scored to yield a total PTSD severity score or individual items can be scored dichotomously to allow for scoring based on DSM-IV criteria (Foa et al., 2001). In order to have more conservative criteria for the presence or absence of a symptom, we only considered a symptom present if youth indicated that a symptom occurred *half of the time* (2) or *all of the time* (3). A probable diagnosis of PTSD was considered present if, as per *DSM-IV-TR* criteria, youth endorsed at least one re-experiencing, three avoidance, and two arousal symptoms.

Depressive symptoms—Depression severity was assessed using the total score from the Children's Depression Inventory (CDI; Kovacs, 1992) included in the patient chart. The CDI is a 27-item self-report measure assessing cognitive, affective, and behavioral symptoms of depression in youth. Each item on the CDI consists of three statements (scored as 0, 1, 2) and respondents were asked to indicate which item best describes them in the past two weeks. The CDI is a widely used measure with good internal consistency, test-retest reliability, and construct validity (Cole, Hoffman, Tram, & Maxwell, 2000; Kovacs, 1992). In the current sample, the CDI $\alpha = .87$.

Suicidality—Information about any past suicidal behavior or attempts (“yes/no”) was obtained from patient's intake records. Current suicidal ideation was considered present if the patient indicated on the CDI (Kovacs, 1992) that they “think about” or “want to” kill themselves.

Diagnosis on admission and discharge—Diagnoses on admission documented in the medical record were provided by child psychiatry residents in consultation with attending child psychiatrists in the emergency department. Diagnoses on discharge documented in the medical record were provided by attending child psychiatrists on the adolescent unit, in consultation with the clinical team. In most cases, we used the specific diagnosis or diagnoses recorded in the chart. Because some specific diagnoses were assigned infrequently, however, for the purpose of the current study we grouped some diagnoses as follows: Anxiety disorders (including generalized anxiety disorder; anxiety disorder, NOS; obsessive compulsive disorder; and panic disorder, but excluding PTSD), psychotic disorders (including schizophrenia; schizoaffective disorder; psychosis, NOS; and delusional disorder); pervasive developmental disorders (including only pervasive developmental disorder, NOS and Asperger's Syndrome, but excluding autism); and impulse control disorders (including impulse control disorder, NOS and intermittent explosive disorder). In addition to coding the individual disorders given to patients, we also tallied the total number of diagnoses given on admission and discharge. Given that 98.6% of youth had five or fewer diagnoses on admission, patients with more extreme scores on this variable were recoded to have a maximum of five admission diagnoses. Diagnoses at discharge were not recoded, given that 5 diagnoses was the maximum number assigned.

Medication regimen on admission and discharge—The medication regimen was coded by reviewing prescriptions for psychotropic medication subjects had prior to admission and at discharge based on all available data in the medical records. The presence or absence of medication from a given class (antidepressant, stimulant, antipsychotic, mood stabilizer, anxiolytic, and other) was coded. Additionally, we tallied the total number of classes of psychotropic medications youth were prescribed prior to admission and at discharge.

Data Analysis

Variables were first examined for skewness and kurtosis and variables with extreme scores (length of stay, number of diagnoses at admission, and number of hospitalizations) were recoded as described above. In the current study, cases with missing data were omitted from analyses through listwise deletion. Minimum covariance coverage was 96%. To address the study aims, we conducted a series of chi-square analyses to examine differences between those coded as having a probable diagnosis of PTSD and those without on categorical variables. In cases where cell sample sizes were less than 5, level of significance was computed using Fisher's exact test. Group differences on continuous variables were assessed using *t*-tests. In instances when the results of Levene's test for homogeneity of variances were significant, the *t*-statistic was computed assuming heterogeneous variances across groups. Cohen's kappa (κ) was used to assess concordance between a probable diagnosis of PTSD based on the CPSS and a diagnosis of PTSD prior to admission, at admission, and at discharge. All tests were 2-tailed, with the significance level set at $p < .05$.

Results

Table 1 shows the overall values of the study variables as well as the differences between those with and without probable PTSD based on the CPSS. Nearly all (96.4%) adolescents reported lifetime exposure to at least one of the seven types of traumatic events assessed. Sexual abuse (24.3%) was endorsed least often whereas witnessed community violence (70%) and personal victimization related to community violence (70%) were endorsed most often. On average, participants reported being exposed to multiple traumatic events ($M=3.26$, $SD=1.69$) in their lifetime. Of the total sample, 28.6% met criteria for a probable diagnosis of PTSD on the CPSS. As expected, cumulative trauma exposure, witnessed community violence, physical abuse, exposure to domestic violence, and sexual abuse were significantly associated with probable PTSD.

Demographics, Service Use, and Clinical Severity

A diagnosis of probable PTSD was not significantly associated with youth age, race/ethnicity, or sex. Youth with probable PTSD reported significantly more past psychiatric hospitalizations and were hospitalized for more days during their current hospitalization. Relative to those without probable PTSD, youth with probable PTSD endorsed current suicidal ideation more often and reported greater depressive symptom severity (see Table 1).

Admission Diagnosis and Medication Regimen

Table 2 presents information about emergency room diagnoses and medication regimen prior to admission for the overall sample and for those with and without probable PTSD. Of the 16 diagnoses coded, probable PTSD was only significantly associated with a diagnosis of bipolar disorder given in the emergency room. A significantly higher proportion of youth with probable PTSD were given a diagnosis of bipolar disorder (27.5%) relative to youth without probable PTSD (9.2%). A probable diagnosis of PTSD was also uniquely associated with being on an antipsychotic medication immediately prior to their hospitalization (47.5% vs. 27.6%). As seen in Table 2, the medication regimen of adolescents with probable PTSD

consisted of more medication classes ($M=1.00$, $SD=1.04$) relative to youth without probable PTSD ($M=0.62$, $SD=0.96$).

Discharge Diagnosis and Medication Regimen

Although the current study is descriptive in nature, we examined discharge diagnoses to explore whether systematically assessing for trauma exposure and PTSD is associated with increased identification of PTSD and changes in medication regimen at discharge. As seen in Table 3, youth with probable PTSD were more likely to be given a chart diagnosis of PTSD (45% vs. 7.1%) at discharge. Furthermore, probable PTSD was also significantly associated with a diagnosis of major depressive disorder at discharge (30% vs. 14.3%). Youth identified as having probable PTSD were also significantly more likely to be prescribed an antidepressant medication (52.5% vs. 33.7%) upon discharge and on average continued to be prescribed more different types of medications ($M=1.58$, $SD=0.98$ vs. $M=1.20$, $SD=0.94$).

PTSD Diagnostic Concordance

To examine the degree of association between a current diagnosis of probable PTSD based on standardized assessment and PTSD diagnoses in prior treatment (based on information available in the entire medical record), in the emergency room psychiatric evaluation, and at discharge we calculated Kappa coefficients. Results revealed that concordance was poor when comparing a current diagnosis of probable PTSD to a diagnosis of PTSD in previous treatment ($\kappa = .14$) or to a diagnosis of PTSD in the emergency room ($\kappa = .01$). Concordance between a diagnosis of probable PTSD based on standardized assessment and a discharge diagnosis of PTSD fell in the fair to moderate range ($\kappa = .43$).

Discussion

With over 96% of youth reporting exposure to at least one potentially traumatic event and youth commonly reporting exposure to multiple types of events, our results underscore the extremely high prevalence of trauma exposure in adolescent psychiatric inpatients. Despite high rates of identified probable PTSD (28.6%) based on standardized assessment, the current study is consistent with previous work suggesting that trauma exposure and PTSD are under-identified in psychiatric inpatient settings (e.g., Allwood, Dyl, Hunt, & Spirito, 2008, Lipschitz et al., 1999). Only one patient out of 40 (2.5%) who met criteria for probable PTSD received an admission diagnosis of PTSD; only 6 patients out of 40 (15%) with probable PTSD had a historical diagnoses of PTSD noted in their admission documentation.

As predicted, adolescents with trauma exposure and PTSD present with complex clinical pictures (e.g., current suicidal ideation) and high service utilization (e.g., longer hospitalizations) relative to youths without probable PTSD. Extending previous work, our findings regarding the lack of recognition of trauma-related symptoms during the admission process suggests that the clinical complexity in this population may complicate appropriate diagnosis and treatment planning. The greater clinical complexity appears to consistently relate to greater medication use and increased likelihood of the prescription of anti-psychotic medications prior to admission as well as increased likelihood of admission diagnoses of bipolar disorder.

PTSD and acute stress disorder are the only diagnoses where knowledge about exposure to a specific event (a trauma) is required to make the diagnosis. Without knowledge about exposure to a traumatic event, it is virtually impossible to evaluate re-experiencing symptoms (e.g., flashbacks, psychological distress associated with the traumatic

experience). Furthermore, evaluation of possible symptoms of avoidance of thoughts and reminders of the traumatic event also require that exposure to the traumatic event is known. Thus, if the youth or their family does not spontaneously disclose exposure to a traumatic event or if the evaluating clinician does not systematically assess exposure to trauma, it will be impossible to obtain information about these central aspects of PTSD. Notably, assessment of symptoms of hyper-arousal (sleep problems, irritability, difficulty concentrating, hyper-vigilance, and an increased startle response) does not require knowledge about exposure to a traumatic event.

When clinicians do not obtain a trauma history and assess for trauma-related problems, the presence of high-risk behaviors, affective volatility, and symptoms of hyper-arousal of adolescents with trauma-related symptoms may lead to diagnostic confusion (including inaccurate diagnosis of bipolar disorder) and increased likelihood of the prescription of second generation anti-psychotics for mood reactivity and interpersonal aggression. While second generation anti-psychotics medications may have a role in the treatment of child and adolescent PTSD (Strawn, Brooks, DelBello, Geraciotti, & Putnam, 2010), there is no evidence that they are efficacious as a single modality treatment. As trauma-focused psychotherapies have the clearest evidence base in child and adolescent PTSD (Silverman et al., 2008), accurate diagnosis of PTSD is an essential foundation for effective treatment.

Previous work conducted in community mental health clinics found poor agreement between results from evidence-based trauma symptom measures available to clinicians and the diagnoses clinicians assign at intake (Osterberg, Jensen-Doss, Cusack, & de Arellano, 2009). In the current study, we found that youth identified as having probable PTSD through standardized assessment were more likely to be assigned a diagnosis of PTSD and prescribed an antidepressant on discharge. As an independent diagnosis of PTSD was not available to provide a “gold standard” for the presence or absence of PTSD, it is not possible to determine the accuracy of these discharge diagnoses and future work will be needed to examine this more closely. However, a PTSD rate of 18.1% on discharge is more consistent with rates of PTSD identified in previous studies of adolescent inpatients (Allwood, Dyl, Hunt, & Spirito, 2008, Lipschitz et al., 1999). Additionally, it is notable that use of a brief assessment instrument provided similar rates of PTSD diagnosis to those of prior studies using structured clinician administered assessments – this has important implications for use in other settings where administration of more lengthy assessments would not be practical.

Reflecting our efforts to target trauma-related symptoms in this population through systematic assessment, patients with probable PTSD were more likely to receive PTSD and major depressive disorder diagnoses on discharge. Although the current study did not employ an experimental design and was not designed to evaluate the implementation of standardized assessment, our descriptive results are consistent with the assertion that routine use of evidence-based assessment instruments improves the identification and treatment of trauma-related problems. Given the high reliance on safety net providers such as emergency rooms and psychiatric hospitals by underserved and racial/ethnic minority youth (Surgeon General, 2001), examining the quality of mental health care in these settings is crucial. The current findings highlight the importance of developing innovations and adaptations in the acute care service delivery model in order to improve the identification of trauma exposure and trauma-related symptomatology, when present. Improved diagnostic accuracy may be one important way of ensuring that the needs of the most vulnerable youth, who encounter the greatest number of barriers to accessing care, receive appropriate and effective services when they do come into contact with the public mental health system.

References

- Ackerman PT, Newton JE, McPherson WB, Jones JG, Dykman RA. Prevalence of post traumatic stress disorder and other psychiatric diagnoses in three groups of abused children (sexual, physical, and both). *Child Abuse & Neglect*. 1998; 22:759–774.10.1016/S0145-2134(98)00062-3 [PubMed: 9717613]
- Allwood MA, Dyl J, Hunt JI, Spirito A. Comorbidity and service utilization among psychiatrically hospitalized adolescents with posttraumatic stress disorder. *Journal of Psychological Trauma*. 2008; 7:104–121.10.1080/19322880802231791
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th, text rev. Arlington, VA: APA; 2000.
- Asarnow JA, Emslie G, Clarke G, Wagner KD, Spirito A, Vitiello B, et al. Treatment of Selective Serotonin Reuptake Inhibitor Resistant Depression in Adolescents: Predictors and Moderators of Treatment Response. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2009; 48:330–339.10.1097/CHI.Ob013e3181977476 [PubMed: 19182688]
- Cloitre M, Morin N, Silva R. The NYU Child and Adolescent Stressors Checklist—Revised (CASC). Unpublished instrument. 2002
- Cohen JA. Practice parameter for the assessment and treatment of children and adolescents with posttraumatic stress disorder (AACAP official action). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010; 49:414–430.10.1097/00004583-201004000-00021 [PubMed: 20410735]
- Cole DA, Hoffman K, Tram JM, Maxwell SE. Structural differences in parent and child reports of children's symptoms of depression and anxiety. *Psychological Assessment*. 2000; 12:174–185.10.1037//1040-3590.12.2.174 [PubMed: 10887763]
- Copeland WE, Keeler G, Angold A, Costello EJ. Traumatic events and posttraumatic stress in childhood. *Archives of General Psychiatry*. 2007; 64:577–584.10.1001/archpsyc.64.5.577
- Craine LS, Henson CE, Colliver JA, MacLean DG. Prevalence of a history of sexual abuse among female psychiatric patients in a state hospital system. *Hospital and Community Psychiatry*. 1988; 39:300–304. [PubMed: 3356438]
- Foa EB, Johnson KM, Feeny NC, Treadwell KRH. The child PTSD symptom scale: A preliminary examination of its psychometric properties. *Journal of Clinical Child Psychology*. 2001; 30:376–384.10.1207/S15374424JCCP3003_9 [PubMed: 11501254]
- Ford, JD. Neurobiological and developmental research. In: Courtois, CA.; Ford, JD., editors. *Treating complex traumatic stress disorders: An evidence-based guide*. New York: Guilford Press; 2009. p. 31-58.
- Giaconia RM, Reinherz HZ, Silverman AB, Pakiz B, Frost AK, Cohen E. Traumas and posttraumatic stress disorder in a community population of older adolescents. *Journal of American Academy of Child Adolescent Psychiatry*. 1995; 34:1369–1380.10.1097/00004583-199510000-00023
- Kovacs, M. *Children's Depression Inventory Manual*. Multi-Health Systems Inc.; North Tonawanda, NY: 1992.
- Lewis CC, Simons AD, Nguyen LJ, Murakami JL, Reid MW, Silva SG, March JS. Impact of Childhood Trauma on Treatment Outcome in the Treatment for Adolescents with Depression Study (TADS). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010; 49:132–140.10.1097/00004583-201002000-00007 [PubMed: 20215935]
- Lipschitz DS, Winegar RK, Hartnick E, Foote B, Southwick SM. Posttraumatic stress disorder in hospitalized adolescents: Psychiatric comorbidity and clinical correlates. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:385–392.10.1097/00004583-199904000-00010 [PubMed: 10199109]
- Miele D, O'Brien EJ. Underdiagnosis of posttraumatic stress disorder in at risk youth. *Journal of Traumatic Stress*. 2010; 23:591–598.10.1002/jts.20572 [PubMed: 20931661]
- Mueser KT, Rosenberg SD, Goodman LA, Trumbetta SL. Trauma, PTSD, and the course of schizophrenia: an interactive model. *Schizophrenia Research*. 2002; 53:123–143. [PubMed: 11728845]

- Mueser KT, Taub JT. Trauma and PTSD among adolescents with severe emotional disorders involved in multiple service systems. *Psychiatric Services*. 2008; 59:627–634.10.1176/appi.ps.59.6.627 [PubMed: 18511582]
- Mullett-Hume E, Anshel D, Guevara V, Cloitre M. Cumulative Trauma and Posttraumatic Stress Disorder Among Children Exposed to the 9/11 World Trade Center Attack. *American Journal of Orthopsychiatry*. 2008; 78:103–108.10.1037/0002-9432.78.1.103 [PubMed: 18444732]
- Osterberg LD, Jensen-Doss A, Cusack KJ, de Arellano MA. Diagnostic practices for traumatized youths: Do clinicians incorporate symptom scale results? *Community Mental Health Journal*. 2009; 45:497–507.10.1007/s10597-009-9258-8 [PubMed: 19957106]
- Shamseddeen W, Asarnowm JR, Clarke G, Vitiello B, Wagner KD, Birmaher B, et al. Impact of Physical and Sexual Abuse on Treatment Response in the Treatment of Resistant Depression in Adolescent Study (TORDIA). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2011; 50:293–301.10.1016/j.jaac.2010.11.019 [PubMed: 21334569]
- Silverman WK, Ortiz C, Visersvaran C, Burns BJ, Kolko DJ, Putnam FW, Amaya-Jackson L. Evidence-Based Psychosocial Treatments for Children and Adolescents Exposed to Traumatic Events. *Journal of Clinical Child & Adolescent Psychology*. 2008; 37:156–183.10.1080/15374410701818293 [PubMed: 18444057]
- Strawn JR, Brooks RK, DelBello MP, Geraciotti TD, Putnam FW. Psychopharmacologic Treatment of Posttraumatic Stress Disorder in Children and Adolescents: A Review. *Journal of Clinical Psychiatry*. 2010; 71:932–941.10.4088/JCP.09r05446blu [PubMed: 20441729]
- Surgeon General. *Mental Health: Culture, Race, and Ethnicity: A Supplement to Mental Health: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health; 2001.

Table 1
Demographic, Service Use, and Mental Health Characteristics Overall and by Probable PTSD Status

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
Age	15.11	1.60	15.16	1.59	14.98	1.63	χ^2 0.63
Sex							
Boy	70	50.0	55	55.0	15	37.5	
Girl	70	50.0	45	45.0	25	62.5	3.50
Race/Ethnicity							
Non-Hispanic White	19	16.1	14	17.3	5	13.5	
African-American	44	37.3	26	32.1	18	48.6	
Hispanic	42	35.6	29	35.8	13	35.1	
Asian American	10	8.5	9	11.1	1	2.7	
Other	3	2.1	3	3.7	-	-	5.58
Service Use							
Past psych hosp	69	50.4	45	46.4	24	60.0	2.10 ^a
No. past psych hosp	1.21	1.58	.98	1.39	1.78	1.89	-2.41 ^a
Past outpatient treatment	76	55.5	56	57.7	20	50.0	0.69 ^a
Past diagnosis PTSD	10	7.2	4	4.1	6	15.0	5.04 ^b
Days hospitalized	23.57	18.64	19.47	12.91	33.63	25.70	-3.32 ^{***b}
Suicidality							
Prior behavior	33	24.1	19	19.6	14	35.0	3.68 ^a
Current ideation	46	34.3	24	25.3	22	56.4	11.90 ^{***c}
Trauma history							
Accident	72	51.4	47	47.0	25	62.5	2.75
Disaster	42	30.0	28	28.0	14	35.0	0.67
Victim com. violence	98	70.0	71	71.0	27	67.5	0.17
Witness com. violence	98	70.0	65	65.0	33	82.5	4.17 [*]
Physical abuse	74	52.9	46	46.0	28	70.0	6.60 ^{**}

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis t/ χ^2
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
Domestic violence	39	27.9	20	20.0	19	47.5	10.75***
Sexual abuse	34	24.3	17	17.0	17	42.5	10.10***
Total no. traumas	3.26	1.69	2.94	1.68	4.08	1.44	-3.75***
Depressive symptoms	11.84	8.15	8.68	5.72	19.51	8.13	-7.59***,c

Note. PTSD = post-traumatic stress disorder, psych hosp = psychiatric hospitalization, com. violence = community violence

^a N=137 due to missing data.

^b N=138 due to missing data.

^c N=134 due to missing data.

* $p < .05$

** $p < .01$

*** $p < .001$.

Table 2
Diagnosis and Medication Regimen at Admission Overall and by PTSD Status

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
Admission diagnosis							
PTSD	3	2.2	2	2.0	1	2.5	0.03
Anxiety	8	5.8	7	7.1	1	2.5	1.12
Mood disorders							
Major depressive d/o	17	12.3	15	15.3	2	5.0	2.79
Dysthymic d/o	6	4.3	3	3.1	3	7.5	1.35
Depressive d/o, NOS	14	10.1	8	8.2	6	15.0	1.46
Bipolar d/o	20	14.5	9	9.2	11	27.5	7.69**
Mood d/o, NOS	38	27.5	27	27.6	11	27.5	<.01
Disruptive behavior							
Conduct d/o	13	9.4	8	8.2	5	12.5	0.62
Opp. Defiant d/o	20	14.5	15	15.3	5	12.5	0.18
Disrupt beh. d/o, NOS	7	5.1	5	5.1	2	5.0	<.01
ADHD	31	22.5	21	21.4	10	25.0	0.21
Impulse control d/o	6	4.3	6	6.1	-	-	2.56
Adjustment d/o	13	9.4	9	9.2	4	10.0	0.02
Substance use d/o	9	6.5	7	7.1	2	5.0	0.21
Pervasive develop. d/o	6	4.3	6	6.1	-	-	2.56
Psychotic d/o	19	13.8	12	12.2	7	17.5	0.66
No. admission diagnoses	1.64	1.11	1.61	1.1.06	1.73	1.22	-0.54
Medication at admission							
Antidepressant	22	15.9	12	12.2	10	25.0	3.45
Stimulant	16	11.6	11	11.2	5	12.5	0.05
Antipsychotic	46	33.3	26	27.6	19	47.5	5.09*
Mood stabilizer	11	8.0	7	7.1	4	10.0	0.32
Anxiolytic	-	-	-	-	-	-	-
Other	6	4.3	4	4.1	2	5.0	0.06

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
No. medication classes	.73	.99	.62	.96	1.00	1.04	t/χ^2 -2.05*

Note. N=138 due to incomplete admission data. PTSD= post-traumatic stress disorder, NOS = not otherwise specified, ADHD = attention deficit and hyperactivity disorder, d/o = disorder, opp. defiant = oppositional defiant, disrupt beh. = disruptive behavior.

* p <.05

** p <.01

Table 3
Diagnosis and Medication Regimen at Discharge Overall and by PTSD Status

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
Discharge diagnosis							
PTSD	25	18.1	7	7.1	18	45.0	27.44***
Anxiety	8	5.8	6	6.1	2	5.0	0.07
Mood disorders							
Major depressive d/o	26	18.8	14	14.3	12	30.0	4.59*
Dysthymic d/o	7	5.1	3	3.1	4	10.0	2.84
Depressive d/o, NOS	19	13.8	13	13.3	6	15.0	0.07
Bipolar d/o	11	8.0	8	8.2	3	7.5	0.02
Mood d/o, NOS	31	22.5	23	23.5	8	20.0	0.20
Disruptive behavior							
Conduct d/o	9	6.5	8	8.2	1	2.5	1.49
Opp. defiant d/o	20	14.5	14	14.3	6	15.0	0.01
Disrupt beh. d/o, NOS	2	1.4	1	1.0	1	2.5	0.44
ADHD	26	18.8	18	18.4	8	20.0	0.05
Impulse control d/o	6	4.3	6	6.1	-	-	2.56
Adjustment d/o	14	10.1	12	12.2	2	5.0	1.64
Substance use d/o	16	11.6	11	11.2	5	12.5	0.05
Pervasive dev. d/o	3	2.2	3	3.1	-	-	1.25
Psychotic spectrum d/o	8	5.8	6	6.1	2	5.0	0.07
No. admission diagnoses	1.67	.88	1.56	1.95	1.95	1.01	-2.39*
Medication at discharge							
Antidepressant	54	39.1	33	33.7	21	52.5	4.23*
Stimulant	33	23.9	25	25.5	8	20.0	0.47
Antipsychotic	53	38.4	34	34.7	19	47.5	1.97
Mood stabilizer	14	10.1	9	9.2	5	12.5	0.34
Anxiolytic	10	7.2	7	7.1	3	7.5	0.01

Variable	Overall N = 140		No PTSD n = 100		Probable PTSD n = 40		Analysis
	N/M	% / SD	n/M	% / SD	n/M	% / SD	
Other	17	12.3	10	10.2	7	17.5	t/χ ² 1.40
No. medication classes	1.31	.97	1.20	.94	1.58	.98	-2.07*

Note. N=138 due to incomplete discharge data. PTSD= post-traumatic stress disorder, NOS = not otherwise specified, ADHD = attention deficit and hyperactivity disorder, d/o = disorder, opp. defiant = oppositional defiant, disrupt beh. = disruptive behavior.

* p < .05

** p < .01

*** p < .001