



Published in final edited form as:

*Obstet Gynecol.* 2013 July ; 122(1): 64–71. doi:10.1097/AOG.0b013e3182955e58.

## Short Interpregnancy Intervals in the United States

Alison Gemmill, MPH, MA and

Department of Demography, University of California, Berkeley, Berkeley, California

Laura Duberstein Lindberg, PhD

Guttmacher Institute, Research Division, New York, New York

### Abstract

**Objective**—To investigate the prevalence and correlates of short interpregnancy intervals in the United States.

**Methods**—We analyzed pregnancy data from a nationally representative sample of 12,279 women from the 2006–2010 National Survey of Family Growth. We limited our sample to second and higher order births within 5 years of the interview. Interpregnancy intervals were calculated as the interval between the delivery date of the preceding live birth and the conception date of the index birth, with short interpregnancy intervals defined as intervals less than 18 months. We used simple and multivariate logistic regression analyses to examine associations between short interpregnancy intervals and maternal demographic and childbearing characteristics, including pregnancy intention.

**Results**—Among the 2,253 pregnancies in our sample, one third (35%) were conceived within 18 months of a prior birth. After adjusting for sociodemographic and childbearing characteristics, women were significantly more likely to have a short interpregnancy interval if they were aged 15–19 years or married at the time of conception of the index pregnancy, initiated childbearing after age 30 years, or reported the pregnancy as unintended. Short interpregnancy intervals were more likely to be intended among more advantaged women (married, non-Hispanic white, college educated, or non-Medicaid delivery). We estimate that preventing unintended pregnancies would reduce the proportion of short interpregnancy intervals from 35% to 23%.

**Conclusion**—Providing counseling about the potential negative consequences of short interpregnancy intervals and improving women's contraceptive use to reduce rates of unintended pregnancy would likely reduce the proportion of short interpregnancy interval pregnancies in the United States.

### Introduction

Short interpregnancy intervals are associated with a number of adverse outcomes for both mother and child, including increased risk of preterm birth, low birth weight, and preeclampsia (1–5), making prevention of short interpregnancy intervals a public health priority in the United States. Specifically, the 2020 Healthy People objectives call for a 10% reduction of pregnancies that occur within 18 months of a previous birth (6).

---

**Corresponding author:** Laura Duberstein Lindberg, Guttmacher Institute, 125 Maiden Lane, 7<sup>th</sup> Floor, New York, NY 10038; llindberg@guttmacher.org; phone: (212) 248-1111.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Financial Disclosure:** The authors did not report any potential conflicts of interest.

There has been little systematic national surveillance of pregnancy interval length in the United States. Instead, most relevant studies focus on single states and key subpopulations (7,8). National-level baseline estimates in Healthy People 2020 relying on unpublished bivariate analyses from the National Survey of Family Growth suggest that about a third of interpregnancy intervals in the United States fall below this 18-month threshold (6). However, to our knowledge, there has been no national-level investigation of the correlates of short interpregnancy intervals in a multivariate framework.

The potential association between pregnancy intentions and interpregnancy intervals is of particular interest. Strategies to reduce unintended pregnancy may indirectly impact the prevalence of short interpregnancy intervals; in particular, reducing mistimed pregnancies (those reported as occurring sooner than a woman desires) would increase interpregnancy interval length. Moreover, a 2001 study in Denmark found that unplanned pregnancy was associated with an increased risk of interpregnancy interval of 9 months or less (9). Given high rates of unintended pregnancy in the United States (10), similar investigation of the relationship between pregnancy intention and short interpregnancy intervals using national-level data is needed.

Our research objective was to identify the characteristics of women associated with short interpregnancy intervals in order to inform programs and policies aimed at reducing the occurrence of short interpregnancy intervals. We use nationally-representative data to analyze the prevalence and correlates of short interpregnancy intervals, paying particular attention to the role of pregnancy intention.

## Materials and Methods

The institutional review board of the second author's organization (Department of Health and Human Services identifier institutional review board 00002197) determined that the project was exempt from Institutional Review Board approval. The data from this study are drawn from the 2006–2010 National Survey of Family Growth, a periodic national probability survey of the noninstitutionalized population aged 15–44 years in the United States, conducted by the National Center for Health Statistics. The survey used a multistage, stratified, clustered sampling frame to collect interviews continuously from June 2006 to June 2010. Methods of data collection and dissemination of the public use dataset are reviewed by the Institutional Review Board at National Center for Health Statistics for protections of human subjects. Further methodological details are available elsewhere (11,12). Face-to-face interviews were conducted with 12,279 women who answered detailed retrospective questions about their pregnancy experiences; the response rate was 78%. Data for each reported pregnancy were contained in a separate data file and linked to the primary respondent file.

The interpregnancy interval was calculated as the time elapsed between the conception date of any second or higher order birth (hereafter labeled index pregnancy) and the date of a prior birth. Pregnancies conceived within 18 months after a previous birth were classified as having a short interpregnancy interval, the convention of prior studies and the Healthy People 2020 objective.

This analysis excluded all pregnancies not ending in live birth (ie, miscarriage, abortion, and stillbirth) due to both likely underreporting of these other pregnancy outcomes and the relevance of birth spacing for perinatal health. We also excluded all multiple births from both the preceding and index pregnancies. To reduce issues of retrospective reporting bias and focus on experiences during a period of current policy interest, we limited our sample to second or higher order singleton births born in the 5 years preceding the interview

(n=2,265); the preceding singleton live births used to calculate the interval could have occurred outside this 5 year window. We also excluded pregnancies with implausible interpregnancy interval lengths due to erroneous reporting (n=5) and pregnancies with missing covariates (n=7). Thus, the final sample size for analysis included 2,253 second and higher order singleton births that occurred within 5 years of the interview date, were preceded by a singleton birth, and had complete data.

Independent variables selected for this analysis include the following measures of mothers' sociodemographic characteristics: race or ethnicity, union status at conception of index pregnancy, Medicaid funded delivery of index pregnancy, and completed education at the time of the interview. Measures related to childbearing include number of births prior to the index pregnancy, age at conception of the index pregnancy, and age at initiation of childbearing, as well as pregnancy intention of the index pregnancy. Pregnancy intention was determined from a series of questions women were asked to assess their feelings right before they got pregnant. Based on their responses, each pregnancy was classified as intended (wanted and on time, later than wanted, didn't care or indifferent), mistimed (wanted but occurring sooner than desired), or unwanted, following conventional measurement approaches for this concept (13).

The present study used three analytical approaches. First, we employed simple and multivariate logistic regression to estimate unadjusted and adjusted odds ratios for the relationship between short interpregnancy intervals and maternal demographic and childbearing characteristics. Multivariate models included all of the measures with a significant bivariate relationship to short interpregnancy intervals ( $\alpha=0.05$ ); union status, parity, and education were retained in the models for theoretical reasons. Because age at first birth and age at most recent conception are essentially a linear combination of the interpregnancy interval length for most pregnancies in the sample, we estimated models controlling for each of these factors separately; Model 1 controls for age at conception of the index birth and Model 2 controls for age at first birth.

The second analytical approach focused on pregnancies occurring from the short interpregnancy interval and used simple logistic regression to examine variation in pregnancy intention by maternal demographics among these pregnancies.

Lastly, we estimated a hypothetical estimate of the share of short interpregnancy intervals if unintended pregnancies (mistimed and unwanted pregnancies) were averted. For this calculation, all pregnancies to women who reported they had not wanted to have any (more) children were removed from both the numerator and the denominator. For pregnancies that were reported by the mother as mistimed (occurring too soon), we recalculated the interpregnancy interval length by adding the number of months by which the pregnancy was reported to be mistimed to the actual interval. For example, if the actual interpregnancy interval was 12 months, but the mother reported that the pregnancy occurred 9 months before desired, then the new interpregnancy interval would be 21 months and would be shifted to a longer interpregnancy interval. Interpregnancy interval lengths for pregnancies reported as intended were unchanged. All analyses were weighted and used the *svy* command prefix in Stata 12.1 (StataCorp, College Station, TX) to adjust for the complex survey design of the National Survey of Family Growth.

## Results

Among the 2,253 pregnancies in our sample, the average interpregnancy interval was 34.0 months. One third (35%) were conceived within 18 months of a prior birth, meeting our criteria of a short interpregnancy interval (Table 1). The majority of pregnancies had an

interpregnancy interval of 18 months or more, with 50% at 18–59 months, and 16% having a length of 60 months or more.

There is some evidence of associations between measures of pregnancy intention, childbearing history, and short interpregnancy intervals (Table 2). Pregnancies reported as mistimed or unwanted were significantly more likely to have short interpregnancy intervals compared to pregnancies reported as intended (unadjusted odds ratios [ORs] 4.3 and 1.8, respectively). Short interpregnancy intervals were significantly inversely associated with age at conception of the pregnancy. In contrast, births to women initiating childbearing before age 30 years were significantly less likely to have shorter interpregnancy intervals than births to women aged 30 years and older at first birth.

There was limited variation in the share of short interpregnancy intervals by other core demographic measures (Table 2). Births to non-Hispanic black women were significantly more likely than births to Hispanic women to have short interpregnancy intervals, as were births whose delivery was paid by Medicaid. However, maternal union status at conception and maternal education were not associated with short interpregnancy intervals at the bivariate level.

Results from multivariate analyses predicting the likelihood of having a short interpregnancy interval are shown in Table 3. Although both models include different measures of age, the results are generally similar to those found in the bivariate results. Both models also indicate that marital status, which was not significant at the bivariate level, is a significant predictor of interpregnancy interval length; the adjusted odds of having a short interpregnancy interval were higher among births to married as compared to single women. Additionally, Model 1 provides evidence that births were more likely to be short interpregnancy interval if born to high parity women or women with a college degree, while Model 2 suggests that interpregnancy intervals are more likely among Medicaid delivery births or births to non-Hispanic black women.

Table 4 further explores the associations between pregnancy intentions and short interpregnancy interval by examining the proportion of all short interpregnancy interval pregnancies (n=791) that were intended. We find that 45% of these pregnancies were reported as intended by the mother and this varied significantly across all of the childbearing and demographic measures examined. Short interpregnancy interval pregnancies to more advantaged mothers were more likely to be intended; 59–70% were intended among births to those aged 30 years or older at first birth, college graduates, and those not using Medicaid to pay for delivery. Similarly, about half of the short interpregnancy interval pregnancies were intended among births to white women, those married at conception, and those aged 30–44 years at the most recent birth.

We calculated the extent to which preventing unintended pregnancies would reduce the share of all pregnancies that were short interpregnancy intervals by assuming that all unwanted pregnancies in our sample were averted and that mistimed pregnancies were appropriately timed by the mother. Based on these counterfactual assumptions, we estimate that the prevention of unintended pregnancies would reduce the proportion of short interpregnancy intervals overall from 35% to 23%.

## Discussion

Using recent nationally representative data, we estimate that more than one in three of second or higher order singleton births occur after a short interpregnancy interval. Indeed, nearly 7% are conceived within 6 months of a prior birth.

Age plays an important role in short interpregnancy intervals for two distinct groups of women. First, we identify short interpregnancy intervals as a correlated and troublesome outcome of second births to teenaged mothers; two-thirds of births to this age group had a short interpregnancy interval. Although teenaged mothers make up only a small share (6%) of all second and higher order births, additional interventions are needed to address suboptimal birth spacing in this population. Second, women with a first birth at age 30 years or older are more likely to experience short interpregnancy intervals than those initiating childbearing earlier, suggesting that closer birth spacing is a response to later initiation of childbearing. This premise is supported by the finding that among pregnancies that had short interpregnancy intervals to women initiating childbearing after age 30 years, nearly three out of four were intended pregnancies. For this group, short interpregnancy intervals appear to be a choice and not an unintended outcome.

With 55% of short interpregnancy interval pregnancies unintended, helping women achieve their desired pregnancy intentions is the low-hanging fruit for public health interventions to reduce the share of short interpregnancy interval pregnancies. Improvements in women's contraceptive use can further reduce rates of unintended pregnancy, and by extension short interpregnancy intervals. Long-acting reversible contraceptives, such as intrauterine devices and implants, seem particularly well-suited to lengthening the interpregnancy interval (17,18). However, this approach will only go so far since we estimated that alleviating all unintended pregnancies among these second and higher order births—an exceptionally lofty goal—would still leave 23% with short interpregnancy intervals.

Further supporting the idea that closely spaced births may be part of a strategy for family building was the finding that more than half of short interpregnancy interval births to more advantaged women were reported as intended. Increasing interpregnancy interval length among intended births is more challenging and likely requires health care providers to educate and counsel patients about the potential negative health consequences of short interpregnancy intervals. Further research is needed to reevaluate the evidence base for negative health consequences of short interpregnancy intervals among the substantial share of intended births to more advantaged women. Indeed, any suggestion of promoting longer pregnancy intervals for these women must weigh benefits against the potential health risks and decreased fecundity associated with increasing maternal age at birth (19,20).

The choice of an 18-month cutoff to define a short interpregnancy interval in this analysis was based on the indicator used in Healthy People 2020. While much literature suggests that interpregnancy intervals less than 18 months are associated with increased risk, it is important to note that even within this 18-month window, the level of risk likely decreases as interval length increases (1). Additionally, while the Healthy People 2020 objective is limited to reduction of short interpregnancy intervals, there is evidence that interpregnancy intervals over 60 months are also detrimental to maternal and child health (1). Considering that 16% of pregnancies in our sample had interpregnancy intervals over 60 months, these data suggest that about half (51%) of interpregnancy intervals in the United States fall outside generally recommended standards.

The National Center of Health Statistics, as part of the Healthy People 2020 objectives, tracks short interpregnancy intervals using the same National Survey of Family Growth data we analyze here. However, National Center of Health Statistics reports as their baseline measure the *share of women* having a short interpregnancy interval during the 5 years preceding the interview, as opposed to the *share of pregnancies*. Since each woman can only provide a single pregnancy to the numerator, their measurement approach is biased away from measuring short interpregnancy intervals; women with shorter intervals may have more than one pregnancy during the period and thus have relevant pregnancy experiences

excluded. Our pregnancy-based measure conceptually parallels the stated objective to reduce the share of short interpregnancy interval pregnancies and is methodologically sound in incorporating all reported pregnancies ending in live birth. Future monitoring should use a pregnancy-based measure as we do here. Although linked vital records are often used to assess the causes and consequences of interpregnancy length, they are unable to provide pregnancy intention status, which, as demonstrated here, is a key determinant of interpregnancy length. Another source of data, the Pregnancy Risk Assessment Monitoring System, provides information on interpregnancy length and pregnancy intention, but not all states participate in data collection efforts. Therefore, this study drew strength from its ability to link interpregnancy intervals and maternal characteristics, including pregnancy intentions, from a recent nationally representative large sample of pregnancies.

While the National Survey of Family Growth data is widely utilized and considered highly reliable and valid, there are always limitations to self-reported data. Relevant to this study, women may potentially misreport birth dates or gestational age of a child, resulting in a miscalculation of the length of the interpregnancy interval. External validation with medical records of this self-reported data was not possible; however, any misreporting is likely random and should not bias the observed relationships. The response rate of the National Survey of Family Growth was 78%, which may have resulted in underrepresentation of certain high-risk groups. Likewise, our exclusion of multiple births may have also impacted our estimate of short interpregnancy interval pregnancies. Lastly, there has been concern about bias in the retrospective reporting of pregnancy intentions, if women adjust their reporting of births towards more intended farther from the actual time of conception (21). Limiting the analyses to a 5-year retrospective period minimizes this concern, and follows methodological approaches established in prior analyses of this measure from the National Survey of Family Growth (10,13).

## Acknowledgments

Supported by awards R01HD059896 and HD07275 from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) of the National Institutes of Health (NIH). The content is solely the responsibility of the authors and does not necessarily represent the official views of NICHD or the NIH.

## References

1. Conde-Agudelo A, Rosas-Bermudez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta analysis. *JAMA*. 2006; 295:1809–1823. [PubMed: 16622143]
2. Conde-Agudelo A, Rosas-Bermudez A, Castaño F, Norton MH. Effects of birth spacing on maternal, perinatal, infant, and child health: a systematic review of causal mechanisms. *Stud Fam Plan*. 2012; 43:93–114.
3. Khoshnood B, Lee KS, Wall S, Hsieh HL, Mittendorf R. Short interpregnancy intervals and the risk of adverse birth outcomes among five racial/ethnic groups in the United States. *Am J Epidemiol*. 1998; 148:798–805. [PubMed: 9786235]
4. Klerman LV, Cliver SP, Goldenberg RL. The impact of short interpregnancy intervals on pregnancy outcomes in a low-income population. *Am J Public Health*. 1998; 88:1182–1185. [PubMed: 9702144]
5. Zhu BP, Rolfs RT, Nangle BE, Horan JM. Effect of the interval between pregnancies on perinatal outcomes. *N Engl J Med*. 1999; 340:589–594. [PubMed: 10029642]
6. U.S. Department of Health and Human Services. Washington, DC: Office of Disease Prevention and Health Promotion. Healthy People 2020. Available at: <http://www.healthypeople.gov/2020/topicsobjectives2020/objectiveslist.aspx?topicId=13>. [Retrieved January 4, 2013]
7. Larder C, Grigorescu V. Michigan Department of Community Health. Interpregnancy intervals, MI PRAMS 2007. MI PRAMS Delivery. 2010; 9:1. Available at: [http://www.michigan.gov/documents/mdch/PRAMS\\_newsletter\\_July\\_2010\\_7\\_8\\_10\\_Draft\\_3\\_Final\\_338644\\_7.pdf](http://www.michigan.gov/documents/mdch/PRAMS_newsletter_July_2010_7_8_10_Draft_3_Final_338644_7.pdf).

8. Bloebaum, L.; Baksh, L.; McGarry, J.; Carapezza, D.; Hossain, S.; Streeter, N. [Retrieved January 4, 2013] Short interpregnancy spacing in Utah. PRAMS PERSPECTIVES. 2006. Available at: <http://health.utah.gov/mihp/pdf/IPI.pdf>.
9. Nabukera SK, Wingate MS, Kirby RS, Owen J, Swaminathan S, Alexander GR, et al. Interpregnancy interval and subsequent perinatal outcomes among women delaying initiation of childbearing. *J Obstet Gynaecol Res*. 2006; 34:941–947. [PubMed: 19012690]
10. Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities, 2006. *Contraception*. 2011; 84:478–485. [PubMed: 22018121]
11. Lepkowski JM, Mosher WD, Davis KE, Groves RM, van Hoewyk J, Willem J. National Survey of Family Growth, Cycle 6: Sample design, weighting, imputation, and variance estimation. *Vital Health Stat 2*. 2006; 142:1–82. [PubMed: 16999003]
12. Groves RM, Mosher WD, Lepkowski J, Kirgis NG. Planning and development of the continuous National Survey of Family Growth. *Vital Health Stat 1*. 2009; 48:1–64. [PubMed: 20141029]
13. Mosher WD, Jones J, Abma JC. Intended and unintended births in the United States: 1982–2010. *Natl Health Stat Report*. 2012; 55:1–28. [PubMed: 23115878]
14. Klerman, L. Another Chance: Preventing Additional Births to Teen Mothers. Washington, DC: National Campaign To Prevent Teen Pregnancy; 2004.
15. Rigsby DC, Macones GA, Driscoll DA. Risk factors for rapid repeat pregnancy among adolescent mothers: a review of the literature. *J Pediatr Adolesc Gynecol*. 1998; 11:115–126. [PubMed: 9704301]
16. Hamilton BE, Ventura SJ. Birth rates for U.S. teenagers reach historic lows for all age and ethnic groups. *NCHS Data Brief*. 2012; 89:1–8. [PubMed: 22617115]
17. American College of Obstetricians and Gynecologists. Increasing use of contraceptive implants and intrauterine devices to reduce unintended pregnancy. *Obstet Gynecol*. 2009; 114:1434–1438. [PubMed: 20134301]
18. Speidel JJ, Harper CC, Shields WC. The potential of long-acting reversible contraception to decrease unintended pregnancy. *Contraception*. 2008; 78:197–200. [PubMed: 18692608]
19. Cleary-Goldman J, Malone FD, Vidaver J, Ball RH, Nyberg DA, Comstock CH, et al. Impact of maternal age on obstetric outcome. *Obstet Gynecol*. 2005; 105:983–990. [PubMed: 15863534]
20. Menken J, Trussell J, Larsen U. Age and infertility. *Science*. 1986; 233:1389–1394. [PubMed: 3755843]
21. Joyce T, Kaestner R, Korenman S. The stability of pregnancy intentions and pregnancy-related maternal behaviors. *Matern Child Health J*. 2000; 4:171–178. [PubMed: 11097504]

**Table 1**

Demographic and Childbearing Characteristics Associated With Pregnancies \* Conceived After a Previous Birth, 2006–2010.

| Characteristic                                  | n=2,253 |
|---|---------|
| Mean interpregnancy interval length, mo         | 34.0    |
| Interpregnancy interval length, mo              |         |
| Less than 18                                    | 35.1    |
| Less than 6                                     | 6.7     |
| 6–11  | 12.4    |
| 12–17   | 16.0    |
| 18–59   | 49.5    |
| 60 or more                                      | 15.5    |
| Pregnancy intention                             |         |
| Intended  | 61.7    |
| Mistimed  | 20.8    |
| Unwanted  | 17.5    |
| Age at conception of most recent pregnancy, y   |         |
| 15–19   | 5.9     |
| 20–29   | 56.3    |
| 30–44   | 37.9    |
| Age at first birth, y                           |         |
| 15–19   | 35.3    |
| 20–29   | 53.9    |
| 30–44   | 10.9    |
| Prior births                                    |         |
| 1   | 53.1    |
| 2   | 28.5    |
| 3 or more                                       | 18.4    |
| Race or ethnicity                               |         |
| Non-Hispanic white                              | 53.4    |
| Hispanic  | 25.1    |
| Non-Hispanic black                              | 14.7    |
| Non-Hispanic other                              | 6.7     |
| Union status at time of conception              |         |
| Not married or cohabiting                       | 13.9    |
| Married   | 65.7    |
| Cohabiting                                      | 20.4    |
| Education                                       |         |
| Less than high school                           | 24.5    |
| High school graduate or equivalency certificate | 27.2    |
| Some college                                    | 23.4    |
| College graduate or above                       | 24.9    |



| Characteristic           | n=2,253 |
|--------------------------|---------|
| Medicaid funded delivery |         |
| Yes                      | 48.3    |
| No                       | 51.8    |

Data are % except unless otherwise specified.

Figures may not add up due to rounding.

\* Among pregnancies conceived within 5 years of the interview date and ending in a live birth.

**Table 2**

Percent of Pregnancies\* Conceived Within 18 Months of a Previous Birth and Unadjusted Odds Ratios and 95% Confidence Intervals for Short Interpregnancy Intervals by Selected Characteristics, 2006–2010.

| Characteristic                                  | Pregnancies Conceived Within 18 Months of a Previous Birth (%) | Unadjusted OR (95% CI)         |
|---|--|--------------------------------|
| All   | 35.0   |                                |
| Pregnancy intention                             |  |                                |
| Intended  | 25.6   | —                              |
| Mistimed  | 59.7   | 4.30 <sup>†</sup> (3.00, 6.18) |
| Unwanted  | 38.6   | 1.82 <sup>‡</sup> (1.24, 2.70) |
| Age at conception of most recent pregnancy, y   |  |                                |
| 15–19   | 67.1   | —                              |
| 20–29   | 37.2   | 0.29 <sup>†</sup> (0.17, 0.48) |
| 30–44   | 26.8   | 0.18 <sup>†</sup> (0.10, 0.31) |
| Age at first birth, y                           |  |                                |
| 15–19   | 36.8   | 0.61 <sup>§</sup> (0.39, 0.92) |
| 20–29   | 31.1   | 0.47 <sup>†</sup> (0.31, 0.71) |
| 30–44   | 49.0   | —                              |
| Prior births                                    |  |                                |
| 1   | 34.5   | —                              |
| 2   | 33.5   | 0.96 (0.72, 1.28)              |
| 3 or more                                       | 38.9   | 1.21 (0.86, 1.71)              |
| Race or ethnicity                               |  |                                |
| Non-Hispanic white                              | 36.5   | 1.31 (0.97, 1.77)              |
| Hispanic  | 30.5   | —                              |
| Non-Hispanic black                              | 39.6   | 1.49 <sup>§</sup> (1.02, 2.18) |
| Non-Hispanic other                              | 29.7   | 0.96 (0.59, 1.56)              |
| Union status at time of conception              |  |                                |
| Not married or cohabiting                       | 32.3   | —                              |
| Married   | 36.0   | 1.18 (0.84, 1.65)              |
| Cohabiting                                      | 33.5   | 1.06 (0.71, 1.58)              |
| Education                                       |  |                                |
| Less than high school                           | 38.9   | 1.03 (0.69, 1.53)              |
| High school graduate or equivalency certificate | 32.9   | 0.79 (0.55, 1.14)              |
| Some college                                    | 29.8   | 0.69 (0.44, 1.06)              |
| College graduate or above                       | 38.3   | —                              |
| Medicaid funded delivery                        |  |                                |
| Yes   | 38.0   | 1.29 <sup>§</sup> (1.01, 1.65) |
| No  | 32.3   | —                              |

OR, odds ratio; CI, confidence interval.

— indicates reference category.

\* Among pregnancies conceived within 5 years of the interview date and ending in a live birth.

<sup>†</sup> Significant difference at  $P < .001$ .

<sup>‡</sup> Significant difference at  $P < .01$ .

<sup>§</sup> Significant difference at  $P < .05$ .

**Table 3**

Adjusted Odds Ratios and 95% Confidence Intervals From Logistic Regression Analysis Investigating the Association Between Pregnancies \* Conceived Within 18 Months of a Previous Birth and Selected Characteristics, 2006–2010.

| Characteristic                                | Model 1                        | Model 2                        |
|---|--------------------------------|--------------------------------|
|   | Adjusted OR (95% CI)           | Adjusted OR (95% CI)           |
| Pregnancy intention                           |                                |                                |
| Intended                                      | —                              | —                              |
| Mistimed                                      | 4.43 <sup>‡</sup> (3.07, 6.39) | 4.78 <sup>‡</sup> (3.36, 6.86) |
| Unwanted                                      | 2.11 <sup>‡</sup> (1.37, 3.27) | 2.21 <sup>‡</sup> (1.43, 3.41) |
| Age at conception of most recent pregnancy, y |                                |                                |
| 15–19   | —                              | —                              |
| 20–29   | 0.27 <sup>‡</sup> (0.15, 0.48) | —                              |
| 30–44   | 0.12 <sup>‡</sup> (0.06, 0.23) | —                              |
| Age at first birth, y                         |                                |                                |
| 15–19   | —                              | 0.51 <sup>§</sup> (0.28, 0.92) |
| 20–29   | —                              | 0.41 <sup>‡</sup> (0.25, 0.68) |
| 30–44   | —                              | —                              |
| Prior births                                  |                                |                                |
| 1   | —                              | —                              |
| 2   | 1.11 (0.79, 1.57)              | 0.99 (0.72, 1.36)              |
| 3 or more                                     | 1.93 <sup>‡</sup> (1.31, 2.86) | 1.34 (0.89, 2.01)              |
| Race or ethnicity                             |                                |                                |
| Non-Hispanic white                            | 1.35 (0.94, 1.94)              | 1.36 (0.95, 1.94)              |
| Hispanic                                      | —                              | —                              |
| Non-Hispanic black                            | 1.40 (0.93, 2.11)              | 1.59 <sup>§</sup> (1.06, 2.39) |
| Non-Hispanic other                            | 1.00 (0.56, 1.79)              | 0.92 (0.53, 1.59)              |
| Union status at time of conception            |                                |                                |
| Not married or cohabiting                     | —                              | —                              |
| Married                                       | 2.13 <sup>‡</sup> (1.42, 3.19) | 1.85 <sup>‡</sup> (1.23, 2.78) |
| Cohabiting                                    | 1.46 (0.93, 2.30)              | 1.38 (0.90, 2.13)              |
| College graduate or above                     | 2.31 <sup>‡</sup> (1.54, 3.48) | 1.42 (0.94, 2.14)              |
| Medicaid funded delivery                      | 1.14 (0.84, 1.56)              | 1.41 <sup>§</sup> (1.04, 1.91) |

OR, odds ratio; CI, confidence interval.

— indicates reference category.

\* Among pregnancies conceived within 5 years of the interview date and ending in a live birth.

<sup>‡</sup> Significant difference at  $P < .001$ .

<sup>‡</sup> Significant difference at  $P < .01$ .

<sup>§</sup> Significant difference at  $P < .05$ .

**Table 4**

Among Pregnancies \* Conceived Within 18 Months of a Previous Birth, Percent Intended and Unadjusted Odds Ratios and 95% Confidence Intervals For Intended Pregnancies by Selected Characteristics, 2006–2010.

| Characteristic                                  | Among Pregnancies<br>Conceived Within 18<br>Months of a Previous<br>Birth, Intended<br>Pregnancies (%) | Unadjusted OR<br>(95% CI)       |
|---|--|---------------------------------|
| All   | 45.2   |                                 |
| Age at Conception of most recent pregnancy, y   |  |                                 |
| 15–19   | 15.4   | —                               |
| 20–29   | 44.1   | 4.33 <sup>†</sup> (2.08, 9.03)  |
| 30–44   | 59.0   | 7.91 <sup>†</sup> (3.356, 17.6) |
| Age at first birth, y                           |  |                                 |
| 15–19   | 30.7   | 0.19 <sup>†</sup> (0.09, 0.39)  |
| 20–29   | 48.4   | 0.39 <sup>§</sup> (0.18, 0.87)  |
| 30–44   | 70.4   | —                               |
| Prior births                                    |  |                                 |
| 1   | 52.2   | —                               |
| 2   | 35.7   | 0.51 <sup>‡</sup> (0.32, 0.82)  |
| 3 or more                                       | 39.7   | 0.60 (0.35, 1.03)               |
| Race or ethnicity                               |  |                                 |
| Non-Hispanic white                              | 52.1   | 1.85 <sup>§</sup> (1.13, 3.05)  |
| Hispanic  | 37.0   | —                               |
| Non-Hispanic black                              | 31.5   | 0.78 (0.45, 1.37)               |
| Non-Hispanic other                              | 47.7   | 1.55 (0.66, 3.68)               |
| Union status at time of conception              |  |                                 |
| Not married or cohabiting                       | 23.5   | —                               |
| Married   | 54.3   | 3.88 <sup>‡</sup> (1.83, 8.24)  |
| Cohabiting                                      | 27.6   | 1.25 (0.55, 2.85)               |
| Education                                       |  |                                 |
| Less than high school                           | 32.3   | 0.21 <sup>†</sup> (0.11, 0.43)  |
| High school graduate or equivalency certificate | 33.7   | 0.23 <sup>†</sup> (0.11, 0.46)  |
| Some college                                    | 45.0   | 0.37 <sup>§</sup> (0.15, 0.88)  |
| College graduate or above                       | 69.0   | —                               |
| Medicaid funded delivery                        |  |                                 |
| Yes   | 32.6   | 0.34 <sup>†</sup> (0.20, 0.56)  |
| No  | 59.0   | —                               |

OR, odds ratio; CI, confidence interval.

— indicates reference category.

\* Among pregnancies conceived within 5 years of the interview date and ending in a live birth.

<sup>†</sup>Significant difference at  $P < .001$ .

<sup>‡</sup>Significant difference at  $P < .01$ .

<sup>§</sup>Significant difference at  $P < .05$ .