

RESEARCH ARTICLE

Open Access

Purple Acid Phosphatase5 is required for maintaining basal resistance against *Pseudomonas syringae* in *Arabidopsis*

Sridhar Ravichandran¹, Sophia L Stone², Bernhard Benkel³ and Balakrishnan Prithiviraj^{1*}

Abstract

Background: Plants have evolved an array of constitutive and inducible defense strategies to restrict pathogen ingress. However, some pathogens still manage to invade plants and impair growth and productivity. Previous studies have revealed several key regulators of defense responses, and efforts have been made to use this information to develop disease resistant crop plants. These efforts are often hampered by the complexity of defense signaling pathways. To further elucidate the complexity of defense responses, we screened a population of T-DNA mutants in Columbia-0 background that displayed altered defense responses to virulent *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000).

Results: In this study, we demonstrated that the *Arabidopsis Purple Acid Phosphatase5 (PAP5)* gene, induced under prolonged phosphate (Pi) starvation, is required for maintaining basal resistance to certain pathogens. The expression of *PAP5* was distinctly induced only under prolonged Pi starvation and during the early stage of *Pst* DC3000 infection (6 h.p.i). T-DNA tagged mutant *pap5* displayed enhanced susceptibility to the virulent bacterial pathogen *Pst* DC3000. The *pap5* mutation greatly reduced the expression of pathogen inducible gene *PR1* compared to wild-type plants. Similarly, other defense related genes including *ICS1* and *PDF1.2* were impaired in *pap5* plants. Moreover, application of BTH (an analog of SA) restored *PR1* expression in *pap5* plants.

Conclusion: Taken together, our results demonstrate the requirement of *PAP5* for maintaining basal resistance against *Pst* DC3000. Furthermore, our results provide evidence that *PAP5* acts upstream of SA accumulation to regulate the expression of other defense responsive genes. We also provide the first experimental evidence indicating the role *PAP5* in plant defense responses.

Keywords: *Arabidopsis*, Plant defense responses, *PAP5*, *Pseudomonas syringae*, Phosphate starvation

Background

Plants are continuously exposed to a diverse array of microorganisms including beneficial mutualists, commensals, and pathogens. To defend against pathogens, plants have evolved an innate immune system to recognize and limit infection (reviewed in [1,2]). Activation of defense responses involves the initial recognition of pathogens by chemical cues (elicitors) or Pathogen Associated Molecular Patterns (PAMPs) that include bacterial lipopolysaccharides, flagellin, fungal chitin and ergosterol [3,4]. Recognition of PAMP by specific Pattern Recognition

Receptors (PRRs) in the plasma membrane leads to activation of defense responses in both non-host and basal disease resistance [5]. Activation of PRRs subsequently induces the calcium-dependent protein kinase (CDPK) and mitogen-activated protein kinase (MAPK) signaling pathways leading to rapid ion fluxes, followed by transcriptional activation of defense responsive genes and synthesis of antimicrobial compounds to restrict infection [6,7].

Primarily, regulation of plant defense responses is mediated through the phytohormones salicylic acid (SA), jasmonic acid (JA) and ethylene (ET) [8,9]. However, in recent years other phytohormones including abscisic acid (ABA), auxins, gibberellins (GA), cytokines (CK) and brassinosteroids (BR) have been shown to mediate specific plant defense responses (reviewed in [2,10]). As

* Correspondence: bprithiviraj@dal.ca

¹Department of Environmental Sciences, Faculty of Agriculture, Dalhousie University, Truro, NS B2N 5E3, Canada

Full list of author information is available at the end of the article

plants are exposed to an array of pathogens with diverse infection strategies, activation of appropriate, pathogen-specific defense responses is vital for plant growth and productivity [11].

Plant pathogens are classified as biotrophs, necrotrophs or hemi-biotrophs based on their life style and infection strategy. Biotrophic pathogens live as obligate parasites that derive nutrients from living host tissues, while necrotrophs feed on dead tissues. Hemi-biotrophs behave as both biotroph and necrotroph depending on the stage of their life cycle [11]. Defense against biotrophs involves SA-dependent responses whereas necrotroph resistance is SA-independent relying primarily on JA/ET-dependent pathways [9]. The SA signaling pathway is associated with transcriptional activation of pathogenesis related (PR) genes and the establishment of systemic acquired resistance (SAR) to provide enhanced, long lasting resistance to secondary infections [12,13]. By contrast, JA/ET signaling pathways are associated with resistance against necrotrophic pathogens and rhizobacteria-mediated induced systemic resistance (ISR), and are not typically associated with PR gene expression [12,14]. However, there are complex signaling and cross talk between the SA-dependent and SA-independent pathways [13].

Genetic screening of mutant plant populations has proved very useful for the functional analysis of defense responses [15-17]. In *Arabidopsis*, genetic screening has revealed a large number of mutants that exhibit altered responses to SA, JA and/or ET and are more susceptible to virulent pathogens [18]. Identification and characterization of enhanced disease susceptibility (*eds*) mutants, including a series of phytoalexin deficient (*pad*) mutants, have helped to elucidate a number of defense signaling pathways involved in both basal and induced defense responses [19-21].

Purple Acid Phosphatases (PAPs) belong to a family of binuclear metalloenzymes that exhibit diverse biological functions in plants, animals and bacterial species [22,23]. While the predominant role of PAPs in plants is regulation of Pi uptake, PAPs also contribute to other biological functions including peroxidation [24], ascorbate recycling [25], mediation of salt tolerance [26] and regulation of cell wall carbohydrate biosynthesis [27]. Plant PAPs share significant sequence similarity with mammalian tartarate-resistant acid phosphatases (TRAPs), which are involved in bone resorption [28], iron transport [29] and also in the generation of reactive oxygen species for microbial killing [30]. In humans, TRAP expression is restricted to activated macrophages where it aids in the generation of free radicals to enhance microbial killing [31]. Although numerous reports have emphasized the importance of PAPs in Pi acquisition, it has been difficult to assign a general physiological role to PAPs due to their diversity [32]. The *Arabidopsis*


genome contains 29 *PAP* encoding genes [33]. Changes in *PAP* gene expression differs in response to Pi concentration where PAP11 and PAP12 are transcriptionally induced while PAP7-PAP10 and *PAP13* remain unchanged in response to Pi deprivation [33]. Kaffarnik and colleagues first reported the accumulation of PAP10 and a decrease in the abundance of PAP14 in the secretome of *Arabidopsis* cell culture following *P. syringae* infection, suggesting a role for PAPs in the host defense response [34]. Recently, Li *et al.*, (2012) also provided the evidence that some soybean PAPs (GmPAPs) are involved in symbiosis under Pi starved conditions. PAPs carry predicted signal peptides and presumably are secreted, however the biological function of these proteins in the extracellular space is unknown [34].

Here we provide evidence that the *Arabidopsis PAP5* is involved in basal resistance against certain plant pathogens. *PAP5* mutant plants exhibited enhanced susceptibility to virulent isolate of *Pseudomonas syringae* pv. *tomato* DC3000. In addition, expression of defense related genes following *Pst* DC3000 infection were impaired in *pap5* plants.

Results

Identification of mutants exhibiting altered defense responses

One thousand two hundred unique *Arabidopsis thaliana* (ecotype Col-0) T-DNA insertion lines were spray inoculated with the virulent isolate of *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) and monitored for altered responses to the pathogen. Mutants exhibiting extensive chlorosis in comparison to wild-type plants, scored by visual examination, were designated as susceptible. Mutants exhibiting reduced chlorosis compared to wild type (Col-0) were designated resistant to *Pst* DC3000. T-DNA insertion lines were also tested for altered root colonization with the plant growth promoting rhizobacterial isolate *Pseudomonas putida* WCS358. Selected T-DNA lines were retested for response to *Pst* DC3000. A total of 24 T-DNA insertion lines exhibited either altered disease susceptibility, root colonization or both compared to wild-type plants (data not shown). The mutant line *salk_126152C* (*pap5-1*), which exhibited enhanced susceptibility to *Pst* DC3000 with extensive chlorosis on leaf tissues, was selected for further analysis (Figure 1A). *salk_126152C* carried a T-DNA insertion in the gene coding for *Purple Acid Phosphatase5* (*PAP5*; At1G52940) (Genome-Wide Insertional Mutagenesis of *Arabidopsis thaliana*, 2003). The enhanced susceptibility phenotype of *pap5-1* plants was confirmed by assessing bacterial growth in leaf tissues post inoculation. As shown in Figure 1B, *pap5-1* plants had greater titers of bacteria at 48 and 72 hours post inoculation (h.p.i) compared to the wild-type plant. To ensure that the altered responses to the pathogen were caused by disruption of



the *PAP5* gene and not by an unlinked mutation, a second knockout mutant line *salk_081481C* (*pap5-2*), carrying a T-DNA insertion on *PAP5* (At1g52940), was tested. *pap5-2* plants also exhibited the extensive chlorosis and higher titer of bacteria similar to that of in *pap5-1* plants (Additional file 1: Figure S1).

Further characterization of *pap5-1* mutant plants

Genotyping via polymerase chain reaction (PCR) confirmed that *pap5-1* (*salk_126152C*) carries a T-DNA insertion within the first intron (Figure 2A and 2B). To determine the impact of T-DNA insertion on transcript levels of *PAP5*, Reverse Transcription-quantitative PCR (RT-qPCR) was performed using gene specific primers (Figure 2A). Most PAPs are reported to be highly inducible under phosphate starvation (Pi). In our experiments, we did not observe an induction of *PAP5* in wild-type seedlings grown in the presence of phosphate (1.25 mM) or under phosphate starved conditions for 5 days (-Pi, 0 mM) (data not shown). We also observed

that the expression of *PAP5* under optimal growing conditions was very low and this was confirmed with *PAP5* expression profile in the comprehensive microarray site <https://www.genevestigator.com/gv/> (Additional file 2: Figure S2). Interestingly, we observed a marked increase in the expression *PAP5* when wild type seedlings were grown under prolonged phosphate starvation (Figure 2C). For prolonged Pi starvation wild-type seedlings were germinated in media containing reduced Pi (0.25 mM) for seven days and then transferred to media with no Pi (0 mM). After 9 days the seedlings were harvested for gene expression analysis. RT-qPCR analysis revealed a ~30 fold increase in transcript levels of *PAP5* in wild-type seedlings grown under prolonged phosphate starvation (-Pi) compared to seedlings grown in the presence of phosphate (+Pi) (Figure 2C). The expression of *PAP5* was not induced in both *pap5-1* (Figure 2C) and *pap5-2* (Additional file 3: Figure S3B) seedlings grown under prolonged phosphate starvation (-Pi). We did not observe any major alteration in germination, growth and development of *pap5* mutant plants compared to wild-type under optimal growth conditions (data not shown).

Mutation in *PAP5* alters expression of host defense responsive genes and ROS production

To explore the enhanced susceptibility of *pap5-1* plants and to determine the role of *PAP5* in host defense responses, plants were spray inoculated with virulent isolate of *Pst* DC3000 (10^8 c.f.u ml⁻¹) and the transcript abundances of selected defense responsive genes, including the pathogenesis-related gene1 (*PR1*), were determined. Infection of wild-type plants with the virulent isolate *Pst* DC3000 resulted in ~10-fold induction of the *PR1* transcript 24 h.p.i, while an increase of only ~2-fold was observed in *pap5-1* plants (Figure 3). The level of *PR1* transcripts in *pap5-1* plants following *Pst* DC3000 infection was variable at 48 h.p.i. However, the expression of *PR1* was a still less induced in *pap5-1* plants compared to wild-type (Figure 3). Expression of *isochorismate synthase1* (*ICS1*) was induced in wild-type plants (~2-fold) while no increase in transcript levels was observed in *pap5-1* plants. Although, expression of *plant defensin1.2* (*PDF1.2*) was induced (~2-fold higher) in wild-type plants, expression of *PDF1.2* was suppressed in *pap5-1* plants (Figure 4A). The expression pattern of these pathogenesis related genes were also confirmed using *Actin* as the internal control (Additional file 4: Figure S4).

A marked increase in the expression of *PAP5* at 6 h.p.i was observed in wild-type plants (Figure 4B). However, this difference did not prolong to 24 and 48 h.p.i. We did not observe induction of *PAP5* in mock infected or *Pst* DC3000 inoculated *pap5-1* plants (Figure 4B). The expression profile of *PAP5* was further verified from the comprehensive microarray site <http://bar.utoronto.ca/> using




Figure 2 Validation of T-DNA insertion in *pap5-1* plants. **A**, Schematic representation of *AtPAP5* (At1G52940); white boxes and solid lines represent exons and introns. T-DNA insertion is represented with a grey arrow and the solid arrows represent the primers used for genotyping and quantitative RT-qPCR. **B**, Location of the T-DNA insertion and homozygosity of *pap5-1* was confirmed by PCR using the gDNA from wild-type and *pap5-1* plants (M, 100 bp marker). A 30 cycle PCR reactions was performed with the primer pairs indicated. **C**, Relative expression of *PAP5* transcripts in response to prolonged Pi starvation; For prolonged Pi starvation wild type and *pap5-1* seedlings were germinated and grown in 0.5X MS media containing reduced Pi (0.25 mM). After seven days the seedlings were washed with sterile water and transferred to 0.5X MS with no Pi (0 mM). After 9 days the seedlings were harvested for gene expression analysis. Total RNA was extracted from wild-type and *pap5-1* plants as described in Methods. Transcript levels of *PAP5* was normalized to the expression of *GAPDH* in the same samples and expressed relative to the normalized transcript levels of Pi starved wild-type plants. The bars represent the mean and standard deviation from two independent experiments. Asterisks represents data sets significantly different from the wild-type data sets ($P < 0.05$ using one-tailed Student's *t*-test).


Arabidopsis eFP Browser (Additional file 5: Figure S5) [35]. Although, *PAP5* was strongly induced only at 6 h.p.i, our results suggest that this level of *PAP5* is required for maintaining resistance against virulent *Pst* DC3000.

To further explore the mechanism of enhanced susceptibility, we studied hydrogen peroxide (H_2O_2) accumulation using 3-3'-Diaminobenzidine (DAB) staining. As shown in Figure 5A, accumulation of H_2O_2 in response to *Pst* DC3000 was reduced in *pap5-1* leaves at 24 and 48 h.p.i. In contrast, there was an accumulation of H_2O_2 in the wild-type plants. The H_2O_2 concentration was quantified in leaf tissues following *Pst* DC3000 infection. The wild-type plants accumulated a higher

concentration of H_2O_2 in response to *Pst* DC3000 inoculation as compared to *pap5-1* plants (Figure 5B).

Resistance to *Botrytis cinerea* is affected in *pap5* plants

Having demonstrated the enhanced susceptibility of *pap5-1* plants to the hemi-biotrophic pathogen *Pst* DC3000, we next tested the level of resistance of *pap5-1* plants to the necrotrophic pathogen *Botrytis cinerea*. Four week old plants were inoculated with spore suspension of *B. cinerea* and lesion size was measured three days later. As shown in Figure 6A, *pap5-1* plants developed a significantly larger lesion (5.4 ± 0.3 mm) than the wild-type (3.9 ± 0.2 mm). The greater lesion size on




pap5-1 plants in response to *B. cinerea* infection, suggests the role of *PAP5* are important in limiting fungal growth.

To identify the role of *PAP5* in the resistance against *B. cinerea*, we assessed the transcript abundance of *PR1* and *PDF1.2*. As shown in Figure 7A, *B. cinerea* strongly induced the expression of *PR1* in both wild-type and *pap5*. In contrast, the level of the *PDF1.2* transcript at 24 h.p.i was only half of that observed in wild-type plants (Figure 7B). By 48 h.p.i., however, the transcript levels of *PDF1.2* were similar in both wild-type and *pap5-1* plants. Similarly, we did not observe any significant differences in *PAP5* transcripts with *B. cinerea* infection (Figure 6B).

Responses to exogenous application of BTH, a salicylic acid analog and methyl jasmonate (MJ) is unaffected in *pap5* plants

Since *pap5-1* plants exhibited enhanced susceptibility to *Pst* DC3000 and *B. cinerea*, we investigated the role of *PAP5* in responses to BTH and MJ. Exogenous application of BTH induced higher levels of *PR1* in wild-type and *pap5-1* (Figure 8A). We also observed a slightly higher increase in the expression *PR1* in *pap5-1* plants 24 h after BTH treatment. Similarly, application of MJ strongly induced the expression of *PDF1.2* in both wild-type and *pap5-1* plants. We did not observe significant differences in expression of *PDF1.2* between wild-type and *pap5-1* plants following application of MJ (Figure 8B). Application of BTH and JA induced expression of *PR1* and *PDF1.2*, respectively, indicative of an intact JA signaling pathway in *pap5* plants. Based on these experiments it was clear that



pap5-1 plant was not defective in responding to exogenously applied BTH or MJ.

Discussion

In this study, we demonstrated the role of *PAP5*, a phosphate responsive gene, and its requirement in maintaining basal disease resistance against virulent *Pst* DC3000. In previous studies *PAP5* transcripts were not detectable under phosphate starvation [36]. Unlike *PAP12* and *PAP26*, *PAP5* is not abundantly expressed under normal phosphate starvation conditions. Our results revealed that *PAP5* is expressed only under prolonged Pi starvation (Figure 2C). Mutation in *PAP26* has been shown to impair






Figure 5 *pap5-1* plants accumulate reduced H₂O₂ in response to *Pst* DC3000 infection. **A**, Histochemical detection of H₂O₂ accumulation using DAB staining. Wild-type and *pap5-1* leaves were excised following *Pst* DC3000 infection and stained with DAB for hydrogen peroxide. **B**, Quantification of H₂O₂ following *Pst* DC3000 infection. The bars represent mean and SD of H₂O₂ accumulation. Asterisks represent significant difference in H₂O₂ production compared to wild type (Student's *t*-test; *P* < 0.05).

growth and increase anthocyanin accumulation in response to Pi starvation [37]. Despite the loss of *PAP5* expression, mutant plants did not show discrete phenotypic differences from that of wild-type plants. Both wild-type and *pap5-1* plants exhibited an increased root/shoot ratio under Pi starvation (data not shown). This finding also indicates that *PAP5* does not play a major role in Pi acquisition and is more likely to regulate other functions. The Arabidopsis genome contains 29 *PAP* encoding genes [33] and this may lead to functional redundancy. This study suggests that the loss of *PAP5* resulted in impairment of defense responsive genes in response to *Pst* DC3000 infection. Further, it appears that other *PAP* genes does not compensate for the loss of *PAP5* function in response to pathogen attack.

Genetic analyses of Arabidopsis mutants have revealed many key regulatory genes in plant defense responses. Enhanced disease susceptibility mutants including *eds5*, *pad4*, *npr1* and *sid2* have previously been reported to exhibit enhanced susceptibility and compromised defense responses to both virulent and avirulent isolates of *Pst*

DC3000 [19,20,38]. It is also evident that most bacterial pathogens including *Pst* DC3000 are inoculated by pressure-infiltration to study plant-bacterial interactions. Although pressure-infiltration is the most commonly used inoculation method, these inoculation procedures may prevent early innate immune responses such as flagellin perception (FLS2 mediated resistance) [39] and stomatal closure [40]. Also FLS2 mediated resistance was effective only when *Pst* DC3000 was sprayed on the leaf surface and not when bacteria was infiltrated in to leaves [39]. Hence, to mimic natural infection and to focus on the early defense responses we sprayed plants with suspension of *Pst* DC3000 containing 10⁸ c.f.u ml⁻¹. We also observed that plants sprayed with 10³ and 10⁵ cells/ml⁻¹ developed reduced symptoms compared to plants sprayed with 10⁸ cells (data not shown). Similar bacterial titers have been previously used for plant-bacterial interaction studies [39,40].


We observed that the expression of *PR1* was slightly induced in *pap5-1* plants following *Pst* DC3000 infection, however the relative transcript level of *PR1* was



several fold lower compared to wild-type (Figure 3). The *PR1* transcripts at 48 h.p.i were slightly lower compared to wild-type (Figure 3). Similar variability has been observed in MPK6 silenced plants that were susceptible to *Pst* DC3000 [41]. We observed that *PAP5* was strongly induced in the early stages of infection (6 h.p.i). This induction was transient as no difference was observed at 24 and 48 h.p.i. One possible explanation of this observation is that the level of *PAP5* induced during the early stages (6 h.p.i) of infection could be sufficient to dephosphorylate signaling proteins required for activation of defense responses downstream of *PAP5*. Thus, it is also possible that *PAP5* might be involved in early responses to pathogens similar to glutathione *S*-transferase (*GST6*) and glucosyltransferase [42]. Moreover, members of the PAP family have been known to exhibit peroxidase activity in addition to Pi acquisition and recycling [24,43].

Although, the role of *PAP5* with regard to peroxidase activity has not been established, we hypothesize that the *PAP5* might mediate generation of reactive oxygen species (ROS) during *Pst* DC3000 infection. ROS was initially proposed to mediate plant defense response especially, during an incompatible interaction [44]. Virulent pathogens, capable of evading pathogen recognition are also known to induce ROS production at latter stages of infection to lower levels [1].

We also identified the importance in *PAP5* in limiting the growth of the necrotrophic fungus, *B. cinerea* at the site of infection. The expression of *PDF1.2* was strongly suppressed in *pap5-1* plants at 24 h.p.i resulting in an increase in lesion size. There were no differences in *PDF1.2* transcripts between *pap5* and wild-type plants at 48 h.p.i. Similarly, *eds4-1* plants have been reported to exhibit enhanced susceptibility to *B. cinerea* despite



comparable expression of *PR1* and *PDF1.2* transcripts [45]. These results also suggest that defense responsive genes other than *PR1* and *PDF1.2* are required to mount wild-type levels of resistance against *B. cinerea*. SA synthesized in response to *B. cinerea* infection has reported to be derived via phenylalanine ammonia lyase (*PAL*) and not via isochorismate synthase (*ICS*) [46]. Since *pap5-1* plants induced comparable levels of *PR1* to wild-type plants following *B. cinerea* infection, it is possible that the effect of *PAP5* is restricted to SA derived via *ICS* and not via *PAL*.

Application of BTH and MJ in wild-type and *pap5* plants induced expression of *PR1* and *PDF1.2*, respectively (Figure 8A and 8B). These results also suggest that *PAP5* is not required for expression of SA dependent *PR1* expression. *PR1* expression in *pap5-1* plants appeared to be slightly higher than wild-type plants after

of BTH treatment (Figure 8A). This slight increase in *PR1* expression and its significance is unclear. Similarly, application of SA on *pad4* plants showed a slight increase in *PR1* expression [47]. Application of MJ induced the expression of *PDF1.2*, indicating the regulatory function of *PAP5* to be upstream of SA and JA.

Although, most plant PAPs are primarily associated with Pi absorption and recycling, PAPs induced under Pi starvation are also known to exhibit peroxidase activity similar to mammalian PAPs [24,48]. All mammalian PAPs characterized exist as monomers of ~35 kDa (Low Molecular Weight, LMW), while plants encode a relatively large family of High Molecular Weight (HMW) homodimeric and oligomeric PAPs (~45-74 kDa). However, a recent study has identified mammalian-like low molecular weight PAP (~34 kDa) from roots of Pi starved bean plants [32]. Moreover, the LMW, 35 kDa plant PAPs are reported to be closely related to the 35 kDa mammalian PAPs than to the large plant PAPs [49]. Thus, from our results we hypothesize that *PAP5* could play a role in both Pi acquisition and in microbial killing during pathogenesis (Figure 9).

Conclusion

We identified the requirement of *PAP5* for maintaining basal defense responses against virulent *Pst* DC3000, suggesting a role for *PAP5* in pathogen triggered immunity (PTI). We further demonstrated that *PAP5* acts upstream of SA to affect the expression of *PR1*, and levels of *PAP5* do not affect BTH and JA perception. Further analysis on *pap5* plants is likely to reveal novel components of signal transduction pathways that regulate defense responsive genes.

Methods

Biological materials and growth conditions

Arabidopsis thaliana (L.) Heynh, ecotype Columbia (Col-0) seed was purchased from Lehle seeds (Round Rock, TX, USA) and T-DNA insertion mutant lines were obtained from Arabidopsis Biological Resource Center (Columbus, OH, USA). Seeds were surface sterilized with NaOCl 2% (v/v), rinsed five times with sterile water and stratified at 4°C for 3 days. Seeds were planted either in Jiffy peat pellets (Halifax seeds, Canada) or on plates with 0.5X MS media [50]. Plants were grown at 22 ± 2°C with a photoperiod of 16 h light at 125 μmol m⁻² s⁻¹ and 8 h dark cycle.

Virulent *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) was kindly gifted by Dr. Diane Cuppels, Agriculture and Agri Food Canada (AAFC), ON, Canada. *Pseudomonas syringae* strain was maintained on King's medium B supplemented with rifampicin (50 μg ml⁻¹). *Botrytis cinerea* was cultured on modified King's medium B (10 g peptone, 1.5 g potassium phosphate monobasic, 15 g dextrose, pH 5.5, 5 ml of 1 M MgSO₄/l).




Figure 9 Model for role of PAP5 during *Pst* DC3000 infection.
 When plants are infected with virulent *Pst* DC3000, PAP5 is required for activation of defense responsive genes including *PR1* and *ICS1*. Recognition of *Pst* DC3000, induce expression of *PAP5* only during the early stages of infection (6 h) and triggers ROS synthesis which subsequently activates other defense related signals down stream for complete resistance.

For plant treatment, Benzothiazolidone (Actigard®; active ingredient 50% w/v BTH) was a gift from Syngenta Corp., USA. Methyl jasmonate and other microbiological media were purchased from Sigma Aldrich, Oakville, Canada.

Mutant screening and pathogen inoculation

Genetic screen was performed on 4 to 5 week old plants by spray inoculation with bacterial suspension of virulent *Pst* DC3000. Plant inoculation and bacterial growth in plant apoplast was determined as described by [39]. In brief, strains of virulent *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) was cultured in King's medium B supplemented with rifampicin (50 µg ml⁻¹) at 28°C until OD₆₀₀ of 0.8. Bacterial cells were collected by centrifugation and resuspended in water containing 0.02% Silwet L-77 (Lehle seeds, USA) to a final concentration of 10⁸ c.f.u ml⁻¹. Plants (4-5 weeks old) were spray inoculated and kept under high humidity for disease development. Leaves were excised (8-10 replicates) from different infected plants and were surface sterilized with ethanol (75% v/v). Four to five samples were made by pooling 2 leaf discs (0.5 cm²) and the samples were ground in sterile water with microfuge tube pestle. The ground tissues were serially diluted and plated on King's B medium containing rifampicin (50 µg/ml). The plates were incubated at 28°C and colonies were counter after 48 hours. For *Pst* DC3000 induced gene expression,

plants were spray inoculated with bacterial suspension (10⁸ c.f.u ml⁻¹) and leaf tissues were frozen in liquid nitrogen at the time points indicated.

For *Botrytis cinerea* (*Bcr*) inoculation, spore suspension (1 × 10⁵ conidia mL⁻¹) was prepared in potato dextrose broth (PDB) as described by [45]. Four to five week old plants were inoculated by placing 5 µl of the spore suspension on either side of the mid vein of fully expanded leaves. Inoculated plants were covered with a transparent plastic dome to maintain high humidity for disease development. For all gene expression analysis, leaf tissues were harvested from four individual plants for each biological replicate and were snap frozen in liquid nitrogen for RNA extraction.

Benzothiazolidone (BTH) and methyl jasmonate (MJ) treatments were performed by spraying 4-5 weeks old plants with solutions containing 0.06% w/v Actigard® (Active ingredient: 50% w/v BTH) or 50 µM methyl jasmonate (MJ) with 0.02% Silwet L-77.

Confirmation of T-DNA insertion

T-DNA insertion and homozygosity of mutant line salk_126152 was confirmed by PCR as described by [51] using *AtPAP5* gene specific primers generated from SALK T-DNA verification primer design tool LP 5'-TTCACGGTTTGTGTTAGACG-3'; RP 5'-TCGTTGAAACTACTCGATTAAAC-3' and left border primer LBb1.3 5'-ATTTTGCCGATTCGGAAC-3'.

Phosphate starvation

Sterile, stratified seeds (20-25 per jar) were dispensed in 50 ml of liquid 0.5X MS medium containing Pi (1.25 mM) or with reduced Pi (0.25 mM). The seedlings were grown under constant shaking (85 rpm) at 22 ± 2°C under continuous illumination at 100 µmol m⁻² s⁻¹. After 9 days the seedlings were rinsed thrice with sterile distilled water and transferred to 0.5X MS medium containing + Pi (1.25 mM) or -Pi (0 mM) [52]. Plants were harvested after 11 days for RNA extraction.

Table 1 Primer sequences used in RT-qPCR experiments

Gene	Locus	Primer sequences (5'-3')
<i>GAPDH</i>	At1g13440	TTGGTGACAACAGGTC AAGCA
		AAACTTGTGCGCTCAATGCAATC
<i>ICS1</i>	At1g74710	GCGTCGTTCCGGTTACAGG
		ACAGCGAGGCTGAATATCAT
<i>PAP5</i>	At1g52940	AACAGGTCGCTCCACTAGACA
		TGGTTAGAGGCATATGTTGTCC
<i>PDF1.2</i>	At5g44420	GTTCTCTTTGCTGCTTTCGAC
		GCAAACCCCTGACCATGT
<i>PR1</i>	At2g14610	TGATCCTCGTGGGAATTATGT
		TGCATGATCACATCACTTTCAT

Whenever Pi was reduced from growth medium, equivalent amounts of sulphate salts were added to maintain the concentration of conjugate cations.

RNA extraction and quantitative Real-time PCR

Total RNA was extracted from frozen tissues using monophasic extraction method [53]. Reverse Transcription was performed with 2 µg of total RNA using Quantiscript RTase (Qiagen, ON, Canada). Relative transcript levels were assayed by Real-Time PCR using gene specific primers (Table 1) on StepOnePlus Real-Time PCR system (Applied Biosystems, ON, Canada) using SYBR Green reagent (Applied Biosystems, ON, Canada). To determine the relative expression levels, the amount of target gene was normalized over the abundance of constitutive *Glyceraldehyde 3-phosphate dehydrogenase (GAPDH)* or *Actin* as endogenous control. Primers were generated using the Roche Universal Probe Library assay design center.

DAB staining

To visualize H₂O₂ production *in situ*, plants were inoculated with suspension of *Pst* DC3000 as described in earlier section. Leaves were excised at 24 and 48 h.p.i and stained with 3-3 Diaminobenzidine (DAB) as described by [54]. Excised leaves were placed in DAB (1 mg/ml) solution for 8-12 hours and the tissues were soaked in ethanol (95%, v/v) to remove chlorophyll. For H₂O₂ quantification, the excised leaf tissues were frozen and ground with liquid nitrogen. To 50 mg of ground frozen tissue 500 µl of phosphate buffer (50 mM, sodium phosphate, pH-7.4) was added. The samples were centrifuged and 50 µl of the aliquot was used for H₂O₂ quantification using an Amplex red hydrogen peroxide/peroxidase assay kit (Molecular Probes, Life Technologies, Canada).

Additional files

Additional file 1: Figure S1. Enhanced susceptibility of *pap5-2* to *Pst* DC3000. A, Phenotype of *pap5-2* plants exhibiting extensive chlorosis. Plants were spray inoculated with 10⁸ c.f.u ml⁻¹ and photographed after 5 days of infection. B, Growth of virulent *Pst* DC3000 in wild type (Col-0) and *pap5-2* mutant leaves. Plants were spray inoculated with *Pst* DC3000 (10⁸ c.f.u ml⁻¹) and bacterial growth in plant apoplast was determined. The bars represent the mean and standard deviation from values of six to eight replicate samples and the experiment was repeated two times with similar results. An asterisk indicates significant increase in *Pst* DC3000 growth compared to wild-type (Student's *t*-test; *P* < 0.05).

Additional file 2: Figure S2. Expression profile of *PAP5* (array element 261341_s_at) in comparison to *PRT1* (array element 266385_at) from Genevestigator Expression Data.

Additional file 3: Figure S3. Validation of T-DNA insertion in *pap5-2* mutant plants. A, Schematic representation of *AtPAP5* (At1G52940); white boxes and solid lines represent exons and introns. T-DNA insertion is represented with a grey arrow and the solid arrows represent the primers used for genotyping and quantitative RT-qPCR. B, Location of the T-DNA insertion and homozygosity of *pap5-2* was confirmed by PCR using the

gDNA from wild-type and *pap5-2* plants (M, 100 bp marker). A 30 cycle PCR reactions was performed with the primer pairs indicated. C, Relative expression of *PAP5* transcripts in response to Pi starvation; Total RNA was extracted from wild-type and *pap5* plants as described in materials and methods. Transcript levels of *PAP5* was normalized to the expression of *GAPDH* in the same samples and expressed relative to the normalized transcript levels of Pi supplemented wild-type plants. The bars represent the mean and standard deviation from two independent experiments. Asterisks represents data sets significantly different from the wild-type data sets (*P* < 0.05 using one-tailed Student's *t*-test).

Additional file 4: Figure S4. Expression of defense related genes in wild-type and *pap5-1* mutant plants after *Pst* DC3000 infection. Transcript levels of *PR1*, *ICS1*, *PDF1.2* and *PAP5* in wild-type and *pap5-1* plants were quantified after spray inoculation with virulent *Pst* DC3000 (10⁸ c.f.u ml⁻¹) was determined. Total RNA was extracted from leaf tissues harvested 24 h.p.i. Transcript levels were normalized to the expression of *Actin* in the same samples. The transcript levels were expressed relative to the normalized transcript levels of mock infected wild-type plants. The bars represent the mean and standard deviation. Significant differences (*P* < 0.05) are indicated by different letters.

Additional file 5: Figure S5. Expression profile of *PAP5* (At1g52940) from the Arabidopsis eFP Browser.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BP conceived the concept of genetic screening and designed all experiments. SR carried out all the experiments and prepared the manuscript. SLS critically evaluated all the experiments and significantly contributed to the manuscript preparation. BB helped with sequencing and contributed to evaluate the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Ms. Kalyani Prithiviraj (Faculty of Agriculture, Dalhousie University) for her assistance with Real-Time PCR analysis and Dr. Hongxia Liu (Department of Biology, Dalhousie University) for her help and advice with our experiments. This work was supported by Natural Science and Engineering Research Council of Canada Discovery grant to B. Prithiviraj.

Author details

¹Department of Environmental Sciences, Faculty of Agriculture, Dalhousie University, Truro, NS B2N 5E3, Canada. ²Department of Biology, Dalhousie University, Halifax, NS B3H 4J1, Canada. ³Department of Plant and Animal Sciences, Faculty of Agriculture, Dalhousie University, Truro, NS B2N 5E3, Canada.

Received: 15 February 2013 Accepted: 24 July 2013

Published: 29 July 2013

References

1. Jones JDG, Dangl JL: **The plant immune system.** *Nature* 2006, **444**(7117):323–329.
2. Pieterse CMJ, Leon-Reyes A, Van de, Van Wees SCM: **Networking by small-molecule hormones in plant immunity.** *Nat Chem Biol* 2009, **5**(5):308–316.
3. Zipfel C, Felix G: **Plants and animals: a different taste for microbes?** *Curr Opin Plant Biol* 2005, **8**(4):353–360.
4. Boller T, Felix G: **A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors.** *Annu Rev Plant Biol* 2009, **60**(1):379–406.
5. Bittel P, Robatzek S: **Microbe-associated molecular patterns (MAMPs) probe plant immunity.** *Curr Opin Plant Biol* 2007, **10**(4):335–341.
6. Chisholm ST, Coaker G, Day B, Staskawicz BJ: **Host-microbe interactions: shaping the evolution of the plant immune response.** *Cell* 2006, **124**(4):803–814.
7. Shan L, He P, Sheen J: **Intercepting host MAPK signaling cascades by bacterial type III effectors.** *Cell Host Microbe* 2007, **1**(3):167.
8. Liu P, Yang Y, Pichersky E, Klessig DF: **Altering expression of benzoic acid/salicylic acid carboxyl methyltransferase 1 compromises systemic**

- acquired resistance and PAMP-triggered immunity in arabidopsis. *Mol Plant Microbe Interact* 2010, **23**(1):82–90.
9. Robert-Seilaniantz A, Navarro L, Bari R, Jones JD: Pathological hormone imbalances. *Curr Opin Plant Biol* 2007, **10**(4):372–379.
 10. Bari R, Jones JDG: Role of plant hormones in plant defence responses. *Plant Mol Biol* 2009, **69**(4):473–488.
 11. Glazebrook J: Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* 2005, **43**(1):205–227.
 12. Grant M, Lamb C: Systemic immunity. *Curr Opin Plant Biol* 2006, **9**(4):414–420.
 13. Vlot AC, Dempsey D, Klessig DF, Vlot AC: Salicylic acid, a multifaceted hormone to combat disease. *Annu Rev Phytopathol* 2009, **47**(1):177–206.
 14. Ton J, Van Pelt J, Van Loon L, Pieterse C: Differential effectiveness of salicylate-dependent and jasmonate/ethylene-dependent induced resistance in *Arabidopsis*. *Mol Plant Microbe Interact* 2002, **15**(1):27.
 15. Ausubel FM, Katagiri F, Mindrinos M, Glazebrook J: Use of *Arabidopsis thaliana* defense-related mutants to dissect the plant response to pathogens. *Proc Natl Acad Sci USA* 1995, **92**(10):4189–4196.
 16. Alonso JM, Ecker JR: Moving forward in reverse: genetic technologies to enable genome-wide phenomic screens in *Arabidopsis*. *Nat Rev Genet* 2006, **7**(7):524–536.
 17. Petersen LN, Ingle RA, Knight MR, Denby KJ: OX11 protein kinase is required for plant immunity against *Pseudomonas syringae* in *Arabidopsis*. *J Exp Bot* 2009, **60**(13):3727.
 18. Volkov SM, Boller T, Ausubel FM: Isolation of new *Arabidopsis* mutants with enhanced disease susceptibility to *Pseudomonas syringae* by direct screening. *Genetics* 1998, **149**(2):537–548.
 19. Glazebrook J, Ausubel FM: Isolation of phytoalexin-deficient mutants of *Arabidopsis thaliana* and characterization of their interactions with bacterial pathogens. *Proc Natl Acad Sci USA* 1994, **91**(19):8955–8959.
 20. Rogers EE, Ausubel FM: *Arabidopsis* enhanced disease susceptibility mutants exhibit enhanced susceptibility to several bacterial pathogens and alterations in PR-1 gene expression. *Plant Cell* 1997, **9**(3):305–316.
 21. Wiermer M, Feys BJ, Parker JE: Plant immunity: the EDS1 regulatory node. *Curr Opin Plant Biol* 2005, **8**(4):383–389.
 22. Schenk G, Elliott TW, Leung E, Carrington LE, Mitic N, Gahan LR, Guddat LW: Crystal structures of a purple acid phosphatase, representing different steps of this enzyme's catalytic cycle. *BMC Struct Biol* 2008, **8**:6.
 23. Mitić NN, Noble CJ, Gahan LR, Hanson GR, Schenk GG: Metal-ion mutagenesis: conversion of a purple acid phosphatase from sweet potato to a neutral phosphatase with the formation of an unprecedented catalytically competent Mn(II)Mn(II) active site. *J Am Chem Soc* 2009, **131**(23):8173.
 24. del Pozo JC, Allona I, Rubio V, Leyva A, de IP, Aragoncillo C, Paz-Ares J: A type 5 acid phosphatase gene from *Arabidopsis thaliana* is induced by phosphate starvation and by some other types of phosphate mobilising/oxidative stress conditions. *Plant J* 1999, **19**(5):579.
 25. Zhang W, Gruszewski HA, Chevone BI, Nessler CL: An *Arabidopsis* Purple Acid Phosphatase with Phytase Activity Increases Foliar Ascorbate. *Plant Physiol* 2008, **146**(2):431–440.
 26. Liao H, Wong F, Phang T, Cheung M, Li WF, Shao G, Yan X, Lam H: *GmPAP3*, a novel purple acid phosphatase-like gene in soybean induced by NaCl stress but not phosphorus deficiency. *Gene* 2003, **318**:103–111.
 27. Kaida R, Satoh Y, Bulone V, Yamada Y, Kaku T, Hayashi T, Kaneko TS: Activation of beta-glucan synthases by wall-bound purple acid phosphatase in tobacco cells. *Plant Physiol* 2009, **150**(4):1822–1830.
 28. Ekrylander B, Flores M, Wendel M, Heinegard D, Andersson G: Dephosphorylation of Osteopontin and Bone Sialoprotein by Osteoclastic Tartrate-Resistant Acid-Phosphatase - Modulation of Osteoclast Adhesion In-Vitro. *J Biol Chem* 1994, **269**(21):14853–14856.
 29. Nuttleman P, Roberts RM: Transfer of iron from uteroferrin (purple acid phosphatase) to transferrin related to acid phosphatase activity. *J Biol Chem* 1990, **265**(21):12192–12199.
 30. Kaija H, Alatalo SL, Halleen JM, Lindqvist Y, Schneider G, Vaananen HK, Vihko P: Phosphatase and oxygen radical-generating activities of mammalian purple acid phosphatase are functionally independent. *Biochem Biophys Res Commun* 2002, **292**(1):128–132.
 31. Hayman AR, Cox TM: Purple acid phosphatase of the human macrophage and osteoclast. Characterization, molecular properties, and crystallization of the recombinant di-iron-oxo protein secreted by baculovirus-infected insect cells. *J Biol Chem* 1994, **269**(2):1294–1300.
 32. Liang C, Tian J, Lam HM, Lim BL, Yan X, Liao H: Biochemical and molecular characterization of PvPAP3, a novel purple acid phosphatase isolated from common bean enhancing extracellular ATP utilization. *Plant Physiol* 2010, **152**(2):854–865.
 33. Li D, Zhu H, Liu K, Liu X, Leggewie G, Udvardi M, Wang D: Purple acid phosphatases of *Arabidopsis thaliana* comparative analysis and differential regulation by phosphate deprivation. *J Biol Chem* 2002, **277**(31):27772–27781.
 34. Kaffarnik FAR, Jones AME, Rathjen JP, Peck SC: Effector proteins of the bacterial pathogen *Pseudomonas syringae* alter the extracellular proteome of the host plant, *Arabidopsis thaliana*. *Molecular & cellular proteomics: MCP* 2009, **8**(1):145–156.
 35. Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ: An "electronic fluorescent pictograph" browser for exploring and analyzing large-scale biological data sets. *PLoS One* 2007, **2**(8):e718.
 36. Zhu H, Qian W, Lu X, Li D, Liu X, Liu K, Wang D: Expression patterns of purple acid phosphatase genes in *Arabidopsis* organs and functional analysis of *AtPAP23* predominantly transcribed in flower. *Plant Mol Biol* 2005, **59**(4):581–594.
 37. Tran HT, Qian W, Hurley BA, She Y, Wang D, Plaxton WC: Biochemical and molecular characterization of *AtPAP12* and *AtPAP26*: the predominant purple acid phosphatase isozymes secreted by phosphate-starved *Arabidopsis thaliana*. *Plant Cell Environ* 2010, **33**(11):1789–1803.
 38. Dewdney J, Reuber TL, Wildermuth MC, Devoto A, Cui J, Stutius LM, Drummond EP, Ausubel FM: Three unique mutants of *Arabidopsis* identify eds loci required for limiting growth of a biotrophic fungal pathogen. *Plant J* 2000, **24**(2):205–218.
 39. Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JDG, Felix G, Boller T: Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature* 2004, **428**(6984):764–767.
 40. Melotto M, Underwood W, Koczan J, Nomura K, He SY: Plant stomata function in innate immunity against bacterial invasion. *Cell* 2006, **126**(5):969–980.
 41. Menke FLH, Van Pelt JA, Pieterse CMJ, Klessig DF: Silencing of the mitogen-activated protein kinase MPK6 compromises disease resistance in *Arabidopsis*. *Plant Cell* 2004, **16**(4):897–907.
 42. Uquillas C, Letelier I, Blanco F, Jordana X, Holuigue L: NPR1-independent activation of immediate early salicylic acid-responsive genes in *Arabidopsis*. *Mol Plant Microbe Interact* 2004, **17**(1):34–42.
 43. Bozzo GG, Raghothama KG, Plaxton WC: Purification and characterization of two secreted purple acid phosphatase isozymes from phosphate-starved tomato (*Lycopersicon esculentum*) cell cultures. *Eur J Biochem* 2002, **269**(24):6278–6286.
 44. Torres MA, Dangl JL: Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Curr Opin Plant Biol* 2005, **8**(4):397–403.
 45. Ferrari S, Galletti R, Denoux C, De Lorenzo G, Ausubel FM, Dewdney J: Resistance to botrytis cinerea induced in *Arabidopsis* by elicitors is independent of salicylic acid, ethylene, or jasmonate signaling but requires *PHYTOALEXIN DEFICIENT*. *Plant Physiol* 2007, **144**(1):367–379.
 46. Ferrari S, Plotnikova JM, De Lorenzo G, Ausubel FM: *Arabidopsis* local resistance to *Botrytis cinerea* involves salicylic acid and camalexin and requires *EDS4* and *PAD2*, but not *SID2*, *EDS5* or *PAD4*. *Plant J* 2003, **35**(2):193–205.
 47. Zhou N, Tootle TL, Tsui F, Klessig DF, Glazebrook J: *PAD4* functions upstream from salicylic acid to control defense responses in *Arabidopsis*. *Plant Cell* 1998, **10**(6):1021–1030.
 48. Schenk G, Mitić N, Hanson GR, Comba P: Purple acid phosphatase: A journey into the function and mechanism of a colorful enzyme. *Coord Chem Rev* 2013, **257**(2):473–482.
 49. Schenk G, Guddat LW, Ge Y, Carrington LE, Hume DA, Hamilton S, de Jersey J: Identification of mammalian-like purple acid phosphatases in a wide range of plants. *Gene* 2000, **250**(1–2):117–125.
 50. Murashige T, Skoog F: A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 1962, **15**:473–497.
 51. Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, Gadrinab C, Heller C, Jeske A, Koesema E, Meyers CC, Parker H, Prednis L, Ansari Y, Choy N: Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 2003, **301**(5633):653.
 52. Morcuende R, Bari R, Gibon Y, Zheng W, Pant BD, Bläsing O, Usadel B, Scheible W: Genome-wide reprogramming of metabolism and regulatory networks of *Arabidopsis* in response to phosphorus. *Plant Cell Environ* 2007, **30**(1):85–112.

53. Chomczynski P, Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987, **162**(1):156–159.
54. Torres MA, Dangl JL, Jones JDG: *Arabidopsis* gp91(phox) homologues *AtrbohD* and *AtrbohF* are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proc Natl Acad Sci USA* 2002, **99**(1):517–522.

doi:10.1186/1471-2229-13-107

Cite this article as: Ravichandran et al.: *Purple Acid Phosphatase5* is required for maintaining basal resistance against *Pseudomonas syringae* in *Arabidopsis*. *BMC Plant Biology* 2013 **13**:107.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

