

Neutralization of Interleukin-10 Significantly Enhances Gamma Interferon Expression in Peripheral Blood by Stimulation with Johnin Purified Protein Derivative and by Infection with *Mycobacterium avium* subsp. *paratuberculosis* in Experimentally Infected Cattle with Paratuberculosis

Joram J. Buza,¹ Hirokazu Hikono,¹ Yasuyuki Mori,² Reiko Nagata,² Sachiyo Hirayama,¹ Abusaleh M. Bari,¹ Aodon-geril,¹ Yujing Shu,¹ Noriko M. Tsuji,³ and Eiichi Momotani^{1*}

*Paratuberculosis and Inflammatory Bowel Disease Research Team*¹ and *Immune System Section*,² *National Institute of Animal Health, and Molecular Immunology Laboratory, National Institute of Agrobiological Sciences*,³ 3-1-5 Kan-nondai, Tsukuba 305-0856, Japan

Received 28 October 2003/Returned for modification 16 December 2003/Accepted 8 January 2004

Monoclonal antibody neutralization of interleukin-10 (IL-10) increased Johnin purified protein derivative-induced whole-blood gamma interferon (IFN- γ) secretion 23-fold and also increased IFN- γ secretion ninefold following *in vitro* *Mycobacterium avium* subsp. *paratuberculosis* infection of peripheral blood mononuclear cells. These results demonstrate the suppressive effect of IL-10 on immune responses to *M. avium* subsp. *paratuberculosis* infection in cattle.

Immune responses to *Mycobacterium avium* subsp. *paratuberculosis* infection in cattle are characterized by a progressive shift from the protective cell-mediated (Th1) response in the subclinical stages of the disease towards the nonprotective antibody-mediated (Th2) response during the clinical stages of the disease (3, 4, 6, 9, 10, 12). It has been suggested that cytokines with an immunosuppressive effect on Th1 responses, in particular interleukin-10 (IL-10), may mediate this change (12, 13). The cytokine IL-10 is produced by Th2 CD4⁺ T cells, B cells, and macrophages and has many functions including inhibition of cytokine production by Th1 cells while promoting B-cell proliferation and differentiation (5, 8). However, the effect of IL-10 on immune responses to *M. avium* subsp. *paratuberculosis* in cattle has never been demonstrated.

Animals used in this study were 40-month-old castrated male Holstein cattle comprising the infected group in the subclinical stage of experimental *M. avium* subsp. *paratuberculosis* infection and an uninfected group, each containing five animals. Details of the experimental infection were described previously (2).

The study first investigated whether IL-10 has an influence on the purified protein derivative (PPD)-induced whole-blood gamma interferon (IFN- γ) response (1). Blood was stimulated as previously described (8) with few modifications. Stimulants and monoclonal antibodies (MAbs) were added into wells of 24-well round-bottomed tissue culture plates (Sumilon; Sumitomo Bakelite, Tokyo, Japan) to a final volume of 1 ml comprising 1/10-diluted blood, 5 μ g of *M. avium* subsp. *paratuber-*

culosis PPD/ml, and different fourfold dilutions of anti-IL-10 MAb CC320 or an immunoglobulin G1 isotype control MAb (Sigma, St. Louis, Mo.). The diluent was RPMI 1640 (Sigma) containing 1% penicillin-streptomycin and L-glutamine. The blood was incubated at 37°C and 5% CO₂ in air. Positive controls included blood stimulated with concanavalin A (ConA; type IV; Sigma) at a final concentration of 10 μ g/ml, and unstimulated cells served as negative control. Supernatants were measured for IFN- γ concentration with the bovine IFN- γ enzyme-linked immunosorbent assay (ELISA) kit (Bio-X SPRL, Marche-en-Famenne, Belgium) after 24 h of incubation. For measurement of cell proliferation, triplicate 100- μ l aliquots of PPD-stimulated blood were pulsed with 0.5 μ Ci of [³H]thymidine at 48 h and plates were incubated for an additional 24 h at 37°C in 5% CO₂ in air. The cells were then harvested, and radioactivity was quantified in a liquid scintillation counter (Tri-Carb 1600TR; Packard, Meriden, Conn.).

To investigate whether PPD induces a differential IL-10 or transforming growth factor β (TGF- β) mRNA expression in the *M. avium* subsp. *paratuberculosis*-infected and uninfected animals, 1 ml of undiluted blood was stimulated with 5 μ g of *M. avium* subsp. *paratuberculosis* PPD/ml and at 0, 1, 3, 6, 9, 12, and 24 h after stimulation, the blood cell fraction was processed for RNA isolation with the QIAamp RNA blood kit (Qiagen GmbH, Hilden, Germany). The RNA was reverse transcribed and subsequently amplified according to the instructions of the manufacturer (Applied Biosystems). The primers and TaqMan probes used are described in Table 1. The PCR conditions were 2 min at 50°C, 10 min at 95°C, and 40 cycles of 95°C for 15 s and 60°C for 1 min.

Finally the study investigated whether IL-10 has an effect on IFN- γ responses to infection with live *M. avium* subsp. *paratuberculosis*. Peripheral blood mononuclear cells were isolated

* Corresponding author. Mailing address: Paratuberculosis and Inflammatory Bowel Disease Research Team, NIAH, 3-1-5 Kan-nondai, Tsukuba 305-0856, Japan. Phone: 81-29-838-7781. Fax: 81-29-838-7781. E-mail: momotani@affrc.go.jp.

TABLE 1. Primers and probes for real-time PCR

Target and type	Designation ^b	Sequence	Reference no.
IL-10			U00799
Forward	F280	GTGATGCCACAGGCTGAGAA	
Reverse	R414	CGCCTTGCTCTTGTTTTCG	
Probe	T364	CGGCTGCGGGCTGTTCATC	
TGF- β			U62110
Forward	F748	GGCCCTGCCCTTACATCTG	
Reverse	R822	CCGGTTGTGCTGGTTGT	
Probe	T770	CCTGGATACACAGTACAGCAAG GTCTGGC	
GAPDH ^a			U85042
Forward	F364	GCGTGAACCACGAGAAGTATAA	
Reverse	R481	CCTCCACGATGCCAAAGTG	
Probe	T393	CCTCAAGATTGTACGAATGCCT CCTG	

^a GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

^b Numbers indicate the sequence positions.

from jugular blood samples with the use of Ficoll-Paque PLUS (Amersham Bioscience, Uppsala, Sweden) density gradient centrifugation per the manufacturer's instructions. The cells were resuspended in RPMI 1640 medium (Sigma) containing 10% fetal calf serum and dispensed into wells of 24-well tissue culture plates (Costar, Cambridge, Mass.) at 3×10^6 cells per well. To each well was added 3×10^6 CFU of *M. avium* subsp. *paratuberculosis* (14) in the presence of 1/1,000-diluted anti-IL-10 MAb CC320 or 5 μ g of isotype control MAb/ml, and after 24, 48, and 96 h of infection, the supernatant was collected for IFN- γ quantification. Data in this study were analyzed by analysis of variance, and a *P* value of <0.05 was considered significant.

Compared to samples stimulated in the absence of anti-IL-10 MAb, the PPD-induced IFN- γ level was significantly increased in the infected group at a MAb dilution of 1/1,600 and below (*P* < 0.05), representing a 23-fold increase (Fig. 1).




FIG. 1. Changes in IFN- γ secretion by whole blood from *M. avium* subsp. *paratuberculosis*-infected (●) and uninfected (○) cattle after 24 h of stimulation with Johnin PPD in the presence of an IL-10-neutralizing MAb.




FIG. 2. Changes in IFN- γ secretion by whole blood from *M. avium* subsp. *paratuberculosis*-infected (●) and uninfected (○) cattle after 24 h of stimulation with ConA in the presence of an IL-10-neutralizing MAb.

The anti-IL-10 MAb, however, did not change the PPD-induced IFN- γ level in the uninfected group, and levels in the infected group were higher at a MAb dilution of 1/6,400 or lower (*P* < 0.01). The neutralization of IL-10 had no effect on IFN- γ secretion by ConA-stimulated (Fig. 2) or nonstimulated (Fig. 3) cells. Unlike the anti-IL-10 MAb, an isotype control MAb had no effect on IFN- γ secretion in response to PPD- or ConA-stimulated or unstimulated cells (data not shown). The PPD-induced proliferation in samples from the infected group was significantly increased at antibody dilutions of 1/400 or lower, representing up to a twofold increase over that in samples stimulated in the absence of IL-10 neutralization (Fig. 4). Parallel to findings for IFN- γ production, the anti-IL-10 MAb had no effect on ConA-stimulated or nonstimulated samples,




FIG. 3. Changes in IFN- γ secretion by whole blood from *M. avium* subsp. *paratuberculosis*-infected (●) and uninfected (○) cattle after 24 h of incubation without stimulation in the presence of an IL-10-neutralizing MAb.




FIG. 4. Changes in Johnin PPD-induced proliferation by whole-blood cells from *M. avium* subsp. *paratuberculosis*-infected (●) and uninfected (○) cattle in the presence of an IL-10-neutralizing MAb. Whole blood diluted 1/10 was stimulated with 5 µg of Johnin PPD/ml, and after 48 h of stimulation, the cells were pulsed with 0.5 µCi of [³H]thymidine and incubated for an additional 24 h. Radioactivity was measured with a liquid scintillation counter.

and similarly, the isotype control MAb had no effect on cell proliferation (data not shown).

The PPD-induced IL-10 mRNA expression rose sharply and was significantly increased at 1 h in the infected and uninfected groups ($P < 0.05$ and $P < 0.01$, respectively) (Fig. 5). After 3 h, however, while expression in the infected group was sustained,




FIG. 5. Changes in IL-10 mRNA expression by whole blood from *M. avium* subsp. *paratuberculosis*-infected (●) and uninfected (○) cattle after stimulation with Johnin PPD. RNA was purified from whole blood by use of the QIAamp kit and reverse transcribed using the TaqMan reverse transcription reagents. The resulting cDNA was amplified for IL-10 and glyceraldehyde-3-phosphate dehydrogenase by real-time PCR with the TaqMan Universal PCR Mastermix (Applied Biosystems). Results are expressed as IL-10 mRNA expression of PPD-stimulated blood as a percentage of that in unstimulated whole blood.




FIG. 6. Changes in IFN- γ secretion by peripheral blood mononuclear cells after in vitro infection with live *M. avium* subsp. *paratuberculosis* in the presence of anti-IL-10 neutralizing MAb or an immunoglobulin G1 isotype control MAb. The different series represent infected cattle cells in the presence of IL-10-neutralizing MAb (●) or isotype control MAb (○) and uninfected cattle cells in the presence of IL-10-neutralizing MAb (■) or isotype control MAb (□). The anti-IL-10 MAb was used at a 1/1,000 dilution while the isotype control MAb was used at 5 µg/ml.

the expression in the uninfected group declined sharply and was significantly lower at 6 ($P < 0.001$), 9 ($P < 0.01$), and 12 ($P < 0.01$) h. The TGF- β expression was not different between the infected and the uninfected groups (data not shown). Live *M. avium* subsp. *paratuberculosis* infection of peripheral blood mononuclear cells under conditions of IL-10 neutralization significantly increased IFN- γ secretion at 48 and 96 h ($P < 0.001$) in the infected and uninfected groups, representing as much as a ninefold and threefold increase, respectively (Fig. 6). However, peak IFN- γ levels were significantly higher ($P < 0.001$) in cells from the infected group to a magnitude of 28-fold.

Results from this study demonstrate for the first time the suppressive effect of IL-10 on Th1 immune responses to *M. avium* subsp. *paratuberculosis* and its antigens in infected cattle and provide an approach for improvement of diagnosis and further studies of the disease's pathogenesis.

The whole-blood IFN- γ ELISA is a useful test for diagnosis of *M. avium* subsp. *paratuberculosis* infection in cattle but becomes less effective late in the disease due to unresponsiveness of the infected animals (6). The amplification of the whole-blood PPD-induced IFN- γ response by the IL-10-neutralizing MAb may help improve diagnosis of *M. avium* subsp. *paratuberculosis* infection, particularly late in the subclinical stage, for animals whose IFN- γ response is below the limit of detection by the IFN- γ ELISA but who have not developed sufficient antibodies for detection by serological tests.

Results from this study showed that *M. avium* subsp. *paratuberculosis* antigens induced a more sustained IL-10 response in infected animals than in uninfected animals, extending previous observations from uninfected animals (13). The suppressive effect of IL-10 on *M. avium* subsp. *paratuberculosis*-specific T-cell IFN- γ and cell proliferation responses may contribute

greatly to the observed immunopathology associated with the disease. The suppression of proliferative responses, if maintained during the infection period, will reduce the numbers of *M. avium* subsp. *paratuberculosis*-specific T cells, leading to the progressive reduction in IFN- γ responses and the decrease in peripheral and intestinal cell-mediated immune responses normally observed (7, 9, 11). Furthermore, since IFN- γ plays a crucial role in antimycobacterial responses (14), its suppression will likely result in suboptimal killing of *M. avium* subsp. *paratuberculosis*, leading to the chronic infections associated with the disease. However, studies with cells from the infected sites are still required for comparison with observations for peripheral blood.

This work was supported by the Japan Bio-oriented Technology Research Advancement Institute (BRAIN).

We thank C. J. Howard of the Institute of Animal Health, Berkshire, United Kingdom, for donating the IL-10-neutralizing MAb. We also thank T. Kamio of the Parasitic Disease Section, National Institute of Animal Health, Tsukuba, Japan, for granting access to some animals used in this study and S. Enoki of the Paratuberculosis and IBD research team for assistance in the preparation of various experiments.

REFERENCES

1. Billman-Jacobe, H., M. Carrigan, and F. Cochram. 1992. A comparison of the interferon gamma assay with absorbed ELISA for the diagnosis of Johne's disease in cattle. *Aust. Vet. J.* **69**:25–28.
2. Buza, J. J., Y. Mori, A. M. Bari, H. Hikono, Aodon-geril, S. Hirayama, Y. Shu, and E. Momotani. 2003. *Mycobacterium avium* subsp. *paratuberculosis* infection causes suppression of RANTES, monocyte chemoattractant protein 1, and tumor necrosis factor alpha expression in peripheral blood of experimentally infected cattle. *Infect. Immun.* **71**:7223–7227.
3. Chiodini, R. J., H. J. Van Kruningen, and R. S. Merkal. 1984. Ruminant paratuberculosis (Johne's disease): the current status and future prospects. *Cornell Vet.* **74**:218–262.
4. Clarke, C. J. 1997. The pathology and pathogenesis of paratuberculosis in ruminants and other species. *J. Comp. Pathol.* **116**:217–261.
5. Fiorentino, D. F., A. Zlotnik, P. Vieira, T. R. Mosmann, M. Howard, K. W. Moore, and A. O'Garra. 1991. IL-10 acts on the antigen presenting cell to inhibit cytokine production by Th1 cells. *J. Immunol.* **146**:3444–3451.
6. Harris, N. B., and R. G. Barletta. 2001. *Mycobacterium avium* subsp. *paratuberculosis* in veterinary medicine. *Clin. Microbiol. Rev.* **14**:489–512.
7. Koets, A., V. Rutten, A. Hoek, F. van Mil, K. Müller, D. Bakker, E. Gruys, and W. van Eden. 2002. Progressive bovine paratuberculosis is associated with local loss of CD4⁺ T cells, increased frequency of $\gamma\delta$ T cells, and related changes in T-cell function. *Infect. Immun.* **70**:3856–3864.
8. Kwong, L. S., J. C. Hope, M. L. Thom, P. Sopp, S. Duggan, G. P. Bembridge, and C. J. Howard. 2002. Development of an ELISA for bovine IL-10. *Vet. Immunol. Immunopathol.* **85**:213–223.
9. Perez, V., J. Tellechea, J. J. Badiola, M. Gutierrez, and J. F. Garcia Marin. 1997. Relation between serologic response and pathologic findings in sheep with naturally acquired paratuberculosis. *Am. J. Vet. Res.* **58**:799–803.
10. Stabel, J. R. 1996. Production of γ -interferon by peripheral blood mononuclear cells: an important diagnostic tool for detection of preclinical paratuberculosis. *J. Vet. Diagn. Investig.* **8**:345–350.
11. Stabel, J. R. 2000. Cytokine secretion by peripheral blood mononuclear cells from cows infected with *Mycobacterium paratuberculosis*. *Am. J. Vet. Res.* **61**:754–760.
12. Sweeney, R. W., D. E. Jones, P. Habecker, and P. Scott. 1998. Interferon- γ and interleukin-4 expression in cows infected with *Mycobacterium paratuberculosis*. *Am. J. Vet. Res.* **59**:842–847.
13. Weiss, D. J., O. A. Evanson, A. Moritz, M. Q. Deng, and M. S. Abrahamsen. 2002. Differential responses of bovine macrophages to *Mycobacterium avium* subsp. *paratuberculosis* and *Mycobacterium avium* subsp. *avium*. *Infect. Immun.* **70**:5556–5561.
14. Zhao, B., M. T. Collins, and C. J. Czuprynski. 1997. Effects of gamma interferon and nitric oxide on the interaction of *Mycobacterium avium* subsp. *paratuberculosis* with bovine monocytes. *Infect. Immun.* **65**:1761–1766.

Editor: S. H. E. Kaufmann