



CrossMark  
click for updates

## Research

**Cite this article:** Calosi P, Rastrick SPS, Lombardi C, de Guzman HJ, Davidson L, Jahnke M, Giangrande A, Hardege JD, Schulze A, Spicer JI, Gambi M-C. 2013 Adaptation and acclimatization to ocean acidification in marine ectotherms: an *in situ* transplant experiment with polychaetes at a shallow CO<sub>2</sub> vent system. *Phil Trans R Soc B* 368: 20120444. <http://dx.doi.org/10.1098/rstb.2012.0444>

One contribution of 10 to a Theme Issue 'Ocean acidification and climate change: advances in ecology and evolution'.

### Subject Areas:

evolution, physiology, ecology, genetics

### Keywords:

adaptation, plasticity, climate change, metabolic rate, ocean acidification, mass extinction

### Author for correspondence:

Piero Calosi

e-mail: [piero.calosi@plymouth.ac.uk](mailto:piero.calosi@plymouth.ac.uk)

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rstb.2012.0444> or via <http://rstb.royalsocietypublishing.org>.

# Adaptation and acclimatization to ocean acidification in marine ectotherms: an *in situ* transplant experiment with polychaetes at a shallow CO<sub>2</sub> vent system

Piero Calosi<sup>1</sup>, Samuel P. S. Rastrick<sup>1</sup>, Chiara Lombardi<sup>2</sup>, Heidi J. de Guzman<sup>3</sup>, Laura Davidson<sup>4</sup>, Marlene Jahnke<sup>4</sup>, Adriana Giangrande<sup>5</sup>, Jörg D. Hardege<sup>4</sup>, Anja Schulze<sup>3</sup>, John I. Spicer<sup>1</sup> and Maria-Cristina Gambi<sup>6</sup>

<sup>1</sup>Marine Biology and Ecology Research Centre, School of Marine Science and Engineering, Plymouth University, Drake Circus, Plymouth PL4 8AA, UK

<sup>2</sup>Marine Ecology Laboratory, Marine Environment and Sustainable Development Unit ENEA, PO Box 224, La Spezia, Italy

<sup>3</sup>Department of Marine Biology, Texas A&M University at Galveston, PO Box 1675, Galveston, TX 77554, USA

<sup>4</sup>Chemical Ecology Group, School of Biological, Biomedical and Environmental Sciences, The University of Hull, Hull HU6 7RX, UK

<sup>5</sup>Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali, University of Salento, Lecce, Italy

<sup>6</sup>Stazione Zoologica Anton Dohrn, Laboratory of Functional and Evolutionary Ecology, Villa Comunale 80121, Napoli, Italy

Metabolic rate determines the physiological and life-history performances of ectotherms. Thus, the extent to which such rates are sensitive and plastic to environmental perturbation is central to an organism's ability to function in a changing environment. Little is known of long-term metabolic plasticity and potential for metabolic adaptation in marine ectotherms exposed to elevated *p*CO<sub>2</sub>. Consequently, we carried out a series of *in situ* transplant experiments using a number of tolerant and sensitive polychaete species living around a natural CO<sub>2</sub> vent system. Here, we show that a marine metazoan (i.e. *Platynereis dumerilii*) was able to adapt to chronic and elevated levels of *p*CO<sub>2</sub>. The vent population of *P. dumerilii* was physiologically and genetically different from nearby populations that experience low *p*CO<sub>2</sub>, as well as smaller in body size. By contrast, different populations of *Amphiglena mediterranea* showed marked physiological plasticity indicating that adaptation or acclimatization are both viable strategies for the successful colonization of elevated *p*CO<sub>2</sub> environments. In addition, sensitive species showed either a reduced or increased metabolism when exposed acutely to elevated *p*CO<sub>2</sub>. Our findings may help explain, from a metabolic perspective, the occurrence of past mass extinction, as well as shed light on alternative pathways of resilience in species facing ongoing ocean acidification.

## 1. Introduction

Metabolic rate is considered the most fundamental of all biological rates [1]. According to the metabolic theory of ecology, metabolic rates set the rates of resource uptake and allocation to life-history traits (such as growth, reproduction and survival [2]), ultimately controlling ecological processes at all levels of organization [1]. Thus, the ability of an organism to preserve sufficient levels of energy metabolism when exposed to environmental challenges is key to a species' ability to preserve positive life-history traits, its Darwinian fitness, and ultimately its distribution and abundance patterns locally and globally [1,3–7].

Investigations of the effects of elevated *p*CO<sub>2</sub> on ectotherms' metabolic rates have revealed a variety of different responses: from differences among phyla at one extreme [8–10], to differences among related species and populations at the other [11,12]. When exposed to elevated *p*CO<sub>2</sub>, a number of taxa exhibit a

marked downregulation of their metabolic rate or ‘metabolic depression’ [11,13–17] but this is not ubiquitous. There are examples of upregulation [18–20], and no change in metabolism in response to elevated  $p\text{CO}_2$  [11,21–24]. It has been proposed that metabolic depression evolved to enable organisms to maintain a balance between energy supply and demand when their physiological machinery may be impaired as a result of environmental challenges [25,26]. Consequently, metabolic depression is considered to be primarily a short-term strategy [27] as, over the long term, it may have high costs in terms of growth, performances, reproductive output and may ultimately affect fitness. Thus, chronic metabolic depression has the potential to limit or prevent colonization of elevated  $p\text{CO}_2$  environments, and in a future, more acidic ocean to increase the risk of local and global taxa extinction. However, moderate forms of metabolic depression may be sustainable and could be achieved, via adaptation (i.e. selection of phenotypes–genotypes with moderately lower metabolic rates) or acclimatization (i.e. via phenotypic plasticity [28]). Nevertheless, adaptation and acclimatization can have different and important ecological consequences. Unfortunately, the nature and significance of physiological plasticity in marine ectotherms during acclimatization to an elevated  $p\text{CO}_2$  environment, as well as their potential physiological adaptation to these conditions, remain virtually unexplored (cf. [29]). Such an understanding of when plastic as opposed to genetic changes occur (and vice versa) is crucial if we are to predict how ocean acidification affects species’ distribution and abundance patterns, and thus predict the likely responses of marine ectotherms to ongoing climate change.

In addition, our interpretation of organisms’ metabolic responses to elevated  $p\text{CO}_2$  is biased by the fact that most species investigated to date are calcifiers. This is important as the overall direction and intensity of metabolic responses to elevated  $p\text{CO}_2$  are potentially affected by the upregulation of calcium carbon deposition [30–32], as well as the need to alter the biomineralization status of the shell, which may require the production of different organic and inorganic components [33], processes which are to date still poorly understood from a metabolic point of view. Furthermore, phylogenetic independence is rarely accounted for [10,34–37, cf. 38]. Clearly, there is an urgent need to investigate the physiological acclimatory ability and potential for physiological adaptation in a range of species within a group of phylogenetically related, non-calcifying ectotherms.

Consequently, we investigated the effect of elevated  $p\text{CO}_2$  on the metabolic rates of a number of species of non-calcifying polychaetes, which live in different proximities to natural  $\text{CO}_2$  vents. Polychaetes, in general, have so far received relatively little attention when compared with other groups (e.g. molluscs and crustaceans) (cf. [39–41]). Here using field transplant experiments, we looked for evidence of physiological acclimatization or adaptation in these species.

The  $\text{CO}_2$  vents of Ischia (Naples, Italy) have been used extensively to investigate the effects of elevated  $p\text{CO}_2$ /low pH conditions on marine communities and to predict possible responses of marine ecosystems to ocean acidification [10,42–45]. Used carefully, areas naturally enriched in  $\text{CO}_2$  make ideal natural laboratories for investigating such evolutionary questions. Cigliano *et al.* [43] found species-specific patterns of settlement in invertebrates along  $\text{CO}_2$  gradients in this vent system, mirrored by patterns of presence/absence and abundance of adults of polychaete species in the hard

bottom community [43,45], potentially indicating the presence of adaptation or acclimatization. In particular, around the  $\text{CO}_2$  vents of Ischia, some polychaete species maintain high densities along  $p\text{CO}_2$  gradients or even increase in density the closer they are to the vents. Such species could be considered as ‘tolerant’ [36]. Others decrease in density progressively along the  $\text{CO}_2$  gradients and are absent from the elevated  $p\text{CO}_2$  areas within the vents. These species could be considered ‘sensitive’ [36]. Given that there are such tolerant and sensitive non-calcifying polychaetes around the vents, the polychaete assemblage in Ischia is an ideal study system to test whether the ability of marine ectotherms to colonize  $\text{CO}_2$  vents depends on their different scope for physiological acclimatization and/or adaptation to elevated  $p\text{CO}_2$  conditions.

To define the potential for metabolic acclimatization and adaptation that allows colonization of elevated  $p\text{CO}_2$  areas, we carried out a series of *in situ* transplant and mutual transplant experiments populated with polychaetes living around the shallow-water  $\text{CO}_2$  vents system off Ischia. Post-transplant, we characterized metabolic rates and responses of the polychaetes, allowing us to infer the potential for metabolic adaptation in tolerant versus sensitive species (experiment 1), as well as between populations of tolerant species found both inside and outside the vent areas (experiment 2). Finally, to explore the evolutionary implications of any potential physiological adaptation of different populations of tolerant species, we used putatively neutral molecular markers to attribute levels of relatedness and phylogeographic pattern in populations of two of the tolerant species collected at increasing distance from the vents.

We predicted that (i) tolerant polychaete species will maintain their metabolism following acclimatization/adaptation to elevated  $p\text{CO}_2$  conditions (type 2 acclimatization/adaptation *sensu* [46]), (ii) sensitive polychaete species will display metabolic depression (type 1 acclimatization/adaptation *sensu* [46]).

## 2. Material and methods

### (a) Selection of sensitive and tolerant species

Two groups of species were identified:

(i) ‘Tolerant’ to low pH/elevated  $p\text{CO}_2$  conditions: species abundant both outside and inside the low pH/elevated  $p\text{CO}_2$  areas of the Castello  $\text{CO}_2$  vents of Ischia, namely *Platynereis dumerilii* (Audouin & Milne-Edwards, 1834; Nereididae), *Amphiglena mediterranea* (Leydig, 1851; Sabellidae), *Syllis prolifera* (Krohn, 1852; Syllidae) and *Polyopthalmus pictus* (Dujardin, 1839; Opheliidae). The first three dominate the most intense venting areas [43,45] and are commonly associated with rocky, shallow, vegetated habitat in the Mediterranean and with *Posidonia oceanica* (L.) Delile, 1813 seagrass beds [47–49]. *P. dumerilii* and *S. prolifera* are mesoherbivores, whereas *A. mediterranea* is a filter feeder. The fourth species, *P. pictus* (Opheliidae), is also a mesoherbivore typical of shallow, vegetated habitats [47]. This species was relatively scarce in the acidified areas in previous studies [43,45], but found (after our work took place) to be abundant on macroalgae and dead matter in a *P. oceanica* bed in high venting areas (E. Ricevuto & M. C. Gambi 2013, personal communication).

(ii) ‘Sensitive’ to low pH/elevated  $p\text{CO}_2$  conditions: species occurring around the vents under higher pH/low  $p\text{CO}_2$  conditions, but largely absent in the vents areas [43], namely *Lysidice ninetta* (Audouin & Milne-Edwards, 1833; Eunicidae), *Lysidice collaris* (Grube, 1870; Eunicidae) and *Sabella spallanzanii* (Gmelin, 1791; Sabellidae). Both *Lysidice* species are associated with vegetated rocky reef and coralligenous formations and are

some of the few species that bore on *P. oceanica* seagrass sheaths [50]. They also occur in the *P. oceanica* meadows at depths of 3–5 m and surrounding the vents, where normal pH/low  $p\text{CO}_2$  conditions exist [51]. Finally, *S. spallanzanii* is one of the most common and conspicuous sabellids in the Mediterranean. It is a filter feeder, tolerant to organic pollution and typical of fouling communities [52,53] outside the Ischia vent area [42]. All the sensitive species occur around the vents, in similar shallow habitat, but do not colonize the low pH/elevated  $p\text{CO}_2$  areas, thus suggesting that these conditions may constrain their distribution.

## (b) Collection details

The four tolerant species live in association with various macroalgae (mainly *Dyctiota* spp., *Halopteris scoparia* and *Cladophora prolifera*), where they were collected by hand, either by means of snorkelling or scuba diving, in both elevated (i) and low (ii)  $p\text{CO}_2$  conditions:

(i) at 1–2 m depth on a rocky reef at the low pH area on the south side of the Castello Aragonese, Ischia, Naples (40°43.84'N, 13°57.08'E). The collection area (Castello S3 in [43]) is approximately 60 × 15 m along a rocky reef and in the nearby *P. oceanica* meadow and dead 'matte'. This area corresponds to the very low pH area described in [45], where the pH ranges between 7.3 and 6.6 [45].

(ii) at 1–2 m depth off San Pietro promontory (approx. 4 km from the vents), where the control site experiments were performed, and around Sant'Anna islets and Cartaromana Bay (about 600 m from the vents). Here, venting activities were absent and water pH values measured were representative of low  $p\text{CO}_2$  conditions (mean pH  $8.17 \pm 0.02$  at S. Pietro and  $8.07 \pm 0.002$  at S. Anna, [45]). At these sites, polychaetes were associated with the same macroalgae species mentioned earlier, as well as with *Corallina* sp. Polychaetes from both localities were mixed and haphazardly allocated to different treatment and stations (see below). Finally, note that *S. prolifera* could not be retrieved from low  $p\text{CO}_2$  areas at the time our work was carried out, despite having been collected during previous studies and surveys [43,45] and in more recent times.

The sensitive species were collected exclusively in low  $p\text{CO}_2$ /high pH conditions. Specifically, *S. spallanzanii* was collected from floating docks in Ischia harbour, where this species form a dense population in the fouling community and thus a sufficient number of individuals for our experiment could be found. The two borer species of *Lysidice* were collected in shallow *P. oceanica* meadow (10 m depth) off Cava beach (approx. 10 km from the vents), a unique site in the area where these species occur in sufficient number in *P. oceanica* shoots [50].

The molecular phylogenetic analyses were conducted only on the tolerant species, *A. mediterranea* and *P. dumerilii*. Specimens of both taxa were collected at two acidified sites (Castello N3 and Castello S3 *sensu* [43]) and at several low  $p\text{CO}_2$  sites at different distances from the vents; *P. dumerilii* was also collected from the Bristol Channel, UK (see the electronic supplementary material, table A1.1 in appendix 1). Individuals of each species ( $n = 7$ –29) were collected at each location for genetic analyses. All specimens of *A. mediterranea* were preserved in 95% ethanol. Specimens of *P. dumerilii* were preserved in 95% ethanol or in dimethyl sulfoxide, frozen at  $-80^\circ\text{C}$  or immediately processed after collection without prior preservation. All *A. mediterranea* specimens were processed at Texas A&M University at Galveston (TAMUG; Galveston, USA); all *P. dumerilii* specimens were processed at the University of Hull (Hull, UK).

## (c) Culture and pre-exposure of polychaetes

To provide material for use in transplant experiments, all specimens of all species were reared in glass bowls (approx. 20 individual per bowl), each containing 300 ml of natural seawater ( $S = 38$ ) at

the original seawater pH/ $p\text{CO}_2$ . All glass bowls were kept in a controlled temperature environment ( $T = 19^\circ\text{C}$ , 12 L:12 D cycle). Each bowl was supplied with a few pieces of macroalgae from the collection site, for the polychaete to attach to and feed upon. Individuals of *S. spallanzanii* were kept under identical conditions except that, owing to their larger body size, the containers used were larger (volume = 14 l, approx. 1.2 individuals per litre), to allow polychaetes to easily open the branchial crown for filtering and respiration.

## (d) Study area and methods

The area where the transplant experiments into acidified conditions (A) were carried out was located in zones of high venting activity (greater than 10 vents  $\text{m}^{-2}$ ), at both south and north sides of the Castello Aragonese d'Ischia (figure 1). Previous studies showed a persistent gradient of low pH conditions in these areas [45]. In particular, three stations (A1, A2, A3—depth 2.5 m, areas approx. 2 × 3 m wide) were selected (figure 1), effectively representing one locality. The control area (C) (San Pietro point, approx. 4 km from the vent area; figure 1) was situated in close proximity to the Benthic Ecology research unit (Villa Dohrn) and was selected because of the persistent high pH/low  $p\text{CO}_2$  conditions and its accessibility [44]. Three stations (C1, C2 and C3—3 m depth) were established approximately 50 m apart from each other (figure 1). Small stony moorings (mass = 6 kg) were deployed in each of the six stations employed. A buoy attached to a nylon rope was used to fix, via a plastic cable tie, the experimental containers ('transplantation chambers') where individuals of the study species were kept during the experiment. Transplantation chambers for all but one species were constructed from white PVC tubes (diameter = 4 cm, length = 11 cm) with a nylon plankton net (mesh = 100  $\mu\text{m}$ ) fixed to both ends. This net mesh size was small enough to prevent polychaete escaping but allow regular water flow through the tube, allowing filtration for filter-feeders and respiration. For mesograzers, some of the algae that the various specimens are found on in the field were introduced in the transplantation chambers to provide both a suitable substratum for the polychaete to attach onto and a source of food to graze upon for the entire duration of the transplant, as for sea urchins in [36]. The large filter feeding *S. spallanzanii* were inserted in larger transplantation chambers constructed as plastic mesh cages (diameter = 15 cm, length = 30 cm, mesh = 1 cm), through which the apical part of their tube protruded by 1 cm allowing the branchial crown to open outside the cage enabling filtering and respiration. In general, while feeding can affect an organism physiology, and in particular metabolic levels [54,55], here we specifically wanted to maintain the polychaetes in conditions as close as possible to those they experience in the field, where they have continuous access to food resources.


## (e) Transplant design

To investigate the potential for metabolic adaptation and acclimatization that may allow or prevent the successful colonization of elevated  $p\text{CO}_2$  areas, the effect of exposure to different  $p\text{CO}_2$  conditions on the metabolic rates of selected polychaete species, collected from either low or elevated  $p\text{CO}_2$  areas around the shallow-water  $\text{CO}_2$  vent system off Ischia, was examined using a two-way orthogonal experimental design (with 'exposure' (exposure to low and elevated  $p\text{CO}_2$ ) and 'species' as factors). The analyses were conducted separately for the transplant experiments ('transplant from control areas'—experiment 1) and the reciprocal or mutual transplant ('transplant from acidified areas'—experiment 2).

### (i) Experiment 1

The first experiment investigated metabolic rates of all species collected from non-acidified conditions in control areas (control—C)





**Figure 1.** Regional (left) and local (right) maps of the study area, showing the low (C1, C2, C3) and elevated (A1, A2, A3)  $p\text{CO}_2$  stations used for the *in situ* transplants of polychaete species in the area around the  $\text{CO}_2$  vents off Ischia (Naples, Italy). The dotted lines represent bathymetric lines.

that were exposed to acidified (C-A) and non-acidified (C-C) conditions *in situ* for five days. For each species separately, and at each station, transplantation chambers (average of four individuals per chamber and so approx. 12 individuals in total per species per treatment) were deployed by scuba diving.

## (ii) Experiment 2

The second experiment investigated whether individuals of tolerant species living inside the  $\text{CO}_2$  vent areas show the same  $\text{CO}_2$ -dependent metabolic rate response among each other and when compared with their conspecifics living outside the  $\text{CO}_2$  vents. We conducted the reciprocal *in situ* transplant experiment on specimens of tolerant species collected from inside the vents (acidified—A), following the same deployment procedure described above in acidified (A-A) and non-acidified (A-C) areas. *S. prolifera* was found only in acidified conditions, while *P. pictus* was not found in sufficient numbers in the acidified area, therefore reciprocal transplants were not possible for these two species.

## (f) Environmental monitoring and profiles

Seawater pH, temperature and salinity were measured at each station daily throughout the three weeks duration of the experiments. For the transplant experiment, environmental monitoring was carried out as follows.  $\text{pH}_{\text{NSB}}$  was measured using a pH microelectrode (Seven Easy pH InLab, Mettler-Toledo Ltd, Beaumont Leys, UK), maintained at ambient seawater temperature, coupled to a pH meter (Sevengo, Mettler-Toledo Ltd), calibrated using pH standards (pH 4.01, 7.00, 9.21 at 25°C, Mettler-Toledo Ltd) and also maintained at ambient seawater temperature. Temperature was measured using a digital thermometer (HH806AU, OMEGA Eng. Ltd, Manchester, UK).

Salinity was measured using a hand-held conductivity meter (TA 197 LFMulti350, WTW, Weilheim, Germany). In addition, to determine total alkalinity (TA), samples of seawater

(volume = 100 ml) were collected at each station daily throughout the duration of the experimental period using glass bottles with a secure tight lid, transported inside a cool box to the laboratory (located approx. 4 km from the vents) and poisoned with  $\text{HgCl}_2$  within approximately 60 min of collection. TA was determined at the Marine Biology and Ecology Research Centre at Plymouth University (Plymouth, UK), using an alkalinity titrator (AS-ALK2, Apollo SciTech, Bogart, USA).

Dissolved inorganic carbon (DIC), partial pressure of  $\text{CO}_2$  ( $p\text{CO}_2$ ), calcite and aragonite saturation ( $\Omega_{\text{calc}}$  and  $\Omega_{\text{ara}}$ ), bicarbonate and carbonate ion concentration ( $[\text{HCO}_3^-]$  and  $[\text{CO}_3^{2-}]$ , respectively) were calculated from pH and TA measurements using the software program CO2SYS [56] with dissociation constants from [57] refit by [58] and  $[\text{KSO}_4]$  using [59]. Our results for the carbonate system (table 1) outside and inside the  $\text{CO}_2$  vent areas are consistent with those previously reported for these areas [42–45,60–62].

During the experimental period, seawater environmental conditions (table 1) were relatively stable with mean values for temperature, salinity and TA being around 23°C, 37 and 2651  $\mu\text{equiv kg}^{-1}$ , respectively. Mean pH values were relatively stable, approximately 8.15 and 7.17 for the control and acidified areas, respectively (table 1). Furthermore, preliminary analysis using the estimate marginal means (EMM) test with Bonferroni correction showed that environmental conditions (salinity, temperature and TA) were comparable across all stations irrespective of the  $p\text{CO}_2$  condition (table 1); while mean pH significantly differed among the low and elevated  $p\text{CO}_2$  treatments, it was, however, comparable among stations of a same treatment (table 1).

## (g) Recovery of transplantation chambers and measurement of metabolic rates

After five days of exposure *in situ* to either acidified or control conditions, transplantation chambers from both experiments 1 and 2 were recovered via snorkelling and diving. They were

**Table 1.** Values (mean  $\pm$  s.e.) for physico-chemical parameters of the seawater used when exposing polychaetes to: (i) current  $p\text{CO}_2$ /pH conditions ('control' stations C1, C2 and C3) and (ii) elevated  $p\text{CO}_2$ /low pH conditions ('acidified' stations A1, A2, A3). Salinity, temperature,  $\text{pH}_{\text{NBS}}$  (Mettler-Toledo pH meter, Luton, UK), TA (AS-ALK2, Apollo SciTech, Bogart, USA), DIC, carbon dioxide partial pressure ( $p\text{CO}_2$ ), bicarbonate and carbonate ion concentration ( $[\text{HCO}_3^-]$  and  $[\text{CO}_3^{2-}]$ ), calcite and aragonite saturation state ( $\Omega_{\text{cal}}$  and  $\Omega_{\text{ara}}$ ) are provided.

parameter	control			acidified			overall
	C1	C2	C3	A1	A2	A3	
salinity	$37.50 \pm 0.15^{(A)}$	$37.55 \pm 0.16^{(A)}$	$37.55 \pm 0.16^{(A)}$	$37.19 \pm 0.07^{(A)}$	$37.14 \pm 0.05^{(A)}$	$37.10 \pm 0.06^{(A)}$	$37.14 \pm 0.03$
temperature ( $^{\circ}\text{C}$ )	$22.83 \pm 0.03^{(A)}$	$22.80 \pm 0.03^{(A)}$	$22.80 \pm 0.03^{(A)}$	$23.33 \pm 0.15^{(A)}$	$23.31 \pm 0.12^{(A)}$	$23.41 \pm 0.18^{(A)}$	$23.34 \pm 0.08$
pH	$8.15 \pm 0.01^{(A)}$	$8.16 \pm 0.01^{(A)}$	$8.15 \pm 0.01^{(A)}$	$7.19 \pm 0.05^{(B)}$	$7.07 \pm 0.07^{(B)}$	$7.24 \pm 0.04^{(B)}$	$7.17 \pm 0.04$
TA ( $\mu\text{equiv kg}^{-1}$ )	$2624.27 \pm 10.34^{(A)}$	$2623.30 \pm 10.76^{(A)}$	$2629.29 \pm 8.91^{(A)}$	$2678.22 \pm 3.78^{(A)}$	$2673.73 \pm 12.57^{(A)}$	$2678.50 \pm 12.90^{(A)}$	$2676.38 \pm 6.41$
DIC ( $\mu\text{mol kg}^{-1}$ ) <sup>a</sup>	$2316.80 \pm 7.68$	$2307 \pm 11.09$	$2311.43 \pm 8.74$	$2781.21 \pm 20.15$	$2767.05 \pm 23.75$	$2892.95 \pm 55.51$	$2815 \pm 22.81$
$p\text{CO}_2$ ( $\mu\text{atm}$ ) <sup>a</sup>	$483.68 \pm 7.19$	$474.70 \pm 8.64$	$447.53 \pm 8.41$	$5337.65 \pm 567.54$	$5267.76 \pm 493.97$	$8847.50 \pm 1364.50$	$6534.16 \pm 564.68$
$[\text{HCO}_3^-]$ ( $\mu\text{mol kg}^{-1}$ ) <sup>a</sup>	$2070.00 \pm 9.50$	$2060.58 \pm 0.05$	$2065.13 \pm 9.29$	$2585.11 \pm 6.87$	$2565.58 \pm 20.72$	$2597.80 \pm 20.28$	$2582.73 \pm 10.29$
$[\text{CO}_3^{2-}]$ ( $\mu\text{mol kg}^{-1}$ ) <sup>a</sup>	$231.18 \pm 2.33$	$232.58 \pm 3.08$	$232.23 \pm 3.14$	$38.30 \pm 2.93$	$45.31 \pm 7.10$	$33.23 \pm 4.92$	$38.98 \pm 3.15$
$\Omega_{\text{cal}}^b$	$5.42 \pm 0.06$	$5.47 \pm 0.07$	$5.46 \pm 0.08$	$0.90 \pm 0.07$	$1.07 \pm 0.17$	$0.78 \pm 0.11$	$0.92 \pm 0.07$
$\Omega_{\text{ara}}^b$	$3.56 \pm 0.04$	$3.59 \pm 0.05$	$3.59 \pm 0.05$	$0.59 \pm 0.05$	$0.70 \pm 0.11$	$0.51 \pm 0.08$	$0.60 \pm 0.05$

Different capital letters (A,B) indicate significant differences for measure parameters in different stations.

<sup>a</sup>Parameters that were calculated using the CO2SYS program [56], using the dissociation constants of Mehrbach *et al.* [57] as refitted by Dickson & Millero [58] and  $[\text{KSO}_4]$  using Dickson [59].

transported within 5 min of harvest, using a bucket filled with seawater, to an aquarium containing seawater (volume = 60 l). Within 30 min of collection, all individuals were introduced individually, and in succession, to the incubation chambers (see below) containing filtered (0.22  $\mu\text{m}$ ) seawater from the appropriate test areas, to avoid 'recovery' of the polychaetes. Preliminary trials showed that this operation did not affect seawater pH, initial  $p\text{O}_2$ , temperature or salinity. Before determination of  $\text{O}_2$  uptake, here used as a proxy for metabolic rate ( $\dot{\text{M}}\text{O}_2$ ) (as in [63–65]), individuals were left for 1 h in the incubation chamber to allow recovery from handling in order to minimize behavioural disturbances, which may affect their physiological responses [66]. For further details on metabolic rate determination, see appendix 2 in the electronic supplementary material.

At the end of the experiment, polychaetes were bagged, placed in a cool box and returned to the laboratory. Here, they were gently blotted to remove excess seawater, before being weighed using a high precision balance (AT 200, Mettler-Toledo Ltd). To maintain environmental seawater chemistry and temperature stable during measurements, seawater was continuously pumped (BE-M 30, Rover Pompe, Polverara, Italy) directly from the exposure site into a large water bath (volume = 64 l) and the overflow drained. To ensure maximum thermal and pH stability, as well as comparability to the exposure sites, a fast flow rate (approx. 85 l  $\text{min}^{-1}$ ) was employed, thereby allowing us to relate metabolic rate responses solely to the effect of field acclimatization to elevated  $p\text{CO}_2$ /low pH. As an extra measure, during the experimental trials, seawater samples were collected at regular intervals from the exposure sites by snorkelling or diving, and from the water bath, to verify that pH, salinity and temperature were relatively stable and comparable. Polychaetes were never directly exposed to high flow rate, avoiding any mechanical damage or disturbance.

### (h) Statistical analysis of the physiological data

As a two-way orthogonal experimental design was employed, separately for both transplant experiments, the effect of exposure to elevated  $p\text{CO}_2$ /low pH on  $\dot{\text{M}}\text{O}_2$  of the species examined was tested using general linear models (GLMs), with 'exposure' (*in situ* incubation to low and elevated  $p\text{CO}_2$ ) and 'species' as fix factors, and using wet body mass as a covariate, in combination with the EMM test with Bonferroni correction. Preliminary analyses showed that the term station had no significant effect on polychaete  $\dot{\text{M}}\text{O}_2$  ( $p > 0.05$ ), and so this term was removed from subsequent analyses.

To investigate the response of populations of tolerant species collected from both inside and outside the  $\text{CO}_2$  vents, a further two-way orthogonal experimental design was employed, separately for each species (*A. mediterranea* and *P. dumerilii*), including the site of collection (collection from low and elevated  $p\text{CO}_2$  site) or 'origin', and the site of *in situ* incubation (to low and elevated  $p\text{CO}_2$ ) or 'exposure' as factors, using GLMs, with 'wet body mass' as covariate, in combination with the EMM test with Bonferroni correction.

$\dot{\text{M}}\text{O}_2$  data for transplant experiments met assumptions of normality of distribution following  $\log_{10}$  transformation (max.  $Z_{92} = 1.109$ ,  $p = 0.280$ , Kolmogorov–Smirnov's test). Assumptions of homogeneity of variance were also met following  $\log_{10}$  transformations for the data from experiment 1 ( $F_{11,84} = 1.904$ ,  $p = 0.05$ , Bartlett's test), whereas assumptions were not met following transformations for data from experiment 2 (minimum  $F_{5,66} = 4.446$ ,  $p < 0.001$ ). However, as our experimental design included a minimum of four treatments per experiment with approximately 11 replicates per treatment, we assumed that the ANOVA design employed should be tolerant to deviation from the assumption of heteroscedasticity [67,68]. However, we also tested the residuals from each analysis against the factor tested, and no significant relationships were detected ( $p > 0.05$ ). Post-hoc

analyses were conducted using the EMM test with  $\alpha = 0.05$ . All statistical analyses were conducted using SPSS v. 19.

### (i) Sequence generation

For *A. mediterranea*, DNA was extracted from each individual using the DNEasy tissue kit (Qiagen), following the manufacturer's instructions. For *P. dumerilii*, DNA was extracted using the hotshot method of Montero-Pau *et al.* [69].

Two gene regions were amplified for both species using polymerase chain reaction (PCR): the mitochondrial cytochrome *c* oxidase subunit I (COI, 658–710 bp) and the nuclear internal transcribed spacer (ITS, alignment of 495 bp). PCRs were performed in 20–25  $\mu\text{l}$  volume using standard chemistry and standard cycles. Some samples of *A. mediterranea* were amplified for COI using a primer cocktail originally designed for fish as described in Ivanova *et al.* [70], whereas other *A. mediterranea* and all *P. dumerilii* were amplified using the universal primers by Folmer *et al.* [71] or specifically designed primers. All primers are listed in the electronic supplementary material, table A1.2 of appendix 1.

PCR products were visualized following gel electrophoresis in ethidium bromide-stained agarose gels and cleaned with ExoSap-IT (Affymetrix) QIAquick PCR purification kit (Qiagen) or with the Illustra GFX PCR DNA purification kit, or purification was carried out by MacroGen Europe before sequencing. For *A. mediterranea*, cycle sequencing was conducted on 10  $\mu\text{l}$  volumes with the BIGDYE TERMINATOR v. 3.1 chemistry. Sequence reactions were cleaned with BIGDYE XTERTMINATOR. Sequences were analysed with an ABI 3130 Genetic Analyzer and edited in SEQUENCHER 4.8 (Genecodes). For *P. dumerilii*, sequences were generated at MacroGen Europe and edited with Codon Code Aligner. All sequences were submitted to GenBank under accession numbers KC591782–KC591950 (see the electronic supplementary material, table A1.3 in appendix 1).

### (j) Sequence analyses

COI sequences were aligned in BIOEDIT [72] using the CLUSTALW algorithm. ITS sequences were aligned in the software MUSCLE as implemented in MEGA 5 [67]. The final alignments had the following lengths: *P. dumerilii*, COI: 568 bp; *A. mediterranea*, COI: 658 bp; *P. dumerilii*, ITS: 608 bp; *A. mediterranea*, ITS: 498. The COI alignments included selected outgroups as available in GenBank. These were used to root the trees. No outgroups were used in the analyses of ITS, because no closely related sequences were available in GenBank. The ITS trees are unrooted.


All tree reconstructions were performed in MEGA v. 5.0 [73] using maximum likelihood as the optimality criterion under a general time reversible model with gamma distribution of substitution rates (five discrete steps) and a proportion of invariable sites (GTR + I + G model). Branch support was calculated by performing 1000 bootstrap replicates. Trees were edited in FIGTREE v. 1.3.1 [74].

## 3. Results

No mortality was observed for any of the test species during collection, transplant or experimentation. All polychaetes appeared in good health, actively moving when transferred from the transplantation chamber to the incubation chambers, and in semi-transparent species food traces were visible in the digestive system.

### (a) $\dot{\text{M}}\text{O}_2$

$\dot{\text{M}}\text{O}_2$  data are presented in the electronic supplementary material, appendix 3, and data on body mass and number of



**Figure 2.** Percentage change of metabolic rates ( $\dot{M}O_2$ ) of non-calcifying marine ‘tolerant’ (T) (*A. mediterranea*, *P. dumerilii* and *P. pictus*) and ‘sensitive’ (S) (*L. collaris*, *L. ninetta* and *S. spallanzanii*) polychaete species collected from low  $pCO_2$ /elevated pH conditions outside the vents and exposed to (i) low  $pCO_2$  conditions outside the vents (C-C), and (ii) elevated  $pCO_2$  conditions within the vents (C-A). Mean  $\dot{M}O_2$  (expressed as  $\log_{10}$  nmol  $O_2$   $mg^{-1}$   $h^{-1}$  STP) measured under control conditions was set as 100% and mean  $\dot{M}O_2$  measured under acidified conditions recalculated accordingly. Asterisk (\*) indicates the presence of a significant difference between the mean  $\dot{M}O_2$  measured under control conditions (C) and acidified conditions (A), according to the EMM test with Bonferroni correction ( $p \leq 0.05$ ).

specimens tested for each treatment are reported in the electronic supplementary material, appendix 4. In summary, mean  $\dot{M}O_2$  varied among species, ranging from (mean  $\pm$  s.e.)  $-0.082 \pm 0.083 \log_{10}$  nmol  $O_2$   $mg^{-1}$   $h^{-1}$  STP in *S. spallanzanii* collected under control conditions and exposed to control conditions, and  $1.632 \pm 0.056 \log_{10}$  nmol  $O_2$   $mg^{-1}$   $h^{-1}$  STP in *S. prolifera* collected inside the vents and exposed to control conditions (see the electronic supplementary material, appendix 3).

### (b) $\dot{M}O_2$ responses of all species collected in low $pCO_2$ areas

The polychaete species collected from low  $pCO_2$  areas differed from each other in the way their mean  $\dot{M}O_2$  responded to exposure to elevated  $pCO_2$  conditions (figure 2), as indicated by the presence of a significant interaction between the terms species and exposure ( $F_{5,95} = 5.611$ ;  $p < 0.0001$ ). The three sensitive species showed significant differences in their  $\dot{M}O_2$  under elevated  $pCO_2$  conditions ( $p < 0.05$ );  $\dot{M}O_2$  increased significantly in *S. spallanzanii*, and decreased significantly in *L. collaris* and *L. ninetta*. The three tolerant species (*A. mediterranea*, *P. dumerilii* and *P. pictus*), on the other hand, showed no significant difference in mean  $\dot{M}O_2$  when exposed to elevated  $pCO_2$  conditions ( $p > 0.05$ ). Finally, wet body mass had no significant effect on mean  $\dot{M}O_2$  ( $F_{1,95} = 1.149$ ;  $p = 0.287$ ).

### (c) $\dot{M}O_2$ responses of tolerant species collected in elevated $pCO_2$ areas

Tolerant species collected from the elevated  $pCO_2$  areas did not differ significantly from each other in the way mean  $\dot{M}O_2$  responded to exposure to low  $pCO_2$  conditions (figure 3), as indicated by the fact that the interaction term ‘species’  $\times$  ‘exposure’ in our analysis was not significant ( $F_{2,71} = 1.635$ ;  $p = 0.203$ ). In fact, all three tolerant species tested (*A.*

*mediterranea*, *P. dumerilii* and *S. prolifera*) showed a significant increase in mean  $\dot{M}O_2$  when exposed to low  $pCO_2$  conditions ( $p < 0.05$ ). Finally, wet body mass had no significant effect on mean  $\dot{M}O_2$  ( $F_{1,71} = 3.545$ ;  $p = 0.064$ ).


### (d) $\dot{M}O_2$ responses of tolerant species from elevated and low $pCO_2$ areas

Individuals of *A. mediterranea* and *P. dumerilii* from both control and elevated  $pCO_2$  areas showed different  $\dot{M}O_2$  responses to exposure to low and elevated  $pCO_2$  conditions (figure 4), as indicated by the presence of a significant three-way interaction between the terms ‘species’, ‘origin’ and ‘exposure’ ( $F_{1,91} = 5.828$ ;  $p = 0.018$ ). *A. mediterranea* collected from acidified areas and re-transplanted into acidified areas showed a significantly lower mean  $\dot{M}O_2$  when compared with all other treatments ( $p < 0.05$ ), while all other comparisons were not statistically significant ( $p > 0.05$ ). By contrast, individuals of *P. dumerilii* collected in acidified areas and exposed to low  $pCO_2$  conditions showed a significantly greater  $\dot{M}O_2$  when compared with all other treatments ( $p < 0.05$ ), while not showing any significant difference in mean  $\dot{M}O_2$  among each other ( $p > 0.05$ ). Finally, wet body mass had no significant effect on mean  $\dot{M}O_2$  ( $F_{1,71} = 3.545$ ;  $p = 0.064$ ).

### (e) Phylogenetic analyses

For both *P. dumerilii* and *A. mediterranea*, the phylogenetic trees generated distinguish multiple distinct evolutionary lineages, with low genetic diversity within them (figure 5). The COI tree for *P. dumerilii* shows that one of these genetic lineages (figure 5a) contains 10 of the 12 sequenced individuals from the acidified site and a single individual from a nearby control site. One individual from the acidified site falls into a clade consisting of individuals from control sites. Another individual forms its own genetic lineage. In





**Figure 3.** Percentage change of  $\dot{M}O_2$  of non-calcifying ‘tolerant’ marine polychaete species collected from elevated  $pCO_2$ /low pH conditions within the vents and exposed to (i) elevated  $pCO_2$  conditions inside the vents (A-A) and (ii) low  $pCO_2$  conditions outside the vents (A-C). Mean  $\dot{M}O_2$  (expressed as  $\log_{10}$  nmol  $O_2$   $mg^{-1}$   $h^{-1}$  STP) measured under acidified conditions was set as 100% and mean  $\dot{M}O_2$  measured under control conditions recalculated accordingly. Asterisk (\*) indicates the presence of a significant difference between the mean  $\dot{M}O_2$  measured under acidified conditions (A) and control conditions (C), according to the EMM test with Bonferroni correction ( $p \leq 0.05$ ). (Online version in colour.)

the ITS tree (figure 5b), the two individuals from the acidified site form a genetic lineage distinct from all other populations.


In contrast to *P. dumerilii*, the COI tree for *A. mediterranea* reveals that all individuals from the acidified site fall into the same clade with individuals from nearby control sites (figure 5c). Likewise, the ITS tree for *A. mediterranea* also does not show genetic separation between acidified and non-acidified sites (figure 5d).

## 4. Discussion and conclusions

Here, we show that a marine metazoan (*P. dumerilii*) is able to physiologically adapt to chronically elevated levels of  $pCO_2$ . However, such adaptation is not ubiquitous among all tolerant species found in the  $CO_2$  vents, even when their ecologies are similar. The physiological plasticity to chronically elevated  $pCO_2$  shown by *A. mediterranea* also appears to be a viable strategy for the successful colonization of elevated  $pCO_2$  environments. Finally, sensitive species, when exposed acutely to elevated  $pCO_2$  conditions, exhibit either considerable up or downregulation of their metabolism. In what follows we discuss species metabolic response to elevated  $pCO_2$  in relation to (i) the current distribution and abundance patterns of different species around the  $CO_2$  vents, (ii) extinctions that occurred during past climate change events, and (iii) species’ alternative pathways of physiological resilience to ongoing ocean acidification.

### (a) Discriminating between acclimatization and adaptation

The effect of  $pCO_2$  on metabolic rate differs consistently between sensitive and tolerant polychaete species, but for




**Figure 4.** Effect of exposure to different  $pCO_2$  conditions on the mean  $\dot{M}O_2$  of the tolerant species *A. mediterranea* and *P. dumerilii* collected from either low  $pCO_2$ /elevated pH conditions outside the vents and exposed to (i) low  $pCO_2$  conditions outside the vents (C-C) and (ii) elevated  $pCO_2$  conditions within the vents (C-A) or collected from elevated  $pCO_2$ /low pH conditions within the vents and exposed to (iii) elevated  $pCO_2$  conditions inside the vents (A-A) and (iv) low  $pCO_2$  conditions outside the vents (A-C). Mean  $\dot{M}O_2$  is expressed as  $\log_{10}$  nmol  $O_2$   $mg^{-1}$   $h^{-1}$  STP. Histograms present mean values  $\pm$  s.e. Significantly different treatments, separately for each species, are indicated by different lower case letters inside the column (according to the EMM test with Bonferroni correction,  $p \leq 0.05$ ). (Online version in colour.)

tolerant taxa the response patterns observed may have been achieved either via acclimatization or adaptation. In fact, organisms may be able to adjust their physiology, via phenotypic plasticity (acclimatization), e.g. during ontogeny in direct developers such as *A. mediterranea* (see [29]), or via the selection of genotypes associated with phenotypes best able to cope with conditions found within the  $CO_2$  vents, as in *P. dumerilii* (see [76]). However, which strategy is adopted (i.e. acclimatization or adaptation) has different genetic, ecological and conservation implications. Therefore, to make predictions on how marine life will respond to future ocean acidification, it is important to discriminate between the strategies.

Here, we show that individuals of *A. mediterranea* living inside the  $CO_2$  vents appear to be acclimatized, but not adapted, to elevated  $pCO_2$ . This is because once removed from elevated  $pCO_2$  (and probably under hypoxaemia [77,78]) their metabolic rates return to a ‘normal’ status: i.e. comparable with that of individuals from low  $pCO_2$  areas. Alternatively, individuals of *P. dumerilii* living inside the  $CO_2$  vents appear physiologically adapted to elevated  $pCO_2$ . When removed from the vents their metabolic rate is approximately 44% more elevated when compared with vent individuals kept inside the vents. The metabolism of vent specimens is thus constantly maintained at high levels, presumably to compensate for (although only in part) the chronic  $pCO_2$ -induced hypoxaemia they are probably subjected to [77,78].

Previous studies show that unicellular organisms can adapt to elevated  $pCO_2$  [79–81]. Our study provides evidence that a marine ectotherm (*P. dumerilii*) has been able to genetically and physiologically adapt to chronic and elevated levels of  $pCO_2$ , and supports those studies that have indicated the potential of marine metazoans to adapt to elevated  $pCO_2$  [76,82–87]. Furthermore, this adaptation may have occurred over a relatively short geological time. In fact, based on archaeological





**Figure 5.** Phylogenetic trees resulting from maximum-likelihood analysis of cytochrome *c* oxidase subunit I (COI) and ITS sequence data. Branch support is indicated as bootstrap percentages (1000 pseudoreplicates); asterisk (\*) indicates bootstrap value greater than 98%. (a) COI tree for *P. dumerilii*. Genbank accession numbers for outgroups: *Nereis zonata*: HQ024403; *Nereis pelagica*: GU672554; (b) ITS tree for *P. dumerilii*. The designations 'clone 1' to 'clone 10' with their respective GenBank accession numbers refer to [75]; (c) COI tree for *Amphiglena mediterranea*. GenBank accession numbers for outgroups: *Pseudopotamilla reniformis*: GU672463; *Eudistylia vancouveri*: HM473371; *Schizobranchia insignis*: HM473778. (d) ITS tree for *A. mediterranea*. (Online version in colour.)

and historical evidences, the CO<sub>2</sub> vent system off Ischia is estimated to be 1850 years old [62].

Although metabolic phenotypic plasticity may be the first 'mechanism' of response to preserve positive function levels

when exposed to environmental disturbance [28], it often comes at a cost [88,89]. Plastic responses can be accompanied by the reallocation of the available energy budget away from growth and reproduction [30], cf. [90,91]. When the cost becomes

too high, the selection of phenotypes better able to cope with elevated  $p\text{CO}_2$  conditions should be favoured as this is a less 'expensive' strategy [76]. Local adaptation leads to the improvement of population physiological performances, thus reducing energy costs of regulation and maintenance, improving an organism's ability to persist locally. However, if adaptation occurs at the expense of genetic diversity, this could lead to a decrease in the performance for other traits (e.g. life-history traits). When evolutionary trade-offs are less costly than phenotypic plastic reshuffling, adaptation should be favoured.

The multiple genetic lineages observed in *P. dumerilii* and *A. mediterranea* suggest that both actually represent complexes of cryptic species, a common phenomenon in polychaetes [92–97]. In our case, the geographical distribution of these lineages provides additional support for our findings on physiological adaptation and acclimatization from the transplant experiments. Having pelagic larval stages [98], *P. dumerilii* might be expected to display a high degree of genetic homogeneity on small geographical scales. However, our molecular analyses show a high degree of genetic differentiation among populations at and near the vents (figure 5a,b). This corroborates the idea that the strain of *P. dumerilii* sampled in the  $\text{CO}_2$  vent is indeed physiologically (and genetically) adapted to low  $p\text{CO}_2$ . Nonetheless, COI analysis indicates that some (but limited) exchange of individuals between the acidified sites and nearby control takes place, as one individual from the low  $p\text{CO}_2$  area was found at the vent sites, and one individual with the genetic make-up of the vent population was found in the control sites. Whether these individuals actually thrive and reproduce in the presumably less suitable habitats remains to be thoroughly examined. Nonetheless, based on our current (limited) evidence, one may infer that some form of ecological competitive exclusion may occur between the two strains.

The average genetic distance between the vent strain and its sister group, consisting of three individuals from Santa Caterina, is 3.8%. Coincidentally, this divergence was chosen by Carr *et al.* [99] to separate molecular operational taxonomic units in polychaetes on the basis of being 10 times greater than intracluster divergence. Estimates of mutation rates in annelids vary tremendously, ranging from 0.2%  $\text{Myr}^{-1}$  [100] to 7%  $\text{Myr}^{-1}$  [94]. Even with a high mutation rate of 7%, the separation of the vent clade from its sister group would date to 542 000 years ago, not compatible with an origin at the same time as the appearance of the island of Ischia, dated approximately 150 kyr [101]. If the emergence of the island of Ischia is used as a calibration point, the resulting mutation rate for *P. dumerilii* would be 12–13% per million years, which is approximately 10 times higher than in most invertebrate taxa [75,102]. Alternatively, we can hypothesize that the 'vent strain' of *P. dumerilii* was already established elsewhere before the vents and the island formed, given the volcanic nature of the whole Gulf of Naples. However, we cannot exclude the possibility of rapid evolution. The distinctiveness of our vent populations makes it unlikely that mating with nearby populations occurs, allowing genetic drift in this population. It is also likely that bottlenecks may have occurred in this population, when the vent was first inhabited by a few individuals, supporting the notion of rapid evolution. Such rapid adaptation along an environmental gradient has been shown to occur on a very fine scale in Trinidadian guppies [103,104]. To resolve the questions of the age of the clade, data from multiple loci and a credible calibration point are required.

On the other hand, in *A. mediterranea*, both the COI and ITS analyses indicate that the vent-inhabiting population is genetically indistinguishable from the nearby control populations. The lack of a strong population structure suggests that local adaptation to the vents may not have taken place in this species, as one would have expected based on the fact that *A. mediterranea* is a brooder [105], and thus more likely to be subjected to the selection pressure of exposure to elevated  $p\text{CO}_2$  during its entire life cycle, from early developmental stages to senescence: see, for example, selective swipes [106,107]. The absence of a distinct 'vent strain' in *A. mediterranea* supports the notion that the ability to colonize areas with elevated  $p\text{CO}_2$  is a result of acclimatization and not adaptation. This suggests that 'phenotypic buffering' *sensu* [108] in some cases may be as good a strategy as adaptation to prevent taxa extinction in the face of elevated  $p\text{CO}_2$  conditions.


### (b) Metabolic responses in $\text{CO}_2$ -tolerant and $\text{CO}_2$ -sensitive species from control areas

We demonstrated that tolerant species, when collected from control areas, maintain their pre-exposure metabolic rate levels during acute exposure to elevated  $p\text{CO}_2$ , thus supporting our initial prediction for these species (type 2 acclimatization/adaptation *sensu* [46]), at least based on the acclimatization regime used here. Our results could be analogous to those of Maas *et al.* [11], who showed that four pteropod species naturally migrating into semi-permanent elevated  $p\text{CO}_2$ /low  $p\text{O}_2$  areas were able to maintain their metabolic rate when exposed to elevated  $p\text{CO}_2$ .

By contrast, the species of polychaetes not found in the  $\text{CO}_2$  vents were unable to maintain the same metabolic rate and displayed either significant upregulation (*S. spallanzani*, +15%—type 3 acclimatization/adaptation *sensu* [46]) or significant downregulation (*Lysidice* spp., approx. –66%—type 1 acclimatization/adaptation *sensu* [46]) of their metabolic rates during acute exposure. Comparably, Maas *et al.* [11] found that the pteropod *Diacria quadridentata* (Blainville, 1821), which does not migrate into semi-permanent elevated  $p\text{CO}_2$ /low  $p\text{O}_2$  areas, responded to the exposure to elevated  $p\text{CO}_2$  by reducing its metabolic rates by approx. 50%. Overall, Maas *et al.*'s study [11], together with this present investigation, suggest that the chemical environment species are acclimatized to *in situ* may influence their resilience to ocean acidification. Our study goes further by supporting the idea that physiological adaptation and phenotypic buffering enable taxa to colonize and persist in chronically elevated  $p\text{CO}_2$  environments.

### (c) The link to past extinctions and future resilience

The physiological ability to preserve metabolic rates to pre-exposure levels while experiencing hypercapnia may be key to species survival in the initial phase of colonization of naturally elevated  $p\text{CO}_2$  areas. Thus, our study supports the idea that taxa possessing well-developed regulatory and homeostasis abilities are most likely to be best able to face future ocean acidification conditions [8,34,36,109,110]. However, the ability to cope with chronic exposure to elevated  $p\text{CO}_2$  conditions appears to be characterized by the acquisition of moderately lower metabolic rates (on average –23% in this study) in both the acclimatized and the adapted species. While metabolic depression in the short term helps an organism to maintain a balance between energy supply and demand [15,16,25,26,111], in the longer term it can



**Figure 6.** Effect of exposure to different  $p\text{CO}_2$  conditions on the mean maximum body mass of the tolerant species *A. mediterranea* and *P. dumerilii* collected from either low  $p\text{CO}_2$ /elevated pH conditions outside the vents or elevated  $p\text{CO}_2$  conditions within the vents. Histograms present mean values  $\pm$  s.e. Asterisk (\*) indicates a significant difference between the mean maximum body mass measured under acidified conditions (A) and control conditions (C), according to a GLM test.

involve the reorganization of its energy budget, often leading to a decrease in its scope for growth and reproduction [2].

We suggest that in the face of elevated  $p\text{CO}_2$  conditions, such as those experienced, for example, during the Permo-Triassic boundary [112–115], the evolution (or development) of moderate metabolic depression may have enabled some marine organisms to persist locally and globally. This may have come, however, at the cost of some life-history traits, such as reduced body size: see [116] for a review on the *Lilliput Effect*. Consistent with this idea, the mean body size of the adult individuals of *P. dumerilii* collected from the  $\text{CO}_2$  vents was approximately 80% lower when compared with that of adult individuals from the non-acidified areas (figure 6;  $F_{1,46} = 14.547$ ;  $p < 0.0001$ ). By contrast, *A. mediterranea* shows no change in body mass (figure 6;  $F_{1,46} = 0.016$ ;  $p = 0.900$ ), against our prediction that acclimatization may come at some cost. In our study, we did not carry out a systematic collection of polychaetes living around the  $\text{CO}_2$  vents, our data requiring further validation; nonetheless, our sampling is the outcome of the haphazard selection of the larger adult individuals we could find. Thus, our measure represents an estimate for the ‘mean maximum body size’ that individuals of *P. dumerilii* and *A. mediterranea* reach when chronically exposed to conditions found inside or outside the  $\text{CO}_2$  vents, including different  $p\text{CO}_2$  levels, altered algal composition and habitat complexity among others [45,117]. Nevertheless, differences in body size patterns observed between *A. mediterranea* and *P. dumerilii*

could be explained by the fact that the strain of *P. dumerilii* from the  $\text{CO}_2$  vents shows overall a higher mean metabolic rate when compared with those of the non-acidified strain. This may suggest an increase in the costs for maintenance and repair in this species, costs that *A. mediterranea* may not incur. As size in several polychaete species defines the maximum number of eggs that a female can produce, it will be important to verify whether individuals from the vent strain could have a reduced reproductive output when compared with individuals from the low  $p\text{CO}_2$  strain.

On the other hand, an extreme increase in metabolism (i.e. *S. spallanzani*) or extreme forms of depression (i.e. *L. collaris* and *L. ninetta*) appear not to support life inside the  $\text{CO}_2$  vents. Thus, past mass extinctions may have stemmed from the inability of some species to maintain their metabolic rate within strict limits and thus their energy budgets. Ultimately, the ability of marine organisms to persist in a rapidly changing ocean [118–120] is largely dependent on the taxa’s ability for rapid physiological adaptation, which could potentially occur, via genetic assimilation of emerging phenotypes [28,103,108,121,122]. However, in the assemblage of polychaetes examined here, phenotypic buffering appears also to be a viable strategy to avoid local extinction. Thus, it appears that both plasticity and adaptation may be key to prevent species’ risk for extinction in the face of ongoing ocean acidification [118], and thus largely determine the fate of marine biodiversity. Nonetheless even within tolerant groups such as the Polychaeta, some species appear sensitive to elevated  $p\text{CO}_2$  and at risk of extinction, as they are unable to cope with ocean acidification [43,45,123–125]. Species extinction will cause shifts in community structure and functions, which may ultimately drive important changes in ecosystem functioning [125,126].

**Acknowledgements.** We thank R. Haslam, M. Hawkins and the staff of the Benthic Ecology research unit (Villa Dorhn, Ischia) for advice and technical support. We particularly thank Captain V. Rando for his outstanding support with all boat operations and for building the *S. spallanzani* transplant chambers. We thank E. Borda for assistance with molecular methods and primer design for *Amphiglena mediterranea*. We are grateful to N.M. Whiteley for the loan of the Oxsense system. We thank C. Ghalambor, F. Melzner, J. Havenhand and an anonymous reviewer for their constructive and useful criticisms on early drafts of this manuscript.

**Data accessibility.** All data are archived with the British Oceanographic Data Centre, <http://www.bodc.ac.uk>.

**Funding statement.** This work was undertaken while P.C. was a recipient of a Research Council UK Research Fellowship to investigate ocean acidification at Plymouth University. J.I.S. was a recipient of RCUK research fund. This project was supported by an ASSEMBLE Grant to P.C. and S.P.S.R, the UKOA NERC grant NE/H017127/1 awarded to J.I.S. and P.C. [Task 1.4 ‘Identify the potential for organism resistance and adaptation to prolonged  $\text{CO}_2$  exposure’ of the NERC Consortium Grant ‘Impacts of ocean acidification on key benthic ecosystems, communities, habitats, species and life cycles’], ENEA internal funding to C.L. and SZN internal funding to M.C.G. The molecular work undertaken on *P. dumerilii* was supported in part by an ASSEMBLE Grant to J.D.H. The molecular work conducted on *A. mediterranea* at TAMUG was funded by the U.S. National Science Foundation (DEB 1036186 to A.S.).

## References

1. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789. (doi:10.1890/03-9000)
2. Sibly RM, Calow P. 1986 *Physiological ecology of animals: an evolutionary approach*. Hong Kong, China: Blackwell Scientific Publications.
3. Calow P, Forbes VE. 1998 How do physiological responses to stress translate into ecological and evolutionary processes. *Comp. Biochem. Physiol.* **120A**, 11–16.

4. Spicer JJ, Gaston KJ. 1999 *Physiological diversity and its ecological implications*. Oxford, UK: Blackwell Science.
5. Gaston KJ. 2003 *The structure and dynamics of geographic ranges*. New York, NY: Oxford University Press Inc.
6. Kooijman SALM. 2010 *Dynamic energy budget theory for metabolic organisation*. Cambridge, UK: Cambridge University Press.
7. Bozinovic F, Calosi P, Spicer JJ. 2011 Physiological correlates of geographical range in animals. *Annu. Rev. Ecol. Syst.* **42**, 155–179. (doi:10.1146/annurev-ecolsys-102710-145055)
8. Melzner F, Gutowska M, Langebuch M, Dupont S, Lucassen M, Thorndyke M, Bleich M, Pörtner H-O. 2009 Physiological basis for high CO<sub>2</sub> tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* **6**, 2313–2331. (doi:10.5194/bg-6-2313-2009)
9. Hendriks IE, Duarte CM, Álvarez M. 2010 Vulnerability of marine biodiversity to ocean acidification: a meta-analysis. *Estuar. Coast. Shelf Sci.* **86**, 157–164. (doi:10.1016/j.ecss.2009.11.022)
10. Kroeker KJ, Kordas RL, Crim RN, Singh GG. 2010 Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecol. Lett.* **13**, 1419–1434. (doi:10.1111/j.1461-0248.2010.01518.x)
11. Maas AE, Wishner KF, Seibel BA. 2012 The metabolic response of pteropods to acidification reflects natural CO<sub>2</sub>-exposure in oxygen minimum zones. *Biogeosciences* **9**, 747–757. (doi:10.5194/bg-9-747-2012)
12. Melatunan S. 2012 'Biochemical, metabolic and morphological responses of the intertidal gastropod *Littorina littorea* to ocean acidification and increased temperature', p. 222. PhD thesis, Marine Biology and Ecology Research Centre, Plymouth University, Plymouth, UK.
13. Reipschläger A, Pörtner H-O. 1996 Metabolic depression during environmental stress: the role of extracellular versus intracellular pH in *Sipunculus nudus*. *J. Exp. Biol.* **199**, 1801–1807.
14. Michaelidis B, Ouzounis C, Palaras A, Pörtner H-O. 2005 Effects of long-term moderate hypercapnia on acid–base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Mar. Ecol. Prog. Ser.* **293**, 109–118. (doi:10.3354/meps293109)
15. Rosa R, Seibel BA. 2008 Synergistic effects of climate-related variables suggest future physiological impairment in a top oceanic predator. *Proc. Natl Acad. Sci. USA* **105**, 20776–20780. (doi:10.1073/pnas.0806886105)
16. Melatunan S, Calosi P, Rundle SD, Moody JA, Widdicombe S. 2011 Exposure to elevated temperature and pCO<sub>2</sub> reduces respiration rate and energy status in the periwinkle *Littorina littorea*. *Physiol. Biochem. Zool.* **84**, 583–594. (doi:10.1086/662680)
17. Nakamura M, Ohki S, Suzuki A, Sakai K. 2011 Coral larvae under ocean acidification: Survival, metabolism, and metamorphosis. *PLoS ONE* **6**, e14521. (doi:10.1371/journal.pone.0014521)
18. Wood HL, Spicer JJ, Widdicombe S. 2008 Ocean acidification may increase calcification rates, but at a cost. *Proc. R. Soc. B* **275**, 1767–1773. (doi:10.1098/rspb.2008.0343)
19. Wood HL, Spicer JJ, Lowe DM, Widdicombe S. 2010 Interaction of ocean acidification and temperature; the high cost of survival in the brittlestar *Ophiura ophiura*. *Mar. Biol.* **157**, 2001–2013. (doi:10.1007/s00227-010-1469-6)
20. Benias E, Ivanina A, Lieb NS, Kurochkin I, Sokolova IM. 2010 Elevated level of carbon dioxide affects metabolism and shell formation in oysters *Crassostrea virginica*. *Mar. Ecol. Prog. Ser.* **419**, 95–108. (doi:10.3354/meps08841)
21. Gutowska MA, Pörtner H-O, Melzner F. 2008 Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO<sub>2</sub>. *Mar. Ecol. Prog. Ser.* **373**, 303–309. (doi:10.3354/meps07782)
22. Marchant HK, Calosi P, Spicer JJ. 2010 Short-term exposure to hypercapnia does not compromise feeding, acid–base balance or respiration of *Patella vulgata* but surprisingly is accompanied by radula damage. *J. Mar. Biol. Assoc. UK* **90**, 1379–1384. (doi:10.1017/S0025315410000457)
23. Thomsen J, Melzner F. 2010 Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*. *Mar. Biol.* **157**, 2667–2676. (doi:10.1007/s00227-010-1527-0)
24. Donohue P, Calosi P, Bates A, Laverock AH, Rastrick SPS, Mark FC, Stroble A, Widdicombe S. 2012 Physiological and behavioural impacts of exposure to elevated pCO<sub>2</sub> on an important ecosystem engineer, the burrowing shrimp *Upogebia deltaura*. *Aquat. Biol.* **15**, 73–86. (doi:10.3354/ab00408)
25. Bishop T, Brand MD. 2000 Processes contributing to metabolic depression in hepatopancreas cells from the snail *Helix aspersa*. *J. Exp. Biol.* **203**, 3603–3612.
26. Seibel BA, Walsh PJ. 2003 Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance. *J. Exp. Biol.* **206**, 641–650. (doi:10.1242/jeb.00141)
27. Guppy M, Withers P. 1999 Metabolic depression in animals: physiological perspectives and biochemical generalizations. *Biol. Rev. Camb. Philos. Soc.* **74**, 1–40. (doi:10.1017/S0006323198005258)
28. Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007 Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct. Ecol.* **21**, 394–407. (doi:10.1111/j.1365-2435.2007.01283.x)
29. Miller GM, Watson S-A, Donelson JM, McCormick MI, Munday PL. 2012 Parental environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature Clim. Change* **2**, 858–861. (doi:10.1038/nclimate1599)
30. Pörtner H-O. 2008 Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar. Ecol. Prog. Ser.* **373**, 203–217. (doi:10.3354/meps07768)
31. Findlay HS, Wood HL, Kendall MA, Spicer JJ, Twitchett RJ, Widdicombe S. 2011 Comparing the impact of high CO<sub>2</sub> on calcium carbonate structures in different marine organisms. *Mar. Biol. Res.* **7**, 565–575. (doi:10.1080/17451000.2010.547200)
32. Melatunan S, Calosi P, Rundle SD, Widdicombe S, Moody JA. 2013 The effects of ocean acidification and elevated temperature on shell plasticity and its energetic basis in an intertidal gastropod. *Mar. Ecol. Prog. Ser.* **472**, 155–168. (doi:10.3354/meps10046)
33. Dove PM, De Yoreo JJ, Weiner S. 2006 *Biomineralization. Reviews in mineralogy and geochemistry, vol. 54* (ed. JJ Rosso). Washington, DC: Mineralogical Society of America.
34. Pane EF, Barry JP. 2007 Extracellular acid–base regulation during short-term hypercapnia is effective in a shallow-water crab, but ineffective in a deep-sea crab. *Mar. Ecol. Prog. Ser.* **334**, 1–9. (doi:10.3354/meps334001)
35. Munday PL, Crawley NE, Nilsson GE. 2009 Interacting effects of elevated temperature and ocean acidification on the aerobic performance of coral reef fishes. *Mar. Ecol. Prog. Ser.* **388**, 235–242. (doi:10.3354/meps08137)
36. Calosi P, Rastrick SPS, Graziano M, Thomas SC, Baggini C, Carter HA, Hall-Spencer JM, Milazzo M, Spicer JJ. In press. Ecophysiology of sea urchins living near shallow water CO<sub>2</sub> vents: investigations of acid–base balance and ionic regulation using in-situ transplantation. *Mar. Pollut. Bull.* (doi:10.1016/j.marpolbul.2012.11.040)
37. Comeau S, Edmunds PJ, Spindel NB, Carpenter RC. 2013 The responses of eight coral reef calcifiers to increasing partial pressure of CO<sub>2</sub> do not exhibit a tipping point. *Limnol. Oceanogr.* **58**, 388–398. (doi:10.4319/lo.2013.58.1.0388)
38. Byrne M, Lamare M, Winter D, Dworjanyn SA, Uthicke S. 2013 The stunting effect of a high CO<sub>2</sub> ocean on calcification and development in sea urchin larvae, a synthesis from the tropics to the poles. *Phil. Trans. R. Soc. B* **368**, 20120439. (doi:10.1098/rstb.2012.0439)
39. Widdicombe S, Needham HR. 2007 Impact of CO<sub>2</sub>-induced seawater acidification on the burrowing activity of *Nereis virens* and sediment nutrient flux. *Mar. Ecol. Prog. Ser.* **341**, 111–122. (doi:10.3354/meps341111)
40. Lane A, Mukherjee J, Chan VS, Thiyagarajan V. In press. Decreased pH does not alter metamorphosis but compromises juvenile calcification of the tube worm *Hydroides elegans*. *Mar. Biol.* (doi:10.1007/s00227-012-2056-9)
41. Godbold JA, Solan M. 2013 Long-term effects of warming and ocean acidification are modified by seasonal variation in species responses and environmental conditions. *Phil. Trans. R. Soc. B* **368**, 20130186. (doi:10.1098/rstb.2013.0186)
42. Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, Turner SM, Rowley SJ, Tedesco D, Buia M-C. 2008 Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* **454**, 96–99. (doi:10.1038/nature07051)



43. Cigliano M, Gambi MC, Rodolfo-Metalpa R, Patti FP, Hall-Spencer JM. 2010 Effects of ocean acidification on invertebrate settlement at natural volcanic CO<sub>2</sub> vents. *Mar. Biol.* **157**, 2489–2502. (doi:10.1007/s00227-010-1513-6)
44. Kroeker KJ, Micheli F, Gambi MC. 2013 Ocean acidification causes ecosystem shifts via altered competitive interactions. *Nat. Clim. Change* **3**, 156–159. (doi:10.1038/nclimate1680)
45. Kroeker KJ, Micheli F, Gambi MC, Martz TR. 2011 Divergent ecosystem responses within a benthic marine community to ocean acidification. *Proc. Natl Acad. Sci. USA* **108**, 14 515–14 520. (doi:10.1073/pnas.1107789108)
46. Precht H. 1958 Concepts of the temperature adaptation of unchanging reaction systems of cold-blooded animals. In *Physiological adaptation* (ed. CL Prosser), pp. 50–78. Washington, DC: American Physiological Society.
47. Gambi MC, Lorenti M, Russo GF, Scipione MB, Zupo V. 1992 Depth and seasonal distribution of some groups of the vagile fauna of *Posidonia oceanica* leaf stratum: structural and feeding guild analyses. *Mar. Ecol.* **13**, 17–39. (doi:10.1111/j.1439-0485.1992.tb00337.x)
48. Giangrande A, Delos AL, Frascchetti S, Musco L, Licciano M, Terlizzi A. 2003 Polychaete assemblages along a rocky shore on the South Adriatic coast (Mediterranean Sea): patterns of spatial distribution. *Mar. Biol.* **143**, 1109–1116. (doi:10.1007/s00227-003-1162-0)
49. Giangrande A, Frascchetti S, Terlizzi A. 2002 Local recruitment differences in *Platynereis dumerilii* (Polychaeta, Nereididae) and their consequences for population structure. *Ital. J. Zool.* **69**, 133–139. (doi:10.1080/1125000209356450)
50. Gambi MC. 2002 Spatio-temporal distribution and ecological role of polychaete borers of *Posidonia oceanica* (L.) Delile scales. *Bull. Mar. Sci.* **71**, 1323–1331.
51. Gambi MC, Cafiero G. 2001 Functional diversity in the *Posidonia oceanica* ecosystem: an example with polychaete borers of the scales. In *Mediterranean ecosystems: structure and processes* (eds FM Faranda, L Guglielmo, G Spezie), pp. 399–405. Milano, Italy: Springer.
52. Bocchetti R, Fattorini D, Gambi MC, Regoli F. 2004 Trace metal concentrations and susceptibility to oxidative stress in the polychaete *Sabella spallanzanii* (Gmelin) (Sabellidae): potential role of antioxidants in revealing stressful environmental conditions in the Mediterranean. *Arch. Environ. Contam. Toxicol.* **46**, 353–361. (doi:10.1007/s00244-003-2300-x)
53. Giangrande A, Licciano M, Pagliara P, Gambi MC. 2000 Gametogenesis and larval development in *Sabella spallanzanii* (Polychaeta: Sabellidae) from the Mediterranean Sea. *Mar. Biol.* **136**, 847–861. (doi:10.1007/s002279900251)
54. Alsop DH, Wood CM. 1997 The interactive effects of feeding and exercise on oxygen consumption, swimming performance and protein usage in juvenile rainbow trout. *J. Exp. Biol.* **200**, 2337–2346.
55. McCue MD. 2006 Specific dynamic action: a century of investigation. *Comp. Biochem. Physiol.* **144A**, 381–394.
56. Pierrot D, Lewis E, Wallace DWR. 2006 *MS Excel program developed for CO<sub>2</sub> system calculations, ORNL/CDIAC-105*. Oak Ridge, TN: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy.
57. Mehrbach C, Culbertson CH, Hawley JE, Pytkowicz RM. 1973 Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol. Oceanogr.* **18**, 897–907. (doi:10.4319/lo.1973.18.6.0897)
58. Dickson AG, Millero FJ. 1987 A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Res.* **34**, 1733–1743. (doi:10.1016/0198-0149(87)90021-5)
59. Dickson AG. 1990 Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K. *Deep-Sea Res.* **37**, 755–766. (doi:10.1016/0198-0149(90)90004-F)
60. Rodolfo-Metalpa R, Lombardi C, Cocito S, Hall-Spencer JM, Gambi MC. 2010 Effects of ocean acidification and high temperatures on the bryozoan *Myriapora truncata* at natural CO<sub>2</sub> vents. *Mar. Ecol.* **31**, 447–456. (doi:10.1111/j.1439-0485.2009.00354.x)
61. Rodolfo-Metalpa R *et al.* 2011 Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat. Clim. Change* **1**, 308–312. (doi:10.1038/nclimate1200)
62. Lombardi C, Gambi MC, Vasapollo C, Taylor AC, Cocito S. 2011 Skeletal alterations and polymorphism in a Mediterranean bryozoan at natural CO<sub>2</sub> vents. *Zoomorphology* **130**, 135–145. (doi:10.1007/s00435-011-0127-y)
63. Spicer JI, Eriksson SP. 2003 Does the development of respiratory regulation always accompany the transition from pelagic larvae to benthic fossorial postlarvae in the Norway lobster *Nephrops norvegicus* (L.)? *J. Exp. Mar. Biol. Ecol.* **295**, 219–243. (doi:10.1016/S0022-0981(03)00296-X)
64. Calosi P, Morrirt D, Chelazzi G, Ugolini A. 2007 Physiological capacity and environmental tolerance in two sandhopper species with contrasting geographical ranges: *Talitrus saltator* and *Talorchestia ugolini*. *Mar. Biol.* **151**, 1647–1655. (doi:10.1007/s00227-006-0582-z)
65. Rastrick SPS, Whiteley NM. 2011 Congeneric amphipods show differing abilities to maintain metabolic rates with latitude. *Physiol. Biochem. Zool.* **84**, 154–165. (doi:10.1086/658857)
66. Kamenos NA, Calosi P, Moore PG. 2006 Substratum-mediated heart rate responses of an invertebrate to predation threat. *Anim. Behav.* **71**, 809–813. (doi:10.1016/j.anbehav.2005.05.026)
67. Sokal R, Rohlf FJ. 1995 *Biometry: the principles and practice of statistics in biological research*. New York, NY: Freeman and Company.
68. Underwood AJ. 1997 *Experiments in ecology: their logical design and interpretation using analysis of variance*. Cambridge, UK: Cambridge University Press.
69. Montero-Pau J, Gómez A, Muñoz J. 2008 Application of an inexpensive and high-throughput genomic DNA extraction method for the molecular ecology of zooplanktonic diapausing eggs. *Limnol. Oceanogr. Meth.* **6**, 218–222. (doi:10.4319/lom.2008.6.218)
70. Ivanova NV, Zemlak TS, Hanner RH, Hebert PDN. 2007 Universal primer cocktails for fish DNA barcoding. *Mol. Ecol. Notes* **7**, 544–548. (doi:10.1111/j.1471-8286.2007.01748.x)
71. Folmer O, Black M, Lutz R, Vrijenhoek R. 1994 DNA primers for amplification of mitochondrial cytochrome c oxidase subunit 1 from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* **3**, 294–299.
72. Hall TA. 1999 BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acid S.* **41**, 95–98.
73. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011 MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **24**, 1596–1599. (doi:10.1093/molbev/msm092)
74. Rambaut A. 2009 *FigTree v.1.3.1. Tree figure drawing tool*. UK: Institute of Evolutionary Biology, University of Edinburgh.
75. Knowlton N, Weigt LA, Solorzano LA, Mills DK, Bermingham E. 1993 Divergence in proteins, mitochondrial DNA, and reproductive compatibility across the isthmus of Panama. *Science* **260**, 1629–1632. (doi:10.1126/science.8503007)
76. Pespeni MH *et al.* 2013 Evolutionary change during experimental ocean acidification. *Proc. Natl Acad. Sci. USA* **110**, 6937–6942. (doi:10.1073/pnas.1220673110)
77. Pörtner H-O, Farrell AP. 2008 Physiology and climate change. *Science* **322**, 690–692. (doi:10.1126/science.1163156)
78. Melzner F, Thomsen J, Koeve W, Oschlies A, Gutowska M, Bange H, Hansen H, Körtzinger A. In press. Future ocean acidification will be amplified by hypoxia in coastal habitats. *Mar. Biol.* (doi:10.1007/s00227-012-1954-1)
79. Lohbeck KT, Riebesell U, Reusch TBH. 2012 Adaptive evolution of a key phytoplankton species to ocean acidification. *Nat. Geosci.* **5**, 346–351. (doi:10.1038/ngeo1441)
80. Benner I, Diner RE, Lefebvre SC, Li D, Komada T, Carpenter EJ, Stillman JH. 2013 *Emiliania huxleyi* increases calcification but not expression of calcification-related genes in long-term exposure to elevated temperature and pCO<sub>2</sub>. *Phil. Trans. R. Soc. B* **368**, 20130049. (doi:10.1098/rsth.2013.0049)

81. Tatters AO *et al.* 2013 Short- and long-term conditioning of a temperate marine diatom community to acidification and warming. *Phil. Trans. R. Soc. B* **368**, 20120437. (doi:10.1098/rstb.2012.0437)
82. Pistevo JCA, Calosi P, Widdicombe S, Bishop JDD. 2011 Will variation among genetic individuals influence species responses to global climate change? *Oikos* **120**, 675–689. (doi:10.1111/j.1600-0706.2010.19470.x)
83. Sunday JM, Crim RN, Harley CDG, Hart MW. 2011 Quantifying rates of evolutionary adaptation in response to ocean acidification. *PLoS ONE* **6**, e22881. (doi:10.1371/journal.pone.0022881)
84. Foo SA, Dworjanyn SA, Poore AGB, Byrne M. 2012 Adaptive capacity of the habitat modifying sea urchin *Centrostephanus rogersii* to ocean warming and ocean acidification: performance of early embryos. *PLoS ONE* **7**, e42497. (doi:10.1371/journal.pone.0042497)
85. Schlegel P, Havenhand JN, Gillings MR, Williamson JE. 2012 Individual variability in reproductive success determines winners and losers under ocean acidification: a case study with sea urchins. *PLoS ONE* **7**, e53118. (doi:10.1371/journal.pone.0053118)
86. Padilla-Gamiño JL, Kelly MW, Evans TG, Hofmann GE. 2013 Temperature and CO<sub>2</sub> additively regulate physiology, morphology and genomic responses of larval sea urchins, *Strongylocentrotus purpuratus*. *Proc. R. Soc. B* **280**, 20130155. (doi:10.1098/rspb.2013.0155)
87. Kelly MW, Padilla-Gamiño JL, Hofmann GE. 2013 Natural variation, and the capacity to adapt to ocean acidification in the keystone sea urchin *Strongylocentrotus purpuratus*. *Glob. Change Biol.* **19**, 2536–2546. (doi:10.1111/gcb.12251)
88. Hoffmann AA. 1995 Acclimation: increasing survival at a cost. *Trends Ecol. Evol.* **10**, 1–2. (doi:10.1016/S0169-5347(00)88949-1)
89. Dewitt TJ, Sih A, Wilson DS. 1998 Costs and limits of phenotypic plasticity. *Trends Ecol. Evol.* **13**, 77–81. (doi:10.1016/S0169-5347(97)01274-3)
90. Deigweier K, Hirse T, Bock C, Lucassen M, Pörtner H-O. 2009 Hypercapnia induced shifts in gill energy budgets of Antarctic notothenioids. *J. Comp. Physiol.* **180B**, 347–359.
91. Stumpp M, Trübenbach K, Brennecke D, Hu MY, Melzner F. 2012 Resource allocation and extracellular acid–base status in the sea urchin *Strongylocentrotus droebachiensis* in response to CO<sub>2</sub> induced seawater acidification. *Aquat. Toxicol.* **110–111**, 194–207. (doi:10.1016/j.aquatox.2011.12.020)
92. Schulze A. 2006 Phylogeny and genetic diversity of palolo worms (*Palola*, Eunicidae, Polychaeta) from the tropical North Pacific and the Caribbean. *Biol. Bull.* **210**, 25–37. (doi:10.2307/4134534)
93. Audzijonyte A, Ovcarenko I, Bastrop R, Väinölä R. 2008 Two cryptic species of the *Hediste diversicolor* group (Polychaeta, Nereididae) in the Baltic Sea, with mitochondrial signatures of different population histories. *Mar. Biol.* **155**, 599–612. (doi:10.1007/s00227-008-1055-3)
94. Barroso R, Klautau M, Solé-Cava A, Paiva PC. 2010 *Eurythoe complanata* (Polychaeta: Amphinomididae), the ‘cosmopolitan’ fireworm, consists of at least three cryptic species. *Mar. Biol.* **157**, 69–80. (doi:10.1007/s00227-009-1296-9)
95. Pires A, Paxton H, Quintino V, Rodrigues AM. 2010 *Diopatra* (Annelida: Onuphidae) diversity in European waters with the description of *Diopatra micrura*, new species. *Zootaxa* **2395**, 17–33.
96. Nygren A, Pleijel F. 2011 From one to ten in a single stroke—resolving the European *Eumida sanguinea* (Phylloclodidae, Annelida) species complex. *Mol. Phylogenet. Evol.* **58**, 132–141. (doi:10.1016/j.ympev.2010.10.010)
97. Nygren A, Norlinder E, Panova M, Pleijel F. 2011 Colour polymorphism in the polychaete *Harmothoe imbricata* (Linnaeus, 1767). *Mar. Biol. Res.* **7**, 54–62. (doi:10.1080/17451001003713555)
98. Fischer A, Henrich T, Arendt D. 2010 The normal development of *Platynereis dumerilii* (Nereididae, Annelida). *Front. Zool.* **7**, 31. (doi:10.1186/1742-9994-7-31)
99. Carr CM, Hardy SM, Brown TM, Macdonald TA, Hebert PDN. 2011 A tri-oceanic perspective: DNA barcoding reveals geographic structure and cryptic diversity in Canadian polychaetes. *PLoS ONE* **6**, e22232. (doi:10.1371/journal.pone.0022232)
100. Chevaldonné P, Jollivet D, Desbruyères D, Lutz RA, Vrijenhoek RC. 2002 Sister-species of eastern Pacific hydrothermal vent worms (Ampharetidae, Alvinellidae, Vestimentifera) provide new mitochondrial COI clock calibration. *Cah. Biol. Mar.* **43**, 367–370.
101. Gillot P-Y, Chiesa S, Pasquaré G, Vezzoli L. 1982 33,000 yr K/Ar dating of the volcano-tectonic horst of the isle of Ischia, Gulf of Naples. *Nature* **229**, 242–245. (doi:10.1038/299242a0)
102. Knowlton N, Weigt LA. 1998 New dates and new rates for divergence across the isthmus of Panama. *Proc. R. Soc. Lond. B* **265**, 2257–2263. (doi:10.1098/rspb.1998.0568)
103. Reznick DN, Ghalambor CK. 2001 The population ecology of contemporary adaptations: what empirical studies reveal about the conditions that promote adaptive evolution. *Genetica* **112–113**, 183–198.
104. Torres-Dowdall J, Handelsman CA, Reznick DN, Ghalambor CK. 2012 Local adaptation and the evolution of phenotypic plasticity in Trinidadian guppies (*Poecilia reticulata*). *Evolution* **66**, 3432–3443. (doi:10.1111/j.1558-5646.2012.01694.x)
105. Rouse GW, Gambi MC. 1998 Evolution of reproductive features and larval development in the genus *Amphiglena* (Polychaeta: Sabellidae). *Mar. Biol.* **131**, 743–754. (doi:10.1007/s002270050365)
106. Schmidt PS, Bertness MD, Rand DM. 2000 Environmental heterogeneity and balancing selection in the acorn barnacle *Semibalanus balanoides*. *Proc. R. Soc. Lond. B* **267**, 379–384. (doi:10.1098/rspb.2000.1012)
107. Gagnaire P-A, Normandeau E, Côté C, Møller Hansen M, Bernatchez L. 2012 The genetic consequences of spatially varying selection in the panmictic American Eel (*Anguilla rostrata*). *Genetics* **190**, 725–736. (doi:10.1534/genetics.111.134825)
108. Waddington CH. 1942 Canalization of development and the inheritance of acquired characters. *Nature* **150**, 563–565. (doi:10.1038/150563a0)
109. Widdicombe S, Spicer JL. 2008 Predicting the impact of ocean acidification on benthic biodiversity: what can animal physiology tell us? *J. Exp. Mar. Biol. Ecol.* **366**, 187–197. (doi:10.1016/j.jembe.2008.07.024)
110. Whiteley NM. 2011 Physiological and ecological responses of crustaceans to ocean acidification. *Mar. Ecol. Prog. Ser.* **430**, 257–271. (doi:10.3354/meps09185)
111. Pörtner H-O, Reipschäger A, Heisler N. 1998 Acid–base regulation, metabolism and energetics in *Sipunculus nudus* as a function of ambient carbon dioxide level. *J. Exp. Biol.* **201**, 43–55.
112. Twitchett RJ, Looy CV, Morante R, Visscher H, Wignall PB. 2001 Rapid and synchronous collapse of marine and terrestrial ecosystems during the end-Permian biotic crisis. *Geology* **29**, 351–354. (doi:10.1130/0091-7613(2001)029<0351:RASCOSM>2.0.CO;2)
113. Benton MJ, Twitchett RJ. 2003 How to kill (almost) all life: the end-Permian extinction event. *Trends Ecol. Evol.* **18**, 358–365. (doi:10.1016/S0169-5347(03)00093-4)
114. Veron JEN. 2008 Mass extinctions and ocean acidification: biological constraints on geological dilemmas. *Coral Reefs* **27**, 459–472. (doi:10.1007/s00338-008-0381-8)
115. Hoegh-Guldberg O. 2012 Coral reefs, climate change, and mass extinction. In *Saving a million species: extinction risk from climate change* (ed. I Hannah), pp. 261–283. Washington, DC: Island Press.
116. Harries PJ, Knorr PO. 2009 What does the ‘Lilliput Effect’ mean? *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **284**, 4–10. (doi:10.1016/j.palaeo.2009.08.021)
117. Porzio L, Buia MC, Hall-Spencer JM. 2011 Effects of ocean acidification on macroalgal communities. *J. Exp. Mar. Biol. Ecol.* **400**, 278–287. (doi:10.1016/j.jembe.2011.02.011)
118. Caldeira K, Wickett ME. 2003 Anthropogenic carbon and ocean pH. *Nature* **425**, 365. (doi:10.1038/425365a)
119. Raven J, Caldeira K, Elderfield H, Hoegh-Guldberg O, Liss P, Riebesell U, Shepherd J, Turley C, Watson A. 2005 Ocean acidification due to increasing atmospheric carbon dioxide. *The Royal Society policy document 12 : 05*. Cardiff, UK: Clyvedon Press.
120. IPCC. 2007 In *Climate change 2007; The physical science basis. Contribution of Working Group I to the 4th Assessment Report of the Intergovernmental Panel on Climate Change* (eds S Solomon, D Qin,

- M Manning, M Marquis, K Averyt, MMB Tignor, HL Miller, Z Chen). Cambridge, UK: Cambridge University Press.
121. Torres Dowdall J, Handelsman CA, Ruell EW, Auer SK, Reznick DN, Ghalambor CK. 2012 Fine-scale local adaptation in life histories along a continuous environmental gradient in Trinidadian guppies. *Funct. Ecol.* **26**, 616–627. (doi:10.1111/j.1365-2435.2012.01980.x)
122. Pigliucci M, Murren CJ, Schlichting CD. 2006 Phenotypic plasticity and evolution by genetic assimilation. *J. Exp. Biol.* **209**, 2362–2367. (doi:10.1242/jeb.02070)
123. Ricevuto E, Lorenti M, Patti FP, Scipione MB, Gambi MC. 2012 Temporal trends of benthic invertebrate settlement along a gradient of ocean acidification at natural CO<sub>2</sub> vents (Tyrrhenian sea). *Biol. Mar. Mediterr.* **19**, 49–52.
124. Hale R, Calosi P, McNeill L, Mieszkowska N, Widdicombe S. 2011 Predicted levels of future ocean acidification and temperature rise could alter community structure and biodiversity in marine benthic communities. *Oikos* **120**, 661–674. (doi:10.1111/j.1600-0706.2010.19469.x)
125. Christen N, Calosi P, McNeill CL, Widdicombe S. In press. Structural and functional vulnerability to elevated pCO<sub>2</sub> in marine benthic communities. *Mar. Biol.* (doi:10.1007/s00227-012-2097-0)
126. Solan M, Cardinale BJ, Downing AL, Engelhardt KAM, Ruesink JL, Srivastava DS. 2004 Extinction and ecosystem function in the marine benthos. *Science* **306**, 1177–1180. (doi:10.1126/science.1103960)