



GROWTH FACTORS, CYTOKINES, AND CELL CYCLE MOLECULES

# Decidual Cell Regulation of Natural Killer Cell—Recruiting Chemokines

## *Implications for the Pathogenesis and Prediction of Preeclampsia*

Charles J. Lockwood,\* S. Joseph Huang,\* Chie-Pein Chen,<sup>†</sup> Yingqun Huang,<sup>‡</sup> Jie Xu,<sup>‡</sup> Saeed Faramarzi,\* Ozlem Kayisli,\* Umit Kayisli,\* Louise Koopman,<sup>§</sup> Dineke Smedts,<sup>¶</sup> Lynn F. Buchwalder,<sup>‡</sup> and Frederick Schatz\*

From the Department of Obstetrics and Gynecology,\* Ohio State University College of Medicine, Columbus, Ohio; the Department of Obstetrics and Gynecology,<sup>†</sup> Mackay Memorial Hospital, Taipei, Taiwan; the Department of Obstetrics, Gynecology and Reproductive Sciences,<sup>‡</sup> Yale University School of Medicine, New Haven, Connecticut; Celgene Avilomics Research,<sup>§</sup> Bedford, Massachusetts; and the Department of Obstetrics and Gynecology,<sup>¶</sup> Erasmus MC Rotterdam, Rotterdam, the Netherlands

Accepted for publication  
May 28, 2013.

Address correspondence to  
Frederick Schatz, Ph.D.,  
Department of Obstetrics and  
Gynecology, Biomedical  
Research Tower room 814, 460  
West 12th Av, Columbus,  
OH 43210. E-mail: [Frederick.Schatz@osumc.edu](mailto:Frederick.Schatz@osumc.edu).

First trimester human decidua is composed of decidual cells, CD56<sup>bright</sup>CD16<sup>-</sup> decidual natural killer (dNK) cells, and macrophages. Decidual cells incubated with NK cell–derived IFN- $\gamma$  and either macrophage-derived TNF- $\alpha$  or IL-1 $\beta$  synergistically enhanced mRNA and protein expression of IP-10 and I-TAC. Both chemokines recruit CXCR3-expressing NK cells. This synergy required IFN- $\gamma$  receptor 1 and 2 mediation via JAK/STAT and NF $\kappa$ B signaling pathways. However, synergy was not observed on neutrophil, monocyte, and NK cell–recruiting chemokines. Immunostaining of first trimester decidua localized IP-10, I-TAC, IFN- $\gamma$ R1, and -R2 to vimentin-positive decidual cells versus cytokeratin-positive interstitial trophoblasts. Flow cytometry identified high CXCR3 levels on dNK cells and minority peripheral CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells and intermediate CXCR3 levels on the majority of CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. Incubation of pNK cells with either IP-10 or I-TAC elicited concentration-dependent enhanced CXCR3 levels and migration of both pNK cell subsets that peaked at 10 ng/mL, whereas each chemokine at a concentration of 50 ng/mL inhibited CXCR3 expression and pNK cell migration. Deciduae from women with preeclampsia, a leading cause of maternal and fetal morbidity and mortality, displayed significantly lower dNK cell numbers and higher IP-10 and I-TAC levels versus gestational age-matched controls. Significantly elevated IP-10 levels in first trimester sera from women eventually developing preeclampsia compared with controls, identifying IP-10 as a novel, robust early predictor of preeclampsia. (*Am J Pathol* 2013, 183: 841–856; <http://dx.doi.org/10.1016/j.ajpath.2013.05.029>)

In normal human pregnancy, blastocyst-derived extravillous cytotrophoblasts (EVTs) traverse the underlying decidua and inner third of the myometrium. As they cross the decidua, EVT's detach from anchoring placental villous columns, then breach spiral arteries and arterioles to mediate replacement of the smooth muscle tunica media and endothelium. This invasive process can occur either from the vessel lumen into the tunica media, mediated by endovascular EVT's, or from the surrounding decidualized stroma into the tunica media, mediated by interstitial EVT's. On entering the vessel, the epithelial cell adhesion molecule

phenotype of trophoblasts is converted to an endothelial cell–like adhesion molecule phenotype,<sup>1</sup> and spiral vessels are transformed into low-resistance, high-capacity conduits that increase uteroplacental blood flow to the developing fetal–placental unit.<sup>1,2</sup>

Preeclampsia, a major cause of maternal and perinatal morbidity and mortality,<sup>3</sup> is frequently associated with shallow trophoblast invasion leading to incomplete uterine

Supported by NIH grants R01HD33937-12 (C.J.L.), Project 2 PO1HD054713-01A1 (C.J.L.), and R01HD056123-04 (S.J.H.).

vascular remodeling.<sup>4</sup> The resulting decreased uteroplacental blood flow can elicit fetal growth restriction and/or elaboration of antiangiogenic and proinflammatory placental factors that mediate the maternal syndrome of hypertension and proteinuria, which usually occurs later in pregnancy and can produce end-organ damage.<sup>5</sup>

At the human implantation site, the decidua is composed primarily of resident decidual cells (50%) and a diverse immune cell population (40%). The latter is dominated by decidual natural killer (dNK) cells (70%), macrophages (20%), and T lymphocytes (10%) with small percentages of dendritic cells and B lymphocytes.<sup>6</sup> Unlike the major antigen-presenting cells, macrophages and dendritic cells, NK cells act as specialized lymphocytes and normally mediate innate immunity by killing tumor and virus-infected cells without prior sensitization before the onset of T- and B-cell-mediated adaptive immunity. In the circulation, NK cells comprise approximately 5% to 15% of the lymphocyte population and consist primarily of two functionally distinct subsets. The majority, CD56<sup>dim</sup>CD16<sup>+</sup> peripheral NK (pNK) cells (90%), exhibit greater cytotoxicity, express high levels of killer cell immunoglobulin-like receptors (KIRs), as well as CD57, and usually do not secrete cytokines. By contrast, the absence of CD16 expression by the minority, less mature, CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells (10%), accounts for their inability to mediate antibody-dependent cell toxicity.<sup>7</sup> These CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells do not display KIRs, but express low levels of perforin and high levels of the CD94/NKG2 receptor and adhesion-mediating L-selectin.<sup>8</sup> They also serve as the major pNK cell source of secreted immunoregulatory cytokines. Chief among these is interferon-gamma (IFN- $\gamma$ ). This prototypic NK cell cytokine is expressed by CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells in response to IL-12 acting in concert with either other cytokines (ie, IL-1, IL-2, IL-15, or IL-18) or engagement of either the CD16 (Fc $\gamma$ RIIIa) or NKG2D pNK cell-activating receptors.<sup>9</sup> Recently, the microRNA (miR155) was also shown to function as a positive regulator of IFN- $\gamma$  expression in pNK cells.<sup>10</sup> Other immunoregulatory cytokines expressed by CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells include tumor necrosis factor- $\beta$  (TNF- $\beta$ ), granulocyte-macrophage colony stimulating factor (GM-CSF), and IL-10 and -13.<sup>7</sup>

Like the minority circulating NK cell population, approximately 80% of dNK cells are also CD56<sup>bright</sup>CD16<sup>-</sup>.<sup>7,11</sup> Extensive investigation indicates that dNK cells represent a unique immune cell subtype that plays a crucial pregnancy-supporting role by fostering immune tolerance of the semiallogeneic fetal-placental unit while promoting EVT invasion and spiral artery and arteriole remodeling via expression of vascular endothelial and placental growth factors.<sup>7,11–13</sup>

The current study postulates that at the human implantation site, targeting of the majority resident decidual cells by paracrine effectors, secreted primarily by dNK cells and macrophages, recruits additional CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells into the decidua. In support of this hypothesis, flow

cytometric analysis singled out high CXCR3 expression by CD56<sup>bright</sup>CD16<sup>-</sup> dNK cells and CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells, and lower expression by CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. Complementing these observations, mRNA and protein expression of IFN- $\gamma$ -induced protein 10 (IP-10/CXCL10) and IFN-inducible T cell- $\alpha$  chemoattractant (I-TAC/CXCL11), chemokines that specifically recruit CXCR3-bearing pNK cells,<sup>14</sup> proved to be synergistically enhanced during incubation of human leukocyte-free first trimester decidual cells with the primary NK cell-derived cytokine, IFN- $\gamma$ ,<sup>12</sup> and either of the primary macrophage-derived cytokines, TNF- $\alpha$  or IL-1 $\beta$ .<sup>15</sup> These observations were augmented by: i) evaluating the potential involvement of both IFN- $\gamma$  receptors (IFN- $\gamma$ R1 and IFN- $\gamma$ R2),<sup>16</sup> as well as signaling pathways in mediating this synergistic up-regulation of IP-10 and I-TAC in decidual cells; ii) functional studies that assessed the individual and combined effects of IP-10 and I-TAC on CXCR3 expression and migration of both subsets of pNK cells; iii) *in situ* studies that assessed cellular localization of IP-10, I-TAC, and IFN- $\gamma$ R1 and -R2 at the human implantation site and compared IP-10 and I-TAC levels and the abundance of CD56<sup>bright</sup>CD16<sup>-</sup> NK cells in preeclamptic versus gestational age-matched control decidua; and iv) measurement of IP-10 and I-TAC levels in first trimester sera to determine whether either chemokine is an early predictor of preeclampsia.

## Materials and Methods

### Tissues and Blood Samples

Decidual samples obtained from patients undergoing termination of pregnancy (6 to 12 weeks gestation) at the New York University Medical Center and Yale-New Haven Hospital, under New York University Institutional Review Board and Yale Human Investigative Committee approval, were used to isolate decidual cells and for flow cytometric analysis of dNK cells. Decidual samples (6 to 12 weeks and 37 to 40 weeks) were also obtained under institutional review board approval at the Mackay Memorial Hospital in Taiwan and used for immunostaining and tissue enzyme-linked immunosorbent assays (ELISAs). Peripheral blood samples from volunteers were obtained from the Harvard Department of Molecular and Cellular Biology for use in flow cytometric analysis, and from the Yale-New Haven Hospital for use in measuring CXCR3 expression by NK cells, as well as their migration in a Transwell assay, under institutional review board and human investigative committee approval, respectively.

### Isolation of Decidual Cells and NK Cells

Decidual tissues derived from first trimester elective terminations were washed in sterile PBS until clear of blood, minced, and then digested with 0.1% collagenase type IV and 0.01% DNase (Sigma-Aldrich, St. Louis, MO) in RPMI

1640 containing 20 µg/mL penicillin/streptomycin, 1 µL/mL fungizone (Invitrogen, Carlsbad, CA) in a 37°C shaking water bath for 30 minutes followed by washing with PBS. The digestate was subjected to consecutive filtration through 100-µm, 70-µm, and 40-µm Millipore filters (Millipore, Billerica, MA). The resulting pellet was resuspended in complete RPMI (Invitrogen), then seeded onto six 10-cm Petri dishes and incubated at 37°C in a humidified 95% O<sub>2</sub>:5% CO<sub>2</sub> incubator for 2 hours to enrich for decidual cells (plastic adherent) and lymphocytes (nonadherent). The leukocyte-enriched supernatants were washed with complete RPMI and cell pellets resuspended in complete RPMI, layered over Ficoll-Hypaque (Pharmacia and Upjohn, Chicago, IL), and centrifuged for 20 minutes at 509 × *g*. The decidual leukocyte band at the interface was washed twice with PBS plus 2% fetal calf serum, and then subjected to flow cytometric analysis (see below).

Adherent decidual cells were detached with 0.1% trypsin-EDTA (Sigma-Aldrich), resuspended in RPMI, grown to confluence on polystyrene tissue culture dishes, harvested using trypsin-EDTA, and analyzed by flow cytometry with anti-CD45 and anti-CD14 monoclonal antibodies (BD Pharmingen, San Diego, CA) to monitor the presence of leukocytes after each passage. After three to four passages, cell cultures were found to be leukocyte-free (<1%). Confluent decidual cells were vimentin-positive and cytokeratin-negative, and displayed progesterone-induced decidualization-related morphological and biochemical changes, including elevated prolactin and plasminogen activator inhibitor-1 (PAI-1), and reduced interstitial collagenase and stromelysin-1 expression (results not shown). Cell aliquots were frozen in fetal calf serum/dimethyl sulfoxide (9:1 ratio) (Sigma-Aldrich) and stored in liquid nitrogen.

As previously described, peripheral blood from anonymous healthy reproductive-age donors was used to isolate NK cells using Ficoll followed by enrichment for NK cells with RosetteSep according to the manufacturer's instructions (Stem Cell Technologies, Vancouver, BC, Canada).<sup>17</sup>

## Experimental Incubations

Thawed decidual cells were incubated in basal medium (BM), a phenol red-free 1:1 v/v mix of Dulbecco's modified Eagle's medium (Invitrogen) and Ham's F-12 (Flow Labs, Rockville, MD), with 100 U/mL penicillin, 100 µg/mL streptomycin, 0.25 µg/mL fungizone supplemented with 10% charcoal-stripped calf serum (BMS). After two additional passages, confluent cultures were incubated in parallel in BMS containing 10<sup>-8</sup> mol/L estradiol (E2) with 10<sup>-7</sup> mol/L medroxyprogesterone acetate (MPA) (Sigma-Aldrich), which was used in place of progesterone because of its greater stability in culture.<sup>18</sup> After 7 days, the cultures were washed twice with HBSS to remove residual serum elements and then switched to a defined medium consisting of BM plus ITS+ (Collaborative Research, Waltham, MA), 5 µmol/L FeSO<sub>4</sub>, 0.5 µmol/L ZnSO<sub>4</sub>, 1 nmol/L CuSO<sub>4</sub>, 20 nmol/L

Na<sub>2</sub>SeO<sub>3</sub>, trace elements (Invitrogen), 50 µg/mL ascorbic acid (Sigma-Aldrich), and 50 ng/mL epidermal growth factor (Becton-Dickinson, Bedford, MA) with steroids alone or with 0.01 to 10 ng/mL IFN-γ, IL-1β, TNF-α, or with IFN-γ plus IL-1β or TNF-α (R&D Systems). After the incubation, cells were harvested by scraping into ice-cold PBS, pelleted, and extracted in ice-cold lysis buffer. Conditioned medium supernatants and cell lysates were stored at -70°C. Total RNA was extracted from parallel incubations with Tri Reagent (Sigma-Aldrich).

## Signaling Pathway Mediation of IP-10 and I-TAC Expression

Confluent first trimester decidual cells were pretreated in defined medium ± 10 µmol/L NFκB inhibitor BAY-11-7082 (BAY; EMD Chemical, San Diego, CA), or 100 µmol/L Janus kinase/signal transducers and activators of transcription (JAK/STAT) inhibitor 5'-deoxy-5'-(methylthio)adenosine (MTA) (EMD Chemical) or BAY + MTA. The cultures were then incubated with control (vehicle) or TNF-α or IFN-γ or TNF-α + IFN-γ in parallel either for 24 hours and then the conditioned medium was collected for ELISAs, or for 30 minutes and then subjected to nuclear extraction for ELISA-based electromobility shift assay (E-EMSA).

## E-EMSA

First trimester decidual cells were treated for 30 minutes as described above, then rinsed in ice-cold PBS containing a protease inhibitor cocktail, and immediately stored at -80°C. A Nuclear Extraction Kit was used according to the manufacturer's instruction (Active Motive, Carlsbad, CA). NFκB transcriptional activity was determined using a TransAM NFκB transcription factor ELISA kit (Active Motive) following the manufacturer's instructions. This kit measures both p65 and p50 NFκB subunits that bind to 96-well plates precoated with NFκB DNA response elements (NFκB-RE). A colorimetric reaction was obtained using a 96-well plate luminometer at 450 nm and an optional reference wavelength of 655 nm.

## Flow Cytometry

Flow cytometry was performed as described previously.<sup>17</sup> Briefly, cell sorting and fluorescence measurements were carried out on a MoFlo high-performance cell sorter (Cytomation, Fort Collins, CO). For fluorescence measurements, the following mouse anti-human monoclonal antibodies were obtained from BD Pharmingen: CD16-fluorescein isothiocyanate (IgG1), CD56-phycoerythrin (IgG1), and CD3-Cyc (IgG1), and from R&D Systems: Phycoerythrin-conjugated mouse monoclonal anti-human *CCR1* (*IgG2b*), *CCR9* (*IgG2a*), *CXCR1* (*IgG2a*), *CXCR3* (*IgG1*), *CXCR4* (*IgG2a*), and *CCR7* (*IgG2a*). The *CCR7* antibody was used together with goat anti-mouse IgG-phycoerythrin as

a secondary fluorochrome conjugate (Caltag, Burlingame, CA). All IgG isotype control antibodies were obtained from BD Biosciences (Los Angeles, CA). For flow cytometric analysis, cells were incubated with 10% human serum to block Fc-receptors before incubation with monoclonal antibodies on ice for 30 minutes, followed by washing twice in PBS with 0.5% bovine serum albumin. Surface expression of chemokine receptors on pNK and dNK cells was detected with a standard FACSCalibur flow cytometer (Immunocytometry Systems; Becton Dickinson, San Jose, CA). In these measurements, data from 10,000 single-cell events were collected and analyzed using CellQuest version 2.0 (Becton Dickinson) or FlowJo version 7.6.5 (TreeStar, San Carlos, CA).

## ELISA

Total cell protein levels were measured by a modified Lowry assay (Bio-Rad Laboratories, Hercules, CA). Commercial ELISA kits measured immunoreactive levels of IP-10, I-TAC, IL-6, IL-8, IL-11, and monocyte chemoattractant protein (MCP)-1 in decidual cell-conditioned medium supernatants according to the manufacturer's instructions (Duoset kits; R&D Systems). The IP-10 and I-TAC ELISAs have sensitivities of 8 and 2 pg/mL, respectively. The intra-assay coefficients of variation (COV) for IP-10 and I-TAC are each <5.0%, and interassay COV are each <10%. The IL-6 ELISA has a sensitivity of 3 pg/mL; intra-assay and interassay COV are 3.1% and 2.7%, respectively. The IL-8 ELISA has a sensitivity of 8 pg/mL, and intra-assay and interassay COV are 4.6% and 6.7%, respectively. The IL-11 ELISA has a sensitivity of 10 pg/mL, and intra-assay and interassay COV are 2.4% and 6.9%, respectively. The MCP-1 ELISA has a sensitivity of 4 pg/mL, and intra-assay and interassay COV are 5.0% and 5.1%, respectively.

## Microarray Analysis

Total RNA from decidual cell cultures was extracted as previously described.<sup>19</sup> For microarray studies, cells harvested with QIAzol lysis reagent (Qiagen, Valencia, CA) were used to prepare total RNA, which was cleaned and precipitated using the RNeasy Mini Kit (Qiagen). The quality of RNA was confirmed by a Bio-Rad bioanalyzer. Array processing was performed at the W. M. Keck Foundation Biotechnology Resource Laboratory at Yale University using HG\_U133 Plus 2.0 chips (Affymetrix, Santa Clara, CA). Raw data without normalization generated from Affymetrix GeneChip Operating Software version 1.2 (GCOS 1.2) (Affymetrix) were analyzed by GeneSpring software version 7.2 (Agilent Technologies/Silicon Genetics, Redwood City, CA). Gene readouts were normalized to the 50th percentile of the distribution of all measurements in each chip. Normalization for each gene across chips was performed using the median value of each gene throughout different chips in the same experimental condition. Normalized data were filtered to eliminate those genes whose mRNA were found to be absent

in all experimental conditions and replicates. A parametric test assuming unequal variance was then used to assess statistical significance. Fold-ratios were derived by comparing normalized data between E2 + MPA versus E2 + MPA + IL-1 $\beta$  or TNF- $\alpha$  groups with a cutoff of  $\geq 2$ -fold. Microarray results were confirmed by quantitative RT-PCR.

## Quantitative RT-PCR

Total RNA from decidual cell cultures derived from eight patients was isolated and purified using a column kit with DNase treatment (Ambion, Austin, TX). Generation of cDNA from purified DNA-free RNA used the SuperScript II Reverse Transcriptase kit (Invitrogen). qPCR was performed using TaqMan gene expression assays for *IP-10*, *I-TAC*, and the reference gene  $\beta$ -*actin* (Applied Biosystems, Foster City, CA). Standard curves were created from serial dilutions of a known sample for each gene assay. All samples were run in duplicate and the average used for each sample. The efficiency for each TaqMan gene assay was >94%. The TaqMan Assay ID#s were: Hs00171042\_m1 (for *IP-10*), Hs00171138\_m1 (for *I-TAC*), and Hs99999903\_m1 (for  $\beta$ -*actin*). The relative standard curve method was used (versus the comparative  $C_T$  method,  $\Delta\Delta C_T$ ). For each standard curve, an independent sample known to contain very high expression of the gene of interest was used. Serial dilutions were made from this independent sample and used in the standard curve to calculate the unknowns according to the manufacturer's instructions.

## Western Blotting

Western blot analysis was performed on conditioned medium supernatants, diluted 1:6 in reducing Laemmli 6 $\times$  sample buffer (Boston Bioproducts, Ashland, MA), then boiled for 5 minutes. Recombinant human IP-10 and I-TAC proteins (R&D Systems) were treated similarly as positive controls. Centrifuged conditioned medium supernatants and positive controls were subjected to SDS-PAGE on 15% Tris-HCl gels (BioRad) with subsequent electroblotting transfer onto 0.20- $\mu$ m nitrocellulose membranes (BioRad). After transfer, membranes were blocked for 3 hours at room temperature in Odyssey blocking buffer (LI-COR Biosciences, Lincoln, NE) and then incubated overnight at 4°C with goat antibodies to IP-10 and I-TAC (R&D Systems). After washing, membranes were incubated in donkey anti-goat IRDye 680 (LI-COR) at room temperature for 1 hour and washed again before visualizing with the Odyssey infrared imaging system (LI-COR).

## siRNA Transfection

Transient transfections used a modified protocol.<sup>20</sup> Briefly, to prepare siRNA/lipid solutions for a 12-well plate scale transfection, 64 pmol (equivalent of 20  $\mu$ L of a 3  $\mu$ mol/L

stock) of each siRNA was diluted in 200  $\mu$ L of OPTI-MEMI (Invitrogen) and incubated at room temperature for 5 minutes. For IFN- $\gamma$ R1 and IFN- $\gamma$ R2 double knockdown, 32 pmol of each siRNA was combined in a final amount of 64 pmol. In a separate tube, 8  $\mu$ L of Lipofectamine 2000 (Invitrogen) was diluted in 200  $\mu$ L of OPTI-MEMI and incubated at room temperature for 5 minutes. The contents of both tubes were mixed by gentle pipetting and then incubated at room temperature for 30 to 50 minutes. The resulting 400- $\mu$ L solution was used to transfect  $2 \times 10^5$  cells (see below). The siRNAs specific for IFN- $\gamma$ R1 (siR1) (ON-TARGETplus SMARTpool L-011057-00-0005) and IFN- $\gamma$ R2 (siR2) (ON-TARGETplus SMARTpool L-012713-00-0005), and control nontarget siRNA (D-001810-10<sup>-05</sup>) were from Thermo Fisher Scientific/Dharmacon Products (Lafayette, CO).

To prepare cell pellets for transfection, first trimester leukocyte-free decidual cells grown in BM were dissociated using 0.25% Trypsin/EDTA (Sigma-Aldrich). Cell pellets were collected by centrifugation at  $700 \times g$  for 2 minutes. Each cell pellet containing about  $2 \times 10^5$  cells was gently resuspended in 400  $\mu$ L of the siRNA/lipid solution prepared as described above, and the cell suspension was incubated at room temperature for 10 minutes (note, incubation for >15 minutes decreases cell viability). At the end of the incubation, 1.6 mL of BM was added and the resulting cell suspension transferred to a well of a 12-well plate, followed by incubation in a CO<sub>2</sub> tissue culture incubator overnight. The medium was replaced on the second day of incubation.

### Immunohistochemistry

First and third trimester deciduae were snap-frozen in liquid nitrogen for 5- $\mu$ m cryosections. Morphology was confirmed after H&E staining. Sections were air-dried and fixed in ice-cold acetone for 10 minutes, then rehydrated with PBS for 5 minutes, subjected to protein block (Dako, Atlanta, GA) for 20 minutes, and incubated in goat anti-human IP-10 (dilution 1:10; R&D Systems), rabbit anti-human I-TAC, or mouse anti-human IFN- $\gamma$ R1/R2 (dilution 1:250; Abcam, Cambridge, MA) antibodies for 1 hour at room temperature. After 3 washes in PBS, sections were incubated with fluorescein isothiocyanate-conjugated rabbit anti-goat (dilution 1:20; Dako), donkey anti-mouse (dilution 1:50; Millipore), or donkey anti-rabbit (dilution 1:50; Millipore) antibodies for 1 hour.

Sections were double stained by washing  $3 \times$  with PBS and incubating in either goat anti-human vimentin antibody (dilution 1:500; Sigma-Aldrich) or mouse anti-human vimentin antibody (dilution 1:400; Millipore), mouse anti-human cytokeratin 7 (CK7) antibody (dilution 1:100; Dako), goat anti-CD16 antibody (dilution 1:100; R&D Systems), or mouse anti-CD56 antibody (dilution 1:100; R&D Systems) for 1 hour at room temperature. After three additional washes in PBS, the sections were incubated with rhodamine-conjugated goat anti-mouse antibody (dilution 1:100; Sigma-Aldrich) or

donkey anti-goat antibody (dilution 1:100; Millipore), or goat fluorescein isothiocyanate-conjugated anti-mouse antibody (dilution 1:100; Sigma-Aldrich) for 1 hour. Staining specificity was confirmed by substituting isotype IgG antibody for the primary antibody. Finally, the sections were washed  $3 \times$  in PBS, stained with DAPI diluted  $1:5 \times 10^5$  in double-distilled H<sub>2</sub>O (Sigma-Aldrich), and mounted in a non-fade mounting medium (Life Sciences International, Basingstoke, UK). Immunofluorescence was assessed at  $\times 400$  with a Zeiss microscope (Carl Zeiss Vision, München-Hallbergmoos, Germany) equipped with a cooled charge-coupled device camera (AxioCam HR; Carl Zeiss Microimaging, Jena, Germany). No cross-reaction was found among these antibodies. Five randomly selected fields from each section (three sections/tissue) from third trimester deciduae were examined. The number of cells per field ( $10^6$  pixel<sup>2</sup>) was counted and calculated as the mean of 15 fields. All of the sections were evaluated by two individuals blinded to the diagnosis.

### Functional pNK Cell Studies

Circulating NK cells were isolated from healthy donors using the human NK Cell Isolation Kit (130-092-657; Miltenyi Biotec, Bergisch Gladbach, Germany) and incubated at  $5 \times 10^5$  cells/mL in BM containing IP-10 or I-TAC or both at a final concentration of 0.1, 1, 10, or 50 ng/mL each as indicated. After 48 hours incubation at 37°C and 5% CO<sub>2</sub>, cells were collected followed by immunofluorescence staining and flow cytometric analysis to assess purity and CXCR3 expression. For the migration assay,  $2 \times 10^5$  pNK cells in 100  $\mu$ L of BM were incubated with either IP-10 or I-TAC or both at a final concentration of 0.1, 1, 10, or 50 ng/mL for 48 hours and then loaded onto Transwell filters (5- $\mu$ m pore size, 24-well cell clusters; Corning, Corning, NY). The filters were then placed in wells containing 600  $\mu$ L of medium supplemented with IP-10 or I-TAC or both. Although the pre-incubation was carried out at various concentrations, the concentration of IP-10 or I-TAC or both in the bottom chamber was 100 ng/mL. To determine nonspecific migration, two wells without chemokine(s) were used as controls. After 150 minutes incubation at 37°C and 5% CO<sub>2</sub>, the upper chambers were removed, and cells in the bottom chamber were collected and counted. The percentage of migrated cells was calculated by subtracting the value of spontaneously migrated cells from the number of migrated cells at a given chemokine concentration, and the result was divided by the total number of cells initially loaded onto the Transwell filter.

### Peripheral Blood IP-10 and I-TAC Concentrations in Patients Developing Preeclampsia Versus Patients with Uncomplicated Pregnancies

Serum samples from consenting pregnant women undergoing first trimester aneuploidy screening were collected under institutional review board approval at Mackay

**Table 1** Characteristics of Pregnant Women with Preeclampsia and of Controls

	Controls (n = 90)	Preeclampsia (n = 30)	P value
Age, years	30.8 ± 2.8	32.1 ± 3.4	0.06
Parity	1.2 ± 0.4	1.3 ± 0.5	0.179
Gestational age of delivery, weeks	38.7 ± 1.1	37.7 ± 2.2	0.001
Birth weight, g	3215.8 ± 284.4	2755.5 ± 658.5	<0.001
Gestational age of sample collection, weeks	12.5 ± 0.5	12.4 ± 0.4	0.402

Memorial Hospital, Taiwan. Among women who eventually gave birth at Mackay Memorial Hospital from April 2010 to February 2011, 30 developed preeclampsia in the third trimester. Preeclampsia was defined according to standard criteria.<sup>21</sup> Gestational age-matched first trimester control (n = 90) blood samples were obtained from women with subsequent uneventful term deliveries. Gestational ages did not differ significantly between preeclamptic patients (12.4 ± 0.4 weeks) and controls (12.5 ± 0.5 weeks) (P = 0.402) (Table 1). Immunoreactive levels of IP-10 and I-TAC in conditioned media were measured by commercial ELISA kits according to the manufacturer's instructions (DuoSet kits; R&D Systems).

#### Tissue ELISA Measurement of IP-10 and I-TAC Concentrations in Preeclamptic Versus Gestational Age-Matched Control Decidua

Frozen deciduae from patients with preeclampsia and gestational age-matched controls were homogenized in radioimmunoprecipitation assay lysis buffer (NP-40 2%, SDS 0.02%, sodium deoxycholate 1%, 1× PBS) with protease inhibitor cocktail (Sigma-Aldrich). Total protein was measured using protein assay dye reagent (PAK500; Strong Biotech, Taipei, Taiwan). Plates coated with 25 µg/mL sample were incubated overnight at 25°C, followed by blocking for 1 hour with PBS containing 2% bovine serum

albumin. Subsequently, anti-human IP-10 or I-TAC (1 µg/mL; R&D Systems) antibody was added and incubated for 1 hour. The plates were incubated with horseradish peroxidase-conjugated secondary antibody (dilution 1:5000; Chemicon/Millipore) for 1 hour, then incubated with NeA-Blue (Clinical, Mansfield, MA) for 1 hour. The reaction was stopped by 2N H<sub>2</sub>SO<sub>4</sub>. Relative chemokine levels were expressed as absorbance.


#### Statistical Analysis

For ELISA and RT-qPCR IP-10 and I-TAC data from decidual cell cultures, nonparametric tests were used because data were not distributed normally. Comparisons of control and treatment groups used the Kruskal-Wallis analysis of variance on ranks test followed by the Student-Newman-Keuls *post hoc* test with a P value <0.05 representing statistical significance. Student's *t*-tests were used for all pNK cell data, with P values <0.01 representing statistical significance. For both serum and tissue ELISA data and fluorescent immunohistochemistry data from preeclamptic versus control patients, the Student's *t*-test was used, with a P value <0.05 representing statistical significance.

## Results

### Flow Cytometric Analysis of NK Cell Chemokine Receptors

Figure 1A displays representative histograms for the expression of six chemokine receptors (CCR1, CCR7, CCR9, CXCR1, CXCR3, and CXCR4) obtained by gating on CD56<sup>bright</sup>CD16<sup>-</sup> dNK cells or on either CD56<sup>bright</sup>CD16<sup>-</sup> or CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. Comparison of mean fluorescence intensities in Figure 1B indicates that compared with the CD56<sup>dim</sup>CD16<sup>+</sup> population of pNK cells, CD56<sup>bright</sup>CD16<sup>-</sup> dNK cells express relatively higher levels of CXCR3, CCR9, CCR1, and CXCR4. Of these chemokine receptors, CXCR3 is also highly expressed on minority circulating CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells. In this subset, CXCR3 is expressed at higher levels than CCR7, and much



**Figure 1** Chemokine receptor expression by decidual and peripheral NK cells. Decidual and peripheral lymphocyte suspensions were stained with fluorescence-conjugated monoclonal antibodies reactive with CD3, CD16, CD56, the indicated chemokine receptors, or with the appropriate isotype control antibodies. Initially, cells were gated by forward scatter/side scatter characteristics. CD56 and CD16 expression were each analyzed after setting a gate on CD3<sup>-</sup> cells. **A:** Representative histograms of chemokine receptor expression in NK cells. Isotype: filled gray curve; anti-chemokine receptor antibody: solid black curve. CD56<sup>bright</sup>CD16<sup>-</sup> dNK cells, minority CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells, and majority CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. **B:** Mean fluorescence intensity (MFI) ± SD of chemokine receptor expression in NK cell from three donors.

**Table 2** Microarray Results for IP-10 and I-TAC mRNA in First Trimester Decidual Cells

First trimester			Fold change	
Chemokine	Cognate receptor	Alternate name	EM vs EMI	EM vs EMTNF
CXCL10	CXCR3	IP-10	239.33	174.86
CXCL11	CXCR3	I-TAC	154.34	58.65

Leukocyte-free first trimester decidual cells were primed in E2 + MPA (Control) and then switched to defined medium with E2 + MPA ± 10 ng/mL IL-1 $\beta$  (EMI) or TNF- $\alpha$  (EMTNF) for 6 hours. mRNA levels were measured by microarray. Results are shown as fold-change versus control.

higher than CXCR4, CCR1, CCR9, and CXCR1. On the majority circulating CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells, only CXCR1, a receptor for IL-8, is highly expressed, whereas CCR7 and CXCR3 are expressed at low-to-intermediate levels, and CCR1 and CXCR4 at low levels (Figure 1B). By indicating that both decidual and peripheral CD56<sup>bright</sup>CD16<sup>-</sup> NK cell subsets express significantly elevated levels of CXCR3, Figure 1 suggests a potential role for CXCR3 in mediating preferential trafficking of circulating CD56<sup>bright</sup>CD16<sup>-</sup> NK cells to the decidua, thereby explaining the unusual abundance of these cells at this site.


#### Effects of TNF- $\alpha$ , IL-1 $\beta$ , and IFN- $\gamma$ on First Trimester Decidual Cell IP-10 and I-TAC Expression

The microarray screening results displayed in Table 2 indicate that in cultured first trimester decidual cells incubated with E2 + MPA, to mimic the hormonal milieu of early pregnancy, addition of either TNF- $\alpha$  or IL-1 $\beta$  significantly up-regulates steady-state levels of mRNAs for the cognate CXCR3 ligands, IP-10, and I-TAC. Figure 2 displays the effects of incubating first trimester leukocyte-free decidual cell monolayers over a physiological range of concentrations up to 10 ng/mL TNF- $\alpha$  or IL-1 $\beta$  or IFN- $\gamma$  on secreted levels of immunoreactive IP-10 (Figure 2A) and I-TAC (Figure 2B). The responses are linear over the entire concentration range for TNF- $\alpha$  and IFN- $\gamma$ , but attain a maximum effect at 1.0 ng/mL for IL-1 $\beta$ .


The individual effects of 1.0 ng/mL each of TNF- $\alpha$  or IL-1 $\beta$  or IFN- $\gamma$  and the combined effects of IFN- $\gamma$  with either TNF- $\alpha$  or IL-1 $\beta$  on mRNA expression are displayed for IP-10 (Figure 3A) and for I-TAC (Figure 3B). For IP-10 mRNA levels (mean ± SEM): IFN- $\gamma$ , TNF- $\alpha$ , and IL-1 $\beta$  produced increases over basal output of 922 ± 430-fold, 581 ± 381-fold, and 508 ± 219-fold, respectively. For I-TAC mRNA levels (mean ± SEM): IFN- $\gamma$ , TNF- $\alpha$ , and IL-1 $\beta$  produced increases over basal output of 897 ± 508-fold, 202 ± 97-fold, and 131 ± 64-fold, respectively. The combination of IFN- $\gamma$  with TNF- $\alpha$  or IL-1 $\beta$  resulted in IP-10 mRNA levels of more than 40,000-fold greater than basal output and about 30-fold greater than the sum of each cytokine used alone. For I-TAC, the combination of IFN- $\gamma$  with TNF- $\alpha$  or IL-1 $\beta$  results in mRNA expression levels

that are about 36,000-fold greater than basal output, and about 40-fold greater when each cytokine was used alone.

Figure 3 also displays the individual effects of 1.0 ng/mL TNF- $\alpha$  or IL-1 $\beta$  or IFN- $\gamma$  and the combined effects of IFN- $\gamma$  with either TNF- $\alpha$  or IL-1 $\beta$  on secreted IP-10 (Figure 3C) and I-TAC (Figure 3D) concentrations. For IP-10 levels (mean ± SEM): IFN- $\gamma$  stimulation induces 3.20 ± 0.84 pg/mL/ $\mu$ g protein, an increase of 468 ± 179-fold over basal output (0.012 ± 0.003 pg/mL/ $\mu$ g protein); TNF- $\alpha$  induces 3.22 ± 1.22 pg/mL/ $\mu$ g protein, an increase of 356 ± 115-fold over basal output; and IL-1 $\beta$  induces 2.26 ± 1.02 pg/mL/ $\mu$ g protein, an increase of 256 ± 104-fold over basal output. For I-TAC (mean ± SEM): IFN- $\gamma$  induces 0.45 ± 0.16 pg/mL/ $\mu$ g protein, an increase of 278 ± 120-fold over basal output (0.002 ± 0.0003 pg/mL/ $\mu$ g protein); TNF- $\alpha$  induces 0.32 ± 0.13 pg/mL/ $\mu$ g protein, an increase of 179 ± 64-fold over basal output; and IL-1 $\beta$  induces 0.10 ± 0.06 pg/mL/ $\mu$ g protein, an increase of 52 ± 29-fold over basal output. Note that basal and induced levels of IP-10 are approximately 5 to 10 times greater than the corresponding levels for I-TAC. Co-incubation of IFN- $\gamma$  with either TNF- $\alpha$  or IL-1 $\beta$  produced striking synergistic



**Figure 2** Concentration-dependent effects of TNF- $\alpha$  or IL-1 $\beta$  or IP-10 and I-TAC protein levels in cultured first trimester decidual cells. Leukocyte-free first trimester decidual cells were primed in E2 + MPA then switched to a defined medium with E2 + MPA ± IL-1 $\beta$  (asterisk) or TNF- $\alpha$  (black circle) or IFN- $\gamma$  (black triangle) at the concentrations indicated on the abscissa for 24 hours. Levels of IP-10 (A) and I-TAC (B) were measured by ELISA in conditioned medium supernatants and normalized to total cell protein (see *Materials and Methods* for details) (average of two separate experiments).



**Figure 3** Separate and interactive effects of TNF-α or IL-1β or IFN-γ on IP-10 and I-TAC protein and mRNA expression in cultured first trimester decidual cells. Leukocyte-free first trimester decidual cells were primed in E2 + MPA (CON). Cultures were then switched to defined medium (DM) with the steroids and either IL-1β or TNF-α or IFN-γ or TNF-α + IFN-γ or IL-1β + IFN-γ for 6 hours for mRNA or 24 hours for protein analysis. The mRNA levels for IP-10 (A) and I-TAC (B) were measured by RT-qPCR on RNA extracts and normalized to β-actin, a reference gene (n = 8, mean ± SEM). Protein levels of IP-10 (C) and I-TAC (D) were measured by ELISA in conditioned medium supernatants and normalized to total cell protein (n = 10, mean ± SEM). See *Materials and Methods* for details. \*P < 0.05 versus CON; †P < 0.05 versus TNF-α alone; ‡P < 0.05 versus IL-1β alone. Western blotting (E) was carried out for IP-10 and I-TAC on 24 hours conditioned medium. Recombinant human IP-10 or I-TAC (2.5 ng per lane) are shown as positive controls (Pos) (see *Materials and Methods* for details).

augmentation of IP-10 and I-TAC outputs. Combinations of IFN-γ with TNF-α or IL-1β produced IP-10 levels of 190.1 ± 32.7 and 158.8 ± 25.3 pg/mL/μg protein, respectively. These represent increases of about 24,000-fold over basal output, and 35-fold greater than the levels induced by each cytokine added individually. In parallel measurements for I-TAC, IFN-γ added with TNF-α or IL-1β produces levels of 38.0 ± 4.9 and 23.5 ± 5.1 pg/mL/μg protein, respectively, which represent a >14,000-fold increase over basal output and an 80-fold increase over each cytokine used alone.


The Western blots depicted in Figure 3E confirm the ELISA results for IP-10 (Figure 3C) and for I-TAC (Figure 3D). Specifically, the antibodies used recognize individual bands in the decidual cell–conditioned medium that migrate with the mobility of IP-10 (8.5 kDa) and I-TAC (8.3 kDa). Compared with the corresponding control incubation, the magnitude of each band is modestly augmented during incubation with TNF-α or IL-1β or IFN-γ, with marked synergistic augmentation evident in co-incubations with IFN-γ plus TNF-α or IL-1β.

**Effects of TNF-α, IL-1β, and IFN-γ on First Trimester Decidual Cell MCP-1, IL-6, IL-8, and IL-11 Expression**

As shown in Figure 4, both TNF-α, and IL-1β significantly increased secreted levels of each of the evaluated

chemokines. Specifically, for MCP-1 compared with basal output of 4.7 ± 1.6 pg/mL/μg cell protein (mean ± SEM), TNF-α elicits a 48.9 ± 13.5-fold increase and IL-1β elicits a 79.6 ± 15.5-fold increase (mean ± SEM), respectively (Figure 4A). Figure 4B indicates basal IL-6 production of 0.06 ± 0.02 pg/mL/μg cell protein, with TNF-α eliciting a 32.5 ± 5.1-fold increase and IL-1β eliciting a 2871 ± 994-fold increase, respectively. Figure 4C shows basal IL-11 output of 0.08 ± 0.02 pg/mL/μg cell protein, with TNF-α eliciting a 7.4 ± 2.1-fold increase and IL-1β eliciting a 618.9 ± 309.6-fold increase, respectively. Figure 4D shows basal IL-8 output of 0.50 ± 0.15 pg/mL/μg cell protein, with TNF-α eliciting a 547.0 ± 116.2-fold increase and IL-1β eliciting a 2850 ± 385-fold increase (mean ± SEM), respectively. Unlike the marked responses to TNF-α and IL-1β, IFN-γ is either ineffective or much less effective at elevating secreted levels of each chemokine. Only for MCP-1 did IFN-γ up-regulation achieve statistical significance (5.9 ± 1.5-fold increase, mean ± SEM, P < 0.05). Moreover, in contrast to the striking synergistic increases in secreted levels of IP-10 and I-TAC displayed in Figure 3, C and D, during co-incubation of IFN-γ with either TNF-α or IL-1β, Figure 4 indicates that co-incubation of IFN-γ with either cytokine produces less than additive effects for MCP-1 (Figure 4A), IL-6 (Figure 4B), and IL-8 (Figure 4D), and a subtractive effect for IL-11 (Figure 4C).






**Figure 4** Separate and interactive effects of TNF- $\alpha$  or IL-1 $\beta$  or IFN- $\gamma$  on monocyte, neutrophil, and NK cell recruiting chemokine expression by first trimester decidual cell monolayers. Leukocyte-free first trimester decidual cells were primed in E2 + MPA [control (CON)]. Cultures were then switched to defined medium with the steroids and either IL-1 $\beta$  or TNF- $\alpha$  or IFN- $\gamma$  or TNF- $\alpha$  + IFN- $\gamma$  or IL-1 $\beta$  + IFN- $\gamma$  for 24 hours. Levels of MCP-1 (A), IL-6 (B), IL-11 (C), and IL-8 (D) measured by ELISA in conditioned medium supernatants and normalized to total cell protein (see *Materials and Methods* for details) ( $n = 8$ , mean  $\pm$  SEM). \* $P < 0.05$  versus CON; † $P < 0.05$  versus IFN- $\gamma$  alone; ‡ $P < 0.05$  versus TNF- $\alpha$  alone; and § $P < 0.05$  versus IL-1 $\beta$  alone.


#### Effects of Knockdown of *IFN- $\gamma$ R1* and *-R2* mRNA on *IP-10* and *I-TAC* mRNA Expression

As shown in **Figure 5A**, 92% knockdown of *IFN- $\gamma$ R1* mRNA was obtained 48 hours after siR1 transfection versus control siRNA. Importantly, the siR1 does not affect the nontargeted  $\beta$ -actin mRNA, confirming the specificity of the *IFN- $\gamma$ R1* knockdown. Similarly, transfection of siR2 leads to a 95% down-regulation of *IFN- $\gamma$ R2* (**Figure 5B**). Furthermore, the combination of siR1 and siR2 (double knockdown) elicits an  $\sim 90\%$  knockdown efficiency with each gene (**Figure 5C**). Consistent with the RT-qPCR results shown in **Figure 3**, A and B, results obtained with control

siRNA indicate that IFN- $\gamma$  induces higher mRNA levels for *IP-10* (**Figure 5D**) and *I-TAC* (**Figure 5E**) than TNF- $\alpha$  added alone and that co-incubation with TNF- $\alpha$  plus IFN- $\gamma$  synergistically enhances both *IP-10* and *I-TAC*. Individual siRNA-mediated knockdown of *IFN- $\gamma$ R1* and *-R2* neutralizes augmentation of *IP-10* (**Figure 5D**) and *I-TAC* (**Figure 5E**) elicited by IFN- $\gamma$ , but only marginally inhibits TNF- $\alpha$ -induced *IP-10* and *I-TAC* mRNA levels. In co-incubation with TNF- $\alpha$  plus IFN- $\gamma$ , siRNA-mediated knockdown of *IFN- $\gamma$ R1* is more effective than *IFN- $\gamma$ R2* in blocking synergistic up-regulation of *IP-10* and *I-TAC*. Simultaneous silencing of both receptors eliminates the synergistic up-regulation of both *IP-10* and *I-TAC* (**Figure 5**, D and E).



**Figure 5** Effects of siRNA-mediated knockdown of IFN- $\gamma$ R1 and IP-10, I-TAC, IFN- $\gamma$ R1, and IFN- $\gamma$ R2 mRNA expression. **A–C**: Gene knockdown of human *IFN- $\gamma$ R1*, or *IFN- $\gamma$ R2*, or both. Levels of  $\beta$ -actin are shown as a nontargeted negative control. **A**: Single *IFN- $\gamma$ R1* gene knockdown. Cells were transfected with control siRNA (siCon) or siR1, and the levels of *IFN- $\gamma$ R1* mRNAs are shown on the ordinate. **B**: Single *IFN- $\gamma$ R2* gene knockdown. Cells were transfected with siCon or siR2, and the levels of *IFN- $\gamma$ R2* mRNAs are shown on the ordinate. **C**: Double knockdown of *IFN- $\gamma$ R1* and *IFN- $\gamma$ R2*. Cells were transfected with siCon, or a combination of siR1 and siR2, and the levels of *IFN- $\gamma$ R1* and *IFN- $\gamma$ R2* mRNAs are shown. The levels of the indicated mRNAs from siCon-transfected cells were arbitrarily set as 1. Error bars indicate mean  $\pm$  SD ( $n = 3$ ). **D** and **E**: Human leukocyte-free first trimester decidual cells were transfected with siRNAs specific for siR1, siR2, or siCon for 24 hours and then incubated with either 1 ng/mL TNF- $\alpha$  or IFN- $\gamma$  or both for 6 hours ( $n = 1$ ). Levels of mRNA were measured using real-time and RT-qPCR for *IP-10* (D) and *I-TAC* (E).



**Figure 6** Immunostaining for IP-10, I-TAC, IFN- $\gamma$ R1, and IFN- $\gamma$ R2 in first trimester decidua. Serial cryosections of first trimester decidua were immunostained with IP-10 (A and B) or I-TAC (D and E) (green fluorescence) and then with either the decidual cell marker vimentin (A and D) or the EVT marker cytokeratin 7 (CK7) (B and E) (red fluorescence). Sections were also immunostained for vimentin (red) and either IFN- $\gamma$ R1 (C) or IFN- $\gamma$ R2 (F) (green). DAPI (blue fluorescence) denotes nuclear immunostaining. Overlap between red and green immunostaining is seen as yellow or orange. **Arrows** indicate colocalization of IP-10 in vimentin-positive decidual cells (A) and indicate CK7-positive EVT cells expressing IP-10 (B). **Arrowheads** indicate decidual cells expressing I-TAC (D) and EVTs expressing I-TAC (E). Original magnification of representative sections,  $\times 400$ .

### Immunostaining of IP-10, I-TAC, IFN- $\gamma$ R1, and -R2 in First Trimester Human Decidua

Cryosections of first trimester deciduae were sequentially immunostained for either IP-10 (Figure 6, A and B) or I-TAC (Figure 6, D and E) and then with either vimentin as a decidual cell marker (Figure 6, A and D) or CK7 as an EVT marker (Figure 6, B and E). The sections were also stained for DAPI to identify nuclei. Overlap between red and green immunostaining appears as yellow-to-orange staining. For IP-10, inspection of Figure 6A indicates that the overwhelming majority of cells demonstrate colocalization of IP-10 in vimentin-positive decidual cells, evident by the resulting yellow-orange staining. The pure green staining in the majority of the cells in Figure 6B indicates that cells expressing IP-10 are mainly non-EVT. Figure 6B indicates that only a few CK7-positive EVT cells express IP-10. Inspection of Figure 6D indicates that virtually all of the decidual cells also express I-TAC, albeit at lower intensity than IP-10. As with IP-10, very few EVTs express I-TAC as seen in Figure 6E. These *in situ* findings are consistent with the higher levels of basal IP-10 than I-TAC mRNA (Figure 3, A and B) and secreted protein (Figure 3, C and D) measured in cultured decidual cells. Cryosections were also immunostained for IFN- $\gamma$ R1 or -R2 and then vimentin. Positive staining for IFN- $\gamma$ R1 (Figure 6C) and -R2 (Figure 6F) was primarily localized to decidual cells, with blue staining denoting nuclei.


### Signaling Pathway Mediation of Synergistic Enhancement of IP-10 and I-TAC Expression

Figure 7 demonstrates that TNF- $\alpha$  + IFN- $\gamma$  significantly enhances IP-10 (Figure 7A) and I-TAC (Figure 7B) expression compared with control or TNF- $\alpha$  alone or IFN- $\gamma$  alone. Pretreatment of cultured decidual cells with either BAY (NF $\kappa$ B inhibitor) or MTA (JAK/STAT inhibitor) alone or in combination blocks the synergistic effects of TNF- $\alpha$  + IFN- $\gamma$  on IP-10 or I-TAC expression.

The E-EMSA analysis displayed in Figure 7C shows that nuclear extracts obtained from decidual cells treated with TNF- $\alpha$  alone or IFN- $\gamma$  alone exhibit higher binding of NF $\kappa$ B subunits p65 and p50 to NF $\kappa$ B-RE compared with control. This binding to NF $\kappa$ B-RE increased further in nuclear extracts from decidual cells treated with TNF- $\alpha$  + IFN- $\gamma$ . Moreover, pretreatment of decidual cells with either BAY or MTA alone or with BAY + MTA for 1 hour significantly reduced NF $\kappa$ B-RE binding of p65 and p50 in nuclear extracts compared with decidual cells treated with TNF- $\alpha$  + IFN- $\gamma$  (Figure 7C).

### The Effects of IP-10 and I-TAC on CXCR3 Expression by and Migration of pNK Cells

Figure 8 demonstrates the effects of incubating CD56<sup>bright</sup> CD16<sup>-</sup> or CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells with IP-10 or I-TAC on CXCR3 expression. Incubation of both NK cell subsets



**Figure 7** Signaling pathway mediation of IP-10 and I-TAC expression. **A:** IP-10 and **B:** I-TAC secretion by decidual cells treated with vehicle, TNF- $\alpha$ , IFN- $\gamma$ , and TNF- $\alpha$  + IFN- $\gamma$  with or without NF $\kappa$ B inhibitor BAY, or JAT/STAT inhibitor (MTA) or BAY + MTA ( $n = 3$ , mean  $\pm$  SEM). \* $P < 0.05$ , \*\* $P < 0.01$ , and \*\*\* $P < 0.005$ . **(C)** Representative binding of decidual cell–derived nuclear NF $\kappa$ B subunits p65 and p50 to NF $\kappa$ B-RE by E-EMSA ( $n = 2$ ).

with either IP-10 (Figure 8A) or I-TAC (Figure 8B) elicits a concentration-dependent increase in CXCR3 that peaks at 10 ng/mL. However, either 50 ng/mL IP-10 or I-TAC down-regulates CXCR3 expression. Figure 8C demonstrates that the combination of equal concentrations of IP-10 and I-TAC elicits a similar concentration-dependent biphasic effect on CXCR3 expression. Parallel concentration-dependent biphasic effects are also observed on the migration of both pNK cell subtypes (Figure 9). Thus, either IP-10 (Figure 9A) or I-TAC (Figure 9B) or the combination of IP-10 and I-TAC (Figure 9C) progressively enhances cell migration at concentrations of 0.1 to 10 ng/mL, whereas 50 ng/mL impedes migration. Importantly, the biphasic effects observed are donor-independent.


### CD56<sup>bright</sup>CD16<sup>-</sup> NK Cells in Preeclamptic Decidua

Decidual sections from women with preeclampsia versus gestational age–matched control specimens were immunostained sequentially for CD56 (green fluorescence) and CD16 (red fluorescence). The absence of CD16 and the


presence of strong staining for CD56 in control and preeclamptic specimens indicate that virtually all NK cells in decidua are CD56<sup>bright</sup>CD16<sup>-</sup>. The more abundant green staining in Figure 10A compared with Figure 10B suggests that preeclamptic decidua contains fewer CD56<sup>bright</sup>CD16<sup>-</sup> NK cells. As indicated in Figure 10C, lower NK cell numbers in preeclamptic versus gestational age–matched control decidua attained statistical significance.

### Measurement of IP-10 and I-TAC Levels in First Trimester Sera and Term Deciduae of Patients With and Without Subsequent Preeclampsia

In Figure 11A, a statistically significant increase in IP-10 levels in first trimester serum predicts later development of preeclampsia. By contrast, no such correlation was observed for the serum levels of I-TAC (Figure 11B). Tissue ELISA analyses for decidual extracts from patients with preeclampsia find significantly higher levels of IP-10 (Figure 11C) and I-TAC (Figure 11D) than age-matched controls.



**Figure 8** Effects of IP-10 and I-TAC on CXCR3 expression by pNK cells. The lymphocyte population from healthy human donor blood was enriched by Ficoll density gradient centrifugation, followed by purification of pNK cells using NK cell isolation kits (see *Materials and Methods*). Enriched (91% to 95%) pNK cells were incubated with 0, 0.1, 1, 10, or 50 ng/mL IP-10 (**A**) or I-TAC (**B**) or IP-10 plus I-TAC (**C**) for 48 hours. After incubation, CXCR3 expression was analyzed on gated CD56<sup>+</sup>CD3<sup>-</sup> NK cells. Data are presented as means  $\pm$  SEM ( $n = 7$ ), with CXCR3 expression levels of cells incubated with 0 ng/mL IP-10 and I-TAC arbitrarily set as 1. The bar groups represent CD56<sup>dim</sup>CD16<sup>+</sup> and CD56<sup>bright</sup>CD16<sup>-</sup> subsets, respectively. \*\* $P < 0.01$  0 ng/mL versus 0.1, 1, 10, or 50 ng/mL IP-10 and I-TAC, by Student's *t*-test.



**Figure 9** Effects of IP-10 and I-TAC on migration of pNK cells. pNK cells isolated from healthy donors were pre-incubated with IP-10 (A), I-TAC (B), or both (C) at a final concentration of 0, 0.1, 1, 10, or 50 ng/mL, each for 48 hours. Migration assays were carried out in the presence of 100 ng/mL IP-10 or I-TAC or both as chemoattractants. Results are presented as the mean  $\pm$  SEM ( $n = 3$ ) of the percentage of migrated cells.  $**P < 0.01$  0 ng/mL versus 0.1, 1, 10, or 50 ng/mL IP-10 and I-TAC, by Student's *t*-test.


## Discussion

In NK cell-deficient mice, the decidual vasculature lacks characteristic mid-gestation dilation, elongation, and loss of smooth muscle associated with increased blood flow, similar to lesions seen in human preeclampsia.<sup>22</sup> These effects are ameliorated by treatment with mouse IFN- $\gamma$  or with the IFN- $\gamma$ -regulated cytokine-binding protein,  $\alpha$ -2 macroglobulin, which enhances decidual remodeling and produces normal histological changes in decidual blood vessels.<sup>23</sup> Ninety percent of mouse uterine IFN- $\gamma$  is dNK cell-derived, indicating that IFN- $\gamma$  is necessary and sufficient to mediate normal uterine vascular remodeling in pregnant mice.<sup>24</sup> Close proximity between dNK cells and blood vessels suggests that dNK cell-derived angiogenic factors may augment the effects of IFN- $\gamma$  to promote normal vascular remodeling.<sup>25</sup> Extrapolation of studies in NK cell-depleted mice to humans<sup>26,27</sup> suggests that the minority


CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells traffic from the circulation to the decidua in early pregnancy.

Transformation of spiral arteries and arterioles into high-capacity vessels with markedly increased uteroplacental blood flow to the developing fetal-placental unit has been attributed to replacement of vascular smooth muscle and endothelial cells by fibrinoid-embedded endovascular EVT.<sup>28</sup> Spiral arterial lining endovascular EVTs express stromal cell-derived factor-1 (SDF-1) (alias CXCL12), which binds to CCR4-expressing CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells to mediate their recruitment to the decidua.<sup>28</sup> In these specimens, infiltrating dNK cells and macrophages were identified in the vascular smooth muscle layers accompanied by the appearance of apoptotic markers in vascular smooth muscle and endothelial cells. This induction of apoptosis was attributed to dNK cell- and macrophage-expressed matrix metalloproteinase (MMP)-7 and MMP-9.<sup>29</sup> Such trophoblast-independent decidualization-associated vascular remodeling is also indicated by early signs of vascular transformation in decidua from ectopic tubal pregnancies or in late luteal phase nonpregnant endometrium.<sup>30,31</sup> In the early pregnant golden hamster, dNK cells invade the vascular wall before trophoblast invasion and are implicated in directly mediating endometrial vascular remodeling.<sup>32</sup>

The origin of dNK cells in humans is alternatively ascribed to: i) differentiation of resident endometrial NK cells<sup>7,11</sup>; ii) self-renewal from local progenitor stem cells<sup>33</sup>; and iii) trafficking of pNK cells.<sup>34</sup> In support of the last, the majority circulating CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells were shown to be recruited to the decidua, then differentiate into CD56<sup>bright</sup>CD16<sup>-</sup> NK cells by decidual cell-derived transforming growth factor- $\beta$ .<sup>35,36</sup> However, abundant evidence points to the minority circulating CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells as the major contributors to the dNK cell population. Their preferential recruitment is suggested by elevated expression of L-selectin, which mediates interactions with vascular endothelium<sup>37</sup> by CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells, but not by CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. Moreover, the current study confirms that CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells express high levels of CXCR3<sup>14</sup> (Figure 1, A and B), a receptor for decidual cell-derived IP-10 and I-TAC that can



**Figure 10** CD56<sup>bright</sup>CD16<sup>-</sup> NK cells in preeclamptic (PE) versus control (CON) decidua. Decidua from patients with preeclampsia (A) and gestational age-matched controls (B) were stained with anti-human CD56 (green) and CD16 (red) antibodies. Arrows indicate CD56<sup>bright</sup>CD16<sup>-</sup> cells. Representative slides from eight independent experiments are shown. Original magnification,  $\times 400$ . C: Comparison of CD56<sup>bright</sup>CD16<sup>-</sup> cells between preeclamptic and control decidua. Results are presented as mean  $\pm$  SEM ( $n = 8$ ) of the CD56<sup>bright</sup>CD16<sup>-</sup> cells.  $***P < 0.001$ .



**Figure 11** Comparison of immunoreactive IP-10 and I-TAC expression in serum from first trimester of controls (CON) versus patients subsequently developing preeclampsia (PE) and decidual tissues from preeclampsia versus gestational age-match third trimester decidua. IP-10 (A) and I-TAC (B) levels in serum obtained from first trimester gestation measured by ELISAs. Results are presented as mean  $\pm$  SEM ( $n = 30$  for preeclampsia,  $n = 90$  for control) of IP-10 and I-TAC serum levels. \* $P < 0.05$ . IP-10 (C) and I-TAC (D) levels in decidua obtained from patients with and without preeclampsia measured by ELISAs. Results are presented as means  $\pm$  SEM ( $n = 7$ ). \* $P < 0.05$ .

preferentially promote trafficking to the decidua. Recently, IP-10 secreted by various human endometrial, but not NK cells, was shown to chemoattract blastocyst-derived CXCR3-expressing trophoblast.<sup>38</sup>

Human maternal–fetal interactions create a mild systemic inflammatory state initiated in the luteal phase of the menstrual cycle characterized by vascular endothelial activation, leukocytosis, increased monocyte activity, and elevated circulating chemokine and cytokine levels that become most apparent in the third trimester of uncomplicated human pregnancies.<sup>39</sup> Decidual cell–mediated recruitment of NK cells in normal pregnancy is indicated by a report that progesterone enhances mRNA expression of NK cell–recruiting chemokines, ie, fractalkine (alias CX3CL1) and MCP-1 (alias CCL2), as well as IP-10 in primary first trimester human decidual cells, and NK cell migration is enhanced by using these decidual cells cultured with progesterone as a substrate.<sup>40</sup> Extending these observations to better mimic the decidual milieu of pregnancy, the current study used E2 + MPA to maintain decidualization of leukocyte-free first trimester decidual cell monolayers in combination with proinflammatory cytokines IFN- $\gamma$ , a primary NK cell product, and/or TNF- $\alpha$  or IL-1 $\beta$ , primary macrophage products. The resulting marked synergistic increase in *IP-10* and *I-TAC* mRNA and protein expression suggests that decidual cell–mediated crosstalk between NK cells and macrophages leads to preferential recruitment and retention of additional CD<sup>bright</sup>CD16<sup>-</sup> NK cells to the decidua.<sup>28</sup>

Previously, expression profiling of several genes revealed striking differences between the minority peripheral versus decidual CD56<sup>bright</sup>CD16<sup>-</sup> NK cell populations, suggesting that dNK cells represent a unique NK cell subset.<sup>17</sup> Evidence that the decidual milieu induces differentiation of recruited CD56<sup>bright</sup>CD16<sup>-</sup> NK cells is provided by observations that incubation of peripheral CD56<sup>bright</sup>CD16<sup>-</sup> NK cells with decidual cell–derived cytokines, particularly IL-15, induces a chemokine receptor repertoire similar to that expressed by dNK cells.<sup>28</sup> Progesterins and NK cell–derived IFN- $\gamma$  enhance decidual cell IL-15 production.<sup>41</sup> These observations complement a report that dNK cells express the IL-2 receptor,<sup>42</sup> which mediates IL-15 activation of NK cells.<sup>43</sup> Like the majority CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells, dNK cells express KIRs and display granules containing cytolytic molecules, such as perforin and granzymes A and B.<sup>17</sup> However, dNK cells exhibit markedly reduced cytotoxicity compared with the majority CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells and are even less cytotoxic than the relatively nontoxic minority CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells.<sup>11,44</sup> Although dNK cells form conjugates and activating immune synapses with target cells, their failure to polarize their microtubule organizing centers and transport perforin-containing granules to the synapse impedes cytotoxicity.<sup>44</sup>

Paradoxically, dNK cells express multiple pregnancy-promoting factors. Like the minority circulating CD56<sup>bright</sup>CD16<sup>-</sup> pNK cells, dNK cells produce growth factors, such as GM-CSF, leukemia inhibitory factor, TNF- $\alpha$ , and IFN- $\gamma$ .<sup>7,17,45–47</sup> In addition, dNK cells express EVT invasion-enhancing IL-8 and IP-10, as well as angiogenic factors, such as vascular endothelial growth factor, placental-derived growth factor, angiopoietin 2, and NKG5.<sup>7,45,46</sup> Taken together, these observations indicate that human dNK cells differ phenotypically from both pNK cells and NK cells in cycling endometrium to form a unique NK cell subtype that supports normal pregnancy by promoting trophoblast invasion, vascular remodeling, and in view of their lack of cytotoxicity, immune tolerance of the fetal semiallograft.


The current study employs a multitiered approach to reveal that targeting of first trimester human decidual cells<sup>48</sup> by paracrine effectors released by the predominant decidual immune cell types, dNK cells and macrophages, recruits additional NK cells from the circulation. Initially, flow cytometric analysis demonstrated high expression of CXCR3 on peripheral and decidual CD56<sup>bright</sup>CD16<sup>-</sup> NK cells with lower CXCR3 levels on the majority circulating CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells. Subsequently, incubation of leukocyte-free first trimester decidual cells with either of the primary macrophage-derived cytokines, TNF- $\alpha$  or IL-1 $\beta$ ,<sup>15</sup> or primary dNK cell–derived cytokine, IFN- $\gamma$ ,<sup>12</sup> markedly elevated IP-10 and I-TAC mRNA and protein levels. Co-incubation of IFN- $\gamma$  with either TNF- $\alpha$  or IL-1 $\beta$  produced striking synergistic increases in IP-10 and I-TAC mRNA and protein levels far exceeding the sum of the individual responses. In parallel incubations, expression of other monocyte, neutrophil, and NK cell chemoattractants and/or

activators, ie, MCP-1, IL-6, -8, and -11 (Figure 4) were at best additive, thereby emphasizing specificity of the synergistic up-regulation of IP-10 and I-TAC expression in human decidual cells. Consistent with *in situ* hybridization observations for IP-10 and I-TAC,<sup>49</sup> we observed preferential immunolocalization of IP-10 and I-TAC to vimentin-positive decidual cells versus CK7-positive trophoblasts in first trimester decidual sections.

The individual and interactive contribution of IFN- $\gamma$  to the marked synergistic enhancement of IP-10 and I-TAC expression by first trimester decidual cells revealed by the current study provides strong evidence that IFN- $\gamma$  is involved in CD56<sup>bright</sup>CD16<sup>-</sup> cell recruitment to the decidua. Two receptors, IFN- $\gamma$ R1 and IFN- $\gamma$ R2, mediate IFN- $\gamma$  effects in target cells. Binding of cell membrane expressed IFN- $\gamma$ R1 initiates IFN- $\gamma$ R1/R2 heterodimerization, which mediates intracellular signaling.<sup>16</sup> Use of specific siRNAs to silence each receptor separately and together neutralized the contribution of IFN- $\gamma$ , but did not affect that of TNF- $\alpha$  to enhanced IP-10 and I-TAC mRNA expression. *In situ* observations complementing *in vitro* results provide the first indication that at the implantation site, immunoreactive IFN- $\gamma$ R1 and -R2 are preferentially localized to vimentin-positive decidual cells (Figure 6).

Elucidation of the potential mechanisms underlying the striking synergistic up-regulation of IP-10 and I-TAC expression induced by TNF- $\alpha$  and IFN- $\gamma$  focused on NF $\kappa$ B and JAK/STAT signaling pathways, which have been shown to mediate cellular effects of TNF- $\alpha$  and IFN- $\gamma$ , respectively.<sup>50,51</sup> Pre-incubation of first trimester decidual cells with either BAY or MTA was sufficient to block the synergistic increase of TNF- $\alpha$  + IFN- $\gamma$ -induced IP-10 and I-TAC expression measured by ELISAs or E-EMSA (Figure 7). Signaling pathway mediation of these effects is depicted by the scheme in Figure 12. Specifically, it indicates that binding of TNF- $\alpha$  to its receptor results in dissociation of NF $\kappa$ B p65 and p50 subunits from NF $\kappa$ B inhibitory molecule I $\kappa$ B, which is degraded by proteasomes. The resulting p65 and p50 homo- and/or heterodimerization is followed by translocation to the nucleus. The p65/p50 heterodimer then binds to NF $\kappa$ B-RE in the IP-10 and I-TAC promoters. IFN- $\gamma$  activates the IFN- $\gamma$  receptor and its downstream signaling JAK/STAT pathway. Activation of JAK/STAT results in nuclear translocation of STAT where it binds to its response element in the IP-10 and I-TAC promoters. Because the JAK/STAT inhibitor MTA also suppresses NF $\kappa$ B binding to NF $\kappa$ B-RE, IFN- $\gamma$  can also activate further NF $\kappa$ B binding via the JAK/STAT pathway, thereby contributing to synergistic up-regulation.

The current observations on primary human first trimester decidual cells and immunostaining of decidual sections are complemented by parallel assessment of CXCR3 expression and *in vitro* migration of CD56<sup>bright</sup>CD16<sup>-</sup> and CD56<sup>dim</sup>CD16<sup>+</sup> pNK cells during incubation with IP-10 or I-TAC or both (Figures 8 and 9). Taken together, our findings have translational implications for normal recruitment of both



**Figure 12** Schematic representation of the contribution of JAK/STAT and NF $\kappa$ B signaling pathways to TNF- $\alpha$  + IP-10 and I-TAC secretion in first trimester decidual cells.

pNK cell subsets to the decidua and aberrant pNK cell trafficking during preeclampsia. Specifically, incubation of both pNK cell subsets with IP-10 and/or I-TAC elicits concentration-dependent enhancement of CXCR3 expression and NK cell migration that peaked at 10 ng/mL, with 50 ng/mL inhibiting both endpoints. These observations suggest that secreted IP-10 and I-TAC levels resulting from co-stimulation of decidual cells by IFN- $\gamma$  with either TNF- $\alpha$  or IL-1 $\beta$  cause the recruitment of additional NK cells to the early decidua, but that overstimulation by these cytokines, as would be observed in the inflammatory milieu of preeclampsia, inhibits CXCR3 expression by, and *in vitro* migration of, both pNK cell subsets. This would account for the reduced number of NK cells observed in preeclamptic decidua (Figure 10) and why elevated IP-10 levels in first trimester serum predict the subsequent development of preeclampsia. Indeed, although elevated circulating levels of angiogenesis mediators, endoglin and soluble fms-like tyrosine kinase receptor-1 (sFlt-1), in second or third trimester serum<sup>52</sup> also predict preeclampsia, IP-10 appears to be an earlier indicator of risk. The importance of our observations is highlighted by a recent review article<sup>53</sup> that states “Currently, no reliable markers exist to identify patients before the development of symptoms. The early identification of women with risk of developing preeclampsia is critical, because careful monitoring and referral to a specialized perinatal care center could substantially improve outcomes for both the mother and fetus.”<sup>53,p957</sup>

The causes of preeclampsia-related decidual inflammation are yet to be fully discerned. Placental oxidative and endoplasmic reticulum stress mediated by trophoblast-derived growth factors, activin-A, corticotrophin-releasing hormone, leptin, and placental microvesicles may play a role.<sup>54,55</sup> Our laboratory and others have observed excess dendritic cells and macrophages in the preeclamptic decidua.<sup>56</sup> Moreover, we found that incubation of first trimester decidual cells with either IL-1 $\beta$  or TNF- $\alpha$  markedly enhances expression of

several monocyte/macrophage-recruiting and -activating cytokines<sup>19</sup>, and that activated macrophages treated with conditioned medium obtained from first trimester decidual cells after incubation with IL-1 $\beta$  or TNF- $\alpha$  induce EVT apoptosis and inhibit EVT invasiveness.<sup>51,57</sup>

Other studies evaluating the relationship between preeclampsia and dNK cell numbers and phenotype have produced conflicting results.<sup>58</sup> However, as noted by Lash and Bulmer,<sup>58</sup> the lack of clarity among various studies as to whether dNK cell numbers are reduced in preeclampsia and intrauterine growth restriction likely reflects differences in sampling, analysis, and severity of the disease. Beyond altered dNK cell numbers, preeclampsia, as well as other adverse pregnancy outcomes, such as fetal growth restriction and recurrent miscarriage, are associated with mismatched KIR haplotype (KIR AA) expression by dNK cells and EVTs.<sup>59</sup>

In summary, we postulate that enhanced periconceptual decidual inflammation contributes to the genesis of shallow placentation in preeclampsia by promoting excess decidual cell IP-10 and I-TAC production, which blunts pNK cell recruitment and leads to deficient dNK cell numbers. Coupled with excess decidual macrophages, these changes impede EVT-mediated remodeling of spiral arteries, the pathological *sine qua non* of preeclampsia.

## Acknowledgments

We thank Drs. Jack L. Strominger for the initial, and crucial, flow cytometry observations made in his laboratory and Mizanur Rahman for his expert cell culture work.

## References

- Zhou Y, Fisher SJ, Janatpour M, Genbacev O, Dejana E, Wheelock M, Damsky CH: Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for successful endovascular invasion? *J Clin Invest* 1997, 99:2139–2151
- Pijnenborg R, Vercruyssen L, Brosens I: Deep placentation. *Best Pract Res Clin Obstet Gynaecol* 2011, 25:273–285
- Young BC, Levine RJ, Karumanchi SA: Pathogenesis of preeclampsia. *Annu Rev Pathol* 2010, 5:173–192
- Steegers EA, von Dadelszen P, Duvetkot JJ, Pijnenborg R: Preeclampsia. *Lancet* 2010, 376:631–644
- Silasi M, Cohen B, Karumanchi SA, Rana S: Abnormal placentation, angiogenic factors, and the pathogenesis of preeclampsia. *Obstet Gynecol Clin North Am* 2010, 37:239–253
- Trundle A, Moffett A: Human uterine leukocytes and pregnancy. *Tissue Antigens* 2004, 63:1–12
- Manaster I, Mandelboim O: The unique properties of uterine NK cells. *Am J Reprod Immunol* 2010, 63:434–444
- Bryceson YT, Chiang SC, Darmanin S, Fauriat C, Schlums H, Theorell J, Wood SM: Molecular mechanisms of natural killer cell activation. *J Innate Immun* 2011, 3:216–226
- Caligiuri MA: Human natural killer cells. *Blood* 2008, 112:461–469
- Trotta R, Chen L, Ciarlariello D, Josyula S, Mao C, Costinean S, Yu L, Butchar JP, Tridandapani S, Croce CM, Caligiuri MA: miR-155 regulates IFN- $\gamma$  production in natural killer cells. *Blood* 2012, 119:3478–3485
- Kalkunte S, Chichester CO, Gotsch F, Sentman CL, Romero R, Sharma S: Evolution of non-cytotoxic uterine natural killer cells. *Am J Reprod Immunol* 2008, 59:425–432
- Lash GE, Robson SC, Bulmer JN: Review: Functional role of uterine natural killer (uNK) cells in human early pregnancy decidua. *Placenta* 2010, 31(Suppl):S87–S92
- Kopcow HD, Karumanchi SA: Angiogenic factors and natural killer (NK) cells in the pathogenesis of preeclampsia. *J Reprod Immunol* 2007, 76:23–29
- Cooper MA, Fehniger TA, Caligiuri MA: The biology of human natural killer-cell subsets. *Trends Immunol* 2001, 22:633–640
- Singh U, Nicholson G, Urban BC, Sargent IL, Kishore U, Bernal AL: Immunological properties of human decidual macrophages: a possible role in intrauterine immunity. *Reproduction* 2005, 129:631–637
- Trinchieri G: Type I interferon: friend or foe? *J Exp Med* 2010, 207:2053–2063
- Koopman LA, Kopcow HD, Rybalov B, Boyson JE, Orange JS, Schatz F, Masch R, Lockwood CJ, Schachter AD, Park PJ, Strominger JL: Human decidual natural killer cells are a unique NK cell subset with immunomodulatory potential. *J Exp Med* 2003, 198:1201–1212
- Arici A, Marshburn PB, MacDonald PC, Dombrowski RA: Progesterone metabolism in human endometrial stromal and gland cells in culture. *Steroids* 1999, 64:530–534
- Huang SJ, Schatz F, Masch R, Rahman M, Buchwalder L, Niven-Fairchild T, Tang C, Abrahams VM, Krikun G, Lockwood CJ: Regulation of chemokine production in response to pro-inflammatory cytokines in first trimester decidual cells. *J Reprod Immunol* 2006, 72:60–73
- Zhang M, Guller S, Huang Y: Method to enhance transfection efficiency of cell lines and placental fibroblasts. *Placenta* 2007, 28:779–782
- ACOG Committee on Practice Bulletins—Obstetrics: ACOG practice bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. *Obstet Gynecol* 2002, 99:159–167
- Greenwood JD, Minhas K, di Santo JP, Makita M, Kiso Y, Croy BA: Ultrastructural studies of implantation sites from mice deficient in uterine natural killer cells. *Placenta* 2000, 21:693–702
- He H, McCartney DJ, Wei Q, Esadeg S, Zhang J, Foster RA, Hayes MA, Tayade C, Van Leuven F, Croy BA: Characterization of a murine alpha 2 macroglobulin gene expressed in reproductive and cardiovascular tissue. *Biol Reprod* 2005, 72:266–275
- Ashkar AA, Croy BA: Functions of uterine natural killer cells are mediated by interferon gamma production during murine pregnancy. *Semin Immunol* 2001, 13:235–241
- Chantakru S, Wang WC, van den Heuvel M, Bashar S, Simpson A, Chen Q, Croy BA, Evans SS: Coordinate regulation of lymphocyte-endothelial interactions by pregnancy-associated hormones. *J Immunol* 2003, 171:4011–4019
- Bilinski MJ, Thorne JG, Oh MJ, Leonard S, Murrant C, Tayade C, Croy BA: Uterine NK cells in murine pregnancy. *Reprod Biomed Online* 2008, 16:218–226
- Guimond MJ, Wang B, Croy BA: Engraftment of bone marrow from severe combined immunodeficient (SCID) mice reverses the reproductive deficits in natural killer cell-deficient tg epsilon 26 mice. *J Exp Med* 1998, 187:217–223
- Hanna J, Wald O, Goldman-Wohl D, Prus D, Markel G, Gazit R, Katz G, Haimov-Kochman R, Fujii N, Yagel S, Peled A, Mandelboim O: CXCL12 expression by invasive trophoblasts induces the specific migration of CD16- human natural killer cells. *Blood* 2003, 102:1569–1577
- Smith SD, Dunk CE, Aplin JD, Harris LK, Jones RL: Evidence for immune cell involvement in decidual spiral arteriole remodeling in early human pregnancy. *Am J Pathol* 2009, 174:1959–1971
- Pijnenborg R, Vercruyssen L, Hanssens M: The uterine spiral arteries in human pregnancy: facts and controversies. *Placenta* 2006, 27:939–958

31. Craven CM, Morgan T, Ward K: Decidual spiral artery remodelling begins before cellular interaction with cytotrophoblasts. *Placenta* 1998, 19:241–252
32. Pijnenborg R, Robertson WB, Brosens I: The arterial migration of trophoblast in the uterus of the golden hamster, *Mesocricetus auratus*. *J Reprod Fertil* 1974, 40:269–280
33. Lynch L, Golden-Mason L, Eogan M, O’Herlihy C, O’Farrelly C: Cells with haematopoietic stem cell phenotype in adult human endometrium: relevance to infertility? *Hum Reprod* 2007, 22: 919–926
34. Male V, Hughes T, McClory S, Colucci F, Caligiuri MA, Moffett A: Immature NK cells, capable of producing IL-22, are present in human uterine mucosa. *J Immunol* 2010, 185:3913–3918
35. Keskin DB, Allan DS, Rybalov B, Andzelm MM, Stern JN, Kopcow HD, Koopman LA, Strominger JL: TGFbeta promotes conversion of CD16+ peripheral blood NK cells into CD16- NK cells with similarities to decidual NK cells. *Proc Natl Acad Sci U S A* 2007, 104:3378–3383
36. Allan DS, Rybalov B, Awong G, Zuniga-Pflucker JC, Kopcow HD, Carlyle JR, Strominger JL: TGF-beta affects development and differentiation of human natural killer cell subsets. *Eur J Immunol* 2010, 40:2289–2295
37. Yamaguchi T, Kitaya K, Daikoku N, Yasuo T, Fushiki S, Honjo H: Potential selectin L ligands involved in selective recruitment of peripheral blood CD16(-) natural killer cells into human endometrium. *Biol Reprod* 2006, 74:35–40
38. Sela HY, Goldman-Wohl DS, Haimov-Kochman R, Greenfield C, Natanson-Yaron S, Hamani Y, Revel A, Lavy Y, Singer O, Yachimovich-Cohen N, Turetsky T, Mandelboim O, Reubinoff B, Yagel S: Human trophoblast apposition is regulated by interferon gamma-induced protein 10 (IP-10) during early implantation. *Placenta* 2013, 34:222–230
39. Sargent IL, Borzychowski AM, Redman CW: NK cells and human pregnancy—an inflammatory view. *Trends Immunol* 2006, 27: 399–404
40. Carlino C, Stabile H, Morrone S, Bulla R, Soriani A, Agostinis C, Bossi F, Mocchi C, Sarazani F, Tedesco F, Santoni A, Gismondi A: Recruitment of circulating NK cells through decidual tissues: a possible mechanism controlling NK cell accumulation in the uterus during early pregnancy. *Blood* 2008, 111:3108–3115
41. Dunn CL, Critchley HO, Kelly RW: IL-15 regulation in human endometrial stromal cells. *J Clin Endocrinol Metab* 2002, 87: 1898–1901
42. Verma S, Hiby SE, Loke YW, King A: Human decidual natural killer cells express the receptor for and respond to the cytokine interleukin 15. *Biol Reprod* 2000, 62:959–968
43. Carson WE, Giri JG, Lindemann MJ, Linett ML, Ahdieh M, Paxton R, Anderson D, Eisenmann J, Grabstein K, Caligiuri MA: Interleukin (IL) 15 is a novel cytokine that activates human natural killer cells via components of the IL-2 receptor. *J Exp Med* 1994, 180:1395–1403
44. Kopcow HD, Allan DS, Chen X, Rybalov B, Andzelm MM, Ge B, Strominger JL: Human decidual NK cells form immature activating synapses and are not cytotoxic. *Proc Natl Acad Sci U S A* 2005, 102: 15563–15568
45. Kalkunte SS, Mselle TF, Norris WE, Wira CR, Sentman CL, Sharma S: Vascular endothelial growth factor C facilitates immune tolerance and endovascular activity of human uterine NK cells at the maternal-fetal interface. *J Immunol* 2009, 182:4085–4092
46. Hanna J, Goldman-Wohl D, Hamani Y, Avraham I, Greenfield C, Natanson-Yaron S, Prus D, Cohen-Daniel L, Arnon TI, Manaster I, Gazit R, Yutkin V, Benharroch D, Porgador A, Keshet E, Yagel S, Mandelboim O: Decidual NK cells regulate key developmental processes at the human fetal-maternal interface. *Nat Med* 2006, 12:1065–1074
47. Hanna J, Mandelboim O: When killers become helpers. *Trends Immunol* 2007, 28:201–206
48. Dunn CL, Kelly RW, Critchley HO: Decidualization of the human endometrial stromal cell: an enigmatic transformation. *Reprod Bio-med Online* 2003, 7:151–161
49. Red-Horse K, Drake PM, Fisher SJ: Human pregnancy: the role of chemokine networks at the fetal-maternal interface. *Expert Rev Mol Med* 2004, 6(11):1–14
50. Horvath CM: The Jak-STAT pathway stimulated by interferon gamma. *Sci STKE* 2004, 260:tr8. <http://dx.doi.org/10.1126/stke.2602004tr8>
51. Li M, Wu ZM, Yang H, Huang SJ: NFkappaB and JNK/MAPK activation mediates the production of major macrophage- or dendritic cell-recruiting chemokine in human first trimester decidual cells in response to proinflammatory stimuli. *J Clin Endocrinol Metab* 2011, 96:2502–2511
52. Hagmann H, Thadhani R, Benzing T, Karumanchi SA, Stepan H: The promise of angiogenic markers for the early diagnosis and prediction of preeclampsia. *Clin Chem* 2012, 58:837–845
53. Zhao Z, Moley KH, Gronowski AM: Diagnostic potential for miRNAs as biomarkers for pregnancy-specific diseases. *Clin Biochem* 2013, 46:953–960
54. Redman CW, Sargent IL: Immunology of pre-eclampsia. *Am J Reprod Immunol* 2010, 63:534–543
55. Burton GJ, Yung HW, Cindrova-Davies T, Charnock-Jones DS: Placental endoplasmic reticulum stress and oxidative stress in the pathophysiology of unexplained intrauterine growth restriction and early onset preeclampsia. *Placenta* 2009, 30(Suppl A):S43–S48
56. Huang SJ, Chen CP, Schatz F, Rahman M, Abrahams VM, Lockwood CJ: Pre-eclampsia is associated with dendritic cell recruitment into the uterine decidua. *J Pathol* 2008, 214:328–336
57. Wu ZM, Yang H, Li M, Yeh CC, Schatz F, Lockwood CJ, Di W, Huang SJ: Pro-inflammatory cytokine-stimulated first trimester decidual cells enhance macrophage-induced apoptosis of extravillous trophoblasts. *Placenta* 2012, 33:188–194
58. Lash GE, Bulmer JN: Do uterine natural killer (uNK) cells contribute to female reproductive disorders? *J Reprod Immunol* 2011, 88:156–164
59. Hiby SE, Apps R, Sharkey AM, Farrell LE, Gardner L, Mulder A, Claas FH, Walker JJ, Redman CW, Morgan L, Tower C, Regan L, Moore GE, Carrington M, Moffett A: Maternal activating KIRs protect against human reproductive failure mediated by fetal HLA-C2. *J Clin Invest* 2010, 120:4102–4110