

# Methylene Blue for Distributive Shock: A Potential New Use of an Old Antidote

David H. Jang · Lewis S. Nelson · Robert S. Hoffman


Published online: 12 April 2013  
© American College of Medical Toxicology 2013

**Abstract** Methylene blue is used primarily in the treatment of patients with methemoglobinemia. Most recently, methylene blue has been used as a treatment for refractory distributive shock from a variety of causes such as sepsis and anaphylaxis. Many studies suggest that the nitric oxide–cyclic guanosine monophosphate (NO–cGMP) pathway plays a significant role in the pathophysiology of distributive shock. There are some experimental and clinical experiences with the use of methylene blue as a selective inhibitor of the NO–cGMP pathway. Methylene blue may play a role in the treatment of distributive shock when standard treatment fails.

**Keywords** Methylene blue · Refractory shock · Cardiovascular drug overdoses · Calcium channel blockers

## Introduction

Methylene blue is a phenothiazine-related heterocyclic aromatic molecule (C<sub>16</sub>H<sub>18</sub>N<sub>3</sub>SCl). It is a solid, odorless, dark green powder at room temperature that yields a blue solution when it is dissolved in water. Methylene blue is used in a wide variety of settings and for many purposes; for example, as a redox indicator or as a dye/stain [1, 2].



Methylene Blue

The medicinal use of methylene blue dates back to the late 1800s as a treatment for malaria, a urinary analgesic, and in the treatment for cyanide and carbon monoxide poisoning [3–6]. Methylene blue is currently utilized as a treatment for ifosfamide neurotoxicity [7, 8]. Methylene blue is most commonly used as a reducing agent in the treatment of patients with methemoglobinemia [9–11]. Most recently, methylene blue has been used as a treatment for refractory distributive shock through a lesser known property: inhibition of the downstream effect of nitric oxide (NO). The objective of this review is to explain the pathophysiological basis of methylene blue for the use in distributive shock and review the current literature.

## Methods

The authors conducted a scientific review of all available literature published over the last 20 years. Our primary objective was to evaluate the use of methylene blue as a treatment for distributive shock from causes such as sepsis and anaphylaxis. Our secondary objective was to summarize the proposed mechanisms for the use of methylene blue based on existing human, animal, and in vitro studies. We initiated a PubMed database search using the MESH terms “methylene blue,” “sepsis,” “anaphylaxis,” “nitric oxide,” “nitric oxide synthase,” and “cGMP.” Articles were selected and agreed upon by the authors based on relevance and impact. Effort was made to include both positive and negative studies where appropriate. Emphasis was placed on well-conducted

D. H. Jang (✉) · L. S. Nelson · R. S. Hoffman  
Division of Medical Toxicology, Department of Emergency  
Medicine, New York University School of Medicine,  
Bellevue Hospital Center, 455 First Avenue, Room 123,  
New York, NY 10016, USA  
e-mail: Jangd01@nyumc.org

L. S. Nelson  
e-mail: lnelsonmd@gmail.com

R. S. Hoffman  
e-mail: bobhoffmd@gmail.com

experimental data, case studies, and controlled trials when possible. Studies were only excluded due to redundancy. After analysis of the available data, this paper concludes with recommendations based on the existing scientific evidence.

### Physiology of Vascular Tone

#### Nitric Oxide Synthase General Structure and Function


NO has many functions, such as serving as a messenger molecule, regulating gene transcription and mRNA translation, as well as assisting in neurotransmission [12, 13]. NO also plays an important role in the regulation of vascular tone (Fig. 1) and pathophysiology of shock when it is excessively produced. Initial evidence that NO plays an important role in the regulation of vascular tone was demonstrated with nitrate production in experimental models [14]. Later work revealed that L-arginine was the substrate and that L-citrulline was a by-product for the production of NO by an enzyme known as nitric oxide synthase (NOS) [15]. NOS, the enzyme that produces NO, is a homodimer with a substrate access channel which extends from the active site toward the dimer interface along with a reductase domain and an oxygenase domain (heme and H4B binding) [16]. In humans, there exist three different isoforms of NOS known as: neuronal NOS (nNOS, NOS1), inducible NOS (iNOS, NOS2), and endothelial NOS (eNOS, NOS3) [17]. All isoforms of NOS utilize L-arginine as a substrate along with oxygen

and reduced nicotinamide-adenine-dinucleotide phosphate as co-substrates [18]. In addition, all isoforms of NOS bind calmodulin and calcium; activated calmodulin is important for the regulation of eNOS and nNOS activity [19]. Production of NO is a two-step process: In the first step, NOS hydroxylates L-arginine to an intermediate known as *N*-hydroxy-L-arginine. In a second step, NOS oxidizes this intermediate to L-citrulline and NO. Inactivation of NO occurs when it combines with a superoxide anion,  $O_2^-$ , to form a very potent oxidant known as peroxynitrite. Peroxynitrite causes DNA damage and oxidation, nitration, and S-nitrosylation of lipids and proteins [20, 21].

Certain neurons of the brain express nNOS that is regulated by calcium and calmodulin [22]. Although nNOS exists in the brain, spinal cord, adrenal glands, and sympathetic ganglia, the largest source of nNOS in mammals is found within skeletal muscle [23]. The physiologic functions of nNOS are varied and include modulation of learning, neurogenesis, and memory [24, 25]. There is strong evidence that nNOS also centrally regulates blood pressure [26, 27]. Certain neurodegenerative diseases such as Parkinson’s and Alzheimer’s disease are associated with increased nNOS function. Increased nNOS function may lead to neuron cell death from large calcium influx mediated by *N*-methyl-D-aspartate receptors. Increased nNOS activity may also result in increased NO production with peroxynitrite formation leading to cell injury and mitochondrial dysfunction [28].

Endothelial NOS (eNOS) exists primarily in endothelial cells, but also occurs in other cell types including platelets and cardiac myocytes [29]. eNOS activity significantly increases

**Fig. 1** Regulation of the relaxation of vascular smooth muscle by nitric oxide. Nitric oxide is produced from L-arginine by nitric oxide synthase leading to the activation of soluble guanylyl cyclase with production of cyclic guanosine 3',5'-monophosphate (cGMP). cGMP-dependent protein kinase is activated, leading to (1) decreased sensitivity of myosin to calcium-induced contractions, (2) activation of calcium-sensitive potassium channels which in turn decreases calcium entry through calcium channels, and (3) inhibition of release of calcium from the sarcoplasmic reticulum. This results in relaxation of smooth muscle



with a rise in intracellular calcium with resultant increased NO production [30]. The physiologic functions of eNOS are varied and include inhibition of platelet aggregation and adhesion and inhibition of leukocyte aggregation, all of which help prevent atherogenesis [31, 32]. The primary role of eNOS appears to be regulation of cardiovascular functions such as blood pressure [33]. Dysfunction of eNOS activity is associated with cardiovascular disease through either inability to generate adequate amounts of NO for proper vasodilation and inhibition of platelet/leukocyte aggregation or excessive production of NO with resultant oxidative stress and endothelial cell dysfunction [34].

Inducible NOS (iNOS) expression is regulated by a variety of factors such as cytokines, interleukins, and bacterial lipopolysaccharide. iNOS was first identified in macrophages. However, further work revealed that expression of iNOS can be stimulated in almost any cell line [35, 36]. The primary physiologic function of iNOS in macrophages is to produce NO as a major cytotoxic agent. Thus, iNOS plays a role in immune defense and mediation of inflammation. NO can also inhibit enzymes that contain iron within their catalytic centers, as is often found in microorganisms. The production of NO has a cytotoxic effect on tumors and various microorganisms, such as parasites [37, 38]. While the production of NO from iNOS plays an important role in host defense, large concentrations of NO may also injure surrounding host cells. Overproduction of NO contributes to allograft rejection and is also implicated in the pathogenesis of various forms of distributive shock such as sepsis and anaphylaxis characterized by arteriolar vasodilation, hypotension, and poor response to vasopressors [39–41].

#### Guanylyl Cyclase and Signaling by Cyclic Guanosine Monophosphate

An important signaling pathway of NO is the activation of soluble guanylyl cyclase (sGC) resulting in the production of cyclic guanosine monophosphate (cGMP) [42]. sGC are a family of proteins composed of membrane-bound and soluble isoforms and expressed in virtually all cell types [43]. sGC catalyze conversion of the purine nucleoside guanosine-5'-triphosphate to cGMP in response to various messengers such as peptide ligands, calcium influx, and NO [44].

Intracellular cGMP gates specific ion channels, modulates nucleotide concentrations by regulating various phosphodiesterases, and activates protein kinases [42]. cGMP has a central role in many other vital processes such as retinal phototransduction, electrolyte regulation, and mediation of vascular smooth muscle tone [45, 46]. Excessive cGMP production, generally through abundant NO availability, is implicated in the pathophysiology of distributive shock such as sepsis and anaphylaxis [47]. Dysfunction within the NO–cGMP pathway has been implicated in certain disease states of distributive shock such as sepsis and anaphylaxis.

#### Distributive Shock

The definition of shock has evolved over centuries. The term “shock” was defined as sudden deterioration of a patient’s condition from major trauma in the 1700s. This term was translated by the English physician Clarke from the writings of a French surgeon Le Dran who is often credited with the origin of the term “shock.” While the term “shock” has taken on many meanings, the current emphasis is on tissue perfusion in relationship to cellular function. Shock is often a syndrome caused by a systemic derangement of perfusion leading to widespread cellular hypoxia and organ dysfunction. While hypovolemic shock from blood loss following trauma was the first form of shock to be recognized, other etiologies of shock are now recognized. These include cardiogenic (e.g., heart failure), obstructive (e.g., tension pneumothorax), and distributive shock. Distributive shock is characterized by loss of vasomotor tone or an overall decrease in systematic vascular resistance, and is often accompanied by an increase in cardiac output. Sepsis and anaphylaxis are the two most common etiologies of distributive shock. Also, drug-induced shock from certain medications such as calcium channel blockers may have a distributive component as well.

#### The NO–cGMP Pathway in Anaphylaxis and the Role of Methylene Blue

Anaphylaxis is a serious allergic reaction with a systemic response that can be life-threatening. The most common causes of anaphylaxis include allergic reaction to medications, food, and hymenoptera venom, although in up to 75 % of cases, there is no previous history of allergy [48]. The signs and symptoms of anaphylaxis manifest in all organ systems, but the life-threatening effects occur in the respiratory (e.g., bronchospasm) and cardiovascular system (e.g., refractory hypotension from peripheral vasodilation) [49].

The pathophysiology of distributive shock from anaphylaxis is complex involving many mediators such as bradykinin and other leukotrienes, but the mediator believed to have the most significant role is histamine [50]. Early experiments demonstrate that histamine-induced vasodilation is dependent on intact endothelium containing eNOS which suggests that histamine’s action is at least partly mediated through nitric oxide [51, 52]. Further evidence to support the role of the NO–cGMP pathway in anaphylaxis is demonstrated by the finding that increased histamine results in upregulation of eNOS gene expression with increased production of NO leading to vasodilation through increased activity of GC. Inhibition of GC reduces vasodilation induced by histamine [53–55]. This evidence provides support that histamine-induced vasodilation is at least partly mediated through the NO–cGMP pathway and that inhibition of this pathway reverses the vasodilation.

The treatment of anaphylaxis-induced hypotension consists of isotonic fluids, antihistamines, steroids, and in severe cases epinephrine. While sufficient in many instances, severe cases of anaphylaxis may not respond to these interventions, and patients may remain persistently hypotensive. There are some experimental and clinical experiences with the use of methylene blue as a selective inhibitor of the NO–cGMP pathway in anaphylaxis. A study of anaphylactic shock in rabbits showed that the use of methylene blue (3 mg/kg) as a single dose increased survival time compared to controls. Plasma nitrate, a marker of NO release, did not differ between groups. This finding suggests that methylene blue does not mediate its action through NOS inhibition or NO scavenging, and suggests that its primary action may be mediated through GC inhibition [56].

The use of methylene blue for the treatment of humans with refractory anaphylactic shock is limited to case reports and case series. Methylene blue was used successfully to treat three patients who experienced anaphylactic shock following radiocontrast injection during coronary angiography. All had prompt improvement and hemodynamic stabilization without obvious adverse effects [57]. The successful use of methylene blue for refractory hypotension secondary to protamine during cardiac bypass is also reported [58]. Table 1 lists reported cases of methylene blue administration for treatment of anaphylactic shock.

#### The Role of the NO–cGMP Pathway in Septic Shock and Treatment with Methylene Blue

Sepsis is defined as a constellation of clinical findings that result from an infection and a subsequent inflammatory response from a combination of pathogenic products and the body’s own immune response. The most severe form of

sepsis is “septic shock,” which is defined as hypotension with signs of tissue hypoperfusion despite fluid resuscitation in the presence of overwhelming infection [59]. There are over 700,000 cases of severe sepsis in the USA annually, and it is the leading cause of ICU mortality [60].

Sepsis is associated with a complex hemodynamic profile when compared to other causes of shock. Septic shock can have features of hypovolemia from capillary leak (i.e., hypovolemic shock) and depressed myocardial function with resultant reduction in cardiac output (i.e., cardiogenic shock). The most characteristic finding with septic shock is profound peripheral vasodilation (i.e., distributive shock). The pathogenesis of septic shock is a complex response of cellular activation that releases a multitude of proinflammatory mediators which includes excessive NO [61]. The overproduction of NO by iNOS in sepsis results in increased production of cGMP contributing to the refractory hypotension observed with septic shock. Numerous animal studies demonstrate large increases of NO after exposure to lipopolysaccharide (i.e., endotoxin) with resultant hypotension [62–67].

The current management of septic shock includes early goal-directed therapy centered on early administration of empiric antibiotics, volume resuscitation, frequent monitoring, and cardiac support with appropriate use of vasopressors. Despite these therapies, the mortality rate remains around 30 % [68]. As a result, the need for additional or alternative therapy for septic shock is clear. One such treatment modality may involve inhibition along the NO–cGMP pathway. Earlier work with nonspecific competitive NOS inhibitors, such as *N*-monomethyl-L-arginine for the treatment of distributive shock in sepsis noted an increase in mean arterial pressure but no mortality benefit. In fact, the use of nonspecific nitric oxide synthase inhibitors resulted in

**Table 1** Summary of publishing cases of methylene blue use for refractory shock from anaphylaxis

Reference	Age/gender	Cause of anaphylaxis	Clinical presentation	Interventions	IV methylene blue dose	Time of effect	Outcome
[57]	57-year-old woman	Radiocontrast	Refractory hypotension, angioedema	Hydrocortisone, epinephrine	1.5 mg/kg bolus	20 min	Survival
[57]	48-year-old woman	Radiocontrast	Refractory hypotension, oxygen desaturation	Intubation, Isotonic fluids, epinephrine, dopamine, hydrocortisone	2 mg/kg bolus followed by 2 mg/kg/h for 2 h	10 min	Survival
[57]	53-year-old man	Radiocontrast	Refractory hypotension, angioedema	Hydrocortisone ×2	1.5 mg/kg infusion	Immediate	Survival
[58]	72-year-old man	Protamine 100 mg	Refractory hypotension, urticaria	Hydrocortisone, pressure, calcium, cardiac bypass	100 mg	15 min	Survival
[58]	72-year-old woman	Aprotinin	Refractory hypotension	Hydrocortisone, pressure, fluids, cardiac bypass	100 mg	10 min	Survival
[98]	23-year-old woman	Unknown	Angioedema	Antihistamine, epinephrine	1.5 mg/kg bolus with 1.5 mg/kg infusion	20 min	Survival
[99]	79-year-old man	Protamine	Urticaria, angioedema, refractory hypotension	Antihistamine pressors, fluids	35 mg	20 min	Survival
[100]	72-year-old man	Protamine	Refractory hypotension	Pressors	2 mg/kg bolus	30 min	Survival

increased mortality in both animal and humans. NO has a vital, positive role in other pathways so nonspecific inhibition of nitric oxide synthase is detrimental [69]. This begs the questions of whether inhibition further along the NO–cGMP pathway might be beneficial. Later studies examined selective inhibition of guanylate cyclase. One of these selective agents is methylene blue, which may restore vascular tone and improve tissue perfusion.

There are multiple experimental models of septic shock that support the use of methylene blue in the treatment of refractory distributive shock [70–76]. The majority of clinical data on the use of methylene blue for such therapy is observational with case reports and small observational studies, but there are also small controlled clinical trials [77–82]. The majority of these studies demonstrate an increase in mean arterial pressure and/or decrease in pressor requirement. The first prospective randomized controlled study using methylene blue in the treatment of septic shock was published in 2001. Patients were randomized 1:1 to receive either methylene blue or isotonic saline as a control in addition to conventional treatment. When compared to the control group, methylene blue reduced pressor requirement. Although 5/10 patients treated with methylene blue survived as opposed to 3/10 patients who received conventional treatment, this study was not powered to find a mortality difference [83].

Another randomized control trial measured plasma concentrations of cytokines in severe sepsis. In this study, 15 patients received methylene blue and 15 patients received isotonic saline for 6 h. Various cytokines such as interleukin-2 and tumor necrosis factor alpha were measured at baseline, immediately after treatment, 24 h posttreatment, and 48 h posttreatment. When compared to the control group, methylene blue resulted in higher mean arterial pressures, but no difference in mortality or in the various cytokines measured. The primary limitation of this study was the short observation period and small sample size [84].

A prospective, randomized, double-blind, single-center study in 15 consecutive, mechanically ventilated patients with septic shock admitted to the intensive care unit compared escalating doses of methylene blue. Methylene blue was infused at 1 mg/kg ( $n=4$ ), 3 mg/kg ( $n=6$ ), or 7 mg/kg ( $n=5$ ) over 20 min. Methylene blue had a dose-dependent effect on cardiac index, mean pulmonary artery and pulmonary artery occlusion pressure as well as oxygen delivery and lactate concentrations. The data suggested that in human septic shock, methylene blue increases mean arterial blood pressure through an increase in cardiac index and systemic vascular resistance [85].

In summary, the clinical studies that evaluate the use of methylene blue for septic shock demonstrate an improvement in hemodynamics and address the mechanism by which methylene blue may work. However, no mortality benefit is shown which may be due to the small sample sizes found in these studies.

The dosing regimen for methylene blue based on experimental and clinical data is not entirely clear but does seem to be similar to what is used for the treatment of methemoglobinemia: 1–2 mg/kg. The dosing used for refractory septic shock has included a single bolus, repeated bolus based on response, low-dose infusion, and infusions followed by a bolus. It is clear based on other studies that the use of high doses of methylene blue, typically doses greater than 7 mg/kg, is associated with adverse effects such as paradoxical induction of methemoglobinemia, acute hemolytic anemia, and detrimental effects on pulmonary function [86, 87].

Based on current data, it appears that the use of a single bolus improves hemodynamics for the duration of a few hours. It is unclear if there is any mortality benefit with temporary improvement of hemodynamics, and for this reason, some authors have advocated using methylene blue as an infusion [78, 79]. However, due to limited experience with this method of administration as well as the lack of data on dosing methylene blue in the setting of liver or kidney dysfunction, the use of methylene blue as an infusion cannot be recommended at this time.

The use of methylene blue has a favorable side effect profile, but it is associated with some adverse effects that are important for the clinician to note. Methylene blue rarely may cause shortness of breath, tremors, vomiting, blue discoloration of body fluids, and acute hemolytic anemia with high doses [88]. Another uncommon adverse effect for clinicians to be aware of is the precipitation of serotonin syndrome [89, 90]. Serotonin syndrome is a potentially fatal condition characterized by autonomic instability that includes agitation, tachycardia, hypertension, and hyperthermia due to excessive stimulation at the 5-HT<sub>2A</sub> receptor. Serotonin syndrome is most commonly precipitated from drug interactions of multiple serotonergic agents or agents that inhibit metabolism of serotonin such as monoamine oxidase inhibitors [91]. Methylene blue may cause serotonin syndrome because both the parent compound and metabolite, azure B, inhibit MAO-A [92]. Methylene blue causing serotonin syndrome is relatively rare occurring mostly with the use of very high doses.

The patient population that may benefit from the use of methylene blue for refractory distributive shock requires careful consideration. Cardiac support with catecholamines and volume expansion should be the initial therapy of choice. In patients who do not respond to standard treatment, methylene blue may be considered. The ideal dose is unknown, but based on the existing literature and safety profile, we recommend a dose of 1–2 mg/kg as a single bolus. Methylene blue should not be used in patients with pulmonary hypertension, underlying G6PD deficiency, and acute lung injury.

## The Use of Methylene Blue for Refractory Shock from Cardiovascular Drug Poisoning

Methylene blue has also been utilized in a case of refractory distributive shock from a cardiovascular drug overdose with a dihydropyridine calcium channel blocker [93]. This case described a confirmed isolated dihydropyridine overdose resulting in profound distributive shock as demonstrated by a Swan-Ganz catheter. The patient remained hypotensive despite multiple therapeutic interventions including isotonic saline, multiple doses of calcium gluconate, glucagon, multiple pressors and high-dose insulin euglycemic (HIE) therapy for several hours. The patient then received methylene blue at 2 mg/kg and soon after showed improvement in hemodynamics with eventual discontinuation of both pressors and HIE. While this case should not be used as sole support for the use of methylene blue as treatment for cardiovascular drug poisoning, this is a potential area of research.

While the primary mechanism of calcium channel blockers involves blockade of L-type calcium channels, experimental studies have demonstrated an additional vasodilatory mechanism through the release of nitric oxide. Although the exact mechanism of NO release is not clear, one of the reasons some dihydropyridine calcium channel blockers increase nitric oxide production is from an increased endothelial nitric oxide synthase activity through phosphorylation of this enzyme [94–97]. It is not clear how much of a role the NO–cGMP pathway may play in an acute overdose with certain calcium channel blockers, but this may explain the mechanism by which methylene blue acts.

### Conclusion

Increased NO and cGMP production in tissues is well documented in various conditions of distributive shock such as sepsis and anaphylaxis. The NO–cGMP pathway plays a central role in the pathophysiology of distributive shock. There is experimental and clinical evidence that support the use of a selective NO–cGMP pathway inhibitor, methylene blue, as a treatment in cases of refractory vasodilatory shock with no response to conventional treatment. Clinicians should be aware of potential adverse effects and drug interactions with serotonergic agents when considering the use of methylene blue.

**Acknowledgments** This study was supported in part by grant 5UL1RR029893 from the National Center for Research Resources, National Institutes of Health.

**Conflict of Interest** The authors have no conflict of interest to declare.

### References

- Spaeth CS, Robison T, Fan JD et al. (2012) Cellular mechanisms of plasmalemmal sealing and axonal repair by polyethylene glycol and methylene blue. *J Neurosci Res* 90(5):955–966 doi:10.1002/jnr.23022
- Sohrabnezhad SH (2011) Study of catalytic reduction and photodegradation of methylene blue by heterogeneous catalyst. *Spectrochim Acta A Mol Biomol Spectrosc* 81(1):228–235
- Kasozi DM, Gromer S, Adler H et al (2011) The bacterial redox signalling pyocyanin as an antiplasmodial agent: comparisons with its thioanalog methylene blue. *Redox Rep* 16(4):154–165
- Meissner PE, Mandi G, Coulibaly B et al (2006) Methylene blue for malaria in Africa: results from a dose-finding study in combination with chloroquine. *Malar J* 8:5–84
- Adjalley SH, Johnston GL, Li T, Eastman RT et al (2011) Quantitative assessment of *Plasmodium falciparum* sexual development reveals potent transmission-blocking activity by methylene blue. *Proc Natl Acad Sci U S A* 108(47):E1214–E1223
- Hanzlik PJ (1933) Subject of this letter: Methylene blue as antidote for cyanide poisoning. *Cal West Med* 38(3):225–226
- Giovanis P, Gama A, Marcante M et al (2009) Ifosfamide encephalopathy and use of methylene blue. A case report of different sequential neurotoxicity. *Tumori* 95(4):545–546
- Richards A, Marshall H, McQuary A (2011) Evaluation of methylene blue, thiamine, and/or albumin in the prevention of ifosfamide-related neurotoxicity. *J Oncol Pharm Pract* 17(4):372–380
- Barclay JA, Ziemba SE, Ibrahim RB (2011) Dapsone-induced methemoglobinemia: a primer for clinicians. *Ann Pharmacother* 45(9):1103–1115
- So TY, Farrington E (2008) Topical benzocaine-induced methemoglobinemia in the pediatric population. *J Pediatr Health Care* 22(6):335–339
- Hahn IH, Hoffman RS, Nelson LS (2004) EMLA-induced methemoglobinemia and systemic topical anesthetic toxicity. *J Emerg Med* 26(1):85–88
- Alderton WK, Cooper CE, Knowles RG (2001) Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357:593–615
- Schuman EM, Madison DV (1991) A requirement for the intercellular messenger nitric oxide in long-term potentiation. *Science* 254:1503–1506
- Llyengar R, Stuehr DJ, Marietta MA (1987) Macrophage synthesis of nitrite, nitrate, and N-nitrosamines: precursors and role of the respiratory burst. *Proc Natl Acad* 84:6369–6373
- Hibbs JB, Taintor RR, Vavrin Z (1987) Macrophage cytotoxicity: role of L-arginine deiminase and imino nitrogen oxidation to nitrite. *Science* 235:473–476
- Garcin ED, Bruns CM, Lloyd SJ et al (2004) Structural basis for isozyme-specific regulation of electron transfer in nitric-oxide synthase. *J Biol Chem* 279:37918–37927
- Daff S (2012) NO synthase: structures and mechanisms. *Nitric Oxide* 23:1–11
- Stuehr D, Pou S, Rosen GM (2001) Oxygen reduction by nitric-oxide synthases. *J Biol Chem* 276:14533–14536
- Tatsumi R, Wuollet AL, Tabata K et al (2009) A role for calcium-calmodulin in regulating nitric oxide production during skeletal muscle satellite cell activation. *Am J Physiol Cell Physiol* 296(4):C922–C929
- Beckman JS, Beckman TW, Chen J et al (1990) Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci U S A* 87(4):1620–1624

21. Beckman JS, Crow JP (1993) Pathological implications of nitric oxide, superoxide and peroxynitrite formation. *Biochem Soc Trans* 21(2):330–334
22. Ma L (1993) Evidence for nitric oxide-generator cells in the brain. *Bull Tokyo Med Dent Univ* 40(3):125–134
23. Forstermann U, Closs EI, Pollock JS, Nakane et al (1994) Nitric oxide synthase isozymes: characterization, purification, molecular cloning, and functions. *Hypertension* 23:1121–1131
24. Salemme E, Diano S, Maharajan P, Maharajan V (1996) Nitric oxide, a neuronal messenger. Its role in the hippocampus neuronal plasticity. *Riv Biol* 89(1):87–107
25. Izumi Y, Clifford DB, Zorumski CF (1992) Inhibition of long-term potentiation by NMDA-mediated nitric oxide release. *Science* 257:1273–1276
26. Handy RL, Wallace P, Moore PK (1996) Inhibition of nitric oxide synthase by isothioureas: cardiovascular and antinociceptive effects. *Pharmacol Biochem Behav* 55(2):179–184
27. Togashi H, Sakuma I, Yoshioka M et al (1992) A central nervous system action of nitric oxide in blood pressure regulation. *J Pharmacol Exp Ther* 262:343–347
28. Lipton SA, Choi YB, Pan ZH et al (1993) A redox-based mechanism for the neuroprotective and neurodestructive effects of nitric oxide and related nitroso compounds. *Nature* 364:626–632
29. Shaul PW (2002) Regulation of endothelial nitric oxide synthase: location, location, location. *Annu Rev Physiol* 64:749–774
30. Venema RC, Sayegh HS, Arnal JF et al (1995) Role of the enzyme calmodulin-binding domain in membrane association and phospholipid inhibition of endothelial nitric oxide synthase. *J Biol Chem* 270(24):14705–14711
31. Zeiher AM, Fisslthaler B, Schray-Utz B et al (1995) Nitric oxide modulates the expression of monocyte chemoattractant protein 1 in cultured human endothelial cells. *Circ Res* 76:980–986
32. Alheid U, Frolich JC, Forstermann U (1987) Endothelium-derived relaxing factor from cultured human endothelial cells inhibits aggregation of human platelets. *Thromb Res* 47:561–571
33. Kurihara N, Alfie ME, Sigmon DH et al (1998) Role of nNOS in blood pressure regulation in eNOS null mutant mice. *Hypertension* 32(5):856–861
34. Triggle CR, Ding H (2010) A review of endothelial dysfunction in diabetes: a focus on the contribution of a dysfunctional eNOS. *J Am Soc Hypertens* 4(3):102–115
35. Nathan CF, Hibbs JB (1991) Role of nitric oxide synthesis in macrophage antimicrobial activity. *Curr Opin Immunol* 3:65–70
36. MacNaul KL, Hutchinson NI (1993) Differential expression of iNOS and eNOS mRNA in human vascular smooth muscle cells and endothelial cells under normal and inflammatory conditions. *Biochem Biophys Res Commun* 196(3):1330–1334
37. Tamir S, deRoja-Walker T, Gal A et al (1995) Nitric oxide production in relation to spontaneous B-cell lymphoma and myositis in SJL mice. *Cancer Res* 55(19):4391–4397
38. Losada AP, Bermúdez R, Faílde LD et al (2012) Quantitative and qualitative evaluation of iNOS expression in turbot (*Psetta maxima*) infected with *Enteromyxum scophthalmi*. *Fish Shellfish Immunol* 32(2):243–248
39. Sade K, Schwartz IF, Etkin S et al (2007) Expression of inducible nitric oxide synthase in a mouse model of anaphylaxis. *J Investig Allergol Clin Immunol* 17(6):379–385
40. Matuschek A, Ulbrich M, Timm S et al (2009) Analysis of parathyroid graft rejection suggests alloantigen-specific production of nitric oxide by iNOS-positive intragraft macrophages. *Transpl Immunol* 21(4):183–191
41. Okamoto I, Abe M, Shibata K et al (2000) Evaluating the role of inducible nitric oxide synthase using a novel and selective inducible nitric oxide synthase inhibitor in septic lung injury produced by cecal ligation and puncture. *Am J Respir Crit Care Med* 162(2 Pt 1):716–722
42. Lucas KA, Pitari GM, Kazerounian S et al (2000) Guanylyl cyclases and signaling by cyclic GMP. *Pharmacol Rev* 52:375–413
43. Wedel B, Garbers D (2001) The guanylyl cyclase family at Y2K. *Annu Rev Physiol* 63:215–233
44. Mujoo K, Sharin VG, Martin E et al (2010) Role of soluble guanylyl cyclase-cyclic GMP signaling in tumor cell proliferation. *Nitric Oxide* 22(1):43–50
45. Schoeffter P, Lugnier C, Demesy-Waeldele F et al (1987) Role of cyclic AMP- and cyclic GMP-phosphodiesterases in the control of cyclic nucleotide levels and smooth muscle tone in rat isolated aorta. A study with selective inhibitors. *Biochem Pharmacol* 36(22):3965–3972
46. Bensinger RE, Podos SM (1975) Cyclic nucleotide metabolism in the retina. *Invest Ophthalmol* 14(4):263–266
47. Kumar A, Brar R, Wang P et al (1999) Role of nitric oxide and cGMP in human septic serum-induced depression of cardiac myocyte contractility. *Am J Physiol* 276:R256–R276
48. Winbery SL, Lieberman PL (2002) Histamine and antihistamines in anaphylaxis. *Clin Allergy Immunol* 17:287–317
49. Lieberman P (1990) The use of antihistamines in the prevention and treatment of anaphylaxis and anaphylactoid reactions. *J Allergy Clin Immunol* 86(4 Pt 2):684–686
50. Enjeti S, Bleecker ER, Smith PL et al (1983) Hemodynamic mechanisms in anaphylaxis. *Circ Shock* 11:297–309
51. Toda N (1984) Endothelium-dependent relaxation induced by angiotensin II and histamine in isolated arteries of dog. *Br J Pharmacol* 81:301–307
52. Krstic MK, Stepanovic RM, Krstic SK et al (1989) Endothelium-dependent relaxation of the rat renal artery caused by activation of histamine H1-receptors. *Pharmacology* 38:113–120
53. Rosenkranz-Weiss P, Sessa WC, Milstien S et al (1994) Regulation of nitric oxide synthesis by proinflammatory cytokines in human umbilical vein endothelial cells: elevations in tetrahydrobiopterin levels enhance endothelial nitric oxide synthase specific activity. *J Clin Invest* 93:2236–2243
54. Li H, Burkhardt C, Heinrich UR et al (2003) Histamine upregulates gene expression of endothelial nitric oxide synthase in human vascular endothelial cells. *Circulation* 107:2348–2354
55. Champion HC, Kadowitz PJ (1997) NO release and the opening of K<sup>+</sup>ATP channels mediate vasodilator responses to histamine in the cat. *Am J Physiol* 273(2 pt 2):H928–H937
56. Buzato MA, Viaro F, Piccinato CE et al (2005) The use of methylene blue in the treatment of anaphylactic shock induced by compound 48/80: experimental studies in rabbits. *Shock* 23(6):582–587
57. Oliveira Neto AM, Duarte NM, Vicente WV et al (2003) Methylene blue: an effective treatment for contrast medium-induced anaphylaxis. *Med Sci Monit* 9(11):CS102–CS106
58. Del Duca D, Sheth SS, Clarke AE et al (2009) Use of methylene blue for catecholamine-refractory vasoplegia from protamine and aprotinin. *Ann Thorac Surg* 87(2):640–642
59. Levy MM, Fink MP, Marshall JC et al (2003) 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 31:1250–1256
60. Angus DC, Linde-Zwirble WT, Lidicker J et al (2001) Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 29:1303–1310
61. Dellinger RP (2003) Inflammation and coagulation: implications for the septic patient. *Clin Infect Dis* 36:1259–1265
62. Kumar A, Brar R, Wang P et al (1999) Role of nitric oxide and cGMP in human septic serum-induced depression of cardiac myocyte contractility. *Am J Physiol* 276(1 Pt 2):R265–R276
63. Symeonides S, Balk RA (1999) Nitric oxide in the pathogenesis of sepsis. *Infect Dis Clin North Am* 13(2):449–63 x

64. Fleming I, Julou-Schaeffer G, Gray GA et al (1991) Evidence that an L-arginine/nitric oxide dependent elevation of tissue cyclic GMP content is involved in depression of vascular reactivity by endotoxin. *Br J Pharmacol* 103(1):1047–1052
65. Schott CA, Gray GA, Stoclet JC (1993) Dependence of endotoxin-induced vascular hyporeactivity on extracellular L-arginine. *Br J Pharmacol* 108(1):38–43
66. Julou-Schaeffer G, Gray GA, Fleming I et al (1990) Loss of vascular responsiveness induced by endotoxin involves L-arginine pathway. *Am J Physiol* 259(4 Pt 2):H1038–H1043
67. Schuller F, Fleming I, Stoclet JC et al (1992) Effect of endotoxin on circulating cyclic GMP in the rat. *Eur J Pharmacol* 212(1):93–96
68. Rivers E, Nguyen B, Havstad S et al (2001) Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345:1368–1377
69. Watson D, Grover R, Anzueto A et al (2004) Cardiovascular effects of the nitric oxide synthase inhibitor NG-methyl-L-arginine hydrochloride (546C88) in patients with septic shock: results of a randomized, double-blind, placebo-controlled multicenter study (study no. 144–002). *Crit Care Med* 32(1):13–20
70. Cheng X, Pang CC (1998) Pressor and vasoconstrictor effects of methylene blue in endotoxaemic rats. *Naunyn Schmiedeberg Arch Pharmacol* 357(6):648–653
71. Fernandes D, Sordi R, Pacheco LK et al (2009) Late, but not early, inhibition of soluble guanylate cyclase decreases mortality in a rat sepsis model. *J Pharmacol Exp Ther* 328(3):991–999
72. Demirbilek S, Sizanli E, Karadag N et al (2006) The effects of methylene blue on lung injury in septic rats. *Eur Surg Res* 38(1):35–41
73. Menardi AC, Viaro F, Vicente WV et al (2006) Hemodynamic and vascular endothelium function studies in healthy pigs after intravenous bolus infusion of methylene blue. *Arq Bras Cardiol* 87(4):525–532
74. Evgenov OV, Sveinbjörnsson B, Bjertnaes LJ (2001) Continuously infused methylene blue modulates the early cardiopulmonary response to endotoxin in awake sheep. *Acta Anaesthesiol Scand* 45(10):1246–1254
75. Evgenov OV, Sager G, Bjertnaes LJ (2001) Methylene blue reduces lung fluid filtration during the early phase of endotoxemia in awake sheep. *Crit Care Med* 29(2):374–379
76. Galili Y, Kluger Y, Mianski Z et al (1997) Methylene blue—a promising treatment modality in sepsis induced by bowel perforation. *Eur Surg Res* 29(5):390–395
77. Dumbarton TC, Minor S, Yeung CK et al (2011) Prolonged methylene blue infusion in refractory septic shock: a case report. *Can J Anaesth* 58(4):401–405
78. van Haren FM, Pickkers P, Foudraïne N et al. (2010) The effects of methylene blue infusion on gastric tonometry and intestinal fatty acid binding protein levels in septic shock patients. *J Crit Care* 25(2):358.e1–7
79. Brown G, Frankl D, Phang T (1996) Continuous infusion of methylene blue for septic shock. *Postgrad Med J* 72(852):612–614
80. Heemskerk S, van Haren FM, Foudraïne NA et al (2008) Short-term beneficial effects of methylene blue on kidney damage in septic shock patients. *Intensive Care Med* 34(2):350–354
81. Park BK, Shim TS, Lim CM et al (2005) The effects of methylene blue on hemodynamic parameters and cytokine levels in refractory septic shock. *Korean J Intern Med* 20(2):123–128
82. Donati A, Conti G, Loggi S et al (2002) Does methylene blue administration to septic shock patients affect vascular permeability and blood volume? *Crit Care Med* 30(10):2271–2277
83. Kirov MY, Evgenov OV, Evgenov NV et al (2001) Infusion of methylene blue in human septic shock: a pilot, randomized, controlled study. *Crit Care Med* 29(10):1860–1867
84. Memis D, Karamanlioglu B, Yuksel M et al (2002) The influence of methylene blue infusion on cytokine levels during severe sepsis. *Anaesth Intensive Care* 30(6):755–762
85. Juffermans NP, Vervloet MG, Daemen-Gubbels CR et al (2010) A dose-finding study of methylene blue to inhibit nitric oxide actions in the hemodynamics of human septic shock. *Nitric Oxide* 22(4):275–280
86. Weingartner R, Oliveira E, Oliveira ES et al (1999) Blockade of the action of nitric oxide in human septic shock increases systemic vascular resistance and has detrimental effects on pulmonary function after a short infusion of methylene blue. *Braz J Med Biol Res* 32(12):1505–1513
87. Gachot B, Bedos JP, Veber B et al (1995) Short-term effects of methylene blue on hemodynamics and gas exchange in humans with septic shock. *Intensive Care Med* 21(12):1027–1031
88. Goluboff N, Wheaton R (1961) Methylene blue-induced cyanosis and acute hemolytic anemia complicating the treatment of methemoglobinemia. *J Pediatr* 58:86–89
89. Héritier Barras AC, Walder B, Seeck M (2010) Serotonin syndrome following methylene blue infusion: a rare complication of antidepressant therapy. *J Neurol Neurosurg Psychiatry* 81(12):1412–1413
90. McDonnell AM, Rybak I, Wadleigh M et al. (2012) Suspected serotonin syndrome in a patient being treated with methylene blue for ifosfamide encephalopathy. *J Oncol Pharm Pract* 18(4):436–9
91. Boyer EW, Shannon M (2005) The serotonin syndrome. *N Engl J Med* 352(11):1112–1120
92. Metzger A, Harvey BH, Wegener G et al (2012) Azure B, a metabolite of methylene blue, is a high-potency, reversible inhibitor of monoamine oxidase. *Toxicol Appl Pharmacol* 258(3):403–409
93. Jang DH, Nelson LS, Hoffman RS (2011) Methylene blue in the treatment of refractory shock from an amlodipine overdose. *Ann Emerg Med* 58(6):565–567
94. Zhang X, Hintze TH (1998) Amlodipine releases nitric oxide from canine coronary microvessels, an unexpected mechanism of action of a calcium channel-blocking agent. *Circulation* 97:576–580
95. Zhang X, Loke KE, Mital S et al (2002) Paradoxical release of nitric oxide by an L-type calcium channel antagonist, the R+ enantiomer of amlodipine. *J Cardiovasc Pharmacol* 39:208–214
96. Lensai H et al (2003) Amlodipine activates the endothelial nitric oxide synthase by altering phosphorylation on Ser and Thr. *Cardiovasc Res* 59:844–853
97. Xu B, Xiao-hung L, Lin G et al (2002) Amlodipine, but not verapamil or nifedipine, dilates rabbit femoral artery largely through a nitric oxide- and kinin-dependent mechanism. *Br J Pharmacol* 136:375–382
98. Rodrigues JM, Pazin Filho A, Rodrigues AJ et al (2007) Methylene blue for clinical anaphylaxis treatment: a case report. *Sao Paulo Med J* 125:60–62
99. Weissgerber AJ (2008) Methylene blue for refractory hypotension: a case report. *AANA J* 76(4):271–274
100. Grayling M, Deakin CD (2003) Methylene blue during cardiopulmonary bypass to treat refractory hypotension in septic endocarditis. *J Thorac Cardiovasc Surg* 125(2):426–427