

Published in final edited form as:

Acta Paediatr. 2013 October ; 102(10): e439–e442. doi:10.1111/apa.12339.

Perinatal systemic inflammatory responses of growth-restricted preterm newborns

T. McElrath^{*}, E. N. Allred⁺, L. Van Marter[°], R. N. Fichorova^{*}, and A. Leviton⁺ for the ELGAN Study Investigators

^{*}Department of Obstetrics and Gynecology, Brigham & Women's Hospital, Harvard Medical School, Boston, MA, USA

⁺Neuroepidemiology Unit, Department of Neurology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

[°]Division of Newborn Medicine, Brigham & Women's Hospital, Harvard Medical School, Boston, MA, USA

Abstract

AIM—To compare the early postnatal pattern of systemic inflammation in growth-restricted infants born before the 28th week of gestation to that of appropriately grown peers.

METHODS—We measured the concentrations of 25 inflammation-related proteins in blood spots collected from 939 newborns during the first 2 postnatal weeks. We calculated the odds ratios (99% confidence intervals) that concentrations would be in the highest quartile.

RESULTS—Severely growth-restricted infants (birth weight Z-score < -2) were not at increased risk of systemic inflammation shortly after birth. On postnatal day 14, however, they were significantly more likely than their peers to have a CRP, IL-1, IL-6, TNF-, IL-8, MCP-4, ICAM-1, ICAM-3, E-SEL, MMP-9, VEGF-R2, and/or IGFBP-1 concentration in the *highest* quartile. These increased risks could not be attributed to delivery indication, bacteremia, or duration of ventilation.

CONCLUSION—Growth-restricted preterm newborns appear to be at increased risk of elevated concentrations of inflammation-associated proteins by postnatal day 14.

Keywords

growth-restricted; inflammation; IUGR; neonate; preterm

Introduction

Growth-restricted preterm newborns are at increased risk of systemic inflammation.(1) Some of this increased risk might be a consequence of their greater tendency to be exposed to inflammatory stimuli, including sepsis/bacteremia (2) and prolonged ventilation.(3) But what if some of the systemic inflammation that characterizes the growth-restricted, preterm newborn is beyond what might be expected in light of obvious inflammatory stimuli? What if the systemic inflammation is an early indicator of the “fetal programming” processes that lead to later-life increased risk of diseases such as obesity, type-2 diabetes, and

Corresponding author: Thomas McElrath, Department of OB/GYN, Brigham & Women's Hospital., 75 Francis St., Boston MA. 02115, USA, Tel: 001-617-732-5452, Fax : 001-617-232-6346, tmcelrath@partners.org.

None of the authors has any financial issue or conflict of interest to disclose

atherosclerotic cardiovascular disease?(4) While each of these diseases is believed to have a prominent systemic inflammatory component in the adult,(5) we are not aware of evidence that the growth-restricted fetus or newborn is at increased risk of systemic inflammation. If the adult consequences of fetal growth restriction are associated with increased systemic inflammation, but the associated intrauterine environment is not characterized by inflammation, then when does the post-natal inflammation first become manifest?

METHODS

Details about the ELGAN Study are presented elsewhere.(6, 7) Inflammation-related proteins were measured only in the blood of children who might have had the neurological dysfunctions that were the focus of the ELGAN Study. Thus, the sample for the analyses presented here consists of the 939 children who had a developmental assessment at age 2 years. Delivery indications and fetal growth parameters were equivalent in both the protein measured and unmeasured groups. Drops of blood from these newborns were collected on filter paper on postnatal day 1 (range: 1-3 days) (N = 861), 7 (range: 5-8 days) (N = 867), and 14 (range: 12-15 days) (N = 786). We evaluated 2 groups of babies whose birth weight was low for gestational age. We identify babies as growth-restricted infants if they had a birth weight Z-score < -2, and as growth-limited if they had a birth weight Z-score ≥ -2 and < -1. We classified infants by whether or not they were mechanically ventilated during every one of the first 14 postnatal days. Infants were classified as having bacteremia if an organism was recovered from blood obtained during the first 14 days.

Statistical Analyses

We evaluated the null hypothesis that growth-restricted and growth-limited infants were no more likely than their appropriately-grown peers (i.e., those who had a birth weight Z-score ≥ -1) to have early postnatal concentrations of 25 inflammation-associated proteins in the highest quartile for gestational age and day of blood specimen collection. Consistent with prior analysis, we present the data with a minimum of reduction.(8)

To assess the potential confounding that might be due to delivery indication, or inflammatory stimuli such as bacteremia and “prolonged” ventilation, we conducted additional analyses limited to four proteins in strata defined by potential confounders. Because bacteremia and prolonged ventilation did not appear to be confounders, we did not adjust for them in the multinomial models, but did adjust for gestational age (23-24, 25-26, 27 weeks). Not presented here are tables of results when we also adjusted for bacteremia and prolonged ventilation. The odds ratios in those analyses are minimally different from those presented.(9) We created multinomial logistic regression models comparing infants in each of two groups to infants whose birth weight Z-score ≥ -1 . This enabled us to calculate odds ratios (and 99% confidence intervals) using the top quartile concentration of each protein as our indicator of elevated concentration.

Results

Sample description (Table S1)

In this sample of 939 newborns, 50 had a birth weight for gestational age that placed them more than 2 standard deviations below the median of a referent sample (left data column), hereafter identified as growth restricted. An additional 122 had a birth weight Z-score that placed them between two and just less than one standard deviations below the median of the referent standard (i.e., ≥ -2 , < -1) (middle data column) hereafter identified as growth limited.

The majority of the growth-restricted infants were born during the 25th and 26th weeks of gestation. While only 6% of infants who were not growth-restricted were born to women who had severe preeclampsia, 39% of the growth-limited, and 58% of the growth-restricted had a preeclamptic mother. Both groups of small babies were more likely than their peers to have been ventilated during the first two weeks.

Odds ratios of elevated protein concentrations associated with fetal growth restriction in strata defined by ventilation status, bacteremia, and pregnancy disorder (Table S2)

Because infants who received mechanical ventilation for 14 days were at increased risk of systemic inflammation in this sample,(10) we explored the relationship between low birth weight Z-scores and inflammation in strata defined by whether the infant was mechanically-ventilated every day through day 14. For ease of presentation, we restricted these analyses to four proteins (*i.e.*, IL-1 , IL-6, TNF- , and IL-8). Regardless of ventilation, gestational age and bacteremia strata, growth-restricted and growth-limited infants were more likely than their peers to have elevated concentrations of the IL-1 , IL-6, TNF- , and IL-8 on postnatal day 14.

The inflammatory response among growth-restricted newborns is more prominent among those born after a spontaneous delivery than after delivery for severe preeclampsia. The inflammatory response is also seen among children whose birth weight for gestational age is not as low, but only if birth followed a spontaneous indication.

Odds ratios of elevated protein concentrations on days 1, 7 and 14 associated with fetal growth restriction (Table S3)

Both growth-restricted and growth-limited infants were at increased risk of elevated concentrations of 3 proteins in day-7 blood. On day 14, the growth-restricted were at increased risk of elevated concentrations of 11 proteins, while the growth-limited infants were at increased risk of elevated concentrations of 7 proteins.

Odds ratios of elevated protein concentrations on at least two days one week apart associated with restricted growth (Table 1)

Growth-restricted newborns were more likely than others to have prominently elevated odds ratios of highest-quartile concentrations of six proteins on two separate days. Elevated concentrations of three of these six proteins on only one day also characterized these growth-restricted newborns. Growth-limited newborns had elevated concentrations of three proteins on two days.

DISCUSSION

This study has three major findings. First, compared to appropriate for gestational age infants, growth-restricted infants tend to have higher circulating concentrations of inflammation-related proteins one to two weeks after birth, but not earlier. Second, the systemic inflammation on postnatal day 14 does not appear to be due to the propensity for growth-restricted newborns to be exposed to such inflammatory stimuli as bacteremia and “prolonged” ventilation. Third, the hyper-inflammation among the growth-restricted was only minimally greater than among the growth-limited.

Inflammation not present at birth

The placentas of infants delivered for maternal or fetal indications are much less likely to harbor bacteria than the placentas of infants delivered for spontaneous indications, and are also much less likely to have histologic inflammation.(11) From these observations, we infer that the intrauterine environment of infants delivered for spontaneous indications can be

characterized as inflammatory.(7) Indeed, growth-restricted infants, who are typically delivered for maternal or fetal indications, tend to have low circulating concentrations of inflammation-related proteins in blood collected during the first four postnatal days.(12-14)

The first strong inflammatory stimulus for growth-restricted infants occurs postnatally

Since the intense inflammatory response appears to begin one to two weeks after delivery, we infer that the stimulus for this response probably also begins after delivery. One plausible stimulus is assisted ventilation.(10, 15) Stratification of our sample by ventilation every day through day 14 (Table S2) and by culture-proven bacteremia documented that ventilation status and bacteremia accounted for some, but not all of the increased propensity of growth-restricted and growth-limited infants to have systemic inflammation two weeks after birth. These observations raise the possibility of other inflammatory stimuli.

Endogenous propensity to postnatal hyper-inflammation

Support for endogenous contributions to postnatal hyper-inflammation comes from a study of rat pups, which found that those who were growth-restricted at birth, tended to respond more vigorously than controls to an inflammatory stimulus.(16) Additional support comes from a study of human infants born before the 32nd week of gestation who were exposed to mechanical ventilation, which showed that those delivered because of severe early-onset preeclampsia (and therefore at prominently elevated risk of severe growth restriction) had more intense systemic inflammatory responses than their peers delivered for other indications (*i.e.*, preterm labor).(17) This latter report of a preeclampsia-postnatal inflammation relationship, and awareness that preeclampsia appears to be characterized by a spectrum of immunologic characteristics, including hyper-responsiveness to inflammatory stimuli,(18) prompted us to stratify our sample by whether delivery was for spontaneous indications or preeclampsia (Table S2). Growth-restricted and growth-limited infants of preeclamptic pregnancies did not show a greater propensity to inflammation than their peers born to women whose delivery was considered spontaneous. These findings support the view that the association between growth restriction and inflammation is not due solely to preeclampsia, or by inference, the aberrant placenta implantation associated with this disorder.(19)

Might hyper-inflammation account for the late putative consequences of growth restriction?

Adult diseases whose risk appears to be influenced by fetal growth restriction (*e.g.*, obesity, type-2 diabetes, and atherosclerotic cardiovascular disease) tend to have a prominent inflammatory component.(5, 20) In light of our finding that the earliest manifestations of hyper-inflammation in growth-restricted very preterm newborns are prominent by the end of the second postnatal week, we raise the possibility that this early hyper-inflammation is the first indicator of the processes that might account for the diseases that occur preferentially in adults who were severely growth-restricted at birth.

Limitations and strengths

The weaknesses of this study are those of all observational studies. We are unable to establish causation and report only associations. Our study has several strengths. First, we included a large number of infants, making it unlikely that we have missed important associations due to lack of statistical power, or claimed associations that might have reflected the instability of small numbers. Second, we selected infants based on gestational age, and not birth weight, in order to minimize confounding due to factors related to fetal growth restriction.(21)

Conclusion

Very preterm growth-restricted infants appear to be at increased risk of systemic inflammation by postnatal day 14.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by a cooperative agreement(5U01NS040069-05) with, and a grant (2R01NS040069 - 06A2) from the National Institute of Neurological Disorders and Stroke, and a center grant award from the National Institute of Child Health and Human Development (5P30HD018655-28). The authors gratefully acknowledge the contributions of their subjects, and their subjects' families, as well as those of their colleagues listed in the supplement.

References

1. Leviton A, O'Shea TM, Bednarek FJ, Allred EN, Fichorova RN, Dammann O. Systemic responses of preterm newborns with presumed or documented bacteraemia. *Acta Paediatr.* 2012; 101:355–9. [PubMed: 22085230]
2. Damodaram M, Story L, Kulinskaya E, Rutherford M, Kumar S. Early adverse perinatal complications in preterm growth-restricted fetuses. *Aust N Z J Obstet Gynaecol.* 2011; 51:204–9. [PubMed: 21631437]
3. Streimish IG, Ehrenkranz RA, Allred EN, O'Shea TM, Kuban KC, Paneth N, et al. Birth weight-and fetal weight-growth restriction: impact on neurodevelopment. *Early Hum Dev.* 2012; 88:765–71. [PubMed: 22732241]
4. Rinaudo P, Wang E. Fetal programming and metabolic syndrome. *Annu Rev Physiol.* 2012; 74:107–30. [PubMed: 21910625]
5. Shirwany NA, Zou MH. Vascular inflammation is a missing link for diabetes-enhanced atherosclerotic cardiovascular diseases. *Front Biosci.* 2012; 17:1140–64.
6. O'Shea TM, Allred EN, Dammann O, Hirtz D, Kuban KC, Paneth N, et al. The ELGAN study of the brain and related disorders in extremely low gestational age newborns. *Early Hum Dev.* 2009; 85:719–25. [PubMed: 19765918]
7. McElrath TF, Hecht JL, Dammann O, Boggess K, Onderdonk A, Markenson G, et al. Pregnancy disorders that lead to delivery before the 28th week of gestation: an epidemiologic approach to classification. *Am J Epidemiol.* 2008; 168:980–9. [PubMed: 18756014]
8. Leviton A, Allred EN, Yamamoto H, Fichorova RN. Relationships among the concentrations of 25 inflammation-associated proteins during the first postnatal weeks in the blood of infants born before the 28th week of gestation. *Cytokine.* 2012; 57:182–90. [PubMed: 22133344]
9. Levine RJ, Maynard SE, Qian C, Lim KH, England LJ, Yu KF, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med.* 2004; 350:672–83. [PubMed: 14764923]
10. Bose CL, Laughon MM, Allred EN, Michael O'Shea T, Van Marter LJ, Ehrenkranz RA, et al. Systemic inflammation associated with mechanical ventilation among extremely preterm infants. *Cytokine.* 2013; 61:315–22. [PubMed: 23148992]
11. Onderdonk AB, Hecht JL, McElrath TF, Delaney ML, Allred EN, Leviton A. Colonization of second-trimester placenta parenchyma. *Am J Obstet Gynecol.* 2008; 199:52 e1–e10. [PubMed: 18313635]
12. McElrath TF, Fichorova RN, Allred EN, Hecht JL, Ismail MA, Yuan H, et al. Blood protein profiles of infants born before 28 weeks differ by pregnancy complication. *Am J Obstet Gynecol.* 2011; 204:418 e1–e12. [PubMed: 21349490]
13. Briana DD, Boutsikou M, Baka S, Papadopoulos G, Gourgiotis D, Puchner KP, et al. Perinatal plasma monocyte chemoattractant protein-1 concentrations in intrauterine growth restriction. *Mediators Inflamm.* 2007; 2007:65032. [PubMed: 18274642]

14. Amarilyo G, Oren A, Mimouni FB, Ochshorn Y, Deutsch V, Mandel D. Increased cord serum inflammatory markers in small-for-gestational-age neonates. *J Perinatol*. 2011; 31:30–2. [PubMed: 20410909]
15. Hillman NH, Moss TJ, Kallapur SG, Bachurski C, Pillow JJ, Polglase GR, et al. Brief, large tidal volume ventilation initiates lung injury and a systemic response in fetal sheep. *Am J Respir Crit Care Med*. 2007; 176:575–81. [PubMed: 17641159]
16. Campbell LR, Pang Y, Ojeda NB, Zheng B, Rhodes PG, Alexander BT. Intracerebral lipopolysaccharide induces neuroinflammatory change and augmented brain injury in growth-restricted neonatal rats. *Pediatr Res*. 2012; 71:645–52. [PubMed: 22337231]
17. Turunen R, Andersson S, Laivuori H, Kajantie E, Siitonen S, Repo H, et al. Increased postnatal inflammation in mechanically ventilated preterm infants born to mothers with early-onset preeclampsia. *Neonatology*. 2011; 100:241–7. [PubMed: 21701213]
18. Redman CW, Sargent IL. Immunology of pre-eclampsia. *Am J Reprod Immunol*. 2010; 63:534–43. [PubMed: 20331588]
19. Redman CW, Sargent IL. Latest advances in understanding preeclampsia. *Science*. 2005; 308:1592–4. [PubMed: 15947178]
20. Kampoli AM, Tousoulis D, Briasoulis A, Latsios G, Papageorgiou N, Stefanadis C. Potential pathogenic inflammatory mechanisms of endothelial dysfunction induced by type 2 diabetes mellitus. *Curr Pharm Des*. 2011; 17:4147–58. [PubMed: 22204375]
21. Arnold CC, Kramer MS, Hobbs CA, McLean FH, Usher RH. Very low birth weight: A problematic cohort for epidemiologic studies of very small or immature neonates. *American J Epidemiology*. 1991; 134:604–13.

Abbreviations

SAA	Serum Amyloid A
MPO	Myeloperoxidase
IL -1	Interleukin-1
IL -6 -6R	Interleukin-6 & Receptor
TNF-	Tumor Necrosis Factor-
TNF-R1 –R2	Tumor Necrosis Factor Receptor-1 & -2
IL-8	Interleukin-8 (CXCL8)
MCP-1 - 4	Monocyte Chemotactic Protein-1 & -4 (CCL2, CCL13)
MIP-1	Macrophage Inflammatory Protein-1 (CCL4)
RANTES	Regulated upon Activation, Normal T-cell Expressed, and Secreted (CCL5)
I-TAC	Interferon-inducible T cell Alpha-Chemoattractant (CXCL11)
ICAM-1	Intercellular Adhesion Molecule -1 (CD54)
ICAM-3	Intercellular Adhesion Molecule -3 (CD50)
VCAM-1	Vascular Cell Adhesion Molecule-1 (CD106)
E-SEL	E-selectin (CD62E)
MMP-1 –9	Matrix Metalloproteinase-1 & -9
VEGF	Vascular Endothelial Growth Factor
VEGF-R1 – R2	Vascular Endothelial Growth Factor Receptor-1 & -2
IGFBP-1	Insulin-like Growth Factor Binding Protein-1

Key notes

Among infants born before 28 weeks, those that were severely growth restricted (Z-score <-2) were not at increased risk of systemic inflammation at birth but demonstrated increased concentrations of inflammation-associated proteins by postnatal day 14.

Table 1

Odds ratios (99% confidence intervals) of a protein concentration sustained in the top quartile on one versus two or more days for newborns in each birth weight Z-score category compared to newborns whose birth weight Z-score was -1 *

	Birth weight Z-score			
	< -2		-2, < -1	
	1 day only	2 days	1 day only	2 days
CRP	1.9 (0.7, 5.1)	3.8 (1.4, 9.9)	1.5 (0.8, 2.7)	2.0 (1.1, 3.9)
SAA	2.1 (0.9, 5.0)	1.7 (0.5, 5.4)	1.2 (0.7, 2.2)	1.9 (0.97, 3.7)
MPO	1.4 (0.6, 3.1)	0.3 (0.1, 1.7)	0.9 (0.5, 1.6)	0.8 (0.4, 1.7)
IL-1	2.1 (0.9, 4.9)	1.5 (0.5, 4.9)	1.1 (0.6, 2.0)	1.3 (0.6, 2.6)
IL-6	1.4 (0.6, 3.5)	2.9 (1.1, 7.8)	1.0 (0.5, 1.8)	1.8 (0.9, 3.6)
IL-6R	1.3 (0.5, 3.0)	0.7 (0.2, 2.3)	1.2 (0.6, 2.1)	1.1 (0.6, 2.2)
TNF-	3.1 (1.2, 8.5)	4.9 (1.7, 14)	1.2 (0.6, 2.1)	1.3 (0.7, 2.7)
TNF-R1	1.8 (0.8, 4.4)	1.7 (0.6, 4.7)	1.8 (0.99, 3.1)	1.5 (0.7, 3.1)
TNF-R2	1.5 (0.6, 3.4)	1.0 (0.3, 3.2)	0.8 (0.4, 1.4)	1.2 (0.6, 2.3)
IL-8	2.7 (1.03, 7.2)	4.7 (1.6, 13)	1.7 (0.96, 3.1)	1.8 (0.9, 3.7)
MCP-1	1.4 (0.6, 3.6)	2.4 (0.9, 6.3)	1.1 (0.6, 2.0)	2.1 (1.1, 4.0)
MCP-4	1.5 (0.6, 3.8)	1.6 (0.6, 4.2)	1.4 (0.7, 2.5)	1.6 (0.9, 3.1)
MIP-1	0.9 (0.3, 2.3)	0.9 (0.3, 2.6)	1.3 (0.7, 2.3)	0.8 (0.4, 1.6)
RANTES	0.7 (0.3, 1.7)	0.3 (0.1, 1.3)	0.8 (0.4, 1.4)	0.3 (0.1, 0.8)
I-TAC	1.8 (0.8, 4.3)	1.5 (0.5, 4.5)	1.1 (0.6, 2.1)	1.3 (0.7, 2.6)
ICAM-1	1.2 (0.5, 3.2)	2.0 (0.8, 5.1)	1.0 (0.6, 2.0)	1.8 (0.96, 3.4)
ICAM-3	1.5 (0.6, 3.6)	0.7 (0.2, 2.3)	0.9 (0.5, 1.6)	0.5 (0.3, 1.1)
VCAM-1	0.9 (0.3, 2.2)	0.6 (0.2, 1.9)	1.2 (0.6, 2.1)	0.7 (0.3, 1.4)
E-SEL	1.7 (0.7, 4.4)	2.5 (0.9, 6.5)	1.5 (0.8, 2.7)	1.2 (0.6, 2.4)
MMP-1	0.7 (0.2, 2.0)	0.7 (0.2, 2.0)	1.1 (0.6, 2.1)	0.5 (0.2, 1.1)
MMP-9	1.0 (0.4, 2.3)	0.4 (0.1, 1.8)	0.8 (0.4, 1.4)	0.4 (0.1, 0.9)
VEGF	1.2 (0.5, 2.9)	0.4 (0.1, 1.5)	1.0 (0.5, 1.8)	0.5 (0.3, 1.1)
VEGF-R1	2.4 (0.9, 6.5)	4.7 (1.7, 13)	1.8 (0.98, 3.1)	1.8 (0.9, 3.5)
VEGF-R2	1.0 (0.4, 2.5)	1.3 (0.5, 3.3)	1.1 (0.6, 2.0)	1.1 (0.5, 2.1)
IGFBP-1	4.2 (1.3, 13)	14 (4.2, 45)	2.0 (1.1, 3.6)	3.4 (1.7, 7.0)

* Adjustment was made for gestational age (23-24, 25-26, 27 weeks).

Bold items are significantly elevated at $p < 0.01$.