



Published in final edited form as:

*Comput Human Behav.* 2014 January ; 30: 13–22. doi:10.1016/j.chb.2013.07.024.

## Giving and Receiving Emotional Support Online: Communication Competence as a Moderator of Psychosocial Benefits for Women with Breast Cancer

Woohyun Yoo<sup>a,\*</sup>, Kang Namkoong<sup>b</sup>, Mina Choi<sup>c</sup>, Dhavan V. Shah<sup>a,d</sup>, Stephanie Tsang<sup>a</sup>,  
Yangsun Hong<sup>a</sup>, Michael Aguilar<sup>a</sup>, and David H. Gustafson<sup>d</sup>

<sup>a</sup>School of Journalism and Mass Communication, University of Wisconsin-Madison, 5115 Vilas Hall, 821 University Avenue, Madison, WI 53706, USA

<sup>b</sup>Department of Community and Leadership Development, University of Kentucky, 500 Garrigus Building, Lexington, KY 40546-0215, USA

<sup>c</sup>Department of Communication Arts, University of Wisconsin-Madison, 6164 Vilas Hall, 821 University Avenue, Madison, WI 53706, USA

<sup>d</sup>Center for Health Enhancement Systems Studies, University of Wisconsin-Madison, 1513 University Avenue, Rm 4120, Madison, WI 53706, USA

### Abstract

This study examines the moderating role of emotional communication competence in the relationship between computer-mediated social support (CMSS) group participation, specifically giving and receiving emotional support, and psychological health outcomes. Data were collected as part of randomized clinical trials for women diagnosed with breast cancer within the last 2 months. Expression and reception of emotional support was assessed by tracking and coding the 18,064 messages that 236 patients posted and read in CMSS groups. The final data used in the analysis was created by merging (a) computer-aided content analysis of discussion posts, (b) action log data analysis of system usage, and (c) baseline and six-month surveys collected to assess change. Results of this study demonstrate that emotional communication competence moderates the effects of expression and reception of emotional support on psychological quality of life and breast cancer-related concerns in both desired and undesired ways. Giving and receiving emotional support in CMSS groups has positive effects on emotional well-being for breast cancer patients with higher emotional communication, while the same exchanges have detrimental impacts on emotional well-being for those with lower emotional communication competence. The theoretical and practical implications for future research are discussed.

### Keywords

Computer-mediated social support groups; Emotional support; Emotional communication competence; Breast cancer

---

© 2013 Elsevier Ltd. All rights reserved.

\*Corresponding author at: 5115 Vilas Communication Hall, 821 University Avenue, Madison, WI 53706, USA, Tel.: +1-608-630-0264, wyoo3@wisc.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## 1. Introduction

Aside from skin cancer, breast cancer is the most frequently diagnosed cancer among American women and approximately 12% of women in the United States develop invasive breast cancer during their lifetime (Ma & Jemal, 2013). As a result of their cancer diagnosis and subsequent treatments, women with breast cancer often face major psychosocial and emotional challenges (Carlsson & Hamrin, 1994; Ganz, 2008; Shapiro et al., 2001). To cope with such stresses, breast cancer patients exchange social support with peer patients in Computer-Mediated Social Support (CMSS) groups (Klemm et al., 2003; Sharf, 1997; Shaw, McTavish, Hawkins, Gustafson, & Pingree, 2000). Accordingly, the participants benefit from the social support of the groups. A growing body of research has found that breast cancer patients' participation in CMSS groups is beneficial to their emotional well-being (Lieberman et al., 2003; Lieberman & Goldstein, 2005; Winzelberg et al., 2003), psychosocial well-being (Gustafson et al., 2001; Rodgers & Chen, 2005), and healthcare competence (van Uden-Kraan et al., 2008).

From the perspective that people exchange social support via text-based messages in virtual communities (Walther & Boyd, 2002), recent research has focused to identify the health benefits of expression and reception of social support in CMSS groups for women with breast cancer (Kim et al., 2012; Namkoong et al., 2010). These studies have found that giving and getting supportive messages benefits women with breast cancer, with the effect of expression often outpacing reception (Pingree, 2007; Han et al., 2011; Winefield, 2006). Other studies have found the individual differences, including demographic characteristics, digital competence, and coping strategies predict giving and receiving emotional support (Kim et al., 2011; Yoo et al., 2013). Yet, despite the importance of individual characteristics in exploring the expression and reception of social support in CMSS groups, relatively little attention has been paid to whether the benefits of the support expression and reception are conditioned by personal characteristics of the communicators. Particularly noteworthy in this context is communication competence, individuals' abilities to send, receive, and interpret messages regarding social support processes (Goffman, 1972; Wentowski, 1981). In other words, communication competence reflects the ability to render and elicit social support (Heller, 1979; Sarason, Sarason, Hacker, & Basham, 1985; Wentowski, 1981). Accordingly, communication competence may affect the health benefits of emotionally supportive expression and reception in breast cancer CMSS groups.

To better understand how the effects of exchanging computer-mediated social support differ according to communication competence, we examine the moderating role of emotional communication competence in the relation between the expression and reception of emotional support and two measures of emotional well-being: (a) psychological quality of life and (b) breast cancer-related concerns. Tracking and coding the messages sent and received by 237 women with breast cancer in CMSS groups was used to assess expression and reception of emotional support. The data used in the analysis was created by merging (a) computer-aided content analysis of discussion posts within online support groups, (b) action log data analysis of use of this system, and (c) baseline and six-month surveys collected to assess change.

## 2. Emotional support for cancer patients

Emotional support is a critical form of social support, facilitating both coping strategies and contributing to sustained well-being. In general, emotional support has been described as messages or actions assuring an individual that he or she is cared for, loved, esteemed, and valued (Cobb, 1976). It centers on the provision of "comfort and security during times of

stress that leads the person to feel he or she is cared for by others” (Cutrona & Russell, 1990, p. 322).

Emotional support can be conceptualized as having four distinct functions. First, emotional support facilitates empathic and sympathetic responding, including the cognitive understanding for the suffering of others (Cutrona & Suhr, 1994; Dakof & Taylor, 1990). Second, emotional support provides encouragement and reassurance, bolstering the recipient’s hope and confidence (Burlinson, 1994; Rook & Underwood, 2000). Third, emotional support offers caring and physical affection, including hugs, kisses, hand-holding, and shoulder patting (Albrecht & Adelman, 1987; Cutrona & Russell, 1990; House, 1981). Finally, emotional support includes universality to emphasize the importance of closeness, bonding, and community (Braithwaite, Waldron, & Finn, 1999; Coursaris & Liu, 2009; Shaw et al., 2000; Yalom, 1970). Most conceptions center on individuals’ efforts to assist one another cope with affective distress.

Emotional support plays an important role in cancer patients’ psychological and physical adjustment to their disease. When confronted with a traumatic event, such as cancer, most individuals have high needs of emotional support to improve their self-esteem or to reduce feelings of personal inadequacy. Indeed, cancer patients identify emotional support as the most helpful kind of support (Dakof & Taylor, 1990; Dunkel-Schetter, 1984; Neuling & Winefield, 1988). Emotional support is linked with greater psychological well-being (Bloom, 1986; Roberts, Cox, Shannon, & Wells, 1994; Slevin et al., 1996; Taylor, Falke, Shoptaw, & Lichtman, 1986), better quality of life (Courten, Stevens, Crebolder, & Philipsen, 1996), and lower rates of cancer morbidity and mortality (Glanz & Lerman, 1992; Reynolds & Kaplan, 1990).

Emotional support has proven particularly beneficial for women with breast cancer. Many studies on breast cancer have shown that emotional support contributes to improve patients’ mental well-being (Alferi, Carver, Antoni, Weiss, & Duran, 2001; Bloom & Spiegel, 1984; Bloom, Stewart, Johnson, Banks, & Fobair, 2001), health-related quality of life and self-efficacy (Arora, Finney Rutten, Gustafson, Moser, & Hawkins, 2007), coping with breast cancer (Northouse, 1988), and survival rates of breast cancer (Ell, Nishimoto, Mediansky, Mantell, & Hamovitch, 1992; Kroenke, Kubzansky, Schernhammer, Holmes, & Kawachi, 2006).

### **3. Computer-mediated social support (CMSS) groups and emotional support**

According to the cancer stigma hypothesis (Cobb, 1976), patients with cancer may feel marked and branded by their illness. The stigma of breast cancer, in particular, with its focus on issues of sexuality and womanhood, leads some patients to locate themselves outside what they see as acceptable criterion. For women with breast cancer, CMSS groups provide anonymous and asynchronous communication that allows participants to be more open and honest about potentially embarrassing topics and stigmatizing concerns without the limits of time, space, and social cues (White & Dorman, 2001; K. B. Wright, 2002). In this regard, CMSS groups have been attractive to breast cancer patients who have feelings of social isolation or unmet social needs. Women with breast cancer cope with varied stressors by exchanging social support with other patients in CMSS groups (Sharf, 1997; van Uden-Kraan et al., 2008; Weinberg, Schmale, Uken, & Wessel, 1996). Among all types of social support, emotional support is exchanged most frequently in CMSS groups for breast cancer patients (Rodgers & Chen, 2005; Winzelberg et al., 2003). As the need for emotional support increases, patients’ family and friends might not be able to offer appropriate support because the situation also causes extensive illness-related concerns among these caregivers

and supporters (Northouse et al., 2002). In response, CMSS groups can serve as an alternative to traditional support channels, with people facing similar challenges sharing practical information with each other and exchanging supportive messages with other participants (McTavish, Pingree, Hawkins, & Gustafson, 2003; Shaw et al., 2000).

#### 4. Health benefits of CMSS groups: Expression and reception of emotional support

Most research on emotionally supportive exchanges investigates the benefits from the recipient's perspective. From this perspective, emotional support has been found to influence the feelings, coping strategies, personal relationships, and even physical health of the recipient (Albrecht & Goldsmith, 2003; Uchino, 2004; Wills & Fegan, 2001). Research has found that emotional support decreases the recipient's emotional distress and results in other desirable health outcomes (Burleson, 2003; Burleson & Goldsmith, 1988; Dunkel-Schetter, Blasband, Feinstein, & Herbert, 1992; Goldsmith, 1994). Similarly, the social and health benefits of emotional support can be achieved through the reception of relevant messages in online cancer support groups (K. B. Wright, 2002; K. B. Wright & Bell, 2003). For example, breast cancer patients consuming more emotional support messages in CMSS groups are likely to report fewer breast cancer-related concerns (Kim et al., 2012; Lieberman & Goldstein, 2005).

Countering the dominant message-reception paradigm, communication scholars have begun to advance a theoretical model of message-expression effects (Pingree, 2007; Shah, Cho, Eveland, & Kwak, 2005; Shah et al., 2007). From the expression-effects perspective, "the act of expression might change the message sender, which expressed ideas often do not exist intact, if at all, in the speaker's mind prior to expression" (Pingree, 2007, p. 439). For this reason, writing therapy has been widely applied to text-based online counseling (J. Wright, 2002). For instance, written expression of specific messages has been shown to improve patients' mental benefits in online health interventions (Han et al., 2008; Lieberman & Goldsmith, 2006; Shaw et al., 2007; Shaw, Hawkins, McTavish, Pingree, & Gustafson, 2006). That is, translating individual concerns into the written word helps patients understand and cope with trauma, which consequently decreases their psychological distress (Pennebaker, 1997). Additionally, expressive writing facilitates social integration by helping patients to connect with those around them in a positive way (Pennebaker & Graybeal, 2001). Among cancer patients, expressive writing interventions have been found to be effective at improving emotional well-being (Gellaitry, Peters, Bloomfield, & Home, 2010). Participants write about cancer-related experiences or coping strategies and share them with others. In the same vein, recent research found the psychological benefits of expressing emotional support in online cancer communities (Han et al., 2011; Kim et al., 2012), revealing that it often outpaces the effects of message reception.

Given that both giving and receiving emotional support online can evoke the psychological benefits of emotional support, such exchanges may contribute to improving emotional health outcomes. Accordingly, this study proposes the following hypotheses:

- H1.** Expression (H1a) and reception (H1b) of emotional support will be positively associated with psychological quality of life.
- H2.** Expression (H2a) and reception (H2b) of emotional support will be negatively associated with breast cancer-related concerns.

## 5. Emotional communication competence as a moderator

Individuals' ability to exchange and interpret messages is critical to healthy functioning and interpersonal interactions (Goffman, 1972; Wentowski, 1981). Such communication competence conditions individuals' ability to render and elicit social support (Heller, 1979; Sarason et al., 1985; Wentowski, 1981). In social support processes, communication competence is defined as the perceived tendency to establish meaningful relationships with others, render support, be relaxed, appreciate others' plights, and take turns appropriately (Query, Parry, & Flint, 1992). Therefore, communication competence plays an integral role in the exchange of social support (Albrecht & Adelman, 1987; Sarason, Sarason, Hacker, & Basham, 1985).

Because both face-to-face communication and computer-mediated communication (CMC) involve the use of relevant communication skills, communication competence can be readily translated into CMSS contexts. Spitzberg (2006) proposed a model of CMC competence that is determined by the motivation, knowledge, and skills of the interaction in relation to the demands of the context and desired outcomes of the discourse. According to the model, competent users of CMC possess four written communication skills: attentiveness, composure, coordination, and expressiveness. These communication skills can be applied to the selection of channels and messages for social support in CMC settings. For example, attentiveness leads to the display of social support and comforting sophistication of message content in CMC. In addition, individuals with expressiveness skills can exchange social support through the use of emoticons and similar paralinguistic features of message content, the use of humor, and even the depth and breadth of self-disclosure. In this sense, it seems reasonable to suggest that communication competence influences the social support processes through CMC.

Communication competence is a multidimensional concept that includes the traditional notion of media literacy as well as interpersonal communication skills (Lee, Shah, & McLeod, 2012). Of the several types of communication competence, this study focuses on the notion of communication competence in relation to emotional communications, reflecting the specific context of exchanging socially supportive messages among breast cancer patients. Riggio and Carney's (2003) communication/social skills model highlights three sets of affective capabilities: skills in encoding emotional messages, skills in decoding emotional messages, and skills in regulating emotions. Accordingly, we define emotional communication competence as one's self-assessment of effectively sending, receiving, and regulating emotional messages.

Our conceptualization of emotional communication competence aligns well with a comprehensive body of research on emotional competence or intelligence. Studies of emotional competence or intelligence often view communicative skill as a core capability, including recognizing others' emotions, knowing the cultural rules for displaying emotions, and using the proper emotional vocabulary in expression (Saarni, 1990; Salovey, Hsee, & Mayer, 1993; Salovey & Mayer, 1990). For example, Mayer and Salovey's (1997) model of emotional intelligence deals with the ability to express emotions as one of three major components along with the abilities to perceive, understand, and manage emotions.

Given this conception of competence as centered on effectively communicating and regulating emotions, the health benefits of giving and receiving emotional support may be contingent on an individual's level of emotional communication competence. Individuals with high levels of emotional communication competence should be able to give and receive emotional support more effectively than those with low levels of emotional communication competence (Bodie & Burleson, 2008). The communicatively competent should not only be

more attentive to emotional cues of communication, but also should be more proficient in (a) sending and receiving emotional support, (b) initiating and maintaining conversations, (c) expressing feelings clearly, and (d) demonstrating to others that they have been heard. On the other hand, those low in this competency are likely insufficient in communication skills that play an integral role in both the provision and reception of emotional support, a deficiency that inhibits their participation in emotionally supportive conversations. Thus, emotional communication competence is hypothesized to condition the emotional benefits of giving and receiving emotional support. Accordingly, the following hypotheses are offered:

**H3.** The positive relationship of expression (H3a) and reception (H3b) of emotional support with psychological quality of life will be greater for those with higher emotional communication competence.

**H4.** The negative relationship of expression (H3a) and reception (H3b) of emotional support with breast cancer-related concerns will be greater for those with higher emotional communication competence.

## 6. Methods

### 6.1. CMSS groups within CHES

The Comprehensive Health Enhancement Support System (CHES) is an online information and social networking system combining vetted medical information with a variety of other web-based resources. The system was developed based on input from health care professionals, patients, family caregivers, web designers, and communication researchers (Hawkins et al., 2010). The CHES system for breast cancer provides three services: (a) information services, (b) support services, and (c) coaching services. The information services offer vetted facts about breast cancer, its treatment and consequences, the medical system, and general life issues. The support services provide an opportunity to communicate with other patients as well as health care professionals. The coaching services enable the patients to cope with their illness more actively by providing tailored responses, guidelines, and simulations.

As the main support service, CMSS groups allow breast cancer patients to anonymously share informational and emotional support with other breast cancer patients. These text-based, asynchronous discussion boards that are accessible 24 hours a day, 7 days a week, and have consistently been the most widely used CHES service (Gustafson et al., 2005; McTavish et al., 2003). A trained facilitator moderates discussions to ensure that discussions are supportive and do not contain inaccurate or harmful information. The data used in this study were collected as part of a larger randomized clinical trial to examine different component parts of the CHES “Living with Breast Cancer” intervention (Gustafson et al., 2001).

### 6.2. Participants

Recruitment for the clinical intervention occurred from April 1, 2005, through May 31, 2007, at the University of Wisconsin Hospital, Hartford Hospital (Connecticut), and MD Anderson (Texas). Patients were recruited via brochures placed at relevant locations and via doctors or nurses during clinic visits. All clinical staff introduced the study to potentially eligible patients and asked if they would be interested in participating. Once patients agreed to participate, a research staff member explained the nature and procedure of the interventions. All participants were offered laptop computers with a CHES browser, which collected user data automatically, and Internet access during the study period.

In total, 661 women with breast cancer were randomized to one of the six treatment conditions. Each condition was provided different numbers and types of support services. The first condition (*Internet Only*,  $n = 112$ ) is the control group that received training on how to navigate and search for relevant information in the Internet. The second condition (*CHES Information Services*,  $n = 118$ ) received only the CHES information services. The third condition (*CHES Information and Support Services*,  $n = 109$ ) received both the information and support services from CHES. The fourth condition (*Full CHES: CHES Information + Support + Coaching Services*,  $n = 111$ ) received all three types of CHES services. The fifth condition (*Human Cancer Mentor Only*,  $n = 106$ ) received customized, confidential services from only cancer experts without the use of any CHES services. The final condition (*Full CHES + Human Cancer Mentor*,  $n = 105$ ) was offered both human cancer mentoring and all CHES services. Among these six conditions, this study focused on the 325 participants of the three conditions — *CHES Information + Support Services*, *Full CHES*, and *Full CHES + Human Cancer Mentor* — in which they could access the CMSS support groups within CHES. Of the 325 participants, we limited our analysis to the 236 women who either wrote or read at least one message via online discussion boards in the groups within the six-month study period.

### 6.3. Data construction

The data used in the current study resulted from a combination of three major data components: (a) computer-aided content analysis of discussion posts within the CMSS groups, (b) action log data analysis of system use, and (c) baseline and six-month surveys collected to assess change. We first analyzed the expression of emotional support within each posted message with the CMSS groups. In this process, we employed a computer-aided content analysis program, InfoTrend (Fan, 1985), to create a coding scheme that had high levels of reliability and validity (Han et al., 2012; Namkoong et al., 2011). We analyzed 18,064 posted messages using this program. Each discrete message post was the unit of analysis.

As mentioned earlier, emotional support was composed of several constructs, such as understanding, empathy, concern, sympathy, reassurance, encouragement, physical affection, and universality. To develop coding rules that accurately reflect the range of emotional support categories, we clustered our coding around four core sub-concepts: (a) empathy/sympathy, (b) encouragement/reassurance, (c) care/physical affection, and (d) universality/interrelationship. Our operationalization of emotional support included most of the sub-concepts identified in previous literature, making it applicable across a wide range of cases and contexts.

After defining these sub-concepts, we created dictionaries of keywords associated with each category. For example, words such as “sorry” served as a starting point for the empathy/sympathy category and “hope” served as a starting point for the encouragement category (see Table 1). Using the keyword dictionaries, coding rules were created by establishing a relationship between multiple terms, phrases, or concepts. Distinct from word-counting content analysis software, the InfoTrend system enabled us to delineate the distance between terms as well as their directionality.

By coding concepts based on the order in which the words appear, we were able to establish the meaning of concepts more accurately. The distance and direction could be set by indicating the space in characters and putting “A (Ahead),” “B (Behind),” or “E (Either Direction)” between the keywords. This function allowed us to capture emotional support as it occurs in natural language without coding different concepts that used the same term. For example, we coded “I am worried about how you are doing,” by making the coding rule, “*Worry A 20 You = Express Concern (Empathy/Sympathy)*,” which means InfoTrend

would code this only if a “Worry” appeared 20 characters ahead of the word “You” (see Fig. 1). This means phrases like “Don’t worry, I know how to get there,” would not be coded as Empathy/Sympathy. In addition, once rules were created, they could be layered with other rules. This way, we could capture complex expression of emotional support that could not be accurately coded using other software.

After setting final coding rules, a reliability test was conducted on a random subset of 200 discussion posts. The reliability test between human and computer coding showed an estimate of 91.0% agreement across the four different categories (empathy/sympathy: 90.9%, encouragement/reassurance: 88.4%, care/physical affection: 92.0%, and universality/relationship: 95.0%). Scott’s pi (1955), which was calculated by comparing chance agreement across the four coding categories with actual agreement, was found to be 86.2% greater than by chance.

As the next step, we integrated these codings with action log data of system use that tracked each “click and keystroke” made by all CHESS participants. MySQL queries were used to match each coded message with the information about who posted the message and who consumed the message. This enabled us to track which CHESS participant expressed and/or received specific messages. This data was then combined with survey data, which was collected at baseline and 6-month follow up surveys, to examine patients’ emotional health outcomes.

## 6.4. Measures

### 6.4.1. Dependent variables

**Psychological quality of life:** Psychological quality of life was assessed using six items in the World Health Organization Quality of Life (WHOQOL) instrument measured using a 5-point scale (0 = not at all to 4 = completely) (Harper & Power, 1998). Respondents were asked to answer the following statements: (a) “How much do you enjoy life?”, (b) “To what extent do you feel your life to be meaningful?”, (c) “How well are you able to concentrate?”, (d) “Are you able to accept your bodily appearance?”, (e) “How satisfied are you with yourself?”, and (f) “How often do you have negative feelings such as blue mood, despair, anxiety, and depression?” ( $M = 2.85$ ,  $SD = .59$ , Cronbach’s  $\alpha = .78$  at pretest;  $M = 2.91$ ,  $SD = .53$ , Cronbach’s  $\alpha = .81$  at posttest).

**Breast cancer-related concerns:** Breast cancer-related concerns were measured with nine items of the Functional Assessment of Cancer Therapy scale (Kay-Lambkin, Baker, Lewin, & Carr, 2009), assessing breast cancer patients’ emotional, physical, and body image concerns related to treatment for breast cancer and its side effects. These were measured using a 6-point scale (0 = *not at all* to 5 = *very much*) on which participants indicated how much they agreed with the following statements: (a) “I was short of breath”, (b) “I was self-conscious about the way I dress”, (c) “I was bothered by swollen or tender arms”, (d) “I felt sexually attractive (reverse coded)”, (e) “My hair loss bothered me”, (f) “I worried about the risk of cancer in other family members”, (g) “I worried about the effect of stress on my illness”, (h) “My change in weight bothered me”, and (i) “I am able to feel like a woman (reverse coded)” ( $M = 1.11$ ,  $SD = .61$ , Cronbach’s  $\alpha = .64$  at pretest;  $M = 1.15$ ,  $SD = .67$ , Cronbach’s  $\alpha = .72$  at posttest). This scale did not have high internal consistency because distinct breast cancer-related concerns can occur independently of each other. Nonetheless, it is still well suited to assess multi-dimensional concerns among women with breast cancer (Baker et al., 2011; Hawkins et al., 2010).



#### 6.4.2. Independent variables

**Expression and reception of emotional support:** To measure the expression and reception of emotional support as independent variables, we used a measure of proportion consistent with previous research (Kim et al., 2012; Kim et al., 2011; Yoo et al., 2013). We constructed these measures by dividing the total count of emotional support expressed or received by the total number of messages written or read, respectively. This approach makes more sense than simply using raw scores because there is significant variance among participants in the total volume of messages written and read. This ratio variable can also rule out the potential confounding effect of verbosity and the writing/reading of other types of message content (Han et al., 2011). Thus, the expression of emotional support was operationalized by the total count of emotionally supportive messages posted divided by the total number of messages posted at CMSS groups for the six-month study period ( $M = .40$ ,  $SD = .54$ ). Reception of emotional support messages was constructed by the total count of emotionally supportive messages read divided by the total number of messages read at CMSS groups for the six-month study period ( $M = 1.21$ ,  $SD = .76$ ). Table 2 shows the descriptive statistics for the relevant components of the ratio measurement.

#### 6.4.3. Moderator variable

**Emotional communication competence:** Emotional communication competence refers to an individual's belief on her abilities to exchange and interpret emotional messages effectively. It served as a moderator variable and was assessed at baseline using a 5-point scale (0 = *I cannot do this at all* to 4 = *I can do this extremely well*) by using four items: (a) "I can identify and label my feelings accurately for others", (b) "When I disagree with someone, I check out my understanding of their viewpoint before I share my own feelings", (c) "I state my feelings and needs clearly and directly", and (d) "I am able to let others know that I understand their feelings even when their feelings are different than my own" ( $M = 2.58$ ,  $SD = .70$ , Cronbach's  $\alpha = .81$ ).

**6.4.4. Control variables—**We included the following variables to control their potentially confounding effects: demographic and disease-related information contained age, education (1 = *did not complete junior/high school* to 8 = *graduate degree*), income (1 = *less than \$20,000* to 6 = *over than \$100,000*), the time interval between breast cancer diagnosis and intervention participation (in days), and surgery experience (1 = *yes* and 0 = *no*). As additional control variables, we used two experimental conditions (1 = *Full CHESS* or *Full CHESS + Mentor* and 0 = *CHESS Information + Support Services* as a reference group), total time spent in CHESS (in minutes), and pretest values of psychological quality of life and breast cancer-related concerns.

### 6.5. Analytic framework

In order to test emotional communication competence as a potential moderator of the effects of emotional support expression and reception on psychological quality of life and breast cancer-related concerns, we used two types of hierarchical ordinary least squares regression models. In each regression model, control variables were entered in the first block. The pretest value of emotional communication competence was entered as moderator in the second block, and the main effect of exchanging emotional support was examined by entering the expression and reception of emotional support in the third block. Finally, in the fourth block, two-way interaction terms between emotional communication competence and expression (or reception) of emotional support were entered into the equation, which multiplies the standardized values of two variables to reduce multicollinearity problems between the interaction terms and their components (Cohen & Cohen, 1983).

## 7. Results

Table 3 presents the demographic and disease-related characteristics of our sample. The participants had a mean age of 51.18 years. The sample also had a diverse educational background with about 65% having at least an associate or bachelor's degree. The median category for annual household income was the range \$60,001 to \$80,000. The participants had an average 75.12 days between breast cancer diagnosis and intervention participation and they used CHESS an average 723.27 minutes for the six-month intervention period.

H1 predicted that the exchange of emotional support would be positively related to psychological quality of life. Table 4 presents the results of the regression analysis that expressing (H1a) and receiving (H1b) emotional support did not have significant effects on psychological quality of life ( $\beta = -.09, .06, p = \text{n.s.}$ , for emotional support expression and reception, respectively). Thus, H1 was not supported.

H2 predicted that the exchange of emotional support would be negatively related to breast cancer-related concerns. As shown in Table 4, the expression (H2a) and reception (H2b) of emotional support was unrelated with breast cancer-related concerns ( $\beta = .001, -.11, p = \text{n.s.}$ , for emotional support expression and reception, respectively). Therefore, H2 was not supported.

H3 predicted emotional communication competence would moderate the relationship of exchanging emotional support on psychological quality of life. The present study found a significant interaction effect between emotional communication competence and the expression of emotional support on psychological quality of life ( $\beta = .17, p < .05$ ). More specifically, the expression of emotional support had a positive relationship with psychological quality of life for breast cancer patients who had higher levels of emotional communication competence, whereas it had a negative relationship on psychological quality of life for those who had lower levels of emotional communication competence (see Fig. 2). However, the interaction between emotional communication competence and the reception of emotional support on psychological quality of life was not significant ( $\beta = -.06, p = \text{n.s.}$ ). Thus, H3 received partial support.

H4 predicted emotional communication competence would moderate the relationship between the exchange of emotional support and breast cancer-related concerns. There was a significant interaction between the reception of emotional support and communication competence on breast cancer-related concerns ( $\beta = -.17, p < .05$ ). That is, the reception of emotional support reduced breast cancer-related concerns for women with higher levels of emotional communication competence, while it increased breast cancer-related concerns for those who had lower levels of emotional communication competence (see Fig. 3). However, emotional communication competence did not moderate the influence of expressing emotional support on breast cancer-related concerns ( $\beta = -.07, p = \text{n.s.}$ ). Thus, H4 received partial support.

## 8. Discussion

The primary focus of this study concerned the moderating role of emotional communication competence on the relationship between the exchange of emotional support and emotional health outcomes. Research on CMSS groups has found that the expression and reception of emotional support promotes participants' emotional well-being (Han et al., 2011; Kim et al., 2012). Prior research also suggests that people with high emotional communication competence are better able to send, receive, and interpret messages concerning emotional support than those with low emotional communication competence (Bodie & Burleson, 2008). Thus, we expected that the psychological benefits of expression and reception of

emotional support would differ depending on the levels of emotional communication competence.

Somewhat surprisingly, the findings reveal that the expression and reception of emotional support were not directly associated with emotional health outcomes. Instead, the impact of exchanging emotional support was moderated by emotional communication competence in both desired and undesired ways. High communication competence amplified the benefits of exchanging emotional support, while low communication competence produced attenuated the benefits of exchanging emotional support. These findings are consistent with a growing body of research indicating that the effects of supportive messages do not unfold in the same way for all individuals (Bodie & Burleson, 2008). Interestingly, the role of emotional support expression was only found to improve psychological quality of life, while emotional support reception was only effective in reducing breast cancer-related concerns.

Building on the past work distinguishing message expression and reception effects, this research suggests that these influences are, for emotional well-being at least, contingent on communication competence. Emotional communication competence is known to foster active engagement that is critical to building supportive relationships; therefore, those who display this capability are more likely to participate in purposeful and goal-directed behaviors. In addition, they are able to interpret another person's situation to render satisfactory social support. Since the expression of emotional support requires cognitive effort and active motivation to engage in CMSS groups, emotionally communication competent patients likely gain self-efficacy by providing emotional support. Given that the regulation of goal-achieving behaviors operates as an important contributor to emotional well-being (Bandura, 1986, 1997), those who have high levels of emotional communication competence appear especially likely to reap the full range of emotional benefits from helping others by providing emotional support. In contrast, patients with low levels of emotional communication skills can experience more uncomfortable and stressful events because poor communication competence is a vulnerability factor (Segrin, 2001). This deficit in emotional communication skills might generate distress, as they tend to express emotional support in CMSS groups.

While breast cancer patients' message expression within CMSS groups improves emotional well-being among those with higher communication competence, it does not influence the worry and concerns related to a given health problem. Instead, the reception of supportive messages appears to have a positive impact on reducing breast cancer-related concerns (Shaw et al., 2000; Kim et al., 2011), though in this study we find this effect is concentrated among those with higher communication competence. The reception of emotional support only helps those with this competency because these individuals can symbolically influence interaction to mobilize their support group or social network (Query & James, 1989). Emotional communication competence makes them feel less isolated and increases their understanding about the illness and treatment processes, thus decreasing breast cancer-related concerns. Scholars examining the Relational Health Communication Competence Model (Kreps, 1988) found that communication competence predicted increased satisfaction from receiving social support (Weathers, Query, & Kreps, 2010; K. B. Wright, Banas, Bessarabova, & Bernard, 2010; K. B. Wright et al., 2013). Additionally, the increased satisfaction of social support resulted in the reduction of perceived stress (K. B. Wright et al., 2010). From this perspective, patients with high emotional communication competence might show low levels of breast cancer-related concerns as they receive emotional support in CMSS groups. However, those with low emotional communication competence might report high levels of breast cancer-related concerns because they are likely to be dissatisfied with emotional support received in CMSS groups.

Although we analyzed secondary data, the data themselves were constructed by integrating computational and conventional social science methods. These insights about message expression and reception are only available because three different types of data were combined: (a) computer-aided content analysis of discussion posts within online support groups, (b) action log data analysis of use of this system, and (c) baseline and six-month surveys collected to assess change. This move toward computational methods – also called “big data” by some — relies on taking tens of thousands of discussion posts, millions of lines of action log data, and multiple waves of survey data to more carefully test theories of influence.

### 8.1. Limitations and directions for future research

Nonetheless, despite these virtues, this study is not without shortcomings. In particular, the measurement of emotional communication competence leaned toward measuring the competence of message expression rather message reception, so it might be less adept at testing the moderating effect of emotional communication competence for getting emotional support. Future research should use a measure of communication competence equally related to both expression and reception. Second, we adopted a measure of proportion for reception of emotional support, such that it was measured by the total counts of emotionally supportive messages read divided by total number of all types of messages opened. However, opening a message does not guarantee that a subject read through the whole message and completely understood its content. Furthermore, the participants of our study used a text-based, asynchronous message system to read emotionally supportive messages in CMSS groups. The environment of online communication enables them to read all the messages that are targeted to others as well as themselves. Hence, if a woman read emotionally supportive messages directed to others, she might be less likely to perceive the benefits of reading those messages. Future work should consider more specified measures of reception for emotional support.

### 8.2. Conclusions and implications

Despite these limitations, it is noteworthy that this study provides further evidence of the importance of moderating factors in investigating the effects of supportive messages in online cancer communities. Numerous studies have found substantial evidence on moderators such as demographic characteristics, personality traits, cognitive factors, and interactional contexts (see reviews by Bodie & Burlison, 2008). This study not only demonstrates the moderating role of emotional communication competence in linking the effects of exchanging emotional support on emotional health outcomes, but it also provides an explanation of how this moderating effect takes place differently according to the features of emotional outcomes. Communication competence moderates the effect of supportive expression on psychological quality of life, but does not affect the impact of emotional support reception. On the contrary, the effect of emotional support reception on breast cancer-related concerns differs by emotional communication competence, but the influence of emotional support expression is unaffected.

These findings have practical implications. E-health practitioners should have different care guidelines in terms of the level of patients' emotional communication competence during the intervention period. For patients with high emotional communication competence, health care professionals may consider developing strategies that can promote more active participation in supportive discussion within CMSS groups. For those lacking emotional communication competence, it appears important to avoid certain types of CMSS interactions or to support the development of emotional communication competence. Hence, it may be worth considering the development of a more integrated e-health system that not

only provides a space for the exchange of emotional support, but helps improve emotional communication competence.

## Acknowledgments

This research was supported by P50 CA095817-05 from the National Cancer Institute. Study sponsors had no involvement in this research.

## References

- Albrecht, TL.; Adelman, MB. Communicating social support: A theoretical perspective. In: Albrecht, TL.; Adelman, MB., editors. *Communicating social support*. Sage; Newbury Park, CA: 1987. p. 18-39.
- Albrecht, TL.; Goldsmith, DJ. Social support, social networks, and health. In: Thompson, TL.; Dorsey, AM.; Miller, KI.; Parrott, R., editors. *Handbook of health communication*. Erlbaum; Mahwah, NJ: 2003. p. 263-284.
- Alferi SM, Carver CS, Antoni MF, Weiss S, Duran RE. An exploratory study of social support, distress, and life disruption among low-income Hispanic women under treatment for early stage breast cancer. *Health Psychology*. 2001; 20:41–46. [PubMed: 11199065]
- Arora NK, Finney Rutten LJ, Gustafson DH, Moser R, Hawkins RP. Perceived helpfulness and impact of social support provided by family, friends, and health care providers to women newly diagnosed with breast cancer. *Psycho-Oncology*. 2007; 16(5):474–486. [PubMed: 16986172]
- Baker TB, Hawkins RP, Pingree S, Roberts LJ, McDowell HE, Shaw BR, Han JY. Optimizing eHealth breast cancer interventions: Which types of eHealth services are effective? *Translational Behavioral Medicine*. 2011; 1(1):134–145. [PubMed: 21709810]
- Bandura, A. *Social foundations of thought and action: A social cognitive theory*. Prentice-Hall; Englewood Cliffs, NJ: 1986.
- Bandura, A. *Self-efficacy: The exercise of control*. Freeman; New York, NY: 1997.
- Bloom, JR. Social support and adjustment to breast cancer. In: Andersen, BL., editor. *Women with cancer: Psychological perspectives*. Springer Verlag; New York, NY: 1986. p. 204-229.
- Bloom JR, Spiegel D. The relationship of two dimensions of social support to the psychological well-being and social functioning of women with advanced breast cancer. *Social Science & Medicine*. 1984; 19(8):831–837. [PubMed: 6505749]
- Bloom JR, Stewart SL, Johnson M, Banks P, Fobair P. Sources of support and the physical and mental well-being of young women with breast cancer. *Social Science & Medicine*. 2001; 53:1513–1524. [PubMed: 11710426]
- Bodie GD, Burleson BR. Explaining variations in the effects of supportive messages: A dual-process framework. *Communication Yearbook*. 2008; 32:355–398.
- Braithwaite DO, Waldron VR, Finn J. Communication of social support in computer-mediated groups for people with disabilities. *Health Communication*. 1999; 11(2):123–151. [PubMed: 16370973]
- Burleson, BR. Comforting communication: Significance, approaches, and effects. In: Burleson, BR.; Albrecht, TL.; Sarason, IG., editors. *Communication of social support: Messages, interactions, relationships, and community*. Sage; Thousand Oaks, CA: 1994. p. 3-28.
- Burleson, BR. Emotional support skills. In: Greene, JO.; Burleson, BR., editors. *Handbook of communication and social interaction skills*. Erlbaum; Mahwah, NJ: 2003. p. 551-594.
- Burleson, BR.; Goldsmith, DJ. How the comforting process works: Alleviating emotional distress through conversationally induced reappraisals. In: Anderson, PA.; Guerrero, LK., editors. *Handbook of communication and emotion: Research, theory, applications, and contexts*. Academic Press; San Diego, CA: 1988. p. 245-280.
- Carlsson M, Hamrin E. Psychological and psychosocial aspects of breast cancer and breast cancer treatment: A literature review. *Cancer Nursing*. 1994; 17(5):418–428. [PubMed: 7954390]
- Cobb S. Social support as a moderator of life stress. *Psychosomatic Medicine*. 1976; 38:300–314. [PubMed: 981490]

- Cohen, J.; Cohen, P. Applied multiple regression/correlation analysis for the behavioral sciences. 2nd ed.. Lawrence Erlbaum Associates; Hillsdale, NJ: 1983.
- Coursaris CK, Liu M. An analysis of social support exchanges in online HIV/AIDS self-help groups. *Computers in Human Behavior*. 2009; 25(4):911–918.
- Courtens AM, Stevens FCJ, Crebolder HFJM, Philipsen H. Longitudinal study on quality of life and social support in cancer patients. *Cancer Nursing*. 1996; 19(3):162–169. [PubMed: 8674024]
- Cutrona, CE.; Russell, DW. Social support: An interactional behaviors. In: Sarason, BR.; Sarason, IG.; Pierce, GR., editors. Types of social support and specific stress: Toward a theory of optimal matching. Wiley; New York, NY: 1990. p. 319-366.
- Cutrona, CE.; Suhr, JA. Social support communication in the context of marriage: An analysis of couples' supportive interactions. In: Burleson, BR.; Albrecht, TL.; Sarason, IG., editors. The communication of social support: Messages, interactions, relationships, and community. Sage; Newbury, CA: 1994. p. 113-135.
- Dakof GA, Taylor SR. Victim's perception of social support: What is helpful from whom? *Journal of Personality and Social Psychology*. 1990; 58:80–89. [PubMed: 2308075]
- Dunkel-Schetter C. Social support and cancer: Findings based on patient interviews and their implications. *Journal of Social Issues*. 1984; 40(4):77–98.
- Dunkel-Schetter, C.; Blasband, D.; Feinstein, L.; Herbert, T. Elements of supportive interactions: When are attempts to help effective?. In: Spacapan, S.; Oskamp, S., editors. Helping and being helped: Naturalistic studies. Sage; Newbury Park, CA: 1992. p. 83-114.
- Ell K, Nishimoto R, Mediansky L, Mantell J, Hamovitch M. Social relations, social support and survival among patients with cancer. *Journal of Psychosomatic Research*. 1992; 36(6):531–541. [PubMed: 1640391]
- Fan DP. Ideodynamics: The kinetics of the evolution of ideas. *Journal of Mathematical Sociology*. 1985; 11(1):1–23.
- Ganz PA. Quality of life across the continuum of breast cancer care. *The Breast Journal*. 2008; 6(5): 324–330. [PubMed: 11348391]
- Gellaity G, Peters K, Bloomfield D, Horne R. Narrowing the gap: The effects of an expressive writing intervention on perceptions of actual and ideal emotional support in women who have completed treatment for early stage breast cancer. *Psycho-Oncology*. 2010; 19:77–84. [PubMed: 19194996]
- Glanz K, Lerman C. Psychosocial impact of breast cancer: A critical review. *Annals of Behavioral Medicine*. 1992; 14:204–212.
- Goffman, E. Relations in public. Harper Colophon; New York, NY: 1972.
- Goldsmith, DJ. The role of facework in supportive communication. In: Burleson, BR.; Albrecht, TL.; Sarason, IG., editors. Communicating of social support: Messages, interactions, relationships, and community. Sage; Thousand Oaks, CA: 1994. p. 29-49.
- Gustafson DH, Hawkins RP, Pingree S, McTavish FM, Arora NK, Mendenhall J, Stewart J. Effect of computer support on younger women with breast cancer. *Journal of General Internal Medicine*. 2001; 16(7):435–445. [PubMed: 11520380]
- Gustafson DH, McTavish FM, Stengle W, Ballard D, Hawkins RP, Shaw BR, Chen WC. Use and impact of eHealth system by low-income women with breast cancer. *Journal of Health Communication*. 2005; 10(S1):195–218. [PubMed: 16377608]
- Han JY, Shah DV, Kim E, Namkoong K, Lee SY, Moon TJ, Gustafson DH. Empathic exchanges in online cancer support groups: distinguishing message expression and reception effects. *Health Communication*. 2011; 26(2):185–197. [PubMed: 21318917]
- Han JY, Shaw BR, Hawkins RP, Pingree S, McTavish FM, Gustafson DH. Expressing positive emotions within online support groups by women with breast cancer. *Journal of Health Psychology*. 2008; 13(8):1002–1007. [PubMed: 18987072]
- Harper A, Power M. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychological Medicine*. 1998; 28:551–558. [PubMed: 9626712]
- Hawkins RP, Pingree S, Shaw BR, Serlin RC, Swoboda C, Han JY, Salner A. Mediating processes of two communication interventions for breast cancer patients. *Patient Education and Counseling*. 2010; 81:S48–S53. [PubMed: 21081261]

- Heller, K. The effects of social support: Prevention and treatment implications. In: Goldstein, AP.; Kanfer, FH., editors. *Maximizing treatment gains: Transfer enhancement in psychotherapy*. Academic; New York, NY: 1979. p. 353-382.
- House, JS. *Work stress and social support*. Addison-Wesley; Reading, MA: 1981.
- Kay-Lambkin FJ, Baker AL, Lewin TJ, Carr VJ. Computer-based psychological treatment for comorbid depression and problematic alcohol and/or cannabis use: A randomized controlled trial of clinical efficacy. *Addiction*. 2009; 104(3):378-388. [PubMed: 19207345]
- Kim E, Han JY, Moon TJ, Shaw BR, Shah DV, McTavish FM, Gustafson DH. The process and effect of supportive message expression and reception in online breast cancer support groups. *Psycho-Oncology*. 2012; 21:531-540. [PubMed: 21416553]
- Kim E, Han JY, Shah DV, Shaw BR, McTavish FM, Gustafson DH, Fan DP. Predictors of supportive message expression and reception in an interactive cancer communication system. *Journal of Health Communication*. 2011; 16(10):1106-1121. [PubMed: 22070449]
- Klemm P, Bunnell D, Cullen M, Soneji R, Gibbons P, Holecek A. Online cancer support groups: A review of the research literature. *CIN: Computers, Informatics, Nursing*. 2003; 21(3):136-142.
- Kreps GL. Relational communication in health care. *Southern Speech Communication Journal*. 1988; 53:344-359.
- Kroenke CH, Kubzansky LD, Schernhammer ES, Holmes MD, Kawachi I. Social networks, social support, and survival after breast cancer diagnosis. *Journal of Clinical Oncology*. 2006; 24(7): 1105-1111. [PubMed: 16505430]
- Lee N-J, Shah DV, McLeod JM. Processes of political socialization: A *Communication* communication mediation approach to youth civic engagement. Research. 2012 Advance online publication.
- Lieberman MA, Golant M, Giese-Davis J, Winzlenberg A, Benjamin H, Humphreys K, Spiegel D. Electronic support groups for breast carcinoma. *Cancer*. 2003; 97(4):920-925. [PubMed: 12569591]
- Lieberman MA, Goldsmith BA. Not all negative emotions are equal: The role of emotional expression in online support groups for women with breast cancer. *Psycho-Oncology*. 2006; 15(2):160-168. [PubMed: 15880627]
- Lieberman MA, Goldstein BA. Self-help on-line: An outcome evaluation of breast cancer bulletin boards. *Journal of Health Psychology*. 2005; 10(6):855-862. [PubMed: 16176962]
- Ma J, Jemal A. Breast cancer statistics. *Breast Cancer Metastasis and Drug Resistance*. 2013:1-18.
- Mayer, JD.; Salovey, P. What is emotional intelligence. In: Salovey, P.; Sluyter, D., editors. *Emotional development and emotional intelligence: Implications for educators*. Basic Books; New York, NY: 1997. p. 3-31.
- McTavish FM, Pingree S, Hawkins RP, Gustafson DH. Cultural differences in use of an electronic discussion group. *Journal of Health Psychology*. 2003; 8(1):105-117. [PubMed: 22113904]
- Namkoong K, Shah DV, Han JY, Kim SC, Yoo W, Fan DP, Gustafson DH. Expression and reception of treatment information in breast cancer support groups: How health self-efficacy moderates effects on emotional well-being. *Patient Education and Counseling*. 2010; 81:S41-S47. [PubMed: 21044825]
- Neuling SJ, Winefield HR. Social support and recovery after surgery for breast cancer: Frequency and correlates of supportive behaviours by family, friends, and surgeon. *Social Science & Medicine*. 1988; 27(4):385-392. [PubMed: 3175721]
- Northouse LL. Social support in patients' and husbands' adjustment to breast cancer. *Nursing Research*. 1988; 37:91-95. [PubMed: 3347526]
- Northouse LL, Mood D, Kershaw T, Schafenacker A, Mellon S, Walker J, Decker V. Quality of Life of Women With Recurrent Breast Cancer and Their Family Members. *Journal of Clinical Oncology*. 2002; 20(19):4050-4064. [PubMed: 12351603]
- Pennebaker, JW. *Opening up: The healing power of expressing emotions*. Guilford Press; New York, NY: 1997.
- Pennebaker JW, Graybeal A. Patterns of natural language use: Disclosure, personality, and social integration. *Current Directions in Psychological Science*. 2001; 10(3):90-93.

- Pingree RJ. How messages affect their senders: A more general model of message effects and implications for deliberation. *Communication Theory*. 2007; 17(4):439–461.
- Query JL, James AC. The relationship between interpersonal communication competence and social support among elderly support groups in retirement communities. *Health Communication*. 1989; 1(3):165–184.
- Query JM, Parry D, Flint LJ. The relationship among social support, communication competence, and cognitive depression for nontraditional students. *Journal of Applied Communication*. 1992; 20:78–94.
- Reynolds P, Kaplan GA. Social connections and risk for cancer: Prospective evidence from the Alameda County study. *Behavioral Medicine*. 1990; 16:101–110. [PubMed: 2224168]
- Riggio, RE.; Carney, D. *Manual for the social skills inventory*. 2nd ed.. MindGarden; Redwood City, CA: 2003.
- Roberts C, Cox C, Shannon V, Wells N. A closer look at social support as a moderator of stress in breast cancer. *Health and Social Work*. 1994; 19:157–164. [PubMed: 7959397]
- Rodgers S, Chen Q. Internet community group participation: Psychosocial benefits for women with breast cancer. *Journal of Computer-Mediated Communication*. 2005; 10(4) <http://jcmc.indiana.edu/vol10/issue4/rodders.html>.
- Rook, KS.; Underwood, LG. Social support measurement and interventions: Comments and future directions. In: Cohen, S.; Underwood, LG.; Gottlieb, BH., editors. *Social support measurement and intervention: A guide for health and social scientists*. Oxford University Press; New York, NY: 2000. p. 311-334.
- Saarni, C. Emotional competence: How emotions and relationships become integrated. In: Thompson, RA., editor. *Nebraska Symposium on Motivation, 1988: Socioemotional development*. University of Nebraska Press; Lincoln, NE: 1990. p. 115-182.
- Salovey, P.; Hsee, C.; Mayer, JD. Emotional intelligence and the self-regulation of affect. In: Wegner, DM.; Pennebaker, JW., editors. *Handbook of mental control*. Prentice-Hall; Englewood Cliffs, NJ: 1993. p. 258-277.
- Salovey P, Mayer JD. Emotional intelligence. *Imagination, Cognition, & Personality*. 1990; 9:185–211.
- Sarason BR, Sarason IG, Hacker A, Basham RB. Concomitants of social support: Social skills, physical attractiveness, and gender. *Journal of Personality and Social Psychology*. 1985; 49:469–480.
- Segrin C. Social skills and negative life events: Testing the deficit stress generation hypothesis. *Current Psychology*. 2001; 20:19–35.
- Shah DV, Cho J, Eveland WP Jr, Kwak N. Information and expression in a digital age: Modeling Internet effects on civic participation. *Communication Research*. 2005; 32(5):531–565.
- Shah DV, Cho J, Nah S, Gotlieb MR, Hwang H, Lee N-J, McLeod DM. Campaign ads, online messaging, and participation: Extending the communication mediation model. *Journal of Communication*. 2007; 57(4):676–703.
- Shapiro SL, Lopez AM, Schwartz GE, Bootzin R, Figueredo AJ, Braden CJ, Kurker SF. Quality of life and breast cancer: Relationship to psychosocial variables. *Journal of Clinical Psychology*. 2001; 57(4):501–519. [PubMed: 11255204]
- Sharf BF. Communicating breast cancer on-line: Support and empowerment on the Internet. *Women & Health*. 1997; 26(1):65–84. [PubMed: 9311100]
- Shaw BR, Han JY, Kim E, Gustafson DH, Hawkins RP, Cleary J, Lumpkins C. Effects of prayer and religious expression within computer support groups on women with breast cancer. *Psycho-Oncology*. 2007; 16(7):676–687. [PubMed: 17131348]
- Shaw BR, Hawkins RP, McTavish FM, Pingree S, Gustafson DH. Effects of insightful disclosure within computer mediated support groups on women with breast cancer. *Health Communication*. 2006; 19(2):133–142. [PubMed: 16548704]
- Shaw BR, McTavish FM, Hawkins RP, Gustafson DH, Pingree S. Experiences of women with breast cancer: Exchanging social support over the CHESS computer network. *Journal of Health Communication*. 2000; 5(2):135–159. [PubMed: 11010346]



- Slevin ML, Nichols SE, Downer SM, Wilson P, Lister TA, Arnott S, Cody M. Emotional support for cancer patients: What do patients really want? *British Journal of Cancer*. 1996; 74:1275–1279. [PubMed: 8883417]
- Spitzberg BH. Preliminary development of a model and measure of computer-mediated communication (CMC) competence. *Journal of Computer-Mediated Communication*. 2006; 11(2): 629–666.
- Taylor SE, Falke RL, Shoptaw SJ, Lichtman RR. Social support, support groups, and the cancer patient. *Journal of Consulting and Clinical Psychology*. 1986; 54(5):608–615. [PubMed: 3771878]
- Uchino, BN. *Social support and physical health: Understanding the health consequences of relationships*. Yale University Press; New Haven, CT: 2004.
- van Uden-Kraan CF, Drossaert CHC, Taal E, Lebrun CEI, Drossaers-Bakker KW, Smit WM, Van de Laar MAFJ. Coping with somatic illnesses in online *Computers in Human* support groups: Do the feared disadvantages actually occur? *Behavior*. 2008; 24:309–324.
- Walther, JB.; Boyd, S. Attraction to computer-mediated social support. In: Lin, CA.; Atkin, D., editors. *Communication technology and society: Audience adoption and uses*. Hampton Press; Cresskill, NJ: 2002. p. 153-188.
- Weathers M, Query JL, Kreps GL. A multivariate test of communication competence, social support, and coping among Hispanic lay caregivers for loved ones with Alzheimer’s disease: An extension of the Relational Health Communication Competence Model. *Journal of Participative Medicine*. 2010; 2:e14.
- Weinberg N, Schmale J, Uken JS, Wessel K. Online help: Cancer patients participate in a computer-mediated support group. *Health and Social Work*. 1996; 21(1):24–29. [PubMed: 8626154]
- Wentowski GJ. Reciprocity and the coping strategies of older people: Cultural dimensions of network building. *The Gerontologist*. 1981; 21(6):600–609. [PubMed: 7333486]
- White M, Dorman SM. Receiving social support online: Implications for health education. *Health Education Research*. 2001; 16(6):693–707. [PubMed: 11780708]
- Wills, TA.; Fegan, MF. Social networks and social support. In: Baum, A.; Revenson, TA.; Singer, JE., editors. *Handbook of health psychology*. Erlbaum; Mahwah, NJ: 2001. p. 209-234.
- Winefield HR. Support provision and emotional work in an Internet support group for cancer patients. *Patient Education and Counseling*. 2006; 62:193–197. [PubMed: 16139466]
- Winzelberg AJ, Classen C, Alpers GW, Roberts H, Koopman C, Adams RE, Taylor CB. Evaluation of an internet support group for women with primary breast cancer. *Cancer*. 2003; 97(5):1164–1173. [PubMed: 12599221]
- Wright KB. Social support within an on-line cancer community: An assessment of emotional support, perceptions of advantages and disadvantages, and motives for using the community from a communication perspective. *Journal of Applied Communication Research*. 2002; 30:195–209.
- Wright KB, Banas JA, Bessarabova E, Bernard DR. A communication competence approach to examining health care social support, stress, and job burnout. *Health Communication*. 2010; 25(4): 375–382. [PubMed: 20512719]
- Wright KB, Bell SB. Health-related support group on the Internet: Linking empirical findings to social support and computer-mediated communication theory. *Journal of Health Psychology*. 2003; 8:39–54. [PubMed: 22113899]
- Wright KB, Rosenberg J, Egbert N, Ploeger NA, Bernard DR, King S. Communication competence, social support, and depression among college students: A model of Facebook and face-to-face support network influence. *Journal of Health Communication*. 2013; 18(1):41–57. [PubMed: 23030518]
- Wright J. Online counseling: Learning from writing therapy. *British Journal of Guidance and Counseling*. 2002; 30(3):285–298.
- Yalom, ID. *The theory and practice of group psychotherapy*. Basic Books; New York, NY: 1970.
- Yoo W, Chih M-Y, Kwon M-W, Yang J, Cho E, McLaughlin B, Gustafson DH. Predictors of the change in the expression of emotional support within an online breast cancer support group: A longitudinal study. *Patient Education and Counseling*. 2013; 90:88–95. [PubMed: 23122429]

- Emotional communication competence moderates the benefits of emotional support.
- Expressing support improves mental well-being for patients with high competence.
- Expressing support decreases mental well-being for patients with low competence.
- Receiving support reduces cancer-related concern for patients with high competence.
- Receiving support increases cancer-related concern for patients with low competence.


```

----->---|<----->---|-----
FSR2      Acting:   No. spaces           R if operator
          Ahead   (-1 means anywhere)   retained Equal
          Behind  |                       | Priority
          Either  |           ADDIF1ST    DELBETWEEN | (E/e)
Nothing or Direction |           INSERT or  DELETED or |__|
Operator---->   (A/B/E) |   <-Target-----><-New target->  | | Rule
-----|-----|-----|-----|-----


NegWords      A    20   Stop           Encouragement2      38
You           A    20   Deserve        Deserving2           39
Worry        A    20   You          ExpressConcern1     40
Trust        A    20   You          TrustYou2            41
-----

```

**Fig. 1.**  
An example of InfoTrend coding rules for emotional support



**Fig. 2.** Interaction between emotional communication competence and expression of emotional support on psychological quality of life (scale ranges only partially displayed on y-axis).



**Fig. 3.** Interaction between emotional communication competence and reception of emotional support on breast cancer-related concerns (scale ranges only partially displayed on y-axis).

**Table 1**

## Subcategories of emotional support and keywords

<b>Emotional support</b>	<b>Keywords</b>
Empathy/sympathy	Empathy, sympathy, understand, sorry, worry, concern, etc.
Encouragement/reassurance	Hope, wish, trust, congratulation, cheer, hang in there, keep stay strong, keep marching, don't give up etc.
Care/physical affection	Take care, hugs, kisses, love, etc.
Universality/interrelationship	Common, isolated, sisterhood, not alone, together, etc.

**Table 2**

Descriptive statistics of messaging activities in CMSS groups for women with breast cancer

	Participants ( <i>N</i> = 236)			
	M	SD	Minimum	Maximum
Total number of messages posted	21.46	54.53	0	426.00
Total number of messages viewed	466.22	934.67	0	7121.00
Total counts of emotional support posted	24.09	70.97	0	459.00
Total counts of emotional support viewed	653.08	1294.11	0	9305.00
Expression of emotional support <sup>a</sup>	.40	.54	0	2.38
Reception of emotional support <sup>b</sup>	1.21	.76	0	6.63

<sup>a</sup>Note: This expression measure is operationalized by the total counts of emotional support posted divided by the total number of messages posted

<sup>b</sup>This reception measure is constructed by the total counts of emotional support viewed divided by the total number of messages viewed.

**Table 3**

## Demographic and disease-related characteristics of study participants

<b>Participants (N = 236)</b>	
Age	
4Mean ( <i>SD</i> )	51.18 (9.05)
Education	
1. Did not complete junior/middle school	0.9%
2. Did not complete high school	1.7%
3. High school degree	15.5%
4. Some college courses	16.7%
5. Associate degree (2 year college)	9.0%
6. Bachelor's degree (4 year college)	26.6%
7. Some graduate courses	7.7%
8. Graduate degree	21.9%
Income	
1. Less than \$20,000	5.5%
2. \$20,001 to \$40,000	14.3%
3. \$40,001 to \$60,000	17.5%
4. \$60,001 to \$80,000	17.1%
5. \$80,001 to \$100,000	19.8%
6. More than \$100,000	25.8%
Surgery experience	
Yes	55.3%
No	44.7%
Time interval between diagnosis to intervention (days)	
Mean ( <i>SD</i> )	75.12 (91.34)
Total time spent in CHES (minutes)	
Mean ( <i>SD</i> )	723.27 (1093.75)



**Table 4**

Hierarchical regression analysis predicting emotional outcomes

Criterion variables	Psychological well-being (posttest)	Breast cancer-related concerns (posttest)
<b>Block 1. Control variables</b>		
Pretest value of each outcome variable	.57***	.43***
Age	-.07	-.01
Education	-.04	-.02
Income	.03	-.01
Time (diagnosis to intervention)	-.09	.06
Surgery experience (yes = 1)	.12	-.20**
Full CHESS group (yes = 1)	-.06	.09
CHESS + mentor group (yes = 1)	-.08	-.002
Total time spent in CHESS	.08	-.16*
$R^2$ (%)	33.7***	20.3***
<b>Block 2. Moderator variable</b>		
Emotional communication competence (pretest)	-.03	.11
$R^2$ (%)	0.1	1.1
<b>Block 3. Main effects</b>		
Expression of emotional support	-.09	.001
Reception of emotional support	.06	-.11
$R^2$ (%)	0.8	1.1
<b>Block 4. Interaction effects</b>		
Emotional communication competence (pretest)	.17*	-.07
× Expression of emotional support		
Emotional communication competence (pretest)	-.06	-.17*
× Reception of emotional support		
$R^2$ (%)	3.0*	2.9*
Total $R^2$ (%)	37.6	25.4

Note: Cell entries are before-entry standardized regression coefficient for Block 4 and final standardized regression coefficients for Blocks 1, 2, and 3.

\*  $p < .05$

\*\*  $p < .01$

\*\*\*  $p < .001$