



Published in final edited form as:

Free Radic Biol Med. 2013 October ; 63: 254–263. doi:10.1016/j.freeradbiomed.2013.05.023.

Oxidative Damage and Cellular Defense Mechanisms in Sea Urchin Models of Aging

Colin Du, Arielle Anderson, Mae Lortie, Rachel Parsons, and Andrea Bodnar
Bermuda Institute of Ocean Sciences, St. George's, Bermuda, GE 01

Abstract

The free radical or oxidative stress theory of aging proposes that the accumulation of oxidative cellular damage is a major contributor to the aging process and a key determinant of species longevity. This study investigates the oxidative stress theory in a novel model for aging research, the sea urchin. Sea urchins present a unique model for the study of aging due to the existence of species with tremendously different natural life spans including some species with extraordinary longevity and negligible senescence. Cellular oxidative damage, antioxidant capacity and proteasome enzyme activities were measured in the tissues of three sea urchin species: short-lived *Lytechinus variegatus*, long-lived *Strongylocentrotus franciscanus* and *Strongylocentrotus purpuratus* which has an intermediate lifespan. Levels of protein carbonyls and 4-hydroxynonenal (HNE) measured in tissues (muscle, nerve, esophagus, gonad, coelomocytes, ampullae) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) measured in cell-free coelomic fluid showed no general increase with age. The fluorescent age-pigment lipofuscin measured in muscle, nerve and esophagus, increased with age however it appeared to be predominantly extracellular. Antioxidant mechanisms (total antioxidant capacity, superoxide dismutase) and proteasome enzyme activities were maintained with age. In some instances, levels of oxidative damage were lower and antioxidant activity higher in cells or tissues of the long-lived species compared to the short-lived species, however further studies are required to determine the relationship between oxidative damage and longevity in these animals. Consistent with the predictions of the oxidative stress theory of aging, the results suggest that negligible senescence is accompanied by a lack of accumulation of cellular oxidative damage with age and maintenance of antioxidant capacity and proteasome enzyme activities may be important mechanisms to mitigate damage.

Keywords

sea urchin; oxidative damage; reactive oxygen species; antioxidants; proteasome; lipofuscin; aging; negligible senescence

Introduction

Aging is a progressive decline in biological function accompanied by an increased risk of degenerative disease and death over time. It is a complex and multifactorial process and there have been many theories proposed to explain this phenomenon at the molecular, cellular, systemic and evolutionary levels [1]. The oxidative stress or free radical theory of

© 2013 Elsevier Inc. All rights reserved.

Corresponding author: Andrea Bodnar, **Tel:** 441-297-1880 ext. 724, **Fax:** 441-297-8143, **E-mail:** andrea.bodnar@bios.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

aging is one of the most studied hypotheses for the molecular basis of aging. This theory proposes that the accumulation of cellular damage caused by reactive oxygen species (ROS) plays a key role in the aging process, as well as in determining organismal longevity [2–4]. ROS are partially reduced derivatives of oxygen that are highly reactive with major components of the cell such as proteins, lipids and DNA. They are continually generated as by-products of a number of cellular processes and are also produced when the cells are exposed to various external stimuli such as ultraviolet light, ionizing radiation and environmental toxins [3]. ROS are required for proper cell function because of their role in cell signalling and the immune response however, when ROS exceed a level of homeostasis, oxidative stress ensues. Oxidative stress results from an imbalance between the production of ROS and the cell's ability to mitigate damage through antioxidant pathways or mechanisms that repair or eliminate damaged molecules.

Over the years, there has been much evidence in support of the oxidative stress theory of aging and numerous studies have shown that oxidative damage increases with age in the cells and tissues of many organisms [5]. In addition, several studies comparing animals with different lifespans suggest that long-lived species tend to show reduced oxidative damage, reduced mitochondrial ROS production, increased antioxidant defenses and/or increased resistance to oxidative stress [6]. However the status of this theory is far from certain as some studies have found a lack of correlation between oxidative damage and lifespan and genetic manipulation of antioxidant pathways in invertebrate models (e.g. the nematode worm *Caenorhabditis elegans* and the fruit fly *Drosophila melanogaster*) have yielded variable results on lifespan while those in higher animals (e.g. the mouse, *Mus musculus*) have generally not supported a role for oxidative stress in modulating longevity [6,7]. Nevertheless there is convincing evidence that a reduction in oxidative stress positively influences biomarkers of health, physiological function and disease resistance and therefore it remains an important aspect of the aging process [6].

Many studies of the oxidative stress theory have focused on humans and short-lived model organisms such as *C. elegans*, *D. melanogaster* and *M. musculus*, all animals that exhibit well characterized organismal senescence, however less is known about the role of oxidative stress in animals that exhibit negligible senescence. Negligible senescence is a term used to describe animals that do not show an increase in mortality rate or a decrease in fertility, physiological function or disease resistance with age [8,9]. In this respect, sea urchins offer an interesting model to explore due to the existence of species with very different natural life spans, including some that display extreme longevity and negligible senescence. The red sea urchin (*Strongylocentrotus franciscanus*) is one of the earth's longest living animals, living in excess of 100 years with no age-related increase in mortality rate or decrease in reproductive capacity [10,11]. In contrast, *Lytechinus variegatus* has an estimated maximum life expectancy of approximately 4 years [12,13] while the maximum life expectancy of *Strongylocentrotus purpuratus* is estimated to be approximately 50 years [14,15]. Thus sea urchins provide a unique model to investigate cellular mechanisms underlying life span determination and negligible senescence. From a phylogenetic viewpoint, sea urchins are also interesting as they are more closely related to humans than other invertebrate models used for aging research (e.g. *C. elegans* and *D. melanogaster*). Sequencing the genome of *S. purpuratus* confirmed the close genetic relationship between sea urchins and humans revealing that sea urchins have an estimated 23,300 genes including representatives of nearly all vertebrate gene families and orthologs of many human disease-associated genes [16]. Age-related gene expression profiles in the tissues of *S. purpuratus* have revealed some interesting distinctions from aging animals. There was no age-related decline in expression of genes involved in energy production and, in some tissues, an increase in expression of genes involved in protein homeostasis, apoptosis and signaling pathways that play a role in

tissue regeneration [17]. No previous studies have investigated oxidative cellular damage and defense mechanisms with respect to age in sea urchins.

When proteins are oxidized by ROS, some amino acid residues are modified which can lead to conformation changes and functional loss. One of the most commonly studied oxidative products is the formation of carbonyl derivatives of amino acids such as lysine, arginine, proline and threonine [18]. When cellular membranes are exposed to ROS, the lipids containing polyunsaturated fatty acids undergo a series of chain reactions to yield end products such as malondialdehyde (MDA) and 4-hydroxynonenal (HNE) [19]. Levels of oxidative DNA damage can be assessed by measuring products of modified nucleosides such as 8-hydroxy-2'-deoxyguanosine (8-OHdG) in tissues or bodily fluids [20–22].

To prevent oxidative damage to proteins, lipids and DNA, the generation of ROS is counterbalanced by various antioxidant systems. These include enzymes such as superoxide dismutase (SOD) and catalase and non-enzymatic antioxidants like glutathione, tocopherols and thioredoxin [19,23]. In addition to antioxidant mechanisms, the degradation of oxidized proteins by the proteasome and autophagy-lysosomal pathways also constitutes an important part of the cell's defense against oxidative stress [24,25]. In particular, the autophagic pathways has been shown to remove damaged mitochondria and oxidatively damaged macromolecules from the cytosol resulting in the accumulation of aggregates of the highly-autofluorescent pigment (lipofuscin) in the lysosomal lumen [26,27]. Lipofuscin accumulation is one of the most recognized hallmarks of aging in a wide range of vertebrate and invertebrate animals.

In order to understand the role of the oxidative stress theory of aging in animals with an alternative life history from humans and short-lived invertebrate models we characterized age-related oxidative cellular damage, antioxidant capacity, as well as proteasome enzyme activities in tissues of three sea urchin species with different lifespans (*L. variegatus*, *S. purpuratus* and *S. franciscanus*).

Materials and Methods

Collection and age determination of sea urchin species

L. variegatus were collected from Flatt's Inlet, Bermuda (32 10.375N and 64 44.216W) in January, 2012. *S. purpuratus* were collected in Mission Bay, San Diego, California (32 46.833N and 117 14.557W) in April, 2012. *S. franciscanus* were collected near Kendrick Island in Gabriola Pass, British Columbia (49 07.554N and 123 41.461W) in July, 2011 and July 2012. Following collection, the sea urchins were kept in flow-through aquaria to ensure all urchins were exposed to a uniform environment prior to tissue dissections.

L. variegatus ages were estimated using test (shell) diameter and comparisons with previously established growth curves generated from size frequency and growth band counting data collected in Florida and Bermuda [12,13]. *S. purpuratus* ages were estimated from test diameter using growth curves previously generated from the Tanaka parameters from tetracycline tagging experiments conducted at Mission Bay, San Diego ($f=1.30757$, $d=2.80184$, $a=0.16314$) [28]. *S. franciscanus* age estimates were based on growth curves generated from tetracycline tagged sea urchins in Oregon and Washington using the Tanaka function ($f=0.22929$, $d=6.07531$, $a=0.19906$) [29,30].

Preparation of tissue and cell extracts

The tissues examined in this study include muscle taken from the jaw structure which is referred to as Aristotle's lantern muscle (Alm), radial nerve (RN), esophagus (Es), gonad (Gon) and ampullae (Amp) which is part of the sea urchins water vascular system. These

tissues were dissected from *L. variegatus*, *S. purpuratus* and *S. franciscanus* and stored at -70°C until analysis. Ampullae were not harvested from *S. franciscanus* collected during July 2011. Coelomocytes were harvested from 1.5–3 ml of coelomic fluid by centrifuging at $6000 \times g$ for 5 minutes and both the cell pellets and cell-free coelomic fluid were frozen until further analysis. Tissue homogenates for the oxidative damage and antioxidant assays were prepared from 30–50 mg of tissue or coelomocyte cell pellets by homogenization in 100 – 200 μl of protein homogenization buffer (50 mM HEPES, 125 mM KCl, 1.25 mM EDTA, 0.6 mM MgSO_4) [31] supplemented with protease inhibitors (40 $\mu\text{g}/\text{ml}$ pepstain A, 5 $\mu\text{g}/\text{ml}$ antipain and 100 $\mu\text{g}/\text{ml}$ phenylmethylsulfonylfluoride) using pestles designed for 1.5 ml microcentrifuge tubes. The tissue homogenates were left on ice for 10 minutes and then were centrifuged at $9,300 \times g$, 4°C for 5 minutes. The supernatant was harvested and protein concentrations determined by the Bradford assay as previously described [32]. The homogenates were stored in aliquots at -70°C prior to subsequent analyses.

Measurement of 8-hydroxy-2'-deoxyguanosine (8-OHdG) in cell-free coelomic fluid

As an indicator of oxidative DNA damage, 8-OHdG was measured in 50 μl cell-free coelomic fluid, prepared as described above, from sea urchins of all three species using the OxiSelect Oxidative DNA Damage ELISA Kit (8-OHdG Quantitation) assay (Cell Biolabs, San Diego, CA, USA). The signals generated by the colorimetric assay were detected at 450 nm on a VERSA max microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Measurement of oxidative damage to proteins and lipids

Protein oxidation in the sea urchin cell and tissue extracts was analyzed by measuring protein carbonyls using the OxiSelect Protein Carbonyl ELISA Kit (Cell Biolabs, San Diego, CA, USA) with 1 μg input protein for each cell or tissue homogenate. The lipid peroxidation by-product, 4-hydroxynonenal histidine protein adduct was measured using the OxiSelect HNE-His adduct ELISA Kit (Cell Biolabs) using 1 μg input protein. The signals generated by the colorimetric assays were detected at 450 nm on a VERSA max microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Measurement of total antioxidant capacity and antioxidant enzyme activities

Total antioxidant capacity in the tissue and cell extracts was assayed using the OxiSelect Total Antioxidant Capacity Assay Kit (Cell Biolabs, San Diego, CA, USA) using 10 μg input protein for each sample. Activities of three antioxidant enzymes in the extracts were measured using the Superoxide Dismutase Activity Assay Kit, the Catalase Activity Assay Kit and the Glutathione Reductase Activity Assay Kit (BioVision, Milpitas, CA, USA), according to the manufacturer's protocols using 5 μg , 20 μg and 20 μg input protein, respectively. The assay reactions were conducted in 96-well plates and the colorimetric signals were detected using a VERSA max microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Measurement of proteasome activity

Proteasome activity was evaluated using the Proteasome-Glo Trypsin-like, Chymotrypsin-like and Caspase-like assays (Promega, Madison, WI, USA), following the manufacturer's guidelines and 5 μg input protein for each sample. For these assays, the tissue or cell homogenates were prepared as described above except that the protease inhibitors were omitted from the homogenization buffer. The assay reactions were placed into a 96-well optical bottom plate (Thermo Fisher Scientific, Waltham, MA, USA) and the luminescence produced was detected and recorded as relative light units (RLU) using a SpectraMax M2 microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Evaluation of lipofuscin by autofluorescence

Tissues (Aristotle's lantern muscle, radial nerve and esophagus) were dissected and placed into 4% paraformaldehyde in phosphate buffered saline and sent to Histopathology Services at the Jackson Laboratory (Bar Harbor, ME, USA) where they were imbedded in paraffin and sectioned (5 μ M) Paraffin was removed from the slides by immersion in toluene for 5 minutes followed by successive treatments with 100%, 95% and 70% ethanol for 5 minutes each. Slides were then rinsed in 1x PBS and left to dry in a 37°C oven for 20 minutes. Autofluorescence was assessed using an Olympus Provis AX70 Electrofluorescence microscope (Olympus, Tokyo, Japan) with a 40x objective by excitation with narrow green light (530–545 nm) and emission at 610–675 nm. Images were captured with a Toshiba IK-TU40A CCD camera (Irvine, CA, USA) or a Retiga EXi Digital camera (Qimaging, Surrey, BC, Canada).

For each tissue section, 12 locations were randomly selected and images captured under bright-field and fluorescent illumination. Autofluorescence levels were quantified using Image Pro Plus, version 4.0 (Media Cybernetics, Bethesda, MD, USA) using the Integrated Optical Density (IOD) measurement tool. In this analysis, lipofuscin granules were discriminated from the surrounding tissues in the digital images by light intensity range selection following background subtraction based on a constant background level set for each tissue. Image Pro's measurement tool was then used to determine the total tissue area in each image using the bright-field image. The average IOD per tissue area for each tissue was calculated after removing the highest and lowest values and averaging the remaining 10 images.

Statistical analysis

Outliers in the data were detected by the Grubbs' test, using QuickCalcs (GraphPad Software, Inc., La Jolla, CA, USA). Species and age specific differences were analyzed using the general linear model for comparison within each tissue type (Statgraphics Centurion, Statpoint Technologies, Inc., Warrenton, VA, USA). Comparisons between the young and old groups within a tissue for each sea urchin species were conducted using Statgraphics Centurion and Excel using t-tests for equal or unequal variances as appropriate after confirming that the data were normally distributed or using the Mann Whitney test if the data were not normally distributed. Species specific differences were evaluated using one way ANOVA for normally distributed data and the Kruskal-Wallis test if the data were not normally distributed, followed by multiple range post-hoc tests (Statgraphics Centurion). Differences of $p < 0.05$ were considered statistically significant. Data are expressed as mean \pm standard error of the mean.

Results

Age estimates for sea urchins

Ages of individual sea urchins were estimated from test (shell) diameter using previously established growth curves generated for sea urchins that were analyzed at or near our collection sites. The sea urchins were divided into two age groups (young and old) and their average test diameter, the number of animals in each group and estimated ages are shown in Table 1. As growth rate can be affected by environmental factors (e.g. food availability) our age estimates serve only as a guideline and not absolute values. However, studying animals within a defined study site should give a reasonable indication of relative age within the population. For our study we selected animals at the upper and lower size range for the populations under study.

8-OHdG levels in cell-free coelomic fluid of *L. variegatus*, *S. purpuratus* and *S. franciscanus*

Cell-free coelomic fluid from *L. variegatus*, *S. purpuratus* and *S. franciscanus* was used to assess the level of 8-OHdG as a marker of oxidative DNA damage. Levels of 8-OHdG in cell-free coelomic fluid showed no significant difference in levels between age groups or between species (age: $F_{1,31}=0.01$, $p=0.93$; species: $F_{2,31}=0.68$, $p=0.51$) (Fig 1).

Protein oxidation and lipid peroxidation

The protein carbonyl content in cell and tissue extracts from the three sea urchin species is shown in Fig. 2. Ampullae were not harvested from *S. franciscanus* collected during July 2011 and so indicators of oxidative damage were not assayed in this tissue. The general linear model showed significant species specific differences in protein carbonyl levels in all tissues and age-specific differences in esophagus and gonads. Post-hoc tests comparing young and old within each species confirmed that most tissues did not show increased protein carbonyls with age. There was no significant age-related change in protein carbonyl content in muscle, esophagus, nerve, ampullae and coelomocytes from *L. variegatus*, however there was a small but statistically significant increase with age in gonad tissue (1.12-fold, $p=0.03$) (Fig. 2, Panel A). For *S. purpuratus*, there was no significant age-related change in protein carbonyl content in muscle, esophagus, nerve, gonad and ampullae tissues but a statistically significant increase with age in protein carbonyl content was found in coelomocytes (1.22-fold, $p=0.02$) (Fig 2, Panel B). There was no significant age-related change in protein carbonyl content in muscle, nerve and coelomocytes from *S. franciscanus* (Fig. 2, Panel C). A statistically significant age-related increase in protein carbonyl content was observed in esophagus (1.76-fold, $p<0.01$) and gonad (2.43-fold, $p<0.01$) of *S. franciscanus* however overall levels in these tissues remained very low. One-way ANOVA or the Kruskal-Wallis tests and post-hoc analyses confirmed that protein carbonyls were significantly higher in all *L. variegatus* tissues (except coelomocytes) compared to the corresponding tissues from the other two long-lived species (>1.7 -fold, $p<0.05$ in each case). Comparisons between the two longer-lived species showed significantly lower levels of protein carbonyls in both young and old muscle tissue (1.3-fold, $p<0.05$) and gonad (>2.5 -fold, $p<0.05$) of *S. franciscanus* compared to *S. purpuratus* while protein carbonyls in coelomocytes of *S. franciscanus* were significantly higher than both *L. variegatus* and *S. purpuratus* (>1.2 -fold, $p<0.05$, in each case).

The same extracts used for protein carbonyl analysis were used to measure levels of HNE-His as an indicator of lipid peroxidation (Fig. 3). The general linear model showed significant species specific differences HNE-His levels in all tissues except gonads and age-specific differences in muscle and radial nerve. Post-hoc tests comparing young and old within each species confirmed that most tissues did not show increased HNE with age. There was no significant age-related change in HNE-His adduct level in all the tissues from *L. variegatus* except the radial nerve, which showed a statistically significant increase of 1.44-fold ($p<0.01$) with age (Fig. 3, Panel A). Similarly, there was no significant age-related change in HNE-His adduct level in all the tissues examined from *S. purpuratus* except radial nerve which showed a 1.54-fold increase ($p=0.03$) (Fig. 3, Panel B). In *S. franciscanus*, a statistically significant age-related increase in HNE-His adduct level in muscle (1.23-fold, $p<0.01$) and nerve (1.50-fold, $p<0.01$) was observed (Fig. 3, Panel C). However, a statistically significant age-related decrease (1.25-fold, $p=0.03$) in HNE-His adduct level in coelomocytes was observed. One-way ANOVA or Kruskal-Wallis tests and post-hoc analyses were used for inter-species comparisons which revealed significantly higher levels of HNE-His adducts in esophagus, radial nerve, coelomocytes and ampullae for *L. variegatus* compared to the corresponding tissues from the long-lived species (>1.9 -fold, $p<0.05$ in all cases). Levels of HNE-His were significantly lower in young and old muscle

tissue of *S. purpuratus* compared to *S. franciscanus* (>1.8-fold and $p<0.05$) and also *L. variegatus* (>1.9-fold and $p<0.05$).

Total antioxidant capacity

Total antioxidant capacity (TAC) was evaluated in tissue extracts from the three sea urchin species by measuring the reduction of Cu^{2+} to Cu^+ . Ampullae were not harvested from *S. franciscanus* collected during July 2011 and therefore antioxidant capacity was not assayed in this tissue. The general linear model showed significant species specific differences in TAC levels in all tissues except esophagus and gonads and age-specific differences in esophagus, radial nerve and coelomocytes. Post-hoc tests revealed no significant age-related change in TAC in all the tissues examined from *L. variegatus* and *S. franciscanus* (Fig. 4, Panel A, C and D). In *S. purpuratus*, there was a statistically significant age-related increase in TAC in muscle (1.19-fold, $p=0.05$), esophagus (2.29-fold, $p<0.01$) and radial nerve (1.77-fold, $p<0.01$) (Fig. 4, Panel B). However, there was a significant age-related decrease in coelomocytes (1.77-fold, $p<0.01$) (Fig 4, Panel D). It was observed that the TAC of coelomocytes was much higher than the other tissues for all three of the sea urchin species examined and thus it was plotted using a different scale (Fig. 4, Panel D). One-way ANOVA or Kruskal-Wallis tests and post-hoc analyses revealed that the coelomocytes from *S. purpuratus* and *S. franciscanus* exhibited statistically higher TAC (> 5-fold, $p<0.05$) than the coelomocytes from *L. variegatus*. While there was no significant difference between TAC in coelomocytes of young *S. franciscanus* and *S. purpuratus*, the coelomocytes from old *S. franciscanus* had higher TAC than old *S. purpuratus* (1.5-fold, $p<0.05$). Interspecies comparisons also revealed a significantly lower level of TAC in young and old muscle of *S. franciscanus* compared to *L. variegatus* and *S. purpuratus* (>1.5-fold, $p<0.05$) and a higher level of TAC in young and old ampullae of *L. variegatus* compared to *S. purpuratus* (>1.6-fold, $p<0.05$).

Antioxidant enzyme activities

Superoxide dismutase (SOD) activity of different tissues from the three sea urchin species is shown in Fig. 5. Ampullae were not harvested from *S. franciscanus* collected during July 2011 and therefore antioxidant enzyme activity was not assayed in this tissue. The general linear model showed significant species specific differences in SOD levels in all tissues except radial nerve and age-specific differences in muscle and radial nerve. Post-hoc tests revealed no significant age-related change in SOD activity in all the tissues examined from *L. variegatus* (Fig. 5, Panel A). In *S. purpuratus*, there was no significant age-related change in SOD activity in all the tissues examined except radial nerve and muscle tissue which each showed a significant age-related increase of 1.19-fold ($p<0.01$ for nerve and $p=0.03$ for muscle) (Fig. 5, Panel B). A significant age-related increase (1.26-fold, $p<0.01$) in SOD activity was observed in the radial nerve from *S. franciscanus*, but no other age-related changes were observed in this species (Fig. 5, Panel C). One-way ANOVA or Kruskal-Wallis tests and post-hoc analyses revealed some species specific differences in all tissues with the most notable being gonad and coelomocyte. SOD activity was significantly higher (>2-fold, $p<0.05$) in young and old gonad of *S. purpuratus* and old gonad of *S. franciscanus* compared to *L. variegatus*. Coelomocytes from both young and old *S. purpuratus* and *S. franciscanus* exhibited statistically higher SOD activity (>2-fold, $p<0.05$) than the coelomocytes from *L. variegatus*.

Catalase and glutathione reductase assays were also conducted with all tissue extracts except coelomocytes. Catalase activity was undetectable in most of the tissues examined in the three sea urchin species when 20 μg of total protein was used in the assay. Catalase activity was detected in the gonad of the three sea urchin species but no significant age-related change was observed (data not shown). Glutathione reductase activity was detectable in all

the tissues examined in *L. variegatus* with no age-related difference (data not shown). In the long-lived species, glutathione reductase activity was undetectable in several tissues when 20 µg of total protein was used in the assay. Glutathione reductase activity was detected in the esophagus and gonad of *S. purpuratus* and the esophagus, gonad and nerve of *S. franciscanus* but no significant age-related change was observed (data not shown).

Proteasome enzyme activity in sea urchin tissues

Proteins that are damaged by ROS are recognized and degraded by the proteasome complex which exhibits trypsin-like, chymotrypsin-like and caspase-like activities. These activities were measured in tissue extracts for the three sea urchin species. Coelomocytes were not assayed for proteasome activity due to a limited amount of material for these samples. Although proteasome activity has been reported to decline with age in other organisms [25] there was no consistent decrease across sea urchin tissues with age (Fig. 6). Trypsin-like activity did significantly decrease in radial nerve of *S. franciscanus* (2.55-fold, $p=0.01$) and the caspase-like activity in esophagus of *S. purpuratus* decreased 1.33-fold ($p=0.02$) with age. Otherwise, there was a significant increase with age for trypsin-like activity in *L. variegatus* muscle tissue (1.22-fold, $p<0.01$). Chymotrypsin-like activity was increased with age in *S. purpuratus* ampullae (2.75-fold, $p=0.02$) and nerve (1.81-fold, $p<0.01$) as well as *S. franciscanus* ampullae (1.61-fold, $p<0.01$) and nerve (2.38-fold, $p=0.01$). Caspase-like activity increased with age in muscle tissue of all three species: *L. variegatus* (1.25-fold, $p=0.04$), *S. purpuratus* (1.44-fold, $p<0.01$) and *S. franciscanus* (1.37-fold, $p<0.01$). Caspase-like activity also increased slightly with age in *L. variegatus* nerve tissue (1.16-fold, $p=0.04$). The general linear model showed significant species specific differences in all tissues except trypsin-like activity of esophagus. One-way ANOVA or Kruskal-Wallis tests and post-hoc analyses revealed the most notable difference in radial nerve and ampullae tissue with the short-lived *L. variegatus* having higher levels of trypsin-like and chymotrypsin-like activities than the longer-lived species (Fig. 6, Panels A, B and C). For ampullae, trypsin-like activity was significantly higher in young and old *L. variegatus* compared to young and old *S. purpuratus* and *S. franciscanus* (>2.5 -fold, $p<0.05$ in all cases) and chymotrypsin-like activity was significantly higher in young *L. variegatus* compared to young tissue from *S. purpuratus* and *S. franciscanus* (>5.7 -fold, $p<0.05$, in all cases). For radial nerve, trypsin-like activity was significantly higher in young and old *L. variegatus* compared to young and old *S. purpuratus* and old *S. franciscanus* (>1.77 -fold, $p<0.05$, in all cases). Also for radial nerve, chymotrypsin-like activity of *L. variegatus* was significantly higher than in young and old *S. franciscanus* and young *S. purpuratus* (>1.8 -fold, $p<0.05$, in each case) and chymotrypsin-like activity of young and old *S. purpuratus* were significantly higher than young and old *S. franciscanus* (>3.82 -fold, $p<0.05$, in each case).

Lipofuscin accumulation in sea urchin tissues

Lipofuscin is an autofluorescent pigment that accumulates in cells and tissues of both vertebrate and invertebrate animals with age. Autofluorescence was evaluated in muscle, esophagus and nerve tissue of young and old sea urchins as an indicator of lipofuscin levels (Fig 7). Analysis of the data using the general linear model revealed both age and species differences for muscle and nerve and species specific differences in esophagus. Although all tissues showed an increase in autofluorescence with age, post-hoc tests revealed this reached significance in muscle tissue of *L. variegatus* ($p=0.02$) and *S. purpuratus* ($p=0.02$), esophagus tissue of *S. purpuratus* ($p=0.02$) and nerve tissue of *L. variegatus* ($p=0.03$), *S. purpuratus* ($p=0.04$) and *S. franciscanus* ($p=0.05$). For young muscle tissue, autofluorescence was significantly higher in *L. variegatus* than both *S. purpuratus* and *S. franciscanus* (Kruskal-Wallis: $H=8.02$, $df=2$, $n=19$, $p=0.02$), however for old muscle, autofluorescence was higher in *S. franciscanus* than *L. variegatus* and *S. purpuratus* (Kruskal-Wallis: $H=11.88$, $df=2$, $n=17$, $p<0.01$). For nerve, there was no species specific

difference between the young tissues ($H=3.07$, $df=2$, $n=19$, $p=0.22$) but autofluorescence was significantly higher in old nerve tissue of *S. franciscanus* compared to that of *L. variegatus* and *S. purpuratus* ($H=11.7$, $df=2$, $n=17$, $p<0.01$). For esophagus, autofluorescence was significantly higher in both young and old sea urchins of *L. variegatus* compared to *S. purpuratus* and *S. franciscanus* (Kruskal-Wallis: young: $H=15.61$, $df=2$, $n=19$, $p<0.01$ and old: $H=7.76$, $df=2$, $n=17$, $p=0.02$). It was evident from these analyses that autofluorescence was not evenly distributed throughout the tissues but appeared in scattered patches. When tissues were treated with the DNA dye DAPI (4',6-diamidino-2-phenylindole), it revealed that areas of autofluorescence did not co-localize with nuclear staining but were found in areas devoid of cellularity. It was also noted that there was a high degree of variability in autofluorescence between individuals as reflected in the large error bars shown in Figure 7.

Discussion

Although many studies have shown that oxidative damage increases with age in the cells and tissues of a variety of organisms [5], a general age-related increase in some classic markers of oxidative damage was not observed in sea urchin tissues. In humans, levels of oxidative DNA damage have been assessed by measuring products of modified nucleosides such as 8-hydroxy-2'-deoxyguanosine (8-OHdG) in tissues and bodily fluids and increased levels of 8-OHdG have been reported with respect to age, smoking, inflammatory diseases and cancer [20–22]. Measuring 8-OHdG in the bodily fluids (e.g. serum or urine) gives an indication of the overall level of damage in the whole body and has the advantage that it avoids artefactual oxidation of DNA that can be introduced during sample preparation [22]. Similar to the blood of higher animals, coelomic fluid of sea urchins is in direct contact with internal cells and tissues and can provide an overall profile of the biological status of the organism. Contrary to what is seen in aging animals, we did not detect an increase in 8-OHdG in cell-free coelomic fluid with age in sea urchins.

Markers of protein oxidation (carbonyls) and lipid peroxidation (HNE) did not show an age-related increase in most cells and tissues of sea urchins. Consistent with these observations, a study on aging in the long-lived ocean quahog clam *Arctica islandica* showed no significant difference in the level of protein carbonyls with age in gill and mantle tissues [31]. A statistically significant increase in levels of HNE was observed in radial nerve tissues of all three sea urchin species examined indicating that certain tissues are susceptible to some forms of oxidative damage. It is interesting that the magnitude of the increase is similar in all three species (approximately 1.5-fold) even though *S. purpuratus* and *S. franciscanus* are estimated to be much longer lived, indicating a lower rate of accumulation of damage in these species. It has been reported that protein carbonyl levels are lower in the tissues of long-lived *A. islandica* compared to tissues of short-lived clams (*Mya arenaria* and *Laternula elliptica*) or scallops (*Aequipecten opercularis* and *Adamassium colbecki*) [31]. Similarly, our results indicated that there was generally less damage in tissues of long-lived species than in the short-lived species of sea urchin. Although interesting, these data should be viewed with caution as the sea urchins used in this study were collected from different geographical locations and it has been shown that variable environmental factors (e.g. temperature, light, pollution) can stochastically modify the production of ROS and the accumulation of damage in animals from different locations [33,34]. Therefore, it is difficult to conclude that differences across species are solely related to lifespan. Further studies could address this point by examining sea urchin species with different lifespans collected from the same geographic location.

To explore potential mechanisms for the lack of accumulation of oxidative damage with age, we investigated the activity of a variety of cellular antioxidant systems: direct ROS scavenging enzymes SOD and catalase as well as glutathione reductase, a rate limiting

enzyme in glutathione recycling. In addition, total antioxidant capacity (TAC) was measured by the ability of the cell and tissue extracts to reduce Cu^{2+} to Cu^+ . In sea urchin tissues, there was generally no age-related change in TAC or antioxidant enzyme activity and little difference between the species with different lifespans. We did observe a high level of antioxidant activity in coelomocytes which may be important to combat high levels of ROS produced during the immune response of the phagocytotic cells present in this cell mixture. It is also interesting that there were much higher levels of TAC and SOD in coelomocytes of the long-lived species compared to *L. variegatus* but whether this reflects a more robust immune response and higher ROS levels in long-lived species remains to be determined. Other studies of age-dependant changes in antioxidant activities have not shown a consistent trend as the activities of some antioxidants decline whereas others remain unchanged or even increase with age depending on enzyme, tissue or species [5]. In addition, long-lived animals often have lower levels of tissue antioxidants than short-lived animals and experimentally altering antioxidants through dietary supplementation or transgenic techniques does not, in most cases, affect maximum life span [23]. It is now thought that the overall level of production of ROS is more important than the modulation of ROS by antioxidant activities in the context of the oxidative stress theory of aging [23] and it will be important to measure ROS production in sea urchin tissues in future studies.

The degradation of oxidized proteins by the proteasome pathway constitutes another important part of the cell's defense against oxidative stress by preventing the accumulation of damaged proteins [24,25]. There is evidence that proteasome activity decreases with age in human tissues and model systems which may be causally related to aging and several age-related diseases [24,25]. In contrast, we found no age-related decrease in the activity of proteasome enzymes except for trypsin-like activity in *S. franciscanus* radial nerve tissue and caspase-like activity in *S. purpuratus* esophagus tissue. In all other tissues trypsin-, chymotrypsin- and caspase-like activities were unchanged or, in some cases, showed a significant increase with age. This suggests a general maintenance in proteasome function with age which may contribute to the lack of increase in protein carbonyls observed in this study. It was noted that some tissues of the short-lived *L. variegatus* had higher proteasomal activities than the longer-lived species and *S. franciscanus* generally had the lowest levels of activity. When proteasome enzyme activities were measured in tissues of clams with different lifespans, there were no inter-species differences between trypsin-like and chymotrypsin-like activity in gills but caspase was significantly lower in gill of long-lived *A. islandica* compared to short-lived *M. mercenaria* [35]. However, a recent study has demonstrated that non-proteasomal proteases in crude cell or tissue extracts can contribute to the activity measured in the proteasomal assay and that this contribution varies between different species and tissue types [36]. Therefore it is essential to distinguish between proteasomal and non-proteasomal activities using proteasome-specific inhibitors before one can make generalizations about interspecies differences.

Another major pathway of catabolism in eukaryotic cells is the autophagy-lysosomal system and it is thought that this pathway is responsible for the accumulation of lipofuscin, one of the most recognized hallmarks of aging in a wide range of vertebrate and invertebrate animals [26,27]. Many studies have reported the negative effects of accumulating lipofuscin on lysosomal activity and cellular function, but more recent studies have suggested that lysosomes can play a protective role in the oxidative stress response and suggest that lipofuscin accumulation is indicative of cellular maintenance through autophagic activity [26,27,37]. In sea urchins, an age-related accumulation of lipofuscin was seen in all tissues examined (muscle, nerve and esophagus) which reaches significance in muscle tissue of *L. variegatus* and *S. purpuratus*, esophagus tissues of *S. purpuratus* and nerve tissue of all three species. Lipofuscin staining was not evenly distributed and not punctate as one would expect if it was confined to lysosomes, but rather appeared as patches of autofluorescence scattered

throughout the tissues in areas devoid of nuclear staining. There was a high level of inter-individual variation in the levels of lipofuscin, consistent with previous studies in which lipofuscin, measured in the gonads of the sea urchin (*Strongylocentrotus intermedius*), was also found to be highly variable between individuals such that no strong correlation was observed between lipofuscin levels and exposure to pollution (DDT, oil hydrocarbons and heavy metals) [38].

Lipofuscin has been shown to accumulate with age in the tissues of other marine invertebrates such as bivalves and crustaceans [31,39–42]. In *A. islandica*, lipofuscin levels were significantly different in three tissues examined (gill > mantle > muscle) and increased significantly with age in all three tissues [31]. The authors report that lipofuscin granules were located primarily in connective tissues and interstitial spaces presumably where they are less likely to impair metabolic processes of the tissues. Lipofuscin accumulation in the neural tissues of lobsters [*Homarus gammarus* (European lobsters) and *Panulirus argus* (Caribbean spiny lobsters)] has been used to determine age and is reported to provide a more accurate estimate of age than carapace length [40–42]. However levels of lipofuscin have been shown to be affected by environmental factors such as seasonal temperature oscillations and metal exposure [34,39,40] and age accumulated lipofuscin can be reversed presumably through exocytosis [43]. Similar to sea urchins, lipofuscin granules were shown to be mainly extracellular in the eye stalk of lobster neural tissue [40], however it remains to be determined if the accumulation of lipofuscin affects tissue function in these animals.

The overall findings of this study show that markers of cellular oxidative damage did not generally increase with age in the tissues of sea urchin species with different lifespans. Antioxidant capacity and proteasomal enzyme activities were maintained with age which may contribute to the lack of accumulation of oxidative damage however future studies should examine if the production of ROS differs between tissues and between species. This study is the first to look at age-related oxidative damage in sea urchins with different lifespans collected from their natural environment. It should be noted that we have only investigated two time points during the lifespan of these animals and therefore may be missing important trends that occur over the lives of these animals. In addition, although the sea urchins investigated represent the oldest and youngest in the populations under study, we cannot definitely say that the oldest animals are approaching their maximum limit. Further studies are required to examine sea urchins with different life spans from the same environment to see if the negative correlation between damage and longevity observed in this study is confirmed. The absence of an age-related increase in oxidative damage in animals that lack senescence and low levels of damage in long-lived species are both consistent with the oxidative stress theory of aging. The data from this study support the idea that negligible senescence is accompanied by maintenance of antioxidant and proteasome activities and mitigation of some forms of oxidative damage and provides a foundation to further explore the connections between oxidative damage, negligible senescence and longevity.

Acknowledgments

The authors would like to thank Dan Leus of the Pacific Biological Station, Nanaimo, B.C. and Phil Zerofski of Scripps Institution of Oceanography in San Diego, CA for collecting *S. fransicanus* and *S. purpuratus*, Thomas Ebert at Oregon State University for assistance with age estimates of sea urchins and the Histopathology Services at the Jackson Laboratory, Bar Harbor, Maine for embedding and sectioning sea urchin tissues. This work was supported by grant number R21AG039761 from the National Institute on Aging and financial support from the Canadian Association for the Bermuda Institute of Ocean Sciences and a Bermuda Charitable Trust. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Weinert BT, Timiras PS. Theories of Aging. *J Appl Physiol*. 2003; 95:1706–1716. [PubMed: 12970376]
2. Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol Rev*. 1998; 78:547–581. [PubMed: 9562038]
3. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of aging. *Nature*. 2000; 408:239–247. [PubMed: 11089981]
4. Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol*. 1956; 11:298–300. [PubMed: 13332224]
5. Martin I, Grotewiel MS. Oxidative damage and age-related functional declines. *Mech Ageing Dev*. 2006; 127:411–423. [PubMed: 16527333]
6. Salmon AB, Richardson A, Pérez VI. Update on the oxidative stress theory of aging: Does oxidative stress play a role in aging or healthy aging? *Free Radic Biol Med*. 2010; 48:642–655. [PubMed: 20036736]
7. Pérez VI, Bokov A, Van Remmen H, Mele J, Ran Q, Ikeno Y, Richardson A. Is the oxidative stress theory of aging dead? *Biochim Biophys Acta*. 2009; 1790:1005–1014. [PubMed: 19524016]
8. Finch, CE. *Longevity Senescence and the Genome*. University of Chicago Press; Chicago: 1990.
9. Finch CE, Austad SN. History and prospects: symposium on organisms with slow aging. *Exp Gerontol*. 2001; 36:593–597. [PubMed: 11295501]
10. Ebert TA. Longevity and lack of senescence in the red sea urchin *Strongylocentrotus franciscanus*. *Exp Gerontol*. 2008; 43:734–738. [PubMed: 18550313]
11. Ebert TA, Southon JR. Red sea urchins (*Strongylocentrotus franciscanus*) can live over 100 years: confirmation with A-bomb ¹⁴carbon. *Fishery Bulletin*. 2003; 101(4):915–922.
12. Beddingfield SD, McClintock JB. Demographic Characteristics of *Lytechinus variegatus* (*Echinoidea: Echinodermata*) from three habitats in North Florida Bay, Gulf of Mexico. *Mar Ecol*. 2000; 21:17–40.
13. Moore HB, Jutare T, Bauer JC, Jones JA. The biology of *Lytechinus variegatus*. *Bull Mar Sci Gulf Caribbean*. 1963; 13:23–53.
14. Ebert, TA. Growth and survival of post-settlement sea urchins. In: Lawrence, JM., editor. *Edible Sea Urchins: Biology and Ecology*. 2. Amsterdam: Elsevier; 2007. p. 95-134.
15. Ebert TA. Demographic patterns of the purple sea urchin *Strongylocentrotus purpuratus* along a latitudinal gradient, 1985–1987. *Mar Ecol Prog Ser*. 2010; 406:105–120.
16. Sodergren. (The Sea Urchin Genome Sequencing Consortium) The genome of the sea urchin *Strongylocentrotus purpuratus*. *Science*. 2006; 314:941–952. [PubMed: 17095691]
17. Loram J, Bodnar A. Age-related changes in gene expression in tissues of the sea urchin *Strongylocentrotus purpuratus*. *Mech Ageing Dev*. 2012; 133:338–347. [PubMed: 22475988]
18. Shringarpure R, Davies KJA. Protein turnover by the proteasome in aging and disease. *Free Radic Biol Med*. 2002; 32:1084–1089. [PubMed: 12031893]
19. Kregel KC, Zhang HJ. An integrated view of oxidative stress in aging: basic mechanisms, functional effects, and pathological considerations. *Am J Physiol Regul Integr Comp Physiol*. 2007; 292:R18–R36. [PubMed: 16917020]
20. Loft S, Poulsen HE. Estimation of oxidative DNA damage in man from urinary excretion of repair products. *Acta Biochim Pol*. 1998; 45:133–144. [PubMed: 9701506]
21. Cooke MS, Henderson PT, Evans MD. Sources of extracellular, oxidatively-modified DNA lesions: Implications for their measurement in urine. *J Clin Biochem Nutr*. 2009; 45:255–270. [PubMed: 19902015]
22. Valavanidis A, Vlachogianni T, Fiotakis C. 8-hydroxy-2'-deoxyguanosine (8-OHdG): A critical biomarker of oxidative stress and carcinogenesis. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev*. 2009; 27:120–139. [PubMed: 19412858]
23. Pamplona R, Costantini D. Molecular and structural antioxidant defenses against oxidative stress in animals. *Am J Physiol Regul Integr Comp Physiol*. 2011; 301:R843–R863. [PubMed: 21775650]

24. Kastle M, Grune T. Protein oxidative modification in the aging organism and the role of the ubiquitin proteasomal system. *Curr Pharm Des.* 2011; 17:4007–4022. [PubMed: 22188451]
25. Löw P. The role of ubiquitin-proteasome system in ageing. *Gen Comp Endocr.* 2011; 172:39–43. [PubMed: 21324320]
26. Kiffin R, Bandyopadhyay U, Cuervo AM. Oxidative stress and autophagy. *Antioxid Redox Sign.* 2005; 8:152–162.
27. Sulzer D, Mosharov E, Talloczy Z, Zucca FA, Simon JD, Zecca L. Neuronal pigmented autophagic vacuoles: lipofuscin, neuromelanin, and ceroid as macroautophagic responses during aging and disease. *J Neurochem.* 2008; 106:24–36. [PubMed: 18384642]
28. Coleman, DA. MS thesis. San Diego State University; CA: 1993. Heterozygosity and growth rate and gonad weight in the purple sea urchin *Strongylocentrotus purpuratus*.
29. Ebert, TA. In: Mooi, R.; Telford, M., editors. An analysis of the importance of allee effects in management of the red sea urchin *Strongylocentrotus franciscanus*; Echinoderms: San Francisco. Proceedings of the ninth international echinoderm conference; Rotterdam: A.A Balkema; 1998. p. 619-627.
30. Ebert TA, Dixon JD, Schroeter SC, Kalvass PE, Richmond NT, Bradbury WA, Woodby DA. Growth and mortality of red sea urchins *Strongylocentrotus franciscanus* across a latitudinal gradient. *Mar Ecol Prog Ser.* 1999; 190:189–209.
31. Strahl J, Philipp E, Brey T, Broeg K, Abele D. Physiological aging in the Icelandic population of the ocean quahog *Arctica islandica*. *Aquat Biol.* 2007; 1:77–83.
32. Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976; 72:248–254. [PubMed: 942051]
33. Lesser M. Oxidative stress in marine environments: Biochemistry and physiological ecology. *Ann Rev Physiol.* 2006; 68:253–278. [PubMed: 16460273]
34. Tully O, O'Dovovan V, Fletcher D. Metabolic rate and lipofuscin accumulation in juvenile European lobster (*Homarus gammarus*) in relation to simulated seasonal changes in temperature. *Mar Biol.* 2000; 137:1031–1040.
35. Ungvari Z, Ridgway I, Philipp EER, Campbell CM, McQuary P, Chow T, Coelho M, Didier ES, Gelino S, Holmbeck MA, Kim I, Levy E, Sosnowska D, Sonntag WE, Austad SN, Csiszar A. Extreme longevity is associated with increased resistance to oxidative stress in *Arctica islandica*, the longest-living non-colonial animal. *J Gerontol A Biol Sci Med Sci.* 2011; 66A(7):741–750. [PubMed: 21486920]
36. Götze S, Bose A, Abele D, Sokolova IM, Saborowski R. Pitfalls in invertebrate proteasome assays. *J Exp Biol.* 2013; 216:1351–1354. [PubMed: 23264486]
37. Stroikin Y, Mild H, Johansson U, Roberg K, Öllinger K. Lysosome-targeted stress reveals increased stability of lipofuscin-containing lysosomes. *AGE.* 2008; 30:31–42. [PubMed: 19424871]
38. Vaschenko MA, Zhadan PM, Aminin DL, Almyashova TN. Lipofuscin-like pigment in gonads of sea urchin *Strongylocentrotus intermedius* as a potential biomarker of marine pollution: A field study. *Arch Environ Contam Toxicol.* 2012; 62:599–613. [PubMed: 22138826]
39. Husman G, Abele D, Monien D, Monien P, Kriews M, Philipp EER. The influence of sedimentation on metal accumulation and cellular oxidative stress markers in the Antarctic bivalve *Laternula elliptica*. *Estuar Coast Shelf S.* 2012; 111:48–59.
40. Maxwell KE, Matthews TR, Sheehy MRJ, Bertelsen RD, Derby CD. Neurolipofuscin is a measure of age in *Panulirus argus*, the Caribbean spiny lobster, in Florida. *Biol Bull.* 2007; 213:55–66. [PubMed: 17679720]
41. Sheehy MRJ, Shelton PMJ, Wickins JF, Belchier M, Gaten E. Ageing the European lobster *Homarus gammarus* by the lipofuscin in its eyestalk ganglia. *Mar Ecol Prog Ser.* 1996; 143:99–111.
42. Uglem I, Belchier M, Svåsand T. Age determination of European Lobsters (*Homarus gammarus* L.) by histological quantification of lipofuscin. *J Crustacean Biol.* 2005; 25:95–99.
43. Fonseca DB, Sheehy MRJ, Blackman N, Shelton PMJ, Prior AE. Reversal of a hallmark of brain ageing: lipofuscin accumulation. *Neurobiol Aging.* 2005; 26:69–76. [PubMed: 15585347]

Abbreviations

8-OHdG	8-hydroxy-2'-deoxyguanosine
HNE	4-hydroxynonenal
ROS	reactive oxygen species
SOD	superoxide dismutase
TAC	total antioxidant capacity

Highlights

- sea urchins are novel model animals for studying aging and negligible senescence
- cellular oxidative damage does not increase with age in most sea urchin tissues
- antioxidant and proteasome activities are maintained with age in sea urchin tissues
- maintenance of defense and mitigation of damage accompanies negligible senescence
- the results are consistent with the oxidative stress theory of aging




Figure 1. 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels in cell-free coelomic fluid of *L. variegatus* (Lv), *S. purpuratus* (Sp) and *S. franciscanus* (Sf). The white bars represent the average 8-OHdG values for the young sea urchins and the black bars represent levels for the old sea urchins. All values are presented as mean \pm standard error of the mean.




Figure 2.

Protein carbonyl content in tissue extracts from the three sea urchin species: *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). The tissues assayed were muscle (Alm), esophagus (Es), radial nerve (RN), gonad (Gon), coelomocytes (Coel) and ampullae (Amp). The white bars represent the average protein carbonyl values for the young sea urchins and the black bars represent levels for the old sea urchins (expressed as nmole per mg protein). All values are mean \pm standard error of the mean. The stars indicate that the protein carbonyl levels were significantly different between the young and old sea urchins ($p < 0.05$).




Figure 3. HNE-Histidine adduct levels in tissue extracts from the three sea urchin species: *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). The tissues assayed were muscle (Alm), esophagus (Es), radial nerve (RN), gonad (Gon), coelomocytes (Coel) and ampullae (Amp). The white bars represent the average HNE-Histidine values for the young sea urchins and the black bars represent levels for the old sea urchins (expressed as µg per µg protein). All values are mean ± standard error of the mean. The stars indicate that the HNE-His adduct levels were significantly different between the young and old sea urchins ($p < 0.05$).




Figure 4.

Total antioxidant capacity (TAC) in tissue extracts from the three sea urchin species: *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). Panels A, B and C show the results for muscle (Alm), esophagus (Es), radial nerve (RN), gonad (Gon) and ampullae (Amp). Panel D shows the results for coelomocytes from the three species. The white bars represent the average TAC values for the young sea urchins and the black bars represent levels for the old sea urchins. TAC is expressed as μM Copper Reducing Equivalents (CRE) per μg of input protein for each tissue extract. All values are mean \pm standard error of the mean. The stars indicate that the TAC levels were significantly different between the young and old sea urchins ($p < 0.05$).




Figure 5. Superoxide dismutase (SOD) activity in tissue extracts from the three sea urchin species: *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). The tissues assayed were muscle (Alm), esophagus (Es), radial nerve (RN), gonad (Gon), coelomocytes (Coel) and ampullae (Amp). The white bars represent the average SOD values for the young sea urchins and the black bars represent levels for the old sea urchins. SOD activity is expressed as activity units (AU) per μg protein in each tissue extract. All values are mean \pm standard error of the mean. The stars indicate that the SOD levels were significantly different between the young and old sea urchins ($p < 0.05$).




Figure 6. Proteasome enzyme activities in tissue extracts from the three sea urchin species: *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). The proteasome enzyme activities [trypsin-like (Try), chymotrypsin-like (Chy) and caspase-like (Cas)] are shown for muscle (Alm), esophagus (Es), radial nerve (RN), gonad (Gon) and ampullae (Amp). The white bars represent the average enzyme activity for the young sea urchins and the black bars represent levels for the old sea urchins. Enzyme activity is expressed as relative light units (RLU) per mg protein in each tissue extract. All values are mean \pm standard error of the mean. The star indicates that the enzyme activity levels were significantly different between the young and old sea urchins ($p < 0.05$).




Figure 7. Autofluorescence levels in muscle (Alm), esophagus (Es) and radial nerve (RN) tissue of *L. variegatus* (Panel A), *S. purpuratus* (Panel B) and *S. franciscanus* (Panel C). Autofluorescence is expressed as Integrated Optical Density (IOD) per tissue area. The white bars represent the average autofluorescence for the young sea urchins and the black bars represent levels for the old sea urchins. All values are mean \pm standard error of the mean. The stars indicate that the autofluorescence levels were significantly different between the young and old sea urchins ($p < 0.05$).

Table 1

Estimated ages of young and old *L. variegatus*, *S. purpuratus* and *S. franciscanus* used for oxidative damage, antioxidant, proteasome and lipofuscin analyses.

Sample	Test Diameter (mm)	Est. Age (years)	Reference
Oxidative damage, antioxidant, proteasome assays:			
Lv-Young	40.2 ± 0.3 (6)	1.3 ± 0.1 (6)	[12]
Lv-Old	73.2 ± 0.4 (6)	4.0 ± 0.1 (6)	
Sp-Young	32.2 ± 0.8 (6)	1.9 ± 0.1 (6)	[28]
Sp-Old	73.7 ± 1.7 (6)	41.0 ± 9.1 (6)	
Sf ^a -Young	48.2 ± 4.1 (6)	3.7 ± 0.2 (6)	[29,30]
Sf ^a -Old	159.5 ± 1.3 (6)	128.1 ± 7.4 (6)	
Sf ^b -Young	34.5 ± 0.1 (7)	3.1 ± 0.1 (7)	[29,30]
Sf ^b -Old	159.0 ± 0.1 (5)	124.7 ± 6.6 (5)	
Lipofuscin:			
Lv-Young	40.8 ± 0.4 (6)	1.4 ± 0.1 (6)	[12]
Lv-Old	73.2 ± 0.4 (6)	4.0 ± 0.1 (6)	
Sp-Young	28.7 ± 1.1 (6)	1.7 ± 0.1 (6)	[28]
Sp-Old	73.7 ± 1.7 (6)	41.0 ± 9.1 (6)	
Sf ^b -Young	39.4 ± 0.2 (7)	3.3 ± 0.1 (7)	[29,30]
Sf ^b -Old	159.0 ± 0.1 (5)	124.7 ± 6.6 (5)	

The estimated ages are expressed as mean ± standard error and the numbers in brackets represent the number of samples (n) in each group

^a *S. franciscanus* collected in July 2011, used for oxidative damage and antioxidant analyses

^b *S. franciscanus* collected in July 2012, used for proteasome and lipofuscin analyses