



Community-Partnered Evaluation of Depression Services for Clients of Community-Based Agencies in Under-Resourced Communities in Los Angeles

Jeanne Miranda, PhD^{1,2,3,12}, Michael K. Ong, MD, PhD⁴, Loretta Jones, MA^{5,6}, Bowen Chung, MD^{1,2,7,8}, Elizabeth L. Dixon, RN, PhD⁹, Lingqi Tang, PhD^{1,2,3}, Jim Gilmore, MBA¹⁰, Cathy Sherbourne, PhD⁷, Victoria K. Ngo, PhD⁷, Susan Stockdale, PhD^{1,2,11}, Esmeralda Ramos, BA^{1,2}, Thomas R. Belin, PhD^{1,2,12}, and Kenneth B. Wells, MD, MPH^{1,2,3,7,12}

¹Department of Psychiatry, David Geffen School of Medicine, UCLA, Los Angeles, USA; ²Department of Biobehavioral Sciences, David Geffen School of Medicine, UCLA, Los Angeles, USA; ³UCLA Jane and Terry Semel Institute for Neuroscience and Human Behavior, Center for Health Services and Society, Los Angeles, USA; ⁴UCLA Division of General Internal Medicine and Health Services Research, Department of Medicine, David Geffen School of Medicine, Los Angeles, USA; ⁵Healthy African American Families II, Los Angeles, USA; ⁶Charles R. Drew University of Medicine and Science, Los Angeles, USA; ⁷RAND Corporation, Los Angeles, USA; ⁸Harbor-UCLA Medical Center, Los Angeles, USA; ⁹QueensCare Health and Faith Partnership, Los Angeles, USA; ¹⁰Behavioral Health Services, Los Angeles, USA; ¹¹Greater Los Angeles Veteran's Affairs, Los Angeles, USA; ¹²Department of Health Services, UCLA Jonathan and Karin Fielding School of Public Health, Los Angeles, USA.

BACKGROUND: As medical homes are developing under health reform, little is known regarding depression services need and use by diverse safety-net populations in under-resourced communities. For chronic conditions like depression, primary care services may face new opportunities to partner with diverse community service providers, such as those in social service and substance abuse centers, to support a collaborative care model of treating depression.

OBJECTIVE: To understand the distribution of need and current burden of services for depression in under-resourced, diverse communities in Los Angeles.

DESIGN: Baseline phase of a participatory trial to improve depression services with data from client screening and follow-up surveys.

PARTICIPANTS: Of 4,440 clients screened from 93 programs (primary care, mental health, substance abuse, homeless, social and other community services) in 50 agencies, 1,322 were depressed according to an eight-item Patient Health Questionnaire (PHQ-8) and gave contact information; 1,246 enrolled and 981 completed surveys. Ninety-three programs, including 17 primary care/public health, 18 mental health, 20 substance abuse, ten homeless services, and 28 social/other community services, participated.

MAIN MEASURES: Comparisons by setting in 6-month retrospective recall of depression services use.

KEY RESULTS: Depression prevalence ranged from 51.9 % in mental health to 17.2 % in social-community programs. Depressed clients used two settings on average to receive depression services; 82 % used any

setting. More clients preferred counseling over medication for depression treatment.

CONCLUSIONS: Need for depression care was high, and a broad range of agencies provide depression care. Although most participants had contact with primary care, most depression services occurred outside of primary care settings, emphasizing the need to coordinate and support the quality of community-based services across diverse community settings.

KEY WORDS: depression services; community-partnered; participatory research; CPPR; CBPR; community-based; under-resourced.

J Gen Intern Med 28(10):1279-87

DOI: 10.1007/s11606-013-2480-7

© Society of General Internal Medicine 2013

BACKGROUND

Current healthcare reforms call for expansions of primary care and integrated medical homes for safety-net populations.^{1,2} Primary care is a “de facto” mental health setting,³ where rates of recognition and treatment for depression are lower than in specialty mental health settings.⁴ While collaborative care is effective for primary care depressed patients,⁵⁻⁷ little is known about how to conceptualize depression management in neighborhoods with high rates of poverty and ethnic minority representation.⁸

Neighborhood poverty is predictive of depression onset,⁹ and while prevalence of depressive disorders is similar for Latinos and African Americans relative to non-Hispanic whites,^{10,11} African Americans may have greater severity when depressed.¹² Ethnic and socioeconomic disparities in

Received December 10, 2012

Revised February 27, 2013

Accepted April 18, 2013

Published online May 14, 2013

access to and quality of mental healthcare exist,^{13,14} and under-served communities with low provider availability may rely on alternative settings, such as faith-based^{15,16} or substance abuse programs.^{17,18} A recent Institute of Medicine report suggests that integrating health and non-health services settings is essential to address chronic health needs.¹⁹ To inform such efforts, this study evaluates the distribution of depression services across diverse health, social and other community-based service settings in two safety-net communities, providing critical information for primary care practices developing community partnerships in an era of medical homes, insurance reform, and parity legislation.¹

Community-based participatory research (CBPR) is recommended for addressing health disparities.^{20–23} While CBPR studies recommend improving depression services by including social and community services settings,^{24–27} we know of no application of CBPR to describe the depression services received by clients across service settings in under-resourced communities. To do so, we use baseline data from a community-partnered, participatory randomized trial that screened clients for depression in healthcare and non-healthcare settings. We used CBPR approaches to identify and recruit a much broader range of settings into this study than would have occurred through traditional methods. We hypothesized that 1) non-healthcare settings would play a substantial role in serving depressed clients,²⁸ posing challenges to coordination; but that 2) most clients across service settings would have at least some access to primary care, which could serve to coordinate care. Given the stigma associated with depression and mental healthcare,²⁹ we sought to identify client attitudinal barriers and facilitators for using depression services in health and non-health related settings.

METHODS

The study uses baseline data from Community Partners in Care (CPIC),³⁰ a community-partnered participatory research (CPPR) initiative to improve depression services in Los Angeles, using a randomized participatory public health demonstration approach.³¹ CPPR is manualized and gives community and academic partners equal authority to develop and evaluate programs through two-way knowledge exchange.³² CPIC was designed and implemented by the CPIC Council of 35 leaders from three academic and 24 community-based agencies. This study was approved by the RAND institutional review board.

Sampling

Communities. South Los Angeles (South LA) and Hollywood-Metropolitan (Hollywood) were selected based

on established partnerships.^{24,33} South LA has a population of 1.5 million (63.3 % Hispanic, 32.4 % African American), high rates of morbidity and mortality, and low rates of educational attainment and insurance coverage.³⁴ Hollywood has a half-million population (56.7 % Hispanic, 5.6 % African American, 20.7 % non-Hispanic white, 16.7 % Asian American), with the majority having less than high school education³⁵ and low rates of insurance or regular source of healthcare.³⁴ Council leaders selected these groups for over-sampling: African Americans and substance abuse clients (South LA), and seniors and homeless clients (Hollywood).

Agencies and Programs

County directories were combined with community member nominations to identify agencies within five settings: 1) outpatient primary care and public health (Primary Health); 2) outpatient mental health (Mental Health); 3) substance abuse residential and outpatient (Substance Abuse); 4) homeless social and housing services (Housing); and 5) other social and community-based services (Social), including family preservation, prisoner re-entry, senior centers, hair salons, exercise clubs, and faith-based. Eligible agencies had to provide services for adults or parents of child clients and be financially stable, i.e., not expecting to close during the study time period. The CPIC Council explained the study through community “kick-off” conferences and telephone and site briefings. From 149 agency names, we used a four-stage process to reach agencies; enumerate their programs and determine eligibility; randomize potentially eligible programs; and conduct site visits to finalize program enrollment.³⁶ To be eligible, programs had to serve at least 15 clients per week, have two or more staff (one for senior centers, businesses, and faith-based programs), and be willing to identify a staff liaison. Across recruitment stages, 19 agencies were not reached or lost to follow-up; 33 were ineligible; 47 refused and 50 consented (52 % (50/97) of reached and eligible agencies). These 50 agencies had 122 programs, of which 16 were ineligible, 11 declined and 95 enrolled (89.6 % of eligible programs in enrolled agencies). The Council excluded programs mainly serving persons with psychotic disorders or delivering home-based services, or that were financially unstable. At two programs, no clients showed at screening, leaving 93 programs, including 17 primary care/public health, 18 mental health, 20 substance abuse, ten homeless services, and 28 social/other community service programs. We used census data to compare average household characteristics at zip code level (age, sex, race, population density, income) for programs that did and did not participate, with no significant differences (each p value > 0.10).

Clients

Within programs, clients were screened in waiting rooms from March 2010 to November 2010. Study screenings were conducted by the RAND Survey Research Group. Staff approached 4,645 adults age 18 and older during 2–3 days per program; 4,440 (95.6 %) agreed to screening. Eligibility was based on having a score ≥ 10 on a modified eight-item Patient Health Questionnaire (PHQ-8),³⁷ which has scoring and operational characteristics of the PHQ-9. The modification was counting a response to the original or alternative version of one item with and without the word “depression”; Pearson correlation of the two items was 0.99. Exclusion criteria were under age 18, gross cognitive disorganization by screener staff assessment or providing no contact information. Of 4,440 screened, 1,322 (29.8 %) were eligible and 1,246 (94.3 %) consented. Between April 2010 and January 2011, 981 enrolled depressed participants (78.7 %) completed a baseline telephone survey; two were deceased; 36 refused; and 227 were not reached. The response rate is acceptable for quality improvement studies.^{7,38–40}

Measures

Demographic and Health Status. From the screener, we used data on sociodemographic factors (age, gender, marital status, family income, education, insurance status, employment status), race/ethnicity (any Latino, African American not Latino, non-Hispanic white, and other); and PHQ-8³⁷ score ≥ 10 to indicate moderate to severe depression. From survey data, we categorized chronic conditions as ≥ 3 versus < 3 , out of 18; and derived an indicator of meeting federal family poverty criteria. Using the Mini-International Neuropsychiatric Interview (MINI),⁴¹ we created indicators for 12-month major depressive or dysthymic disorder, and alcohol abuse or use of illicit drugs in 12 months.

Service Use

We used retrospective, self-report data to develop indicators of any service use in the past 6 months for overnight hospital stays for mental health or substance abuse (ADM) issues; emergency room (ER) visits for ADM issues, outpatient visits to mental health providers and self/family groups, calls to mental health hotlines, and use of outpatient primary care or public health clinics, substance abuse or social services programs, parks and community centers, and faith-based and other community locations counting as depression-related visits for which the client reported receiving information, referral, counseling, or medication management for depression or emotional problems. We developed indicators for any use and counts of contacts.

We coded other outpatient contacts (primary care, substance abuse, social services, community centers, homeless, faith-based) as including “depression” if the client reported any provider: a) talked about depression, stress or emotions or gave information like a brochure; b) suggested visiting a specialist or program for depression, stress or emotions; c) suggested taking medication or encouraged the respondent to stay on a treatment plan for depression, stress or emotions; or d) spent > 5 min counseling about these issues, or gave suggestions about how to cope or encouragement to do things the respondent enjoyed. Client reports of “other” services for depression were assigned to specific categories using provider names and addresses, internet information and phone calls. For each setting, we measured any outpatient service use, visit counts, and proportion of visits mental health-related, for five sectors of outpatient services: 1) Primary Health; 2) specialty Mental Health; 3) self-help or family support groups for people with emotional or mental health problems (Self Help); 4) Substance Abuse including self-help meetings; and 5) other Social services. We developed an indicator of any mental health-related outpatient use and a count of settings visited.

Treatment Acceptability

We created binary indicators for acceptable/not acceptable for use of anti-depressant drugs; one-on-one counseling from a mental health specialist; and waiting to get over feelings of depression naturally.

Analyses

We conducted univariate analyses to describe the sample and bivariate analyses to compare types of screening locations (Primary Health; Mental Health; Substance Abuse; Housing; Social) for sociodemographic factors and probable depression; and among depressed clients at baseline, in clinical characteristics, services utilization, and satisfaction and treatment acceptability. We present means or percentages with standard errors and significance based on Chi-square tests from bivariate analyses. We account for intra-class correlation within program using SUDAAN 10.0.⁴² To control for potential response bias, attrition weights were constructed by fitting logistic regression models stratified by intervention condition to predict enrollment status and baseline completion from screener predictors (for enrollment: age, community, and screener program category; for baseline: age, sex, ethnic group, family income $< \$10$ K, homeless, community, and screener program category).^{43,44} For item level missing data, we used an extended hot-deck multiple imputation based on the predictive mean matching method.^{45,46} We imputed five data sets, averaged results and adjusted standard errors for

uncertainty due to imputation.⁴⁷ All variables had missingness rates of less than 5 %, except for income and MINI variables, which had 10–15 %.

We conducted three sensitivity analyses: 1) multiple imputation for missing surveys to the full eligible sample (1,322); 2) raw data; and 3) sex and age adjustment. Conclusions from the imputed main model were nearly identical with raw data and sex-adjustment; several findings were not significant with multiple imputation to the eligible sample owing to reduced precision; we focus on the main model and show exact *p* values. We use *p* < 0.05 to denote significance, but focus on the overall pattern of results.

RESULTS

Sociodemographic Characteristics. All sociodemographic characteristics differed significantly by screening program location at *p* < 0.05 (Table 1). While the sample overall is

half from each community, those from South LA varied from 85.9 % in Substance Abuse programs (a sampling priority for South LA) to 24.8 % Housing programs (a priority for Hollywood). Average age was highest in social community (Social). While the majority of women participants were in Mental Health, Housing and Social programs, the majority of men were in Primary Health and Substance Abuse. Those married were highest (38.0 %) in Social and lowest (14.8 %) in Housing programs. Those with less than a high school education ranged from 47.6 % in Mental Health to 27.8 % in Housing. Those clients who self-identified as Latino were 51.4–52.0 % across Primary Health, Mental Health and Social programs, but 23.1–36.2 % in Housing and Substance Abuse, where the majority (54.1–59.1 %) was African American. Overall, 22.5 % of clients worked. Those uninsured ranged from 61.2 % in Primary Health to 28.6–33.2 % in Social and Mental Health. Overall, 64.5 % of clients had less than \$10 K annual family income, with 78.4 % in Housing to 48.4 % in Social programs. The prevalence of probable moderate to severe depression (PHQ-8 score ≥ 10) was

Table 1. Characteristics of Clients Screened for Depression by Program Location

Characteristic	Overall (N=4440)	Primary Care (N=1399)	Mental Health (N=458)	Substance Abuse (N=799)	Homeless Services (N=656)	Social Community Services (N=1128)	χ^2 (df) [*]	<i>P</i>
	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)		
South Los Angeles (%)	49.9 (7.0)	35.3 (14.4)	60.5 (15.4)	85.9 (8.1)	24.8 (17.3)	52.7 (13.2)	9.5 (4)	0.049
Mean Age (years)	46.6 (1.4)	43.7 (1.1)	42.3 (1.5)	41.5 (2.2)	46.5 (1.2)	55.5 (3.9)	15.5 (4)	0.004
Female (%)	53.8 (3.7)	41.8 (7.4)	69.9 (4.8)	38.7 (5.6)	62.3 (11.1)	67.7 (3.3)	28.0 (4)	< 0.001
Married or living with partner (%)	28.4 (2.3)	30.3 (5.1)	27.8 (4.1)	23.0 (1.7)	14.8 (3.2)	38.0 (5.0)	18.0 (4)	0.001
Less than high school (%)	38.8 (2.1)	38.9 (4.2)	47.6 (3.6)	39.6 (2.2)	27.8 (4.8)	41.0 (4.8)	9.9 (4)	0.041
Race/Ethnicity (%)							55.1 (12)	< 0.001
Latino	44.6 (3.5)	51.7 (6.9)	52.0 (6.6)	36.2 (6.6)	23.1 (2.5)	51.4 (7.3)	NA	
African American	40.3 (3.6)	33.8 (6.2)	39.5 (5.6)	54.1 (7.5)	59.1 (5.6)	27.8 (7.0)	NA	
Non-Hispanic white	10.6 (1.7)	9.9 (2.6)	6.1 (1.8)	6.1 (1.6)	11.5 (2.6)	16.0 (5.0)	NA	
Other ethnicity	4.5 (0.7)	4.5 (1.0)	2.4 (0.8)	3.5 (0.7)	6.3 (2.0)	4.8 (1.8)	NA	
Working status (%)							50.7 (8)	< 0.001
Working	22.5 (2.0)	26.4 (3.3)	21.0 (4.2)	15.5 (3.2)	13.7 (3.1)	28.3 (4.6)	NA	
Unemployed, in work force	27.7 (1.7)	30.5 (2.4)	25.7 (2.0)	38.2 (2.9)	33.5 (3.7)	14.4 (2.5)	NA	
Unemployed, not in work force	49.8 (2.3)	43.2 (2.8)	53.3 (4.8)	46.3 (4.4)	52.8 (4.4)	57.3 (5.8)	NA	
No health insurance (%)	49.6 (2.8)	61.2 (2.4)	33.2 (5.9)	62.7 (5.6)	56.2 (4.3)	28.6 (5.7)	35.5 (4)	< 0.001
Family income from work ≤ \$10,000/year (%)	64.5 (2.1)	66.8 (4.4)	68.2 (6.0)	70.0 (2.4)	78.4 (1.9)	48.4 (3.7)	53.0 (4)	< 0.001
Probable depression [†] (%)	33.2 (1.6)	35.3 (2.4)	51.7 (3.6)	34.9 (3.1)	39.0 (1.5)	18.5 (2.1)	67.1 (4)	< 0.001
Probable depression [‡] (%) age-sex adjusted		36.2 (2.5)	51.9 (3.7)	36.3 (3.0)	38.7 (1.8)	17.2 (2.2)	15.2 (4)	< 0.001

Multiple imputation at the item level; sample completing screener

^{*}To compare differences across five service settings, Chi-square tests were calculated using SUDAAN software and take into account clustering (clients within programs)

[†]PHQ-8 (modified) ≥ 10

[‡]Estimates and test statistic for comparison of differences across five service settings, adjusted for sex and age

33.2 %, varying from 51.7 % in Mental Health to 18.5 % in Social programs, and was not substantially altered by sex-age adjustment ($\chi^2=15.2, p < 0.001$).

Health and Mental Health Conditions. As shown in Table 2, over half of depressed clients (54.9 %) reported three or more chronic medical conditions, from 64.4 % in Housing to 46.7 % in Substance Abuse programs. Those in poverty ranged from 80.8 % in Housing to 57.6 % in Social programs. Mean physical and mental health quality of life scores were about 40 (not shown in Table 2), a standard deviation (SD) below national norms.³⁷ Those with 12-month

depressive disorder was 61.9 %, ranging from 47.2 % in Social to 66.5–71.0 % in Housing and Mental Health programs. 12-month alcohol abuse or illicit drug use affected 19.9 % in Social but 71.0 % in Substance Abuse programs.

Service Use. Overall, 69.2 % had a Primary Health visit, varying from 76.9 % for those screened in Primary Health programs to 54.3 % in Substance Abuse programs (some screened in Primary Health accompanied someone and did not have a visit). Those having a mental health-related Primary Health visit, 41.7 % overall, did not differ significantly by program type. Those with a Mental Health visit varied from

Table 2. Depressed Client Sample Characteristics and Past 6-Month Self-Report of Service Use for Alcohol, Drug, or Mental Health (ADM) or Depression*

Background Characteristics	Overall (N=981)	Primary Care (N=278)	Mental Health (N=187)	Substance Abuse (SA) (N=223)	Homeless Services (HO) (N=159)	Social-Community (SC) (N=134)	$\chi^2(df=4)^{\dagger}$	P value
	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)	% or mean (SE)		
Age, Mean	45.7 (0.9)	46.0 (0.8)	42.9 (1.7)	42.3 (1.9)	47.7 (0.9)	51.4 (3.7)	12.5	0.014
Female (%)	56.9 (3.5)	48.7 (6.3)	72.6 (4.7)	45.3 (6.9)	62.1 (10.3)	66.1 (5.4)	15.0	0.005
≥ 3 chronic conditions (%)	54.9 (2.7)	56.2 (5.2)	55.4 (5.0)	46.7 (6.3)	64.4 (5.4)	53.1 (5.5)	4.7	0.32
Income < federal poverty (%)	73.6 (2.1)	75.9 (4.1)	73.9 (3.8)	74.9 (4.5)	80.8 (3.1)	57.6 (6.4)	10.6	0.032
12-month depressive disorder (%)	62.0 (2.3)	58.2 (5.0)	71.6 (3.5)	66.0 (3.6)	65.8 (5.3)	47.5 (3.8)	25.2	< 0.001
Alcohol abuse or use of illicit drugs 12 months (%)	39.1 (3.3)	36.4 (5.0)	25.9 (4.1)	71.0 (4.0)	32.2 (5.8)	19.9 (5.0)	66.6	< 0.001
Services use								
Any primary care or public health visit	69.2 (2.5)	76.9 (3.5)	71.0 (2.6)	54.3 (5.9)	69.3 (3.0)	73.7 (3.6)	12.2	0.016
Primary care or public health visit with Mental Health service (%)	41.7 (2.4)	45.7 (4.3)	47.2 (4.5)	31.2 (4.6)	43.0 (4.7)	40.7 (5.3)	6.9	0.14
Outpatient Mental Health (%)	57.5 (3.5)	52.2 (7.1)	80.6 (4.4)	57.6 (3.6)	64.8 (8.8)	30.2 (4.1)	49.8	< 0.001
# of outpatient Mental Health visits if any (N=568), Mean	13.4 (1.0)	10.2 (1.2)	16.9 (2.2)	16.5 (2.7)	11.5 (1.7)	9.0 (1.3)	14.5	0.006
Mental Health, Self-Help (%)	18.9 (2.3)	13.6 (3.2)	36.1 (7.4)	19.5 (2.0)	17.6 (4.4)	8.4 (2.9)	15.7	0.003
Any substance abuse visits	33.1 (4.2)	25.9 (6.2)	14.2 (4.1)	78.4 (6.3)	28.8 (6.5)	7.1 (2.2)	65.2	< 0.001
Substance Abuse visit with Mental Health service (%)	24.0 (3.0)	19.4 (4.1)	12.0 (3.3)	57.1 (5.1)	18.3 (4.0)	4.0 (1.9)	71.8	< 0.001
Any Social-Community services visits	85.6 (1.3)	84.5 (2.7)	84.2 (2.8)	84.8 (2.8)	88.4 (2.6)	88.0 (3.3)	2.0	0.74
Social-Community services visit with Mental Health service (%)	47.0 (2.0)	44.7 (3.3)	44.2 (3.1)	50.7 (4.5)	56.8 (5.3)	38.2 (4.0)	9.2	0.057
Any outpatient Mental Health service (%)	81.9 (2.3)	76.6 (4.7)	94.1 (1.8)	87.1 (3.0)	83.2 (5.1)	67.6 (4.9)	31.3	<0.001
# outpatient settings visited for Mental Health service [‡] , Mean	2.0 (0.1)	1.8 (0.2)	2.3 (0.2)	2.3 (0.1)	2.1 (0.2)	1.3 (0.1)	76.2	< 0.001
Overnight hospital stay for alcohol, drug or mental health (ADM) (%)	13.8 (1.3)	9.9 (1.7)	14.6 (2.4)	20.1 (3.0)	20.0 (3.7)	4.2 (1.3)	29.4	< 0.001
ER visit for ADM (%)	26.5 (1.9)	23.2 (3.1)	29.3 (5.0)	34.5 (3.5)	30.9 (4.0)	12.7 (2.7)	22.8	< 0.001

*Multiple imputation at the item level, weighted to characteristics of eligible sample (for enrollment offer and baseline response)

[†]To compare differences across five service settings, Chi-square tests were calculated using SUDAAN software and take into account clustering (clients within programs), and weighting

[‡]Five settings: 1) primary care or public health; 2) outpatient specialty mental health; 3) mental health self-help group; 4) substance abuse including self-help; 5) other social-community services

Table 3. Total Contacts for Depression/Mental Health in 6 Months, If Any Contact (Baseline Depressed Clients)* by Type of Program for Screening and Setting for Services Received*

Contacts	Type of Program for Screening																		
	Overall		Primary Care		Mental Health		Substance Abuse		Homeless		Social								
	N	%	N	%	N	%	N	%	N	%	N	%							
Total	808	29,072	100	7.7	212	7,130	100	176	6,826	100	194	9,031	100	133	4,919	100	93	1,167	100
Primary Care or ER	509	2,229	7.7	142	695	9.8	112	433	6.3	103	546	6.0	91	363	7.4	61	192	16.5	
Mental Health Specialty	568	7,728	26.6	145	1,474	20.7	151	2,552	37.4	129	2,145	23.8	103	1,194	24.3	41	363	31.1	
Mental Health, Self-help or family support group	190	5,085	17.5	37	916	12.8	69	2,185	32.0	43	1,232	13.6	29	654	13.3	11	98	8.4	
Substance Abuse	238	7,739	26.6	52	2,108	29.6	22	629	9.2	128	4,007	44.4	30	931	18.9	5	64	5.5	
Social or community services	462	6,291	21.6	123	1,936	27.2	83	1,027	15.1	113	1,101	12.2	90	1,776	36.1	53	450	38.6	

*Excludes overnight stays in a hospital or residential substance abuse programs, but includes services in homeless housing programs. Substance abuse services include self-help for substance abuse, but Mental Health specialty excludes self-help for mental health. Social or community services includes social services such as family preservation or prisoner reentry services, faith-based services, use of parks and recreation community centers, telephone hotlines, and personal services such as hair salons or exercise clubs.
 N Number of clients who reported one or more contacts of that type; Sum total number of contacts; ER emergency room;

80.6 % for those screened in Mental Health to 30.2 % in Social programs. Those with a Self-Help visit varied from 8.4 % in Social to 36.1 % in Mental Health. Those with a Substance Abuse visit varied from 78.4 % in Substance Abuse to 7.1 % in Social programs, and with a mental health-related Substance Abuse visit varied from 57.1 % in Substance Abuse to 4.0 % in Social programs. Those with a Social visit were 85.6 %, and with a mental health-related Social visit were 47.0 %; these percentages did not vary significantly by program type. The percent with any mental health-related visit in any setting varied from a low of 67.6 % in Social to 94.1 % in Mental Health. The 808 users with any mental health-related visit had a total of 29,162 visits within 6 months, of which 7.6 % were Primary Health, 26.4 % Mental Health, 17.3 % in Mental Health support group, 26.5 % in Substance Abuse treatment or self-help, and 22.2 % in other Social services. The mean number of settings visited for depression varied from 1.3 in Social to 2.3 in Mental Health and Substance Abuse. Those with any ADM hospital visit varied from 20.1 % for Substance Abuse to 5.0 % for Social programs, and those with any ER ADM visit varied from 34.5 % for Substance Abuse to 12.7 % for Social programs. Mean visits for those with any ER ADM use (2.9 [SE=0.2]) did not differ significantly by program type ($\chi^2=5.8, p> 0.05$) (Table 3).

Acceptability of Services. Those viewing one-on-one counseling from a specialist as acceptable ranged from 94.3 % in Mental Health to 85.5 % in Social programs. Those finding anti-depressant medication acceptable ranged from 62.7 to 68.3 % in Substance Abuse, Mental Health and Housing to 45.7–55.4 % in Social and Primary Health. Those finding waiting to get over depression acceptable varied from 59.4 to 61.8 % for Primary Health and Social to 34.2 % in Mental Health programs.

CONCLUSIONS

This “portrait” of depression services in two Los Angeles communities was developed using rigorous CPPR and framed within each community’s views of relevant services for depression.³⁰ We found it feasible to implement a rigorous partnered evaluation of depression services, and had relatively high cooperation among programs and clients in enrolled agencies. In terms of substantive findings, clients had a high prevalence of depression, 33 % overall and 35 % in primary care/public health settings with even higher prevalence in other screening locations; by comparison, depression rates in primary care are typically 6–20 %.^{7,40,48–50} Thus, clients in safety net populations within under-resourced communities have a high burden of depression meriting evaluation. The clients were primarily African American or Latino and most were uninsured, met federal poverty criteria, or were not

employed. Over half of the depressed at baseline had 12-month depressive disorder and moderate to high rates of substance abuse and medical comorbidity.

We found that depressed clients reported use of multiple types of mental health-related services in 6 months. Over 40 % of clients reported primary care/public health visits for mental health/depression and nearly 70 % reported contact with primary care/public health services. However, primary care/public health settings had only 8 % of the total volume of depression contacts reported; most depressed clients reported that they received most of their depression services elsewhere, with an average of two settings. With only one-quarter of depression services reported to be in mental health specialty care, most contacts, 67 %, were outside of healthcare settings. These findings suggest that primary care/public health services are appropriate for organizing depression care through partnering with a broader set of programs.

The majority of clients viewed individual counseling as acceptable, and as expected from prior studies,²⁹ antidepressant medication was acceptable to over half of clients. The acceptability of watch and wait, a component of “stepped care” for mild depression in primary care, was highest for clients in primary care and social-community programs; however, this strategy requires outcomes monitoring,^{51,52} which may be hard to implement in safety-net programs.⁵³ In developing medical homes for under-served populations, partnering with other community programs for counseling could potentially be acceptable to clients and clinically effective. For example, group cognitive behavioral therapy, provided in substance abuse treatment settings can improve depression and reduce substance abuse.⁵⁴ Although counseling services could be co-located in primary care, given our findings, it may be necessary to coordinate services in any case with multiple settings, raising the issue of whether it is more effective and efficient to support quality of existing alternative settings that are visited anyway or create internal capacities.

Our findings are limited to clients of programs in two under-resourced communities. Our response rates for clients and programs were acceptable (78 to over 90 %), but low to moderate (50 %) for agencies relative to the field. Many studies of quality improvement in specific sites or applying CBPR to health rely on convenience samples,^{7,38,55–61} and use integrated systems or households or settings in government districts.^{39,49,62–66} The study findings are based on client self-report and may not accurately reflect actual service use due to problems such as client recall. In addition, our findings are limited to financially stable, community-based safety-net programs in Los Angeles.

While the study’s limitations are clear, our CPPR approach supported community involvement in all aspects of the research.⁶⁷ Our findings highlight the extensive provision of depression services in non-healthcare locations, where staff are unlikely to be reimbursed for or receive training in depression services. Exploring how to collaborate to organize

and support diverse settings in addressing depression is an important direction for future work. As we move into healthcare reform, with an emphasis on accountability, efficiency, and patient-centeredness, mechanisms are needed to involve all stakeholders, but especially for vulnerable populations, in the process of understanding the needs of their community. This study represents an approach for working with vulnerable communities in a participatory way to understand community strength and needs for support services for depression. Community input was particularly important in identifying and recruiting the relevant agencies and obtaining high recruitment rates in vulnerable populations.

Acknowledgements: We thank the 25 participating agencies of the Council and their representatives: QueensCare Health and Faith Partnership; COPE Health Solutions; UCLA Center for Health Services and Society; Cal State University Dominguez Hills; RAND; Healthy African American Families II; Los Angeles Urban League; Los Angeles Christian Health Centers; Los Angeles County Department of Mental Health and West Central Mental Health Center; Homeless Outreach Program/Integrated Care System; National Alliance on Mental Illness (NAMI) Urban Los Angeles; Behavioral Health Services, Inc.; Avalon Carver Community Center; USC Keck School of Medicine Department of Psychiatry and Behavioral Sciences; Kaiser Watts Counseling and Learning Center; People Assisting the Homeless; Children’s Bureau; Saban Free Clinic; New Vision Church of Jesus Christ; Jewish Family Services of Los Angeles; St. John’s Well Child and Family Center; Charles Drew University of Medicine and Science; City of Los Angeles Department of Recreation and Parks; To Help Everyone Clinic; QueensCare Family Clinics and the National Institute of Mental Health (funder). We thank the participating Los Angeles programs, their providers and staff, and the clients who participated. We thank the RAND Survey Research Group and trained community members who conducted client data collection. We also thank Robert Brook and Jurgen Unutzer for helpful comments on earlier drafts.

Funding/Support: Community Partners in Care was funded by Award Numbers R01MENTAL HEALTH078853, P30MENTAL HEALTH082760, and P30MENTAL HEALTH068639 from the National Institute of Mental Health, and the Robert Wood Johnson Foundation (64244). The content is the responsibility of the authors and does not necessarily represent the views of the funders.

Conflict of Interest: The authors declare that they do not have a conflict of interest.

Corresponding Author: Jeanne Miranda, PhD; 10920 Wilshire Blvd, Suite 300, Los Angeles, CA 90024, USA (e-mail: jmmiranda@mednet.ucla.edu).

REFERENCES

1. Barry CL, Huskamp HA. Moving beyond parity — mental health and addiction care under the ACA. *New Engl J Med.* 2011;365(11):973–5.
2. Dietrich AJ, Oxman TE, Williams JW, Schulberg HC, Bruce ML, Lee PW, et al. Re-engineering systems for the treatment of depression in primary care: cluster randomised controlled trial. *BMJ.* 2004;329(7466):602.
3. Burns B, Wagner H, Gaynes B, Wells K, Schulberg H. General medical and specialty mental health service use for major depression. *Int J Psychiatry Med.* 2000;30(2):127–43.
4. Wells KB, Sturm R, Sherbourn CD, Meredith LS. *Caring for Depression.* Cambridge, Mass: Harvard University Press; 1996.

5. **Miranda J, Duan N, Sherbourne C, Schoenbaum M, Lagomasino I, Jackson-Triche M, et al.** Improving care for minorities: can quality improvement interventions improve care and outcomes for depressed minorities? Results of a controlled randomized trial. *Health Serv Res.* 2003;38(2):613-30.
6. **Wells K, Sherbourne C, Schoenbaum M, Ettner S, Buan N, Miranda J, et al.** Five-year impact of quality improvement for depression. *Arch Gen Psychiatry.* 2004;61(4):378-86.
7. **Wells KB, Sherbourne C, Schoenbaum M, Duan N, Meredith L, Unutzer J, et al.** Impact of disseminating quality improvement programs for depression in managed primary care: a randomized controlled trial. *JAMA.* 2000;283(2):212-20.
8. **Hadley J, Cunningham P.** Availability of safety net providers and access to care of uninsured persons. *Health Serv Res.* 2004;39(5):1527-46.
9. **Galea S, Ahern J, Nandi A, Tracy M, Beard J, Vlahov D.** Urban neighborhood poverty and the incidence of depression in a population-based cohort study. *Ann Epidemiol.* 2007;17(3):171-9.
10. **Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al.** The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA.* 2003;289(23):3095-105.
11. **Wang PS, Berglund PA, Kessler RC.** Patterns and correlates of contacting clergy for mental disorders in the United States. *Health Serv Res.* 2003;38(2):647-73.
12. **Williams DR, Gonzalez HM, Neighbors H, Nesse R, Abelson JM, Sweetman J, et al.** Prevalence and distribution of major depressive disorder in African Americans, Caribbean blacks, and non-Hispanic whites: results from the National Survey of American Life. *Arch Gen Psychiatry.* 2007;64(3):305-15.
13. **Alegria M, Canino G, Rios R, Vera M, Calderon J, Rusch D, et al.** Inequalities in use of specialty mental health services among Latinos, African Americans, and non-Latino whites. *Psychiatr Serv.* 2002;53(12):1547-55.
14. **Miranda J, Azocar F, Organista KC, Dwyer E, Areane P.** Treatment of depression among impoverished primary care patients from ethnic minority groups. *Psychiatr Serv.* 2003;54(2):219-25.
15. **Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC.** Twelve-month use of mental health services in the United States: results from the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005;62(6):629.
16. **Wang PS, Demler O, Olfson M, Pincus HA, Wells KB, Kessler RC.** Changing profiles of service sectors used for mental health care in the United States. *Am J Psychiatry.* 2006;163(7):1187.
17. **Druss BG, Goldman HH.** New Freedom Commission Report: introduction to the special section on the president's New Freedom Commission Report. *Psychiatr Serv.* 2003;54(11):1465-6.
18. U.S. Department of Health and Human Services. In: Rosenthal RN, ed. *Managing Depressive Symptoms in Substance Abuse Clients During Early Recovery: Treatment Improvement Protocol.* Darby, PA: DIANE Publishing; 2008.
19. Institute of Medicine. *Living Well with Chronic Illness: A Call for Public Action.* Washington, DC: Institute of Medicine of the National Academies; 2012.
20. **Washington AE, Lipstein SH.** The Patient-Centered Outcomes Research Institute - promoting better information, decisions, and health. *New Engl J Med.* 2011;365(15).
21. **Israel BA, Coombe CM, Cheezum RR, Schulz AJ, McGranaghan RJ, Lichtenstein R, et al.** Community-based participatory research: a capacity-building approach for policy advocacy aimed at eliminating health disparities. *Am J Public Health.* 2010;100(11):2094.
22. **Smedley BD, Syme SL.** *Promoting health: intervention strategies from social and behavioral research.* National Academies Press; 2000.
23. **Tunis SR, Stryer DB, Clancy CM.** Practical clinical trials. *JAMA.* 2003;290(12):1624.
24. **Bluthenthal R, Jones L, Fackler-Lowrie N, Ellison M, Booker T, Jones F, et al.** Witness for wellness: preliminary findings from a community-academic participatory research mental health initiative. *Ethn Dis.* 2006;16(Suppl):S18-34.
25. **Chung B, Jones L, Jones A, Corbett C, Booker T, Wells K, et al.** Using community arts events to enhance collective efficacy and community engagement to address depression in an African American community. *Am J Public Health.* 2009;99(2):237.
26. **Dobransky-Fasiska D, Nowalk M, Pincus H, Castillo E, Lee B, Walnoha A, et al.** Public-academic partnerships: improving depression care for disadvantaged adults by partnering with non-mental health agencies. *Psychiatr Serv.* 2010;61(2):110.
27. **Springgate BF, Allen C, Jones C, Lovera S, Meyers D, Campbell L, et al.** Rapid community participatory assessment of health care in post-storm New Orleans. *Am J Prev Med.* 2009;37(6):S237-S43.
28. **Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al.** The epidemiology of major depressive disorder. *JAMA.* 2003;289(23):3095-105.
29. **Dwight Johnson M, Sherbourne CD, Liao D, Wells KB.** Treatment preferences among depressed primary care patients. *J Gen Intern Med.* 2000;15(8):527-34.
30. **Chung B, Dixon EL, Miranda J, Wells K, Jones L.** Using a community-partnered participatory research approach to implement a randomized controlled trial: planning Community Partners in Care. *J Health Care Poor Underserved.* 2010;21(3):780-95.
31. **Katz DL, Murimi M, Gonzalez A, Njike V, Green LW.** From controlled trial to community adoption: the Multisite Translational Community Trial. *Am J Public Health.* 2011;101(8).
32. **Wells K, Jones L.** Commentary: "research" in community-partnered, participatory research. *JAMA.* 2009;302(3):320-1.
33. **Wells KB, Klap R, Fuentes S, Dosssett E.** Brief reports: obstacles and opportunities in providing mental health services through a faith-based network in Los Angeles. *Psychiatr Serv.* 2005;56(2):206-8.
34. Los Angeles County Department of Health Services. *LA County Department of Health Services Key Health Indicators.* Los Angeles, CA; 2009.
35. U.S. Census Bureau. *United States Census 2000.* Washington, D.C.; 2000.
36. **Wells KB, Jones L, Chung B, Dixon E, Tang L, Gilmore J, et al.** Community-Partnered Cluster-Randomized Comparative Effectiveness Trial of Community Engagement and Planning or Program Technical Assistance to Address Depression Disparities. Under review. 2012.
37. **Kroenke K, Strine TW, Spitzer RL, Williams JBW, Berry JT, Mokdad AH.** The PHQ-8 as a measure of current depression in the general population. *J Affect Disord.* 2009;114(1-3):163-73.
38. **Bruce ML, Ten Have TR, Reynolds CF, Katz II, Schulberg HC, Mulsant BH, et al.** Reducing suicidal ideation and depressive symptoms in depressed older primary care patients. *JAMA.* 2004;291(9):1081.
39. **Datto CJ, Thompson R, Horowitz D, Disbot M, Oslin DW.** The pilot study of a telephone disease management program for depression. *Gen Hosp Psychiatry.* 2003;25(3):169-77.
40. **Unutzer J, Katon W, Williams JW Jr, Callahan CM, Harpole L, Hunkeler EM, et al.** Improving primary care for depression in late life: the design of a multicenter randomized trial. *Medical Care.* 2001;39(8):785.
41. **Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al.** The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry.* 1998;59:22-33.
42. RTI International. SUDAAN 10.0. Research Triangle Park, NC. 2012. <http://www.rti.org/SUDAAN/>. Accessed 04/18/2013.
43. **Korn EL, Graubard BI.** *Analysis of Health Surveys.* Hoboken, New Jersey: Wiley-Interscience; 1999.
44. **Groves RM, Dillman D, Eltinge JL, Little RJA.** *Survey Nonresponse.* Hoboken, NJ: Wiley-Interscience; 2002.
45. **Little RJA.** Missing-data adjustments in large surveys. *J Bus Econ Stat.* 1988;287-96.
46. **Wells KB, Sherbourne C, Schoenbaum M, Duan N, Meredith L, Unutzer J, et al.** Impact of disseminating quality improvement programs for depression in managed primary care. *JAMA.* 2000;283(2):212.
47. **Rubin DB.** *Multiple Imputation for Nonresponsive in Surveys.* New York: Wiley; 1987.
48. **Pyne JM, Rost KM, Zhang M, Williams DK, Smith J, Fortney J.** Cost-effectiveness of a primary care depression intervention. *J Gen Intern Med.* 2003;18(6):432-41.
49. **Rost K, Nutting PA, Smith J, Werner JJ.** Designing and implementing a primary care intervention trial to improve the quality and outcome of care for major depression. *Gen Hosp Psychiatry.* 2000;22(2):66-77.
50. **Whooley MA, Stone B, Soghikian K.** Randomized trial of case - finding for depression in elderly primary care patients. *J Gen Intern Med.* 2000;15(5):293-300.
51. **Katon W, VonKorff M, Lin E, Simon G, Walker E, Unutzer J, et al.** Stepped collaborative care for primary care patients with persistent symptoms of depression: a randomized trial. *Arch Gen Psychiatry.* 1999;56(12):1109-15.

52. **Meredith L, Cheng W, Hickey S, Dwight-Johnson M.** Factors associated with primary care clinicians' choice of a watchful waiting approach to managing depression. *Psychiatr Serv.* 2007;58(1):72–8.
53. **Unützer J, Katon W, Callahan CM, Williams JW, Hunkeler E, Harpole L, et al.** Collaborative care management of late-life depression in the primary care setting. *JAMA.* 2002;288(22):2836.
54. **Watkins KE, Hunter SB, Hepner KA, Paddock SM, de la Cruz E, Zhou AJ, et al.** An effectiveness trial of group cognitive behavioral therapy for patients with persistent depressive symptoms in substance abuse treatment. *Arch Gen Psychiatry.* 2011;68(6):577.
55. **Corbie-Smith G, Thomas SB, Williams MV, Moody-Ayers S.** Attitudes and beliefs of African Americans toward participation in medical research. *J Gen Intern Med.* 2001;14(9):537–46.
56. **Eli K, Xie B, Guon B, Quinn DI, Dwight-Johnson M, Lee PJ.** Randomized controlled trial of collaborative care management of depression among low-income patients with cancer. *J Clin Oncol.* 2008;26(27):4488–96.
57. **Blumenthal DS, Smith SA, Majett CD, Alema-Mensah E.** A trial of 3 interventions to promote colorectal cancer screening in African Americans. *Cancer.* 2010;116(4):922–9.
58. **Siegel JM, Prelip ML, Erasquin JT, Kim SA.** A worksite obesity intervention: results from a group-randomized trial. *Am J Public Health.* 2010;100(2):327.
59. **Katzelnick DJ, Simon GE, Pearson SD, Manning WG, Helstad CP, Henk HJ, et al.** Randomized trial of a depression management program in high utilizers of medical care. *Arch Fam Med.* 2000;9(4):345.
60. **Rollman BL, Belnap BH, LeMenager MS, Mazumdar S, Houck PR, Counihan PJ, et al.** Telephone-delivered collaborative care for treating post-CABG depression a randomized controlled trial. *JAMA.* 2009;302(19):2095–103.
61. **Cordasco KM, Asch SM, Bell DS, Guterman JJ, Gross-Schulman S, Ramer L, et al.** A low-literacy medication education tool for safety-net hospital patients. *Am J Prev Med.* 2009;37(6):S209–S16.
62. **Dick J, Clarke M, Van Zyl H, Daniels K.** Primary health care nurses implement and evaluate a community outreach approach to health care in the South African agricultural sector. *Int Nurs Rev.* 2007;54(4):383–90.
63. **Perry CL, Stigler MH, Arora M, Reddy KS.** Prevention in translation: tobacco use prevention in India. *Health Promot Pract.* 2008;9(4):378–86.
64. **Brook O, van Hout H, Nieuwenhuysen H, Heerdink E.** Impact of coaching by community pharmacists on drug attitude of depressive primary care patients and acceptability to patients: a randomized controlled trial. *Eur Neuropsychopharmacol.* 2003;13(1):1–9.
65. **Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, et al.** Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *Lancet.* 2010;376(9758):18–31.
66. **Wilson TE, Feldman J, King G, Coll B, Homel P, Browne R, et al.** Hair salon stylists as breast cancer prevention lay health advisors for African American and Afro-Caribbean women. *J Health Care Poor Underserved.* 2008;19(1):216–26.
67. **Jones L, Wells K.** Strategies for academic and clinician engagement in community-participatory partnered research. *JAMA.* 2007;297(4):407–10.