

Assessment of Transcriptional Activity of *Borrelia burgdorferi* and Host Cytokine Genes During Early and Late Infection in a Mouse Model

Emir Hodzic,^{1,2} Sunlian Feng,¹ and Stephen W. Barthold¹

Abstract

Differential gene expression by *Borrelia burgdorferi* spirochetes during mammalian infection facilitates their dissemination as well as immune evasion. Modulation of gene transcription in response to host immunity has been documented with the outer surface protein C, but the influence of transcription of other genes is largely unknown. A low-density array (LDA) was developed to study transcriptional activity of 43 *B. burgdorferi* genes and 19 host genes that may be involved in various host-agent interactions. Gene transcription in heart, joint, and muscle tissue was compared in immunocompetent C3H and immunodeficient C3H-*scid* mice during early (3 weeks) and late (2 months) *B. burgdorferi* infection. Among all tissue types, levels of relative transcription of over 80% of *B. burgdorferi* genes tested were one- to nine-fold less in C3H mice compared to C3H-*scid* mice. At the later time point, all genes were transcribed in C3H-*scid* mice, whereas transcription of 16 genes out of 43 tested was not detected in analyzed tissues of C3H mice. Our data suggest that during infection of immunocompetent mice, a majority of *B. burgdorferi* genes tested are downregulated in response to acquired host immunity. LDA revealed variable patterns of host gene expression in different tissues and at different intervals in infected mice. Higher levels of relative expression for IL-10 during both early and late infection were detected in heart base, and it was unchanged in the tibiotarsal joint. Comparative analysis of *B. burgdorferi* and host genes transcriptional activity revealed that increased *flaB* mRNA during early infection was followed by increases of CCL7, CCL8, interleukin-10 (IL-10), and tumor necrosis factor- α (TNF- α) in all assessed tissue types. LDA represents a valuable approach for sensitive and quantitative gene transcription profiling and for understanding Lyme borreliosis.

Key Words: *Borrelia burgdorferi*—Cytokine—Transcription—Mouse—Low-density array.

Introduction

THE ETIOLOGIC AGENT OF LYME BORRELIOSIS, *Borrelia burgdorferi*, maintains a complex life cycle in which it must infect, populate, and persist within resting ticks from widely variable environments, feeding ticks that transmit *Borrelia* to new hosts, and within mammalian, avian, and reptilian hosts. This remarkable adaptability is associated with alteration of gene expression in response to different environmental cues, including temperature, pH, and host environment (Pal and Fikrig 2003). In the mammalian host, *B. burgdorferi* is able to persist and effectively evade host immunity (Barthold 2000, Malawista 2000, Bubeck-Martinez 2005, Pal et al. 2008, Palmer, et al. 2009). Dissemination and persistence of *B. burgdorferi* among host tissues are facilitated by its ability to

adhere to a variety of cell types and ligands within the extracellular matrix (ECM) (Backenson et al. 1995, Coleman et al. 1997, Grab et al. 1998, Guo et al. 1998, Leong, et al. 1998, Parveen and Leong 2000, Coburn et al. 2002, Cooke et al. 2005). The functions of many *B. burgdorferi* genes, as well as the *in vivo* cues that modulate their transcription, remain unknown,

Numerous studies have proposed several possible mechanisms of *B. burgdorferi* immune evasion, including binding and invasion of ECM and connective tissue (Liang et al. 2004a, Zambrano et al. 2004), antigenic variation (Bankhead and Chaconas 2007, Palmer et al. 2009), suppression of harmful immune responses (Diterich et al. 2003, Hovis et al. 2006, Bhide et al. 2009), and selective expression of surface antigens (Liang et al. 2002a, Liang et al. 2004a, 2004b). Lipidated spirochetal outer surface proteins are recognized by Toll-like

¹Center for Comparative Medicine, Schools of Veterinary Medicine and Medicine, ²Real-Time PCR Research and Diagnostic Core Facility, Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California at Davis, Davis, California.

receptor 2 (TLR2) (Lien et al. 1999), which leads to activation of monocytes and neutrophils, with subsequent secretion of inflammatory cytokines (Lazarus et al. 2006, Strle et al. 2009, Shin et al. 2010).

Host immune pressure and associated spirochetal immune evasion are dynamic processes, so that many genes are likely to be variably expressed as a consequence of host immunity. In addition, expression of surface antigens may confer tissue specificity and guide dissemination within the host and may therefore be differentially expressed in various tissues and different stages of infection. For example, several ligand-binding lipoproteins, including BmpA outer surface protein and its three paralogous proteins, BmpB, BmpC, and BmpD, play a role in infection by binding to laminin (Verma et al. 2009), and it has been shown that *bmpA*- or *bmpB*-deleted mutants are unable to persist in mouse joint tissues (Pal et al. 2008). Decorin-binding protein (Dbp) A and B bind to tissues rich in decorin, including joints, heart base, and skin. These collagen-rich sites seem to favor persistence of spirochetes with higher *dbpA* expression (Liang et al. 2004a). Deletion of *dbpBA* impedes dissemination of spirochetes and alters long-term survival (Weening et al. 2008). Other ECM ligand-binding outer surface proteins include fibronectin-binding protein (Fbp), *Borrelia* glycosaminoglycan-binding protein (Bgp), RevA, ErpX, P66 ($\alpha_{IIb}\beta_3$ integrin), and an unidentified type I collagen-binding adhesion, among others (Probert and Johnson 1998, Defoe and Coburn 2001, Parveen et al. 2006, Cabello et al. 2007, Brissette et al. 2009a, b).

Pathogen profiling of *B. burgdorferi* obtained by techniques such as serotyping (Wilske et al. 1996), immunohistochemistry (Miklosy et al. 2008), multilocus enzyme electrophoresis (Boerlin et al. 1992), *in situ* hybridization (Fuchs et al. 1992), ribotyping (Postic et al. 1996), pulsed-field gel electrophoresis (Belfaiza et al. 1993), plasmid fingerprinting (Xu and Johnson 1995), randomly amplified polymorphic DNA (Wang et al. 1998), species-specific PCR and PCR-based restriction fragment length polymorphism (RFLP) analysis (Le Fleche et al. 1997, Marconi and Garon 1992), and sequence analysis (Postic et al. 1998) have all been used to help understand the pathogenesis of Lyme borreliosis, but transcriptional activity of selected genes is becoming an essential part of such analysis. Several new molecular techniques such as membrane-based array (Hyde et al. 2006), and microarray (Adusumilli et al. 2010), show promise in that they are objective, quantitative, robust, and allow high throughput.

Advances in gene quantification technologies allow quantitative, sensitive, and robust expression profiling of multiple genes in single samples simultaneously. One such technology, low-density array (LDA), is a medium-throughput method based on a quantitative real-time reverse transcription PCR (qRT-PCR) platform (Abruzzo et al. 2005). LDA allows the simultaneous measurement of expression of up to 384 genes in a single sample with minimal handling of the samples, thereby decreasing contamination and operator-induced errors and allowing standardization of method and data interpretation across laboratories (de Cremoux et al. 2004).

Despite these recent advances in gene transcription detection technology, there has been limited transcriptional analysis of large clusters of *B. burgdorferi* genes (Anderton et al. 2004, Tokarz et al. 2004, Hyde et al. 2006), especially in the infected host (Liang et al. 2002b). In an initial survey to understand the global effect of host immune response on gene

transcription of *B. burgdorferi* during infection in the mouse model, expression profiles of 43 different genes were selected, including those presumed to be involved in attachment, cell envelope, metabolism, complement regulation, cellular processes, and replication. We used a qRT-PCR-based method on the LDA platform to compare transcriptional levels of selected *B. burgdorferi* genes and 19 mouse chemokines and cytokines in mice during early (3 weeks) and late (2 months) infection.

Materials and Methods

Mice

Specific pathogen-free, 3- to 5-week-old C3H/HeN (C3H) mice were purchased from the Frederick Cancer Research Center (Frederick, MD); C3H.C-*Prkdc*^{scid}/*IcrSmnHsd* (C3H-*scid*) mice were purchased from Harlan Sprague-Dawley, Inc. (Indianapolis, IN). The maintenance and care of mice complied with the *Guide for the Care and Use of Laboratory Animals* under an approved University of California Davis Institutional Animal Care and Use protocol. Mice were necropsied following carbon dioxide narcosis and exsanguination by cardiocentesis.

Borrelia burgdorferi

A clonal strain of *B. burgdorferi* sensu stricto (cN40) was grown in modified Barbour-Stoener-Kelly (BSK) II medium (Barbour 1984). Spirochetes were grown to mid-log phase, harvested, and resuspended to 10^9 cells/mL in BSK II medium. Eight C3H mice and eight C3H-*scid* mice were infected by intradermal inoculation at the dorsal thoracic midline with 10^4 spirochetes each and then randomly divided into groups of four mice each. Eight C3H mice were sham-inoculated with BSK II medium and randomly divided into two equal groups. At 3 weeks and 2 months after inoculation, four infected C3H mice, four infected C3H-*scid* mice, and four sham-inoculated C3H mice were necropsied at each time point. To confirm infection, the urinary bladder and the inoculation site were cultured from each mouse, as described previously (Barthold et al. 1993).

Tissue collection and nucleic acid isolation

Heart base, ventricular myocardium, quadriceps muscle, and tibiotarsal joint were collected for analysis. Samples for DNA and cDNA analysis were immediately weighed, snap-frozen in liquid nitrogen, and stored at -80°C before nucleic acid extraction. For isolation of template DNA for quantitative analysis of spirochetes and total RNA to assess transcriptional activity of selected genes, we used ~ 10 mg of tissue, as been described previously (Hodzic et al. 2003).

DNA amplification

For quantitative analysis of DNA isolated from tissue samples, *flaB* qPCR was performed as previously described (Hodzic et al. 2002). Absolute sensitivity and reproducibility of the qPCR was established by limiting dilution assay of cultured spirochetes and of positive tissue samples. The analytical sensitivity was in the range of $1-10^9$ spirochetes, with a yield of detection close to 90% of the calculated amount of known target in each sample. Specificity of the qPCR was confirmed by the lack of amplification from negative control

samples, and amplification was confirmed from positive control samples.

cDNA synthesis

For transcriptional activity analysis of the selected *B. burgdorferi* and host genes, first-strand cDNA was synthesized from total RNA using the QuantiTect Reverse Transcription Kit (Qiagen) in 50- μ L reactions, according to the manufacturer's instructions. Because limited numbers of spirochetes and spirochetal RNA (cDNA) in tissue samples restricted the analytical sensitivity, the Advantage[®] 2 Polymerase Mix (Clontech Laboratories) was used to increase fidelity, efficiency, and greater yield of cDNA, according to the manufacturer's instructions. The preamplification reaction involved activation at 95°C for 15 sec, amplification for 25 cycles at 95°C for 15 sec, 55°C for 15 sec, and 70°C for 45 sec, followed by elongation at 70°C for 5 min. The preamplified products were diluted at a ratio of 1:10 and used as templates for qPCR analysis. All samples were analyzed for the presence of 18S rRNA to determine the efficiency of the nucleic acid extraction, amplification, and as an indicator of inhibition.

LDA assays

LDA assays were designed on the microfluidic card (Applied Biosystems) that contains eight sample-loading ports, each connected by a microchannel to 48 miniature reaction chambers for a total of 384 wells per card. A total of 43 *B. burgdorferi* N40 genes were selected from the National Center for Biotechnology Information (NCBI) GenBank (Schutzer et al. 2011). Genes were selected from different locations throughout the *B. burgdorferi* N40 genome (chromosome, linear plasmids, and circular plasmids), and among 12 different categories according to their putative functions. The selected genes encode proteins with a variety of functions, including attachment, cell envelope, motility, cell division, metabolism, cellular processes (toxin production, resistance, adaptation, response), DNA metabolism, complement regulation, persistence, general functions, protein synthesis, and hypothetical (Table 1).

Another LDA was designed to assess transcriptional activity of host genes, including MyD88 (5'-AGCTGCTGGCCTTGTTAGACC, 3'-TCTGGCAGTCCTCCTCGATG, probe #17 from Roche, Pleasanton, CA), cytokines interleukin-1 β (IL-1 β ; Life Technologies, Grand Island, NY), IL-2 (Life Technologies), IL-4 (Life Technologies), IL-6 (5'-TTCACAAGTCGGA GGCTTAATTACA, 3'-AAGTGCAT CATCGTTGTTCCATA CA, probe FAM-TGAGAAAAGAGTT GTGCAATGGCAAT TCTG), IL-10 (Life technologies), IL-12p40 (Life Technologies), IL-17 α 5'ACTCCCTTG GCGCAAAAGT, 3'-AGGGT CTTTCATTGCGGTGG, probe #50 from Roche), IL-21 (Life Technologies), interferon- γ (IFN- γ) (5'-TGAATAACTATIT TAACCTCAAGTGGCATAG, 3'-ATA ATCTGGCTCTGCAG GATTTTC, probe FAM-TCTTGGATATCTGGAGGAACTG GCAAAAGG), tumor necrosis factor- α (TNF- α) (5'-TGG CCTCCCTCTCATCAGTT, 3'-GCTACAGGCTTGTCCTC GAATT, probe FAM-CCCAGACCCTCAC ACTCAGAT CATCTTCT), and chemokines CCL2 (5'-ACCA GCAAGA TGATCCCAATG, 3'-GAGCTTGGTGACAAAACTACAGC, probe #19 from Roche), CCL7 (5'-GATCTCTGCCACGCTT CTGTG, 3'-GCATTGGGCCATCTGG, probe #89 from Roche), CCL8 (5'-CAACATGAAGATCTACGCAG TGC, 3'-

AGCCTTATCTGGCCCAGTCA, probe #26 from Roche), CCL12 (5'-ACCCCAGTCACGTGCTGTTAT, 3'-TGGTCCTGA AGATCACAGCTTC, probe #2 from Roche), CXCL12 (Life Technologies), CXCL13 (Life Technologies), CCL19 (5'-GGGTGCTAATGATGCGGAAG, 3'-GGTGAACACAACAGC AGGCA, probe # 17 from Roche), and CCL21 (Life Technologies). Gene-specific primers and hydrolysis probes were embedded in each well. The synthesized cDNA template from each sample (30 μ L) was added to 50 μ L of 2 \times Universal PCR Master Mix (Applied Biosystems) in a 100- μ L reaction mixture. The mixture was added to each line of the microfluidic card after vortexing and brief centrifugation. The system allows analysis of four different samples at the same time, in duplicate. The PCR amplifications were performed in the microfluidic card sample block of an ABI Prism 7900HT sequence detection system (Applied Biosystems). The amplification was done under the following thermal cycle conditions: 2 min at 50°C to 10 min at 94.5°C, followed by 40 cycles of denaturation at 97°C for 30 sec, and annealing and extension at 59.7°C for 1 min.

For each of 43 *B. burgdorferi* N40 and 19 mouse target genes, two specific primers and one internal, fluorescence-labeled probe were designed with Primer Express software (Applied Biosystems). The amplification efficiency (E) of all assays was calculated from the slope of a standard curve generated on a 10-fold dilution in triplicate for every cDNA sample using the formula $E = 10^{(-1/\text{slope})} - 1$. To obtain accurate and reproducible results, all assays were determined to have an efficiency of >95%. On the basis of the amplification efficiencies, detection limits were approximately 10 copies of cDNA per reaction. The coefficient of variability of the qPCR determined for 10 replicates was 15% or less. The specificity of each assay was determined using DNA or cDNA templates in duplicate from uninfected mouse tissues and other pathogens (*Leptospira* spp., *Escherichia coli*, *Listeria monocytogenes*, *Staphylococcus aureus*, *Helicobacter* spp.).

Relative quantification of transcriptional activity of each target gene was calculated using the equation $2^{-\Delta\Delta C_q}$. The method uses the transcriptional activity of a target gene based on E and C_q deviation (ΔC_q) of tissue samples from infected C3H mice versus same sample types obtained from infected C3H-*scid* mice, and expressed in comparison to a reference gene (Pfaffl 2001). C_q values of target genes of all samples were normalized to a reference gene (16S rRNA) to compensate sample-to-sample and run-to-run variations and to ensure experimental reliability. Additional calculation was performed using Relative Expression Software Tool (REST[®]), a mathematical model that uses PCR efficiencies and the mean C_q point deviation between experimental and control groups (Pfaffl et al. 2002).

The stability of six potential *B. burgdorferi* reference genes (16S rRNA, *flaB*, *gapDH*, *p13*, *p66*, *dbpA*) was assessed between individual mice and mouse groups, between different tissue samples (heart, quadriceps muscle, and tibiotarsus), and at the different time points by using BestKeeper software (<http://qpcrnews.gene-quantification.info>), and following the guidelines for selection of the best reference gene described by Radonic et al. (2004). To normalize the expression levels of mouse genes, mouse 18S rRNA, β -actin (*Actb*), β_2 -microglobulin (*B2M*), and glyceraldehyde-3-phosphate dehydrogenase (*gapDH*) genes were assessed for transcriptional stability to be used as reference genes.

TABLE 1. LIST OF *B. BURGDORFERI* GENES, OF WHICH TRANSCRIPTIONAL ACTIVITY WAS ASSESSED IN HEART BASE, VENTRICULAR MUSCLE, TIBIOTARSAL JOINT, AND QUADRICEPS MUSCLE OF IMMUNOCOMPETENT C3H AND IMMUNODEFICIENT C3H-*scid* MICE DURING EARLY (2 WEEKS) AND LATE (2 MONTHS) INFECTION^a

Annotation	Accession number	N40 gene symbol	N40 gene location	Forward primer	Reverse primer	Probe
16 ribosomal RNA (16S rRNA)	CP002228.1	BBUN40_165E	Ribosome	GCTTCGCTGTAGATGAGTCTGC	TTCAGTGTGACCCGTTCCACC	TACCAAGGCAATGATAAGTA
Acetate kinase (<i>ackA</i>)	CP002228.1	BBUN40_0622	Chromosome	GAAGCGTGATATAGACCACATCA	AATTCAGATGCCGGCTTTC	TGATGAGC ^a
Alanin racemase (<i>alr</i>)	CP002228.1	BBUN40_0158	Chromosome	CCATACTCTAAITCGGGACACATT	AGAAATA GGCATTCACAGTAAACCGTAA	CAATTTAATAATGTTAAGACCCGGG
Acriflavine resistance protein (<i>acrB</i>)	CP002228.1	BBUN40_0140	Chromosome	TAAGCCTTGGATTACAAAGTTCGG	CCTTCATTAGTGGCAATGCTT	CTCCACCA ^a
Basic membrane protein A (<i>bmpA</i>)	CP002228.1	BBUN40_0383	Chromosome	GGTAGAAGCGTTGCAACTAGAAATG	ATGTAATGCCAGAA ^a CCAAGTTC	ACATTATTCATCATGCTGCAG GCCTTGG
Basic membrane protein B (<i>bmpB</i>)	CP002228.1	BBUN40_0382	Chromosome	CGGTATTTTGTGACTTCTTCTTTAG	TCGTCAAGAACACCATCTACCAA	AGAAATGGAATAGAAATCTAG
Basic membrane protein C (<i>bmpC</i>)	CP002228.1	BBUN40_0384	Chromosome	AAGGGTGAATTAATTAATGATCGG	CCTATTTGTTAAACATCGGGATCC	GGAAAGAG ^a
Basic membrane protein D (<i>bmpD</i>)	CP002228.1	BBUN40_0385	Chromosome	GATACTTGGCTCGAAGGCTTC	GTTTGCATCTTAGCACGACGCTTC	TGGAGGAG ^a
Cell division protein FtsZ	CP002228.1	BBUN40_0299	Chromosome	GATATGGTGTATTACTGCTGGTATGG	CCTGCTCAGCAAGTCTCAACTTC	CAGCTCCAGTTATTGGCCA AGTTGCA
Colicin V production protein (<i>colV</i>)	DQ662922	BBUN40_0766	Chromosome	TAGCGGATTTGCTGAAAGTTTTTG	GGTTCAAACAACCCTTCTAAA TTCTTCAG	TCCTGCTG ^a
Flagellin B (<i>flaB</i>)	CP002228.1	BBUN40_0147	Chromosome	TGCAGCTAATCTTGGAAAATCTTTTTTC	GCAGGTGCTGGGCTGTGTA	AAACTGCTCAGGCTGCACC GGTTC
Glyceraldehyde 3-phosphate dehydrogenase (<i>gapDH</i>)	CP002228.1	BBUN40_0057	Chromosome	GCAAAGCTTGGAAATGATGTTG	TCTCGTCTTTAGCGGGA ^a ACTGT	CTGGGGCT ^a
Glycerol-3-phosphate dehydrogenase (<i>glpA</i>)	CP002228.1	BBUN40_0243	Chromosome	GTTTGTGGAAGCACACTGACATCC	CAATCTCTTCTCAAAATCTTTTGGG	AGAAGAAGA ^a
Glycosaminoglycan-binding protein (<i>lyp</i> , <i>lps-2</i>)	CP002228.1	BBUN40_0588	Chromosome	TGTCATTAATTTCTGGCGTTGCT	CACCACCA ^a CAATCTCCCACITTTAATA	TGGCGTTGTAGTGCTAA
Guanosine-3',5'-bis(diphosphate) 3'-pyrophosphohydrolyase (<i>relA/spot</i>)	CP002228.1	BBUN40_0198	Chromosome	TGGAGAGCCTTACATTAATTCACCC	AAGTACGTCATGCAGTAGTCCAGC	TTCTTGCC ^a
Membrane-associated protein P66	CP002228.1	BBUN40_0603	Chromosome	CTGTAAATTCAGAAACACCTTTTGA	TGATGAATTAATGAAATGTCGTTCCC	CAGGAGCC ^a
Membrane fusion protein (<i>mitrC</i>)	CP002228.1	BBUN40_0141	Chromosome	TCTTGAAGGATCAAGAGTTGCAAC	CGCTGTGGCAAAATAGACTTC	GCTCCAAG ^a
Outer membrane efflux protein	CP002228.1	BBUN40_0142	Chromosome	TTCGCCITTTCAATITCCAACAT	CCTTCTGCTTTGAGTAGCTTTC	CCCAACTCCAAACCCAAA ACCCCA
Outer membrane protein P13	CP002228.1	BBUN40_0034	Chromosome	AATGTTGCCITTTGGGATGA	CAAAGGAGCCTATTC ^a AAACCC	TGCTGGAG ^a
Outer membrane spanning protein (<i>oms38</i>)	CP002228.1	BBUN40_0418	Chromosome	CACCATTGGCATACTTATGAAG	GAAGGCCAAGTITAAAGGTTTGA	GGCAAAGA ^a
Queuine tRNA-ribosyltransferase (<i>tgr</i>)	EF537570	BBUN40_0809	Chromosome	CATTGGCAGTCTTATGGGATTGA	GCTGTAGCCCAATTTGGTTG	GAAGCCAA ^a

(continued)

TABLE 1. (CONTINUED)

Annotation	Accession number	N40 gene symbol	N40 gene location	Forward primer	Reverse primer	Probe
Periplasmic oligopeptide-binding protein (<i>oppA-2</i>)	CP002228.1	BBUN40_0329	Chromosome	AACGAAGAATGGACAACATA CTTAAACA	AATGTCA AAGGATCAGCATAATCG	ATCCATCTGCTCTTGCTAT TTCATAAATTTCCATTT
Response regulatory protein (<i>rrp-1</i>)	FJ643425	BBUN40_0419	Chromosome	CACAGATGCTCAGCTTG AAGGA	GTAAGCCCATCAATTTAGACAAGC	TGGTGGAG ^a
Recombinase A (<i>recA</i>)	NC_001318	BBUN40_0131	Chromosome	GCGCTC AAGCGAATGATGTAAT	CCAGCTTCTCTGAAATACCTTTCC	CTCCACCT ^a
S2 lipoprotein	CP002228.1	BBUN40_0158	Chromosome	CATATTAATGCTGAGTTGGCTTT	TGGTTTGCCAGTAATGCTGCT	AGAAACCCACAATCTGTACATC
Exported protein A (<i>expA</i>)	CP002231.1	BBUN40_C05	cp9	CAAAGCGTGTCCAGAAATAGG	ACGAGGCCCTAAAGCAAAC	TGATATGTGGTATATCTG
Outer surface protein C (<i>ospC</i>)	CP002239.1	BBUN40_B19	cp26	TGTTAAAGGCTTAATCTACAG AAATAA	TACCAAATAGCTTTGGTAAGTTTCAT GTTCTGGCCGT	TACAGAAATCTAACCGAGTT GTTCTGGCCGT
OspE related gene 23 (<i>erp23</i>)	CP002235.1	BBUN40_R18	cp32-4	GTGGTGTGGAGTGC AAGCTTT	ACTGCCGCCAATAGAAATAACTTACA	AGTGTGCCAATGAAT
OspE related gene 24 (<i>erp24</i>)	CP002235.1	BBUN40_R19	cp32-4	CTTCAAAATGGTCCAAAAGTAG ATGAAG	GACTATTAITTAGGATCATCACCCGTCA	AGAAGAAGA ^a
OspE related protein 25 (<i>erp25</i>)	AY142085	BBUN40_V37	cp32-5	GGTTTGGCATGATAAGTTCCTGG	TGGGCTTGCATTAATTTCTCAAC	TTGGTGGT ^a
Outer surface protein E (<i>ospE</i>)	CP002229.1	BBUN40_026	cp32-7	TGTTTATGCTGTTTTATACTTAT AGGTGC	TGACTCACCACTACTTTGCTCATCA	TGCAAGAATTCATACTTCAT
OspE related gene 22 (<i>erp22</i>)	CP002242.1	BBUN40_N45	cp32-9	GGAAGAAGCTAGGGCTAAAGCAG	CTTCTGCTCCTATGCTAGTTTGACC	AGAAGAAGA ^a
Protein 21 (<i>p21</i>)	CP002242.1	BBUN40_N44	cp32-9	GGAAGACTTAGGAACCTTTGGTT GTAAG	GCAGAGTATCCCTTCCCAACAI	AGATGGTATITGATACAGGGTTA
OspE related protein 26 (<i>erp26</i>)	AY142087	BBUN40_Q42	cp32-10	CGCTCAAAATGCTCCAACA	CATCCGCCTGCCATTAATTCITC	AGAAGAAGA ^a
P23-12, conserved hypothetical protein	CP002240.1	BBUN40_Q32	cp32-10	AGATAAGAANAATACACGCAACAA AACATAAC	CTTAGTGTAGTGGGAGCTAGACCAATC	CTACTGCTTTAAACGAAAAT
OspE related protein 27 (<i>erp27</i>)	AY142088	BBUN40_X36	cp32-12	AAGAGCAGATTGGTAGTCTAGTGG TGAAGATAATGTGTTATCGA	ATACCAAGCCCTAATGAATTAACACA CTGCAGATAACCTGTGCGCTT	ATTGGACTCCAGGGCTT CAGCTTATAAATTCCTGAGCTT
<i>Borrelia</i> persistence in ticks (<i>bptA</i>)	AY894345	BBUN40_E05	lp25	TGAAGTAAATGTTGTTATCGA TGAATC	CTGCAGATAACCTGTGCGCTT	
Arthritis related protein (<i>arp</i>)	DQ903071	BBUN40_Y16	lp28-5	CTTCGCCCATGTATTGGAT	ATCCGGAAAAGAAGTTGAAGAAAA	TCCGGGATTAATACCGCTT
Fibronectin-binding protein (<i>fbp</i>)	CP002230.1	BBUN40_K30	lp36	AACAACAAGATGAATATAAAG GGATGACT	TTTGGCTTAAATCAGAAATCTATA GTAAGA	TCACCCACTTCCACCGCTAAG GGAATTTAAACT
Outer surface protein VisE	CP002230.1	BBUN40_K37	lp36	TGATATGAAGAAAGGAGATAA GGTTGCT	TTTAGTACATCCTCCTCTTACCAACAG	AGTAACCCGAAAACCTTCCCAT CTTAGGCCAA
Complement regulator acquiring protein 1 (<i>crisp1</i>)	CP001651.1	BBUN40_A68	lp54	CCCCGGATTGGTGAATTTT	CATGGCCACCTTTTAGCAAA	CTTAGTGTTCGCAATTTGC
Decorin binding protein A (<i>dbpA</i>)	CP001651.1	BBUN40_A24	lp54	GATAACAATGTA AATTTTGCT GCCITT	GTAGCTGCACCTTTTGGCTTCAAG	GTGAAACAGGTAGCAAGGT ATCAGAAAAATTCAT
Outer surface protein A (<i>ospA</i>)	CP001651.1	BBUN40_A15	lp54	TGAAGCGTAAAAGCTGACAAAA	TTCTGTGATGACTGTCTTTGGAA	CAATTTCTGACGATCTAGGTC AAACCAACACTTGA

^aLocked nucleic acid (LNA) probe (Roche, Pleasanton, CA).

Pathology

Knee and tibiotarsal joint (the other knee and tibiotarsal joint were used for nucleic acid extraction) were fixed in 10% neutral-buffered formalin, decalcified, paraffin embedded, sectioned, and stained with Hematoxylin & Eosin using standard technique. Slides were examined blindly for presence of arthritis, and severity of inflammation was scored on a scale of 0 (negative) to 3 (severe), as previously described (Barthold et al. 2006).

Statistical analysis

All statistical analyses were conducted with PASW Statistics 18 (IBM). To evaluate the transcriptional level differences of each target gene between tissues, heart base, ventricular muscle, tibiotarsal joint, quadriceps muscle during early (3 weeks), and late (2 months) infection in C3H compared to immunodeficient C3H-*scid* mice or infected C3H compared to uninfected C3H mice, one-way analysis of variance (ANOVA) followed by a least-squares difference *post hoc* test was performed. The significance level for this test was considered $p < 0.05$. The transcriptional activity ratio of genes was also tested by REST[®] software using pairwise fixed reallocation randomization test.

Results

Infection status

All C3H and C3H-*scid* mice inoculated with *B. burgdorferi* cN40 were confirmed to be infected by culture (all urinary bladders and all inoculation sites positive). None of the sham-

inoculated mice was culture-positive. *B. burgdorferi flaB* DNA was detected in all tissues of all infected mice, but not in sham-inoculated mice. The total bacterial loads, based on *flaB* DNA copy numbers, in heart base, ventricular muscle, tibiotarsal joint, and quadriceps muscle, were significantly higher in C3H-*scid* mice compared to C3H mice at 3 weeks and 2 months after inoculation (Fig. 1, $p < 0.05$). The mean bacterial loads in tissues of both C3H and C3H-*scid* mice decreased over time (3 weeks compared to 2 months), but not significantly ($p = 0.08$, $p = 0.26$, respectively).

Histology

C3H-*scid* mice developed tibiotarsal arthritis (2+ severity at 3 weeks and 3+ severity at 2 months). Inflammation was mild (1+) in only one of four tibiotarsal joints of C3H mice at 3 weeks and absent in C3H mouse joints at 2 months. Knee inflammation was not scored, but present in all C3H-*scid* mice at both intervals, and in none of the infected C3H mice. Heart base, which is consistently inflamed during early infection (Armstrong et al. 1992), was used for DNA and cDNA extraction, and therefore was not available for histology.

LDA analysis

To determine the reproducibility of the LDA assay, five different dilutions of *B. burgdorferi* culture, as well as mouse tissues, were analyzed for all *B. burgdorferi* and host target genes. Intra-assay variations were found to be 1.6–3.2%, and interassay variations of the same sample in three separate runs were 4.1–5.3%. The data also demonstrated that using the LDA assay, it was possible to accurately detect two-fold




FIG. 1. Copy numbers of *flaB* DNA (mean \pm standard deviation [SD]) in heart base, ventricular muscle, tibiotarsal joint, and quadriceps muscle of immunocompetent C3H and immunodeficient C3H-*scid* mice during early (3 weeks) and late (2 months) infection.

changes for every target gene, and small changes could be detected at both low C_q (high expression) and high C_q (low expression). Total RNA was also extracted from all tissue samples from sham-inoculated mice and was subjected to RT-qPCR. Transcriptional activity was detected for host 18S rRNA, but not for any of the *B. burgdorferi* genes.

Selection of reference genes

The C_q values were plotted to compare transcriptional activity of 16S rRNA, *flaB*, *gapDH*, *p13*, *p66*, and *dbpA* of *B. burgdorferi* (Fig. 2). Transcription of *dbpA* mRNA was detected in all tissue samples obtained from both C3H and C3H-*scid* mice during early and late infection. Therefore, this gene was included as a potential reference gene for normalization against the other reference genes, as suggested by Bustin et al. (2010). The evaluated genes had different patterns of transcription within the same sample type, different sample types, and different mouse groups. Genes with higher coefficients of variation (CV) included *p66* (17.08%), *p13* (23.62%), and *gapDH* (33.34%), and were thus characterized as the least stable genes. The most stable reference gene identified with high transcriptional activity and invariant of the condition studied was 16S rRNA (8.33%), followed by *flaB* (15.24%) and *dbpA* (16.05%). A good indicator of constant RNA transcription over all tissue samples is a transcriptional range. The lowest range was observed for 16S rRNA (7.59), followed by *p66* (9.72), *flaB* (9.92), *p13* (12.88), *dbpA* (13.61), and *gapDH* (14.96). The reference gene candidates evaluated for transcriptional stability for normalization of mouse gene transcription included 18S rRNA, *Actβ*, *B2M*, and *gapDH*. The

most stable host reference genes detected for this study were 18S rRNA and *gapDH*.

Transcriptional activity of *B. burgdorferi*

LDA analysis of heart base, ventricular muscle, tibiotarsal joint, and quadriceps muscle revealed that all analyzed *B. burgdorferi* genes were transcriptionally active in all early and late infected C3H-*scid* mice (data not shown). In contrast to C3H-*scid* mice, *B. burgdorferi* in C3H mice did not transcribe detectable amounts of *ospA*, *relA/spoT*, *oms38*, *acrB*, and *ackA* mRNA during either early or late infection. Transcription was not detected for *vlxE*, *eppA*, *bptA*, *crasp1*, *glpA*, *gapDH*, *S2*, *alr*, *colV*, and *p23-T2* at 2 months, whereas *BB0142* transcription was not detected at 3 weeks of infection in C3H mice. During early infection of C3H mice, *ospC* transcription was detected in nine of 12 tissues at early infection, but only in one of 11 tissues at the later infection interval. As expected, *ospA* transcription was not detected in any of the C3H mouse samples, but low levels of *ospA* transcription were detected in C3H-*scid* mice (3/16 during early infection and 7/12 during late infection)(data not shown).

The spirochetal response to host immunity was complex. Transcriptional activity of the analyzed *B. burgdorferi* genes was significantly ($p < 0.05$) higher in C3H-*scid* mice compared to C3H mice at each time point (data not shown). Comparing transcription of genes within each mouse group related to time of infection, no significant differences were found for C3H or C3H-*scid* mice ($p = 0.08$ and $p = 0.11$, respectively). However, at the 2-month interval of infection, transcription of several genes was downregulated compared to early infection in C3H mice.




FIG. 2. Validation of transcriptional mRNA stability of 16S rRNA, *flaB*, *gapDH*, *p13*, *p66*, and *dbpA* as candidates for selecting reference genes. Absolute C_q values (mean \pm standard deviation [SD]) were plotted within the same sample type, different sample types, and different mouse groups during early and late infection.

Tissue samples from infected C3H and C3H-*scid* mice were analyzed for a relative *B. burgdorferi* transcriptional activity profile between the two types of mice to compare the global effects of the immune response on *B. burgdorferi* gene expression. C_q values obtained from every gene and all tissue types were normalized to *16S rRNA*, and transcription level

differences of ΔC_q values were calculated between tissues of C3H versus C3H-*scid* mouse samples. Analysis showed higher levels of transcription for the majority of assessed genes in immunodeficient compared to immunocompetent mice. The transcription of *B. burgdorferi* genes that had changes (up- and/or downregulation) in tissues of C3H mice




FIG. 3. RNA transcription levels (means \pm standard deviation [SD]) of 42 *B. burgdorferi* genes in the heart base of C3H mice relative to gene transcription levels in C3H-*scid* mice during early and late infection. Samples from C3H mice were normalized to samples from C3H-*scid* mice, expressed in comparison to a reference gene (*16S rRNA*), and plotted as fold-changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in C3H mice compared to C3H-*scid* mice. Zero relative expression indicates no expression in C3H mice.

compared to C3H-*scid* were plotted and depicted (Figs. 3–6). Levels of relative expression for over 80% of genes were one- to nine-fold lower in C3H mice compared to C3H-*scid* mice, or no signal was detected. During early infection, only *fbp* and *flaB* were upregulated in heart base of C3H mice compared to

C3H-*scid* mice (Fig. 3), whereas eight genes involved in attachment (*bgp*, *dbpA*), motility (*flaB*), DNA metabolism (*recA*), and complement regulation (*p21*, *erp22*, *erp23*, *erp27*) were upregulated in ventricular muscle (Fig. 4). In the tibiotarsal joint, *flaB*, *tgt*, *recA*, and *erp27* were upregulated during early




FIG. 4. RNA transcription levels (means \pm standard deviation [SD]) of 41 *B. burgdorferi* genes in ventricular muscle of C3H mice relative to gene transcription levels in C3H-*scid* mice during early and late infection. Samples from C3H mice were normalized to samples from C3H-*scid* mice, expressed in comparison to a reference gene (*16S rRNA*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in C3H mice compared to C3H-*scid* mice. Zero relative expression indicates no expression in C3H mice. Transcriptional activity was not detected for *crasp1* in C3H or C3H-*scid* mice during early or late infection, and was therefore not plotted.




FIG. 5. RNA transcriptional levels (mean \pm standard deviation [SD]) of 42 *B. burgdorferi* genes in tibiotalar joint of C3H mice relative to gene transcriptional levels in C3H-*scid* mice during early infection. Samples from C3H mice were normalized to samples from C3H-*scid* mice, expressed in comparison to a reference gene (*16S rRNA*), and plotted as fold-changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in C3H mice compared to C3H-*scid* mice. Zero relative expression indicates no expression in C3H mice. *B. burgdorferi* mRNA was not detected in tibiotalar joint samples collected at late infection, so the transcriptional activity was not assessed.

infection (Fig. 5), whereas in quadriceps muscle, *flaB*, *p66*, *S2*, and *ospE* were upregulated (Fig. 6). The results indicated that *B. burgdorferi* in tibiotalar joints and quadriceps muscle transcribed fewer genes compared to heart base and ventricular muscle during both early and late infection in C3H mice.

At 2 months after inoculation, transcription of 16 genes was not detected in analyzed tissues of C3H mice (Figs. 3–6). Among 29 genes with detectable transcription, 14 were upregulated (one- to seven-fold increase) in heart base, including genes involved in attachment (*bgp*, *dbpA*, *p66*), cell envelope (*ospE*, *bmpA*, *p13*), DNA metabolism (*recA*), complement




FIG. 6. RNA transcriptional levels (mean \pm standard deviation [SD]) of 39 *B. burgdorferi* genes in the quadriceps muscle of C3H mice relative to gene transcriptional levels in C3H-*scid* mice during early infection. Samples from C3H mice were normalized to samples from C3H-*scid* mice, expressed in comparison to a reference gene (*16S rRNA*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in C3H mice compared to C3H-*scid* mice. Zero relative expression indicates no expression in C3H mice. Transcriptional activity was not detected for *crasp1*, *vlsE*, and *oms38* genes in both infected C3H and C3H-*scid* mice during early and late infection, and was therefore not plotted.

regulation (*p21*, *erp23*, *erp24*, *erp26*, *erp27*), cellular processes (*mtrc*), and unknown function (*arp*) (Fig. 3). Fewer genes were upregulated in ventricular muscle of C3H mice, including genes involved in attachment (*bgp*), cell envelope (*bmpA*), DNA metabolism (*recA*), complement regulation (*erp23*, *erp24*,

erp26), and cellular processes (*mtrc*) (Fig. 4). In quadriceps muscle at late infection, transcription was detected for 13 genes, of which 7 (*oppA-2*, *ftsZ*, *bmpA*, *bmpB*, *p13*, *erp22*, *erp25*) were upregulated in C3H mice compared to C3H-*scid* mice (Fig. 6). Due to inadvertent loss of samples, only one

tibiotalar joint at later infection (positive for *flaB* DNA) was available for LDA analysis, but transcription was not detected for any of 43 genes. Interestingly, the *ftsZ* gene, which has been shown to play role in cell division of *B. burgdorferi* (Dubytska et al. 2006), was downregulated in all samples during both early and late infection in C3H mice compared to C3H-*scid* mice.

In summary, all genes were transcriptionally active in C3H-*scid* mice in all tissues. In contrast, the majority of *B. burgdorferi* genes were downregulated in C3H mice relative to C3H-*scid* mice, with upregulation of various genes, depending upon tissue type and interval of infection.

Transcriptional activity of host genes

To compare transcription of host immunoregulatory genes in response to infection, gene transcription levels were compared between infected C3H mice relative to uninfected C3H mice at 3 weeks and 2 months of infection. Comparative analysis of the heart base and ventricular muscle during early infection revealed similar significantly ($p < 0.05$) higher levels of relative expression for CCL7, CXCL13, IL-10, and TNF- α , but not significantly CCL8 ($p = 0.15$), whereas IL4 was downregulated in ventricular muscle (Figs. 7 and 8). At the later infection interval, more cytokines and chemokines were upregulated compared to early infection. During late infection, IL-2, IL-12p40, and TNF- α were downregulated in




FIG. 7. RNA transcription levels (means \pm standard deviation [SD]) of 19 mouse genes in the heart base of infected C3H mice relative to gene transcription levels in uninfected (sham inoculated) C3H mice during early and late infection. Samples from infected C3H mice were normalized to samples from sham-inoculated mice, expressed in comparison to reference genes (*18S rRNA*, *Actb*, *B2M*, and *gapDH*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in infected C3H mice compared to sham-inoculated mice. Zero relative expression indicates no expression in infected C3H mice. Mouse genes not assessed in mice during early infection were CXCL12, CXCL13, CCL21, and IL-21.




FIG. 8. RNA transcription levels (means \pm standard deviation [SD]) of 19 mouse genes in the ventricular muscle of infected C3H mice relative to gene transcription levels in uninfected (sham inoculated) C3H mice during early and late infection. Samples from infected C3H mice were normalized to samples from sham-inoculated mice, expressed in comparison to reference genes (*18S rRNA*, *Actb*, *B2M*, and *gapDH*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in infected C3H mice compared to sham-inoculated mice. Zero relative expression indicates no expression in infected C3H mice. Mouse genes not assessed in mice during early infection were CXCL12, CXCL13, CCL21, and IL-21.

heart base, while IL-4 ($p=0.07$) and CXCL12 ($p<0.05$) were downregulated in ventricular muscle.

Detectable levels of mouse target mRNA during both early and late infection in tibiotarsal joint and quadriceps muscle were fewer (12 samples for both tissues) compared to heart base and ventricular muscle (19 and 17 samples, respectively) (Figs. 9 and 10). In the tibiotarsal joint during early infection, CCL2, CCL7, and TNF- α were upregulated, while CCL12, CCL19, and MyD88 were downregulated, but not significantly. At late infection in the tibiotarsal joint, only CXCL13




FIG. 9. RNA transcription levels (means \pm standard deviation [SD]) of 19 mouse genes in the tibiotarsal joint of infected C3H mice relative to gene transcription levels in uninfected (sham inoculated) C3H mice during early and late infection. Samples from infected C3H mice were normalized to samples from sham-inoculated mice, expressed in comparison to reference genes (*18S rRNA*, *Actb*, *B2M*, and *gapDH*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in infected C3H mice compared to sham-inoculated mice. Zero relative expression indicates no expression in infected C3H mice. Mouse genes not assessed in mice during early infection were CXCL12, CXCL13, CCL21, and IL-21. Transcriptional activity was not detected for chemokines interleukin-2 (IL-2), IL-17a, IL-21, and interferon- γ (IFN- γ) in infected C3H and sham-inoculated mice during early and late infection and was not plotted.




FIG. 10. RNA transcription levels (means \pm standard deviation [SD]) of 19 mouse genes in the quadriceps muscle of infected C3H mice relative to gene transcription levels in uninfected (sham inoculated) C3H mice during early and late infection. Samples from infected C3H mice were normalized to samples from sham inoculated mice, expressed in comparison to reference genes (*18S rRNA*, *Actb*, *B2M*, and *gapDH*), and plotted as fold changes for each gene. Bars to the left of the center line indicate reduced relative expression and bars to the right indicate increased relative expression in infected C3H mice compared to sham-inoculated mice. Zero relative expression indicates no expression in infected C3H mice. Mouse genes not assessed in mice during early infection were CXCL12, CXCL13, CCL21, and IL-21. Transcriptional activity was not detected for chemokines interleukin-17a (IL-17a) and interferon- γ (IFN- γ) in infected C3H and sham-inoculated mice during early and late infection and was not plotted.

was upregulated ($p<0.05$), while CCL2, CCL7, CCL8, CCL12, CXCL12, CCL19, CCL21, IL-1 β , IL-10, TNF- α , and MyD88 were downregulated (Fig. 9). Figure 10 depicts transcriptional activity of mouse genes in quadriceps muscle during early and late infection. During early infection, all detected cytokines were upregulated, while at late infection CCL2, CCL7, CCL8, CXCL12, CXCL13, CCL19, IL-10, and TNF- α were relatively upregulated and CCL12 and MyD88 were downregulated.

In summary, LDA revealed variable patterns of host gene expression in different tissues and at different intervals in infected mice.

Discussion

In this study, responses of the pathogen and the host during early and late infection were investigated, comparing relative gene expressions in selected tissues of immunocompetent versus immunodeficient mice, and infected versus uninfected immunocompetent mice. This was performed with LDA assays that included genes of *B. burgdorferi* cN40 known to be involved in attachment, cell envelope, metabolism, complement regulation, cellular processes, and replication, as well as selected host innate and acquired immune response genes. LDA proved to be sensitive and reproducible and revealed complex patterns of relative response among treatment groups and tissues. Therefore, LDA should be useful for more incisive analysis of *B. burgdorferi* infection in both the vector and the host.

To study the transcriptional activity of a gene's mRNA, it is appropriate to choose a valid internal control for monitoring intersample variation. Genes that are transcribed with the target genes, but are not regulated by the experimental design, can be used as reference genes (Bustin 2000, Huggett et al. 2005, Gilsbach et al. 2006). For the identification of a suitable reference gene, comparison of transcriptional activity among six selected *B. burgdorferi* genes (*16S rRNA*, *flaB*, *gapDH*, *p13*, *p66*, *dbpA*) was evaluated. Analysis revealed that transcription of most of these possible reference genes is highly variable. In particular, some of the potential reference genes had tissue-specific patterns. The most stable reference gene selected was *16S rRNA*, which appeared to be constantly transcribed at similar range among different tissue types and at the different phases of infection. In several studies, *flaB* was used as a reference gene for relative quantification, as in nonhuman primates necropsied 2–3 months postinjection (Miller et al. 2005, Miller and Stevenson 2006), in ticks (Gilmore et al. 2001, Koci et al. 2006), and in culture (Hyde et al. 2007). This gene has generally been regarded to be constitutively expressed, but results suggest otherwise. Using RT-qPCR, we reported previously that *flaB* mRNA transcription was not detected in all tissues and not in all samples, especially those collected during late infection (Hodzic et al. 2003). This study supports our previous finding that *flaB* is not always constitutively expressed, especially during infection, suggesting that *flaB* is not an ideal reference gene for studying *B. burgdorferi* during infection.

Although the *16S rRNA* gene was determined to be the most constantly transcribed gene for use as a reference gene, a problem in transcription of ribosomal genes in prokaryotes that are used as reference genes can arise on the mRNA before translation is terminated, causing a ribosome to stall. This can occur when a mutation in the DNA changes a termination codon that specifies an amino acid, or when transcription terminates prematurely. In these cases, the ribosome reaches the end of the mRNA without encountering a termination codon and stalls, still attached to mRNA (Pierce 2008). Vandecasteele et al. (2001) have reported that expression of the *16S rRNA* gene in *Staphylococcus aureus* decreased significantly more rapidly and earlier than the expression of other reference genes. Ideally, expression of a good reference gene

should not be altered by the experimental conditions, but many studies have reported that not a single gene is expressed invariantly. The most stable *B. burgdorferi* reference gene identified for this study was *16S rRNA*, with high transcriptional activity and invariant of the condition studied.

LDA is a technological approach capable of measuring transcriptional activity of many genes in a single sample as in microarray, while featuring the sensitivity and quantification attributes to real-time qPCR. In recent years, LDA has been used to study complex gene expression changes under various environmental conditions in a host (Goulter et al. 2006, Steg et al. 2006, Cai et al. 2007). One of the aims of this study was to assess the LDA approach for study of Lyme borreliosis, and the ability of LDA to detect and measure transcriptional activity of 43 *B. burgdorferi* genes simultaneously in single samples, as well as 19 host genes. Hence, LDA is suitable for examining changes in gene expression that are involved in pathogenesis and persistence, as well as in response to changing environmental and physiological conditions and pharmacological stimuli (Goulter et al. 2006, Steg et al. 2006, Lu et al. 2008, Sanchez-Espiridion et al. 2009). Searching available literature, no publications were found that described the use of LDA to study *B. burgdorferi* gene expression.

The amounts of *B. burgdorferi* RNA and cDNA available from collected mouse tissues are limited, especially in immunocompetent mice and during late infection, which restricts the number of analyzable genes. To overcome this problem, a preamplification technique was used to enhance the sensitivity and fidelity of the RT-qPCR, especially for low-abundance target genes, which increases the fidelity of target genes that can be analyzed. Identical primers were used for both the preamplification reaction and the succeeding RT-qPCR. The preamplification approach for gene transcription gave a high yield of cDNA, an increase in sensitivity, low interassay variation, and consistent results. This confirms that preamplification is a suitable tool for investigating gene expression analyses with the limited RNA/cDNA amounts derived from mice with persistent infection and fewer spirochete burdens. Narasimhan et al. (2003) have used a preamplification approach to study *B. burgdorferi* gene expression in the central nervous system and heart of nonhuman primates. The authors used a different methodology (membrane arrays) than that which was employed here, which makes comparison of results difficult. Ciotti et al. (2009) have reported that preamplification is a powerful and reproducible approach that can be performed even using poorer RNA quality samples.

The present study focused on a variety of genes located throughout the *B. burgdorferi* genome that encode proteins associated with a variety of functions, during early (3 weeks) and late (2 months) infection in immunocompetent and immunodeficient mice. We used the well-characterized C3H and C3H-*scid* mouse models, in which inflammation is consistently present in the heart base, but minimal in the ventricular muscle at 3 weeks of infection; inflammation subsides at 2 months of infection (Armstrong et al. 1992). Thus, the model allowed testing the same tissue types during the inflammatory and resolved (persistent) phases of infection. Development of arthritis, however, can be variable, depending upon mouse strain, age, and infectious dose (Barthold et al. 2010). In the current study, arthritis was minimal in C3H mice and was found in only one mouse at the 3-week interval.

Host responses may play roles in the pathogenesis of *Borrelia* infection, such as inducing expression of lipoproteins on the cell surface, which facilitates dissemination and inflammation. Expression of surface-exposed lipoproteins results in cytokine/chemokine activation that may induce host inflammatory reactions at the pathogen's predilection sites (skin, heart, joint, central nervous system), while their downregulation permits the survival of spirochetes for prolonged periods of time despite the host immune responses (Norris et al. 2010). Comparative analysis of *B. burgdorferi* gene transcription revealed higher activity in tissues of immunodeficient compared to immunocompetent mice at both early and late infection in all tissues, and for all genes except *acrB* (not transcribed at either time point) and *crasp1* (not transcribed during late infection). These observations support previous findings that transcriptional activity of *B. burgdorferi* genes have been influenced by the host immune system as a strategy of immune evasion (Liang et al. 2002a, Brooks et al. 2003).

Notably, transcription of two genes in particular, *ospA* and *ospC*, in tissues of mice in the current study validated the LDA approach. *OspA* is expressed by spirochetes within the midgut of ticks, and *ospA* is downregulated during mammalian infection. One signal for downregulation of *ospA* in the tick is a natural antibody that is acquired during the blood meal (Belperron and Bockenstedt 2001). In contrast, *ospA* is transcribed in immunodeficient C3H-*scid* mice, and it has been found that passive transfer of normal mouse serum in infected C3H-*scid* mice results in *ospA* downregulation (Hodzic et al. 2005). *OspA* also tends to be expressed in the context of inflammation (Crowley and Huber 2003). These patterns of *ospA* transcription in infected C3H-*scid* mice were observed in the current study. *OspC* has been shown to be essential to establishing initial infection, but is vulnerable to a specific antibody, and therefore is downregulated as infection proceeds in immunocompetent hosts (Liang et al. 2004b, Seemanapalli et al. 2010). As expected, *ospC* was transcribed in C3H-*scid* mice, but was downregulated in immunocompetent C3H mice as infection progressed. These known patterns of transcription serve as validation of the LDA approach.

Differential expression of *B. burgdorferi* genes was assessed between genes located on the chromosome versus plasmid. No differences were noted during early infection, whereas during late infection most of the upregulated genes that were identified on plasmids encode proteins involved in complement regulation, and those identified on the chromosome encode proteins involved in attachment, cell envelope, cell processes, and DNA metabolism. Brooks et al. (2003) have reported that almost all of the upregulated genes of *B. burgdorferi* cultured under different conditions were chromosomally located, and the vast majority of downregulated genes were encoded on plasmids. Revel et al. (2002) reported similar results, in which cultured as well as host-adapted spirochetes in dialysis membrane chambers implanted in rats were used. Liang et al. (2002c) have found that during early infection five of six of the assessed lipoprotein genes were transcribed, while at later infection four of five of those transcribed during early infection were downregulated. Of the downregulated genes, two of three were plasmid-encoded, and one of three was located on a chromosome. A microarray technique has been used to study differential expression of borrelial genes

cultivated in different environments (Revel et al. 2002) and exposed to different temperatures (Ojaimi et al. 2003) or to mammalian host factors (Brooks et al. 2003, Tokarz et al. 2004). Our data are difficult to compare with those cited above because they used a different technique, *in vitro*-grown spirochetes, and different tissue samples and different stages of infection. Nevertheless, LDA offers the advantage of quantitatively assessing an array of genes simultaneously.

Many studies have shown that cytokines play an important role in the pathogenesis of Lyme borreliosis in the initial phase of infection (Miller et al. 2008, Rupprecht et al. 2009, Bachmann et al. 2010), while at late infection expression of some, especially Th1-associated cytokines, waned (Sjowall et al. 2011). One of the most investigated cytokines is IL-10, which was slightly overexpressed (Benhnia et al. 2005, Lazarus et al. 2008), and has been shown to be a main cytokine to regulate inflammatory response in *B. burgdorferi* infection (Gautam et al. 2011). Comparative analysis of the experimental data of heart tissue revealed higher levels of relative expression for IL-10 during both early and late infection. In the tibiotarsal joint, expression of IL-10 during early and late infection was unchanged when compared to sham-infected control mice, whereas in the quadriceps muscle one out of five samples had higher detectable levels of this cytokine. Lazarus et al. (2006) reported that clearance of *B. burgdorferi* by innate immunity is more efficient in the absence of IL-10, which could explain our finding of IL-10 downregulation and lack of inflammation in the tibiotarsal joint.

Major proinflammatory cytokines that are involved in induction of inflammatory responses include IFN- γ and TNF- α , which were upregulated and correlated with a greater severity of Lyme disease (Christopherson et al. 2003, Dame et al. 2007). The synergistic effect of *B. burgdorferi* and IFN- γ induces upregulated expression of numerous genes, of which seven encode chemokines (CCL7, CCL8, CXCL2, CX3CL1, CXCL9, CXCL10, CXCL11) that attract T lymphocytes and neutrophils (Dame et al. 2007). In our study, we demonstrated a significant increase of IFN- γ in heart tissue during early and late infection, along with CCL7 and CCL8, whereas TNF- α was upregulated in heart tissues, except in heart base samples, during late infection.

Comparative analysis of the transcriptional activity of *B. burgdorferi* and cytokines revealed that increased *flaB* mRNA during early infection was followed by an increase of CCL7, CCL8, IL10, and TNF- α in all assessed tissue types. Also, during early infection in all tissue tested, most complement regulation genes tested were downregulated. In ventricular muscle during both early and late infection, IL-4 was downregulated as well as *B. burgdorferi* genes involved in cellular processes (*acrB*, *relA/spot*) and cell envelope (*bmpC*, *bmpD*, *ospC*, *eppA*).

The main aim of this study was to assess the feasibility of LDA to study multiple gene expression of *B. burgdorferi* during infection and host responses to infection. This work shows complex expression of *B. burgdorferi* genes that correspond to expression of certain host cytokines and chemokines, of which immunoregulatory effects seem to be optimized for bacterial persistence in the host. These results will pave the way for a more comprehensive study of multiple genes in selected tissues under different conditions and treatments during different stages of infection in this important vector-borne disease.

Acknowledgments

This work was supported by PHS grant R01 AI026815 from the National Institute of Allergy and Infectious Diseases. The technical assistance of Kimberly Olsen, Edlin Escobar, Samantha Mapes, and Alicja Omanska is gratefully acknowledged.

Author Disclosure Statement

The authors have no conflicts of interest in connection with the submitted manuscript.

References

- Abruzzo LV, Lee KY, Fuller A, Silverman A, et al. Validation of oligonucleotide microarray data using microfluidic low-density arrays: A new statistical method to normalize real-time RT-PCR data. *Biotechniques* 2005; 38:785–792.
- Adusumilli S, Booth CJ, Anguita J, Fikrig E. Passage through *Ixodes scapularis* ticks enhances the virulence of a weakly pathogenic isolate of *Borrelia burgdorferi*. *Infect Immun* 2010; 78:138–144.
- Anderton JM, Tokarz R, Thill CD, Kuhlow CJ, et al. Whole-genome DNA array analysis of the response of *Borrelia burgdorferi* to a bactericidal monoclonal antibody. *Infect Immun* 2004; 72:2035–2044.
- Armstrong AL, Barthold SW, Persing DH, Beck DS. Carditis in Lyme disease susceptible and resistant strains of laboratory mice infected with *Borrelia burgdorferi*. *Am J Trop Med Hyg* 1992; 47:249–258.
- Bachmann M, Horn K, Rudloff I, Goren I, et al. Early production of IL-22 but not IL-17 by peripheral blood mononuclear cells exposed to live *Borrelia burgdorferi*: The role of monocytes and interleukin-1. *PLoS Pathogens* 2010; 6:e1001144.
- Backenson PB, Coleman JL, Benach JL. *Borrelia burgdorferi* shows specificity of binding to glycosphingolipids. *Infect Immun* 1995; 63:2811–2817.
- Bankhead T, Chaconas G. The role of VlsE antigenic variation in the Lyme disease spirochete: Persistence through a mechanism that differs from other pathogens. *Mol Microbiol* 2007; 65:1547–1558.
- Barbour AG. Isolation and cultivation of Lyme disease spirochetes. *Yale J Biol Med* 1984; 57:521–525.
- Barthold SW. Lyme borreliosis. In: Nataro JP, Blaser MJ, Cunningham-Rundles S, eds. *Persistent Bacterial Infection*. Washington, DC: ASM Press, 2000:281–304.
- Barthold SW, de Souza MS, Janotka JL, Smith AL, et al. Chronic Lyme borreliosis in the laboratory mouse. *Am J Pathol* 1993; 143:959–971.
- Barthold SW, Hodzic E, Tunev S, Feng S. Antibody-mediated disease remission in the mouse model of Lyme borreliosis. *Infect Immun* 2006; 74:4817–4825.
- Barthold SW, Cadavid D, Philipp MT. Animal models of borreliosis. In: Samuels DS, Radolf J, eds. *Borrelia, Molecular Biology, Host Interaction and Pathogenesis*. Norfolk, UK: Caister Academic Press, 2010:353–405.
- Belfaiza J, Postic D, Bellenger E, Baranton G, et al. Genomic fingerprinting of *Borrelia burgdorferi* sensu lato by pulsed-field gel electrophoresis. *J Clin Microbiol* 1993; 31:2873–2877.
- Belperron AA, Bockenstedt LK. Natural antibody affects survival of the spirochete *Borrelia burgdorferi* within feeding ticks. *Infect Immun* 2001; 69:6456–6462.
- Benhnia MR, Wroblewski D, Akhtar MN, Patel RA, et al. Signaling through CD14 attenuates the inflammatory response to *Borrelia burgdorferi*, the agent of Lyme disease. *J Immunol* 2005; 174:1539–1548.
- Bhide MR, Escudero R, Camafeita E, Gil H, et al. Complement factor H binding by different Lyme disease and relapsing fever *Borrelia* in animals and human. *BMC Research Notes* 2009; 2.
- Boerlin P, Peter O, Bretz AG, Postic D, et al. Population genetic analysis of *Borrelia burgdorferi* isolates by multilocus enzyme electrophoresis. *Infect Immun* 1992; 60:1677–1683.
- Brisette CA, Bykowski T, Cooley AE, Bowman A, et al. *Borrelia burgdorferi* RevA antigen binds host fibronectin. *Infect Immun* 2009a; 77:2802–2812.
- Brisette CA, Verma A, Bowman A, Cooley AE, et al. The *Borrelia burgdorferi* outer-surface protein ErpX binds mammalian laminin. *Microbiology* 2009b; 155:863–872.
- Brooks CS, Hefty PS, Jolliff SE, Akins DR. Global analysis of *Borrelia burgdorferi* genes regulated by mammalian host-specific signals. *Infect Immun* 2003; 71:3371–3383.
- Bubeck-Martinez S. Immune evasion of the Lyme disease spirochetes. *Front Biosci* 2005; 10:873–878.
- Bustin SA. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* 2000; 25:169–193.
- Bustin SA, Beaulieu JF, Huggett J, Jaggi R, et al. MIQE precis: Practical implementation of minimum standard guidelines for fluorescence-based quantitative real-time PCR experiments. *BMC Mol Biol* 2010; 11:74.
- Cabello FC, Godfrey HP, Newman SA. Hidden in plain sight: *Borrelia burgdorferi* and the extracellular matrix. *Trends Microbiol* 2007; 15:350–354.
- Cai JH, Deng S, Kumpf SW, Lee PA, et al. Validation of rat reference genes for improved quantitative gene expression analysis using low density arrays. *Biotechniques* 2007; 42:503–512.
- Christopherson JA, Munson EL, England DM, Croke CL, et al. Destructive arthritis in vaccinated interferon gamma-deficient mice challenged with *Borrelia burgdorferi*: Modulation by tumor necrosis factor alpha. *Clin Diagnostic Lab Immunol* 2003; 10:44–52.
- Ciotti P, Garuti A, Ballestrero A, Cirmena G, et al. Reliability and reproducibility of a RNA preamplification method for low-density array analysis from formalin-fixed paraffin-embedded breast cancer samples. *Diagn Mol Pathol* 2009; 18:112–118.
- Coburn J, Medrano M, Cugini C. *Borrelia burgdorferi* and its tropisms for adhesion molecules in the joint. *Curr Opin Rheumatol* 2002; 14:394–398.
- Coleman JL, Gebbia JA, Piesman J, Degen JL, Bugge TH, Benach JL. Plasminogen is required for efficient dissemination of *B. burgdorferi* in ticks and for enhancement of spirochetemia in mice. *Cell* 1997; 89:1111–9.
- Cooke BM, Mohandas N, Cowman AF, Coppel RL. Cellular adhesive phenomena in apicomplexan parasites of red blood cells. *Vet Parasitol* 2005; 132:273–295.
- Crowley H, Huber BT. Host-adapted *Borrelia burgdorferi* in mice expresses OspA during inflammation. *Infect Immun* 2003; 71:4003–4010.
- Dame TM, Orenzoff BL, Palmer LE, Furie MB. IFN-gamma alters the response of *Borrelia burgdorferi*-activated endothelium to favor chronic inflammation. *J Immunol* 2007; 178:1172–1179.
- de Cremoux P, Bieche I, Tran-Perennou C, Vignaud S, et al. Inter-laboratory quality control for hormone-dependent gene expression in human breast tumors using real-time reverse transcription-polymerase chain reaction. *Endocr Relat Cancer* 2004; 11:489–495.
- Defoe G, Coburn J. Delineation of *Borrelia burgdorferi* p66 sequences required for integrin alpha(IIb)beta(3) recognition. *Infect Immun* 2001; 69:3455–3459.

- Diterich I, Rauter C, Kirschning CJ, Hartung T. *Borrelia burgdorferi*-induced tolerance as a model of persistence via immunosuppression. *Infect Immun* 2003; 71:3979–3987.
- Dubytska L, Godfrey HP, Cabello FC. *Borrelia burgdorferi* ftsZ plays a role in cell division. *J Bacteriol* 2006; 188:1969–1978.
- Fuchs R, Jauris S, Lottspeich F, Preac-Mursic V, et al. Molecular analysis and expression of a *Borrelia burgdorferi* gene encoding a 22 kDa protein (pC) in *Escherichia coli*. *Mol Microbiol* 1992; 6:503–509.
- Gautam A, Dixit S, Philipp MT, Singh SR, et al. Interleukin-10 alters effector functions of multiple genes induced by *Borrelia burgdorferi* in macrophages to regulate Lyme disease inflammation. *Infect Immun* 2011; 79:4876–4892.
- Gilmore RD, Jr., Mbow ML, Stevenson B. Analysis of *Borrelia burgdorferi* gene expression during life cycle phases of the tick vector *Ixodes scapularis*. *Microbes Infect* 2001; 3:799–808.
- Gilsbach R, Kouta M, Bonisch H, Bruss M. Comparison of in vitro and in vivo reference genes for internal standardization of real-time PCR data. *Biotechniques* 2006; 40:173–177.
- Goulter AB, Harmer DW, Clark KL. Evaluation of low density array technology for quantitative parallel measurement of multiple genes in human tissue. *BMC Genomics* 2006; 7:34.
- Grab DJ, Givens C, Kennedy R. Fibronectin-binding activity in *Borrelia burgdorferi*. *Biochim Biophys Acta* 1998; 1407:135–145.
- Guo BP, Brown EL, Dorward DW, Rosenberg LC, et al. Decorin-binding adhesins from *Borrelia burgdorferi*. *Mol Microbiol* 1998; 30:711–723.
- Hodzic E, Feng S, Freet KJ, Borjesson DL, et al. *Borrelia burgdorferi* population kinetics and selected gene expression at the host-vector interface. *Infect Immun* 2002; 70:3382–3388.
- Hodzic E, Feng S, Freet KJ, Barthold SW. *Borrelia burgdorferi* population dynamics and prototype gene expression during infection of immunocompetent and immunodeficient mice. *Infect Immun* 2003; 71:5042–5055.
- Hodzic E, Tunev S, Feng S, Freet KJ, et al. Immunoglobulin-regulated expression of *Borrelia burgdorferi* outer surface protein A in vivo. *Infect Immun* 2005; 73:3313–3321.
- Hovis KM, Tran E, Sundry CM, Buckles E, et al. Selective binding of *Borrelia burgdorferi* OspE paralogs to factor H and serum proteins from diverse animals: Possible expansion of the role of OspE in Lyme disease pathogenesis. *Infect Immun* 2006; 74:1967–1972.
- Huggett J, Dheda K, Bustin S, Zumla A. Real-time RT-PCR normalisation; strategies and considerations. *Genes Immun* 2005; 6:279–284.
- Hyde JA, Seshu J, Skare JT. Transcriptional profiling of *Borrelia burgdorferi* containing a unique bosR allele identifies a putative oxidative stress regulon. *Microbiology* 2006; 152:2599–2609.
- Hyde JA, Trzeciakowski JP, Skare JT. *Borrelia burgdorferi* alters its gene expression and antigenic profile in response to CO₂ levels. *J Bacteriol* 2007; 189:437–445.
- Koci J, Derdakova M, Peterkova K, Kazimirova M, et al. *Borrelia afzelii* gene expression in *Ixodes ricinus* (Acari: Ixodidae) ticks. *Vector Borne Zoonotic Dis* 2006; 6:296–304.
- Lazarus JJ, Meadows MJ, Lintner RE, Wooten RM. IL-10 deficiency promotes increased *Borrelia burgdorferi* clearance predominantly through enhanced innate immune responses. *J Immunol* 2006; 177:7076–7085.
- Lazarus JJ, Kay MA, McCarter AL, Wooten RM. Viable *Borrelia burgdorferi* enhances interleukin-10 production and suppresses activation of murine macrophages. *Infect Immun* 2008; 76:1153–1162.
- Le Fleche A, Postic D, Girardet K, Peter O, et al. Characterization of *Borrelia lusitaniae* sp. nov. by 16S ribosomal DNA sequence analysis. *Int J Syst Bacteriol* 1997; 47:921–925.
- Leong JM, Wang H, Magoun L, Field JA, et al. Different classes of proteoglycans contribute to the attachment of *Borrelia burgdorferi* to cultured endothelial and brain cells. *Infect Immun* 1998; 66:994–999.
- Liang FT, Jacobs MB, Bowers LC, Philipp MT. An immune evasion mechanism for spirochetal persistence in Lyme borreliosis. *J Exp Med* 2002a; 195:415–422.
- Liang FT, Nelson FK, Fikrig E. DNA Microarray assessment of putative *Borrelia burgdorferi* lipoprotein genes. *Infect Immun* 2002b; 70:3300–3303.
- Liang FT, Nelson FK, Fikrig E. Molecular adaptation of *Borrelia burgdorferi* in the murine host. *J Exp Med* 2002c; 196:275–80.
- Liang FT, Brown EL, Wang T, Iozzo RV, et al. Protective niche for *Borrelia burgdorferi* to evade humoral immunity. *Am J Pathol* 2004a; 165:977–985.
- Liang FT, Yan J, Mbow ML, Sviat SL, Gilmore RD, Mamula M, Fikrig E. *Borrelia burgdorferi* changes its surface antigenic expression in response to host immune responses. *Infect Immun* 2004b; 72:5759–5767.
- Lien E, Sellati TJ, Yoshimura A, Flo TH, et al. Toll-like receptor 2 functions as a pattern recognition receptor for diverse bacterial products. *J Biol Chem* 1999; 274:33419–33425.
- Lu B, Xu J, Chen J, Yu J, et al. TaqMan low density array is roughly right for gene expression quantification in colorectal cancer. *Clin Chim Acta* 2008; 389:146–151.
- Malawista SE. Resolution of Lyme arthritis, acute or prolonged: A new look. *Inflammation* 2000; 24:493–504.
- Marconi RT, Garon CF. Development of polymerase chain reaction primer sets for diagnosis of Lyme disease and for species-specific identification of Lyme disease isolates by 16S rRNA signature nucleotide analysis. *J Clin Microbiol* 1992; 30:2830–2834.
- Miklosy J, Kasas S, Zurn AD, McCall S, et al. Persisting atypical and cystic forms of *Borrelia burgdorferi* and local inflammation in Lyme neuroborreliosis. *J Neuroinflammation* 2008; 5:40.
- Miller JC, Stevenson B. *Borrelia burgdorferi* *erp* genes are expressed at different levels within tissues of chronically infected mammalian hosts. *Int J Med Microbiol* 2006; 296(Suppl 40): 185–194.
- Miller JC, Narayan K, Stevenson B, Pachner AR. Expression of *Borrelia burgdorferi* *erp* genes during infection of non-human primates. *Microb Pathog* 2005; 39:27–33.
- Miller JC, Ma Y, Bian J, Sheehan KC, et al. A critical role for type I IFN in arthritis development following *Borrelia burgdorferi* infection of mice. *J Immunol* 2008; 181:8492–8503.
- Narasimhan S, Caimano MJ, Liang FT, Santiago F, et al. *Borrelia burgdorferi* transcriptome in the central nervous system of non-human primates. *Proc Natl Acad Sci USA* 2003; 100:15953–15958.
- Norris DE, Coburn J, Leong JM, Hu LT, et al. Pathobiology of Lyme disease borrelia. In: Samuels DS, Radolf J, eds. *Borrelia: Molecular Biology, Host Interaction and Pathogenesis*. Norfolk, UK: Caister Academic Press, 2010:295–325.
- Ojaimi C, Brooks C, Casjens S, Rosa P, et al. Profiling of temperature-induced changes in *Borrelia burgdorferi* gene expression by using whole genome arrays. *Infect Immun* 2003; 71:1689–1705.
- Pal U, Fikrig E. Adaptation of *Borrelia burgdorferi* in the vector and vertebrate host. *Microbes Infect* 2003; 5:659–666.
- Pal U, Wang P, Bao F, Yang X, et al. *Borrelia burgdorferi* basic membrane proteins A and B participate in the genesis of Lyme arthritis. *J Exp Med* 2008; 205:133–141.

- Palmer GH, Bankhead T, Lukehart SA. 'Nothing is permanent but change'—antigenic variation in persistent bacterial pathogens. *Cell Microbiol* 2009; 11:1697–1705.
- Parveen N, Leong JM. Identification of a candidate glycosaminoglycan-binding adhesin of the Lyme disease spirochete *Borrelia burgdorferi*. *Mol Microbiol* 2000; 35:1220–1234.
- Parveen N, Cornell KA, Bono JL, Chamberland C, et al. Bgp, a secreted glycosaminoglycan-binding protein of *Borrelia burgdorferi* strain N40, displays nucleosidase activity and is not essential for infection of immunodeficient mice. *Infect Immun* 2006; 74:3016–3020.
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001; 29:e45.
- Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res* 2002; 30:e36.
- Pierce BA. The Genetic Code and Translation. In: *Genetics: A Conceptual Approach*. New York: W.H. Freeman & Company, 2008:395–424.
- Postic D, Assous M, Belfaiza J, Baranton G. Genetic diversity of borrelia of Lyme borreliosis. *Wien Klin Wochenschr* 1996; 108:748–751.
- Postic D, Ras NM, Lane RS, Henderson M, et al. Expanded diversity among Californian borrelia isolates and description of *Borrelia bissettii* sp. nov. (formerly *Borrelia* group DN127). *J Clin Microbiol* 1998; 36:3497–3504.
- Probert WS, Johnson BJ. Identification of a 47 kDa fibronectin-binding protein expressed by *Borrelia burgdorferi* isolate B31. *Mol Microbiol* 1998; 30:1003–1015.
- Radonic A, Thulke S, Mackay IM, Landt O, et al. Guideline to reference gene selection for quantitative real-time PCR. *Biochem Biophys Res Commun* 2004; 313:856–862.
- Revel AT, Talaat AM, Norgard MV. DNA microarray analysis of differential gene expression in *Borrelia burgdorferi*, the Lyme disease spirochete. *Proc Natl Acad Sci USA* 2002; 99:1562–1567.
- Rupprecht TA, Plate A, Adam M, Wick M, et al. The chemokine CXCL13 is a key regulator of B cell recruitment to the cerebrospinal fluid in acute Lyme neuroborreliosis. *J Neuroinflammation* 2009; 6:42.
- Sanchez-Espiridion B, Sanchez-Aguilera A, Montalban C, Martin C, et al. A TaqMan low-density array to predict outcome in advanced Hodgkin's lymphoma using paraffin-embedded samples. *Clin Cancer Res* 2009; 15:1367–1375.
- Schutzer SE, Fraser-Liggett CM, Casjens SR, Qiu WG, et al. Whole-genome sequences of thirteen isolates of *Borrelia burgdorferi*. *J Bacteriol* 2011; 193:1018–1020.
- Seemanapalli SV, Xu Q, McShan K, Liang FT. Outer surface protein C is a dissemination-facilitating factor of *Borrelia burgdorferi* during mammalian infection. *PLoS One* 2010; 5:e15830.
- Shin JJ, Strle K, Glickstein LJ, Luster AD, et al. *Borrelia burgdorferi* stimulation of chemokine secretion by cells of monocyte lineage in patients with Lyme arthritis. *Arthritis Res Ther* 2010; 12:R168.
- Sjowall J, Fryland L, Nordberg M, Sjogren F, et al. Decreased Th1-type inflammatory cytokine expression in the skin is associated with persisting symptoms after treatment of erythema migrans. *PLoS One* 2011; 6:e18220.
- Steg A, Wang W, Blanquicett C, Grunda JM, et al. Multiple gene expression analyses in paraffin-embedded tissues by TaqMan low-density array: Application to hedgehog and Wnt pathway analysis in ovarian endometrioid adenocarcinoma. *J Mol Diagn* 2006; 8:76–83.
- Strle K, Drouin EE, Shen S, Khoury JE, et al. *Borrelia burgdorferi* stimulates macrophages to secrete higher levels of cytokines and chemokines than *Borrelia afzelii* or *Borrelia garinii*. *J Infect Dis* 2009; 200:1936–1943.
- Tokarz R, Anderton JM, Katona LI, Benach JL. Combined effects of blood and temperature shift on *Borrelia burgdorferi* gene expression as determined by whole genome DNA array. *Infect Immun* 2004; 72:5419–5432.
- Vandecasteele SJ, Peetermans WE, Merckx R, Van Eldere J. Quantification of expression of *Staphylococcus epidermidis* housekeeping genes with Taqman quantitative PCR during in vitro growth and under different conditions. *J Bacteriol* 2001; 183:7094–7101.
- Verma A, Brisette CA, Bowman A, Stevenson B. *Borrelia burgdorferi* BmpA is a laminin-binding protein. *Infect Immun* 2009; 77:4940–4946.
- Wang G, van Dam AP, Spanjaard L, Dankert J. Molecular typing of *Borrelia burgdorferi* sensu lato by randomly amplified polymorphic DNA fingerprinting analysis. *J Clin Microbiol* 1998; 36:768–776.
- Weening EH, Parveen N, Trzeciakowski JP, Leong JM, et al. *Borrelia burgdorferi* lacking DbpBA exhibits an early survival defect during experimental infection. *Infect Immun* 2008; 76:5694–5705.
- Wilske B, Busch U, Fingerle V, Jauris-Heipke S, et al. Immunological and molecular variability of OspA and OspC. Implications for *Borrelia* vaccine development. *Infection* 1996; 24:208–212.
- Xu Y, Johnson RC. Analysis and comparison of plasmid profiles of *Borrelia burgdorferi* sensu lato strains. *J Clin Microbiol* 1995; 33:2679–2685.
- Zambrano MC, Beklemisheva AA, Bryksin AV, Newman SA, et al. *Borrelia burgdorferi* binds to, invades, and colonizes native type I collagen lattices. *Infect Immun* 2004; 72:3138–3146.

Address correspondence to:

Emir Hodzic

Center for Comparative Medicine

School of Veterinary Medicine

University of California Davis

One Shields Avenue

Davis, CA 95616

E-mail: ehodzic@ucdavis.edu