

Original Article

Mitochondrial and DNA damage in bovine somatic cell nuclear transfer embryos

In-Sun Hwang, Hyo-Kyung Bae, Hee-Tae Cheong*

College of Veterinary Medicine and Institute of Veterinary Science, Kangwon National University, Chuncheon 200-701, Korea

The generation of reactive oxygen species (ROS) and subsequent mitochondrial and DNA damage in bovine somatic cell nuclear transfer (SCNT) embryos were examined. Bovine enucleated oocytes were electrofused with donor cells and then activated by a combination of Ca-ionophore and 6-dimethylaminopurine culture. The H₂O₂ and [•]OH radical levels, mitochondrial morphology and membrane potential ($\Delta\Psi$), and DNA fragmentation of SCNT and *in vitro* fertilized (IVF) embryos at the zygote stage were analyzed. The H₂O₂ (35.6 ± 1.1 pixels/embryo) and [•]OH radical levels (44.6 ± 1.2 pixels/embryo) of SCNT embryos were significantly higher than those of IVF embryos (19.2 ± 1.5 and 23.8 ± 1.8 pixels/embryo, respectively, $p < 0.05$). The mitochondria morphology of SCNT embryos was diffused within the cytoplasm. The $\Delta\Psi$ of SCNT embryos was significantly lower ($p < 0.05$) than that of IVF embryos (0.95 ± 0.04 vs. 1.21 ± 0.06 , red/green). Moreover, the comet tail length of SCNT embryos was longer than that of IVF embryos (515.5 ± 26.4 μm vs. 425.6 ± 25.0 μm , $p < 0.05$). These results indicate that mitochondrial and DNA damage increased in bovine SCNT embryos, which may have been induced by increased ROS levels.

Keywords: cattle, cellular damage, DNA fragmentation, ROS generation, somatic cell nuclear transfer

Introduction

The somatic cell nuclear transfer (SCNT) technique has several serious problems that must be resolved to improve its efficiency [7]. These problems are considered to be a result of abnormal reprogramming of SCNT embryos [14,15]. The reprogramming of SCNT embryos can be influenced by a number of factors involving nucleo-cytoplasmic interactions [4]. Mechanical and physiochemical treatments such as

enucleation, electrofusion and activation can also cause cellular stresses in the SCNT embryos. We recently reported that cellular stress that can be caused by the manipulations during SCNT procedures, resulting in generation of reactive oxygen species (ROS) in bovine SCNT embryos [13].

High levels of ROS in the SCNT embryos might result in serious damage to cell membranes [1] and nuclear and mitochondrial DNA of the cells [9, 20], including apoptosis [29], DNA fragmentation [16], abnormal metabolism [19] and aberrant gene expression [20]. Such effects can restrict reprogramming of SCNT embryos. Therefore, the present study was conducted to examine the ROS generation level and subsequent damage to mitochondrial DNA in bovine SCNT embryos.

Materials and Methods

In vitro maturation of oocytes

Bovine cumulus-oocyte complexes (COCs) were collected from the follicles (2- to 7-mm diameter) of ovaries. About ten COCs were cultured in 50 μL droplets of *in vitro* maturation (IVM) medium overlaid with paraffin oil for 20 ~ 22 h at 39°C under 5% CO₂ in air. The IVM medium was Tissue Culture Medium 199 (TCM199; Gibco BRL, USA) supplemented with 10% fetal bovine serum (FBS; Gibco BRL), 0.02 U/mL follicle-stimulating hormone (Sigma-Aldrich, USA), 1 $\mu\text{g}/\text{mL}$ estradiol (Sigma-Aldrich), 50 $\mu\text{g}/\text{mL}$ gentamicin (Sigma-Aldrich), and 0.2 mM Na-pyruvate (Sigma-Aldrich).

Preparation of donor cells

Bovine ear skin fibroblasts (4 ~ 6 passages) from a Korean native cow were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco BRL) supplemented with 10% FBS, 0.2 mM Na-pyruvate (Sigma-Aldrich), and 1% penicillin/streptomycin for 2 ~ 3 days to achieve about 70%

*Corresponding author: Tel: +82-33-250-8659; Fax: +82-33-244-2367; E-mail: htcheong@kangwon.ac.kr

confluency. The cells were then cultured for an additional 5 days in DMEM containing 0.5% FBS. Prior to use, the cells were trypsinized and washed by centrifugation in TCM199 medium supplemented with 3 mg/mL bovine serum albumin (BSA; Sigma-Aldrich).

Nuclear transfer

SCNT was carried out in Hepes-buffered TCM199 (Gibco BRL) supplemented with 3 mg/mL BSA and 5 μ g/mL cytochalasin B (Sigma-Aldrich) according to the methods reported previously [17]. The cumulus cells of *in vitro* matured COCs were removed by vortexing for 5 min in phosphate-buffered saline (PBS) supplemented with 0.1% (w/v) hyaluronidase (Sigma-Aldrich) and 0.1% (w/v) polyvinyl alcohol (PVA; Sigma-Aldrich). Prior to the enucleation, oocytes were cultured in TCM199 containing 0.4 μ g/mL demecolcine (Sigma-Aldrich) for 40 min to extrude their metaphase II (MII) chromosome mass. Enucleation of oocytes was undertaken by removing the MII chromosome mass and the 1st polar body. Donor cells were then injected into the perivitelline space of enucleated oocyte recipients.

Electrofusion and activation

Reconstructed oocytes were manually aligned between two wire electrodes (1-mm apart) of a fusion chamber overlaid with 0.3 M mannitol solution containing 0.1 mM MgSO_4 , 0.05 mM CaCl_2 , and 0.1% BSA. A single direct-current pulse of 1.3 kV/cm was applied for 30 μ sec using a BTX Electro Cell Manipulator 200 (BTX, USA). After fusion treatment, the reconstituted oocytes were placed in CR1aa [21] containing 3 mg/mL BSA and checked for fusion. The fused oocytes were then activated using 10 μ M Ca-ionophore (A23187; Sigma-Aldrich) for 5 min and cultured in CR1aa containing 3 mg/mL BSA and 2 mM 6-dimethylaminopurine (DMAP; Sigma-Aldrich) for 3 h.

In vitro fertilization (IVF)

Bovine COCs matured *in vitro* for 22 h were coincubated with frozen-thawed spermatozoa (2×10^6 spermatozoa/mL) in a 50 μ L drop of BO medium [3] supplemented with 5 mM caffeine (Sigma-Aldrich), 10 μ g/mL heparin (Sigma-Aldrich), and 3 mg/mL BSA at 39°C under 5% CO_2 in air for 6 h.

In vitro culture of embryos

After activation or insemination culture, the SCNT and IVF embryos were further cultured in a 50 μ L drop of CR1aa containing 3 mg/mL BSA and 50 μ g/mL gentamicin at 39°C and 5% CO_2 in air prior to analysis of ROS levels or cellular damage (18~20 h post fusion or insemination).

Analysis of ROS products

The ROS levels of SCNT and IVF embryos were measured as previously described [13]. Briefly, SCNT and IVF embryos at the zygote stage were stained with 10 μ M dichlorodihydrofluorescein diacetate (H_2DCFDA ; Molecular Probes, USA) or 10 μ M 3'-(p-hydroxyphenyl) fluorescein (HPF, Molecular Probes) for 30 min at 39°C to determine the H_2O_2 level [11] or the $\cdot\text{OH}$ radical level, respectively [23]. After washing in PBS, the embryos were mounted onto glass slides and examined under a fluorescent microscope (BX-50; Olympus, Japan) with filters at 450~480 nm for excitation and 515 nm for emission. The fluorescent images from the samples were recorded as JPEG files using a digital camera (ESO 600D; Canon, Japan) and analyzed using ImageJ software 1.37 (NIH) based on the intensity of fluorescence in each embryo.

Laser-scanning confocal microscopic visualization and determination of mitochondrial membrane potential ($\Delta\Psi$)

To assess the morphology and distribution patterns of the mitochondria, SCNT and IVF embryos were stained with MitoTracker Red CMXRos dye (Molecular Probes; Invitrogen, USA) at 18~20 h after electrofusion or insemination for 30 min at 37°C. The fluorescent images of equatorial sections were obtained using a laser-scanning confocal microscope (LSM510; Zeiss, Germany) with excitation at 543 nm and emission at 560 nm.

To assess the $\Delta\Psi$, SCNT and IVF embryos were stained with 20 μ g/mL 5,5',6,6'-tetrachloro-1,1,3,3'-tetraethylbenzimidazolylcarbocyanine iodide (JC-1) (Molecular Probes) at 18~20 h after electrofusion or insemination for 30 min at 37°C. The embryos were then mounted in Hepes-buffered TCM199 and visualized using a laser-scanning confocal microscope (LSM510; Zeiss) with excitation at 488 nm and emission at 500~550 nm (green) and 560 nm (red). The images of individual embryos were captured and the ratio of red to green fluorescence intensity was analyzed using the LSM Image Browser (ver. 4.20; Zeiss).

Assessment of apoptosis

Two analyzing methods were used to assess the apoptosis of SCNT embryos. A TUNEL assay was used to determine the apoptosis of SCNT and IVF embryos. The one-cell stage embryos at 18~20 h after electrofusion or insemination were fixed for 1 h in PBS (Gibco BRL) containing 4% (w/v) paraformaldehyde (Sigma-Aldrich) and then permeabilized for 30 min in PBS containing 0.5% Triton X-100 and 0.3% (w/v) polyvinylpyrrolidone (PVP) at 37°C. The fixed embryos were then incubated in TUNEL reaction medium (*In Situ* Cell Death Detection Kit, TMR red; Hoffmann-La Roche, Germany) in the dark for 1 h at 37°C. The embryos were then re-permeabilized for 5 min in

PBS containing 0.5% Triton X-100 and 0.3% (w/v) PVP at room temperature, after which they were stained for 30 min with 10 $\mu\text{g}/\text{mL}$ of Hoechst 33342 (Sigma-Aldrich) at 37°C and mounted on a glass slide in Vecta-Shield anti-fade (Vector Laboratories, USA) under a cover slip. The apoptosis of embryos was determined by fluorescence microscopy.

A caspase-3 activity assay was performed using a Caspase-3 Colorimetric Activity Assay Kit (EMD Millipore, USA). Briefly, 20 individual one-cell stage SCNT and IVF embryos were put into 1-mL tubes containing 50 μL cell lysis buffer (1 \times) and lysed for 15 min on ice at 18~20 h after electrofusion or insemination. Cell lysates were centrifuged for 5 min at 7,000 \times g, after which supernatants were transferred into 100 μL Assay Mixture containing 10 μL Caspase-3 substrate prepared in a 96-well plate and incubated for 1 h at 37°C. The enzyme reactions were then colorimetrically monitored at 405 nm with a microtiter plate reader (Power Wave XS; BioTek, USA) by reading the OD values using the Gene 5 software.

Comet assay

The DNA damage in SCNT and IVF embryos was analyzed by comet assay at 18~20 h after electrofusion or insemination. A comet assay was performed using an OxiSelect Comet Assay Kit (Cell Biolabs, USA). Prior to assay, OxiSelect comet agarose (Assay kit no. 235002; Cell Biolabs) was melted at 90°C for 20 min and then cooled at 37°C for 20 min. About 75 μL of agarose was then dropped onto an OxiSelect 3-well comet slide (Assay kit no. 235001; Cell Biolabs), after which approximately ten zona-free embryos were transferred to agarose drops and chilled at 4°C for 20 min. The samples were then lysed in lysis buffer [250 mM NaCl, 20% EDTA solution (Assay kit no. 235004; Cell Biolabs), 10% DMSO, 10% 10 \times lysis solution (Assay kit no. 235005; Cell Biolabs), pH ~10.0] at 4°C for 3 h. Next, the slides were carefully transferred to chilled alkaline solution (300 mM NaOH, 1 mM EDTA) and immersed for 30 min at 4°C. Subsequently, the slides were transferred to a horizontal electrophoresis chamber (Mupid α ; Advance Electric, Japan) filled with cold TAE buffer (ELPIS Biotech, Korea) and electrophoresed for 20 min at 50 V, after which the slides were stained with 1 \times Vista Green DNA Dye (Assay kit no. 235003; Cell Biolabs) for 20 min and then examined using a fluorescent microscope with a FITC filter. All steps after agarose treatment were conducted in the dark to prevent additional DNA damage. The comet tail lengths were measured in individual embryos using CASP (ver. 1.2.2; Zbigniew Koza, Poland).

Statistical analysis

Data were analyzed by ANOVA using the Statistical Analysis System software package (SAS Institute, USA).

Significant differences between means were identified by Fisher's least-significant-difference (LSD) test.

Results

ROS levels of SCNT and IVF embryos

The H_2O_2 and $\cdot\text{OH}$ radical levels of SCNT embryos were significantly higher ($p < 0.05$) (35.6 ± 1.1 and 44.6 ± 1.2 pixels/embryo, respectively) than those of the IVF embryos (19.2 ± 1.5 and 23.8 ± 1.8 pixels/embryo, respectively, Fig. 1).

Mitochondrial morphology and membrane potential ($\Delta\Psi$)

The mitochondrial morphology of IVF embryos exhibited a granular or punctate pattern, which is typical of normal mitochondria (Fig. 2A). Conversely, the mitochondrial morphology of SCNT embryos was characterized by a diffuse and non-granularity cytoplasmic staining pattern (Fig. 2B). IVF embryos showed intense red JC-1 staining with active $\Delta\Psi$, whereas SCNT embryos showed weak red JC-1 staining with reduced or inhibited $\Delta\Psi$ (Fig. 2C and D). Quantitative analysis revealed that mitochondria $\Delta\Psi$ (red/green) was significantly lower ($p < 0.05$) in SCNT embryos (0.95 ± 0.04) than IVF embryos (1.21 ± 0.06 , Fig. 3).

Apoptosis

TUNEL Assay (Fig. 2E and F) revealed that no apoptosis had occurred in SCNT embryos or IVF embryos at the one-cell stage (a total of 50~55 embryos in each group, data not shown). Caspase-3 activity assay also confirmed




Fig. 1. Levels of reactive oxygen species (ROS) in bovine somatic cell nuclear transfer (SCNT) and *in vitro* fertilized (IVF) embryos. Five replicates were performed ($n = 50 \sim 55$ embryos in each group). The SCNT and IVF embryos were analyzed on the same day, enabling direct comparisons between SCNT and IVF groups. Data are presented as the mean \pm SEM (bars). *Significantly different from IVF within each treatment ($p < 0.05$).




Fig. 2. Fluorescence microscopic evidence of mitochondrial and DNA damage in one-cell stage IVF (A, C, E and G) and SCNT (B, D, F and H) embryos. (A and B) MitoTracker Red staining images of embryos. (C and D) JC-1 stained embryos. (E and F) TUNEL images of embryos. (G and H) Comet images of fragmented DNA migration of embryos. Scale bars = 50 μm .




Fig. 3. Mitochondrial membrane potential ($\Delta\Psi$) of IVF and SCNT embryos (n = 40 ~ 50 in each group). Data are presented as the mean \pm SEM (bars). *Significantly different from IVF ($p < 0.05$).




Fig. 4. Tail moment length of IVF (n = 37) and SCNT (n = 38) embryos. Data are presented as the mean \pm SEM (bars). *Significantly different from IVF ($p < 0.05$).

the results of the TUNEL assay. There was no difference in the OD values of SCNT (OD 0.056, n = 101) and IVF (OD 0.057, n = 107) embryos, which were similar to that of the standard blank (OD 0.055).

DNA fragmentation

Comet assay results revealed that both IVF and SCNT

embryos exhibited a clearly defined comet tail (Fig. 2G and H). However, the length of migrated DNA fragments in SCNT embryos (515.5 \pm 26.4 μm) was found to be significantly ($p < 0.05$) longer than that of IVF embryos (425.6 \pm 25.0 μm , Fig. 4).

Discussion

The results of our previous study suggested that cellular stress might occur through micromanipulation, resulting in generation of a high level of ROS production [13]. In the present study, the ROS level of SCNT embryos at the zygote stage was higher than that of IVF embryos, which confirmed the results of the previous study, in which micromanipulation procedures were found to produce ROS in the cytoplasm of SCNT embryos [13]. In the present study, we focused on evaluation of cellular damage that may be induced by excessive ROS generated during the SCNT procedure; thus, we did not evaluate the development and ROS level in later stage embryos.

ROS induces morphological and metabolic damage to mitochondria, indicative of mitochondrial dysfunction, followed by apoptotic degeneration [28]. Lipid peroxides that are induced by ROS affect vital mitochondrial functions, such as respiration and oxidative phosphorylation, inner membrane barrier properties, and maintenance of $\Delta\Psi$ [2,30]. It has also been suggested that ROS activates mitochondrial permeability transition (mPT) pores that allow the release of proapoptotic factors such as mitochondrial cytochrome C [8]. In this study, the mitochondrial morphology of SCNT embryos was characterized by a diffuse and non-granular cytoplasmic staining pattern. Unlike the granular pattern of intact mitochondria, the diffuse and non-granular pattern was interpreted to indicate the release of mitochondrial CMXRos back into the cytoplasm and as evidence of an mPT [8]. In this study, $\Delta\Psi$ of SCNT embryos was lower relative to IVF embryos, which is also suggestive of an mPT [28]. It has been reported that the metabolic consequence of sustained mPT is a phenomenon of mitochondrial uncoupling [27]. Uncoupled mitochondria are characterized by decreased inner mitochondrial membrane integrity and proton gradient, and subsequent reduced or abolished ATP synthesis [28].

It has been reported that higher levels of $\cdot\text{OH}$ radical derived from H_2O_2 induce lipid peroxidation [24] and DNA damage [9]. DNA damage induced by ROS is involved in apoptosis [29] and DNA fragmentation [12,16], which can lead to abnormal gene expression as well as suppression of further embryonic development [16]. Although no apoptosis was detected in SCNT embryos at the one-cell stage by TUNEL and caspase-3 assays in this study, apoptotic nuclei might increase during *in vitro* development of the SCNT embryos when compared with IVF embryos. In a previous study, apoptotic nuclei were detected from the four-cell stage and increased greatly during further *in vitro* culture of SCNT bovine embryos when compared to those of IVF embryos [6]. In another study, no TUNEL-positive nuclei were observed in IVF or SCNT porcine early embryos prior to day 5 of *in vitro* culture; however, the apoptotic rate of

porcine SCNT embryos increased with culture time from day 5 to 8 and was significantly higher than in the IVF embryos [10].

The $\cdot\text{OH}$ radical assaults the deoxyribose moiety of DNA to trigger strand breaks [22,26], which induces mispairing during DNA polymerization [18] and irreversible double-strand blocks at the time of DNA replication [5]. DNA fragmentation can be detected by comet assay [25], which has been widely accepted as a sensitive tool for assessing DNA damage in animal cells. In the present study, although no apoptosis was detected by TUNEL and Caspase-3 assays, comet tail movement was observed in both SCNT and IVF one-cell embryos. The comet tail length of SCNT embryos was found to be significantly longer than that of IVF embryos, indicating that the DNA fragmentation of SCNT embryos is induced with high frequency. Such DNA fragmentation in the zygote stage SCNT embryos would result in further increases in irreversible double-strand breaks of DNA through DNA replications during embryo development [5].

In conclusion, the results of the present study show that cellular stress during micromanipulation procedures can generate ROS and subsequently induce mitochondrial and DNA damage in bovine SCNT embryos, which could lead to abnormal nuclear reprogramming of SCNT embryos.

Acknowledgments

We thank Dr. Seung-Hae Kwon of the Korea Basic Science Institute, Chuncheon, Korea, for his assistance with confocal microscopy. The results of this study were presented at the 37th IETS annual conference, 2011. This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2008-313-F00067) and Institute of Veterinary Science, Kangwon National University, Korea.

References

1. **Aitken RJ, Clarkson JS, Fishel S.** Generation of reactive oxygen species, lipid peroxidation, and human sperm function. *Biol Reprod* 1989, **40**, 183-197.
2. **Albano E, Bellomo G, Parola M, Carini R, Dianzani MU.** Stimulation of lipid peroxidation increases the intracellular calcium content of isolated hepatocytes. *Biochimica et Biophysica Acta* 1991, **1091**, 310-316.
3. **Brackett BG, Oliphant G.** Capacitation of rabbit spermatozoa *in vitro*. *Biol Reprod* 1975, **12**, 260-274.
4. **Choi JY, Kim CI, Park CK, Yang BK, Cheong HT.** Effect of activation time on the nuclear remodeling and *in vitro* development of nuclear transfer embryos derived from bovine somatic cells. *Mol Reprod Dev* 2004, **69**, 289-295.
5. **Cox MM, Goodman MF, Kreuzer KN, Sherratt DJ, Sandler SJ, Mariani KJ.** The importance of repairing

- stalled replication forks. *Nature* 2000, **404**, 37-41.
6. **Fahrudin M, Otoi T, Karja NWK, Mori M, Murakami M, Suzuki T.** Analysis of DNA fragmentation in bovine somatic nuclear transfer embryos using TUNEL. *Reproduction* 2002, **124**, 813-819.
 7. **Garry FB, Adams R, McCann JP, Odde KG.** Postnatal characteristics of calves produced by nuclear transfer cloning. *Theriogenology* 1996, **45**, 141-152.
 8. **Green DR, Reed JC.** Mitochondria and apoptosis. *Science* 1998, **281**, 1309-1312.
 9. **Halliwell B, Aruoma OI.** DNA damage by oxygen-derived species: its mechanism and measurement in mammalian systems. *FEBS Lett* 1991, **281**, 9-19.
 10. **Hao Y, Lai L, Mao J, Im GS, Bonk A, Prather RS.** Apoptosis and in vitro development of preimplantation porcine embryos derived in vitro or by nuclear transfer. *Biol Reprod* 2003, **69**, 501-507.
 11. **Hashimoto S, Minami N, Yamada M, Imai H.** Excessive concentration of glucose during in vitro maturation impairs the developmental competence of bovine oocytes after in vitro fertilization: relevance to intracellular reactive oxygen species and glutathione contents. *Mol Reprod Dev* 2000, **56**, 520-526.
 12. **Henle ES, Linn S.** Formation, prevention, and repair of DNA damage by iron/hydrogen peroxide. *J Biol Chem* 1997, **272**, 19095-19098.
 13. **Hwang IS, Bae HK, Park CK, Yang BK, Cheong HT.** Generation of reactive oxygen species in bovine somatic cell nuclear transfer embryos during micromanipulation procedures. *Reprod Dev Biol* 2012, **36**, 49-53.
 14. **Inoue K, Kohda T, Lee J, Ogonuki N, Mochida K, Noguchi Y, Tanemura K, Kaneko-Ishino T, Ishino F, Ogura A.** Faithful expression of imprinted genes in cloned mice. *Science* 2002, **295**, 297.
 15. **Kang YK, Koo DB, Park JS, Choi YH, Chung AS, Lee KK, Han YM.** Aberrant methylation of donor genome in cloned bovine embryos. *Nat Genet* 2001, **28**, 173-177.
 16. **Kitagawa Y, Suzuki K, Yoneda A, Watanabe T.** Effect of oxygen concentration and antioxidants on the in vitro developmental ability, production of reactive oxygen species (ROS), and DNA fragmentation in porcine embryos. *Theriogenology* 2004, **62**, 1186-1197.
 17. **Kwon DJ, Lee YM, Hwang IS, Park CK, Yang BK, Cheong HT.** Microtubule distribution in somatic cell nuclear transfer bovine embryos following control of nuclear remodeling type. *J Vet Sci* 2010, **11**, 93-101.
 18. **Marnett LJ.** Oxyradicals and DNA damage. *Carcinogenesis* 2000, **21**, 361-370.
 19. **Raha S, Robinson BH.** Mitochondria, oxygen free radicals, disease and ageing. *Trends Biochem Sci* 2000, **25**, 502-508.
 20. **Rhoads DM, Umbach AL, Subbaiah CC, Siedow JN.** Mitochondrial reactive oxygen species. Contribution to oxidative stress and interorganellar signaling. *Plant Physiol* 2006, **141**, 357-366.
 21. **Rosenkrans CF Jr, First NL.** Effect of free amino acids and vitamins on cleavage and developmental rate of bovine zygotes in vitro. *J Anim Sci* 1994, **72**, 434-437.
 22. **Salgo MG, Stone K, Squadrito GL, Battista JR, Pryor WA.** Peroxynitrite causes DNA nicks in plasmid pBR322. *Biochem Biophys Res Commun* 1995, **210**, 1025-1030.
 23. **Setukinai KI, Urano Y, Kakinuma K, Majima HJ, Nagano T.** Development of novel fluorescence probes that can reliably detect reactive oxygen species and distinguish specific species. *J Biol Chem* 2003, **31**, 3170-3175.
 24. **Slater TF.** Free-radical mechanisms in tissue injury. *Biochem J* 1984, **222**, 1-15.
 25. **Takahashi M, Saka N, Takahashi H, Kanai Y, Schultz RM, Okano A.** Assessment of DNA damage in individual hamster embryos by comet assay. *Mol Reprod Dev* 1999, **54**, 1-7.
 26. **Takahashi M, Nagai T, Hamano S, Kuwayama M, Okamura N, Okano A.** Effect of thiol compounds on in vitro development and intracellular glutathione content of bovine embryos. *Biol Reprod* 1993, **49**, 228-232.
 27. **Thompson JG, McNaughton C, Gasparrini B, McGowan LT, Tervit HR.** Effect of inhibitors and uncouplers of oxidative phosphorylation during compaction and blastulation of bovine embryos cultured in vitro. *J Reprod Fertil* 2000, **118**, 47-55.
 28. **Thouas GA, Trounson AO, Wolvetang EJ, Jones GM.** Mitochondrial dysfunction in mouse oocytes results in preimplantation embryo arrest in vitro. *Biol Reprod* 2004, **71**, 1936-1942.
 29. **Yang HW, Hwang KJ, Kwon HC, Kim HS, Choi KW, Oh KS.** Detection of reactive oxygen species (ROS) and apoptosis in human fragmented embryos. *Hum Reprod* 1998, **13**, 998-1002.
 30. **Zhang Y, Marcillat O, Giulivi C, Ernster L, Davies KJ.** The oxidative inactivation of mitochondrial electron transport chain components and ATPase. *J Biol Chem* 1990, **265**, 16330-16336.