

Bioactivity studies on *Musa seminifera* Lour

Sanjib Saha, Jamil A. Shilpi, Himangsu Mondal, Royhan Gofur¹, Morsaline Billah², Lutfun Nahar³, Satyajit D. Sarker⁴

Pharmacy Discipline, Khulna University, Khulna 9208, ¹Department of Animal Husbandry and Veterinary Science, Rajshahi University, Rajshahi 6205, ²Department of Biotechnology and Genetic Engineering Discipline, Khulna University, Khulna 9208, Bangladesh, ³Leicester School of Pharmacy, De Montfort University, The Gateway, Leicester LE1 9BH, ⁴Department of Pharmacy, School of Applied Sciences, University of Wolverhampton, Wolverhampton WV1 1LY, West Midlands, UK

Submitted: 05-09-2012

Revised: 04-11-2012

Published: 07-09-2013

ABSTRACT

Background: *Musa seminifera* Lour is a tree-like perennial herb that has been used in folk medicine in Bangladesh to heal a number of ailments. **Objective:** To evaluate the antioxidant, analgesic, antidiarrheal, anthelmintic activities, and general toxicity of the ethanol extract of the roots. **Materials and Methods:** The extract was assessed for free-radical-scavenging activity by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay, total phenolic content (TPC) by the Folin Ciocalteu reagent, antioxidant activity by the ferric reducing power assay, analgesic activity by the acetic acid-induced writhing and hot-plate tests, antidiarrheal activity by the castor oil-induced diarrhea model in mice, anthelmintic activity on *Paramphistomum cervi* and *Haemonchus contortus*, and general toxicity by the brine shrimp lethality assay. **Results:** The extract showed free-radical-scavenging activity with an IC_{50} value of 44.86 $\mu\text{g/mL}$. TPC was 537.89 mg gallic acid equivalent/100 g of dried plant material. It showed concentration-dependent reducing power, and displayed 42.11 and 69.32% writhing inhibition at doses of 250 and 500 mg/kg body weight, respectively. The extract also significantly raised the pain threshold at the above-mentioned dose levels. In vivo antidiarrheal property was substantiated by significant prolongation of latent period and decrease in total number of stools compared with the control. The LC_{50} against brine shrimp nauplii was 36.21 $\mu\text{g/mL}$. The extract exhibited dose-dependent decrease in paralysis and death time of the helminths. **Conclusion:** The above results demonstrated that the plant possesses notable bioactivities and somewhat supports its use in folk medicine.

Key words: Analgesic activity, anthelmintic activity, brine shrimp lethality, 2,2-diphenyl-1-picrylhydrazyl, Folin Ciocalteu's reagent, reducing power

INTRODUCTION

Musa seminifera Lour (Synonym: *M. sylvestris*, fam: Musaceae) is native to Asia, grows in 5-9 m in height with tuberous rhizome, hard pseudo-stem, and a crown of large extended green leaves. The inflorescence can be consumed as vegetable. The ripe fruits are sweet, juicy, and full of seeds. The peel is thicker than other species of the genus *Musa*. In Bangladesh, this plant is commonly known as Bichi kola, Aitta kola, or Doia kola and distributed throughout the country. The fruit is used in the treatment of diarrhea, dysentery, and excess menstruation.^[1] The flower is used to treat diarrhea, diabetes, and menorrhagia. Stem juice is used to treat cholera, hemoptysis, otalgia, and dysentery.^[2]

Address for correspondence:

Prof. Satyajit D. Sarker, Department of Pharmacy, School of Applied Sciences, University of Wolverhampton, Wulfruna Street, Wolverhampton WV1 1LY, West Midlands, UK.
E-mail: s.sarker@wlv.ac.uk

There are no reports available on any bioactivity studies on the roots of this species. However, other species of this genus, e.g., *Musa acuminata*, *Musa paradisiaca*, and *Musa sapientum*, were studied before. The antidiarrheal, antioxidant, and antimicrobial activities of the seeds^[3] and hypoglycemic activity of the flowers^[4] of *M. sapientum* were studied. The fruits of *M. sapientum* revealed antibacterial, wound-healing, and antiallergic properties.^[5-7] Phytochemical analysis on *M. sapientum* indicated the presence of triterpenes (β -amyrin, cyclomusalenone, cyclomusalenol, 24-methylenecycloartanol stigmast-7-en-3-ol, stigmast-7-methylenecycloartanol), flavonoids (quercetin and its 3-O-glucoside, 3-O-galactoside and 3-O-rhamnosyl glucoside),^[8] 4-epicyclomusalenone, and 4-epicycloeucalenone.^[9] Antihyperglycemic effect of the roots of *M. paradisiaca* was investigated in streptozotocin-induced diabetic male Albino rats.^[10] Inhibition of dehydrogenase activity in pathogenic bacteria was studied for aqueous fruit peel and leaf extracts of *M. paradisiaca*.^[11] Chemical investigation on *M. acuminata* led to the isolation

Access this article online

Website:

www.phcog.com

DOI:

10.4103/0973-1296.117827

Quick Response Code:



of 4,4'-dihydroxy-anigorootin, 4'-hydroxy-anigorootin, 3,3'-bishydroxyanigorufone,^[12] (S)-(+)-6-methoxy- α -methyl-2-naphthaleneacetic acid,^[13] and oxabenzochrysenones.^[14]

As part of our ongoing studies on medicinal plants from the Bangladeshi flora,^[15-30] we now, for the very first time, report on the antioxidant, analgesic, antidiarrheal, anthelmintic activity, and general toxicity of the ethanol extract of the roots of *M. seminifera*.

MATERIALS AND METHODS

Plant material

Roots of *M. seminifera* were collected from Khulna University area, Khulna, Bangladesh, in November 2011. A voucher specimen (DACB 37523) was lodged at the Bangladesh National Herbarium, Dhaka.

Extraction

The roots were washed with water, shed dried, ground, and soaked in ethanol for 3 days with occasional shaking. The extract was filtered through a cotton plug and dried using a rotary evaporator at 50°C to obtain crude extract (yield 1.83%).

Test animals

Young Swiss Albino mice aged 4-5 weeks, average weight 20-25 g, procured from the International Centre for Diarrheal Disease and Research, Bangladesh (ICDDR, B) were used. The mice were provided with ICDDR, B formulated rodent food, and water *ad libitum*. The experiments were conducted strictly in accordance with the animal ethics guidelines^[31] and the ethical guidelines and permission provided by the ICDDR, B.

Test pathogens

Live parasites *Paramphistomum cervi* (Trematoda) and *Haemonchus contortus* (Nematode) were collected from freshly slaughtered cattle at local abattoirs and identified by Dr. Md. Royhan Gofur (Rajshahi University, Rajshahi). After cleaning, parasites were stored in 0.9% phosphate-buffered saline (PBS) of pH 7.4 prepared with 8.01 g NaCl, 0.20 g KCl, 1.78 g Na₂HPO₄, and 0.27 g KH₂PO₄ in 1 L of distilled water at 37 ± 1°C.

Chemicals and reagents

Folin Ciocalteu's reagent and 2,2-diphenyl-1-picrylhydrazyl (DPPH) were purchased from Sigma-Aldrich, USA. Gallic acid, ascorbic acid, trichloroacetic acid, potassium ferricyanide, sodium carbonate, acetic acid, dimethyl sulfoxide (DMSO), and ferric chloride were purchased from Merck, Germany. Castor oil and tween-80 were procured from Loba Chemie Pvt., Ltd, India. Solvents and all other chemicals used in the study were of analytical

grade. Standard vincristine sulphate was obtained from Cipla Pharmaceuticals, India. Standard diclofenac sodium and albendazole were from Beximco Pharmaceuticals Ltd, Bangladesh. Standard morphine was collected from Popular Pharmaceuticals Ltd, Bangladesh.

Determination of DPPH radical scavenging activity

In vitro antioxidant activity was determined by the DPPH assay.^[32] Stock solution (512 µg/mL) of the root extract was prepared in ethanol, and was serially diluted with ethanol to obtain concentrations of 256, 128, 64, 32, 16, 8, 4, 2, and 1 µg/mL. Serially diluted solutions (1 mL) were mixed with 3 mL DPPH solution (0.004% in ethanol) and kept for 30 min to complete any reaction that occurs. The absorbance was recorded at 517 nm. Ascorbic acid was used as the positive standard. Percent inhibition was calculated using the formula $(1 - A_1/A_0) \times 100$, where A₀ is the absorbance of control and A₁ is the absorbance of sample or standard. IC₅₀ was determined from % inhibition versus concentration (µg/ml) plot.

Determination of total phenolic content (TPC)

Ground roots (0.5 g) were mixed with 80% aq. MeOH (50 mL), sonicated for 20 min, and a portion (2 mL) was centrifuged for 15 min at 14,000 rpm. TPC was determined by the Folin Ciocalteu reagent.^[33] Gallic acid solution in MeOH was prepared at 500, 250, 125, 62.5, 31.25, and 15.62 mg/L concentrations. From each of the concentrations, and from the extract, a volume of 1 mL was transferred to a 25 mL volumetric flask and 9 mL distilled water was added. Diluted (1 mL reagent in 9 mL distilled water) Folin Ciocalteu's reagent (1 mL) was added and mixed by continuous shaking. After 5 min, 7% Na₂CO₃ (10 mL) was added to each flask and distilled water was added to make the final volume of 25 mL. Keeping for 30 min at room temperature, absorbance was measured at 750 nm against blank, which was prepared by following all the above-mentioned steps except for the addition of extract or gallic acid. Standard curve was prepared by plotting absorbance versus respective concentration of gallic acid. TPC of the extract was determined from the standard curve and expressed as mg gallic acid equivalent (GAE)/100 g dried plant material.

Reducing power assay

Reducing power was determined by the method described by Oyaizu (1986).^[34] The root extract (1 mL) at different concentrations (15.62-500 µg/mL) was mixed with phosphate buffer (2.5 mL, 0.2 M, pH 6.6) and 2.5 mL of 1% potassium ferricyanide [K₃Fe(CN)₆]. The mixture was incubated at 50°C for 20 min. Trichloroacetic acid (10%, 2.5 mL) was added, and centrifuged at 3000 rpm for 10 min. The supernatant (2.5 mL) was mixed with distilled water (2.5 mL) and FeCl₃ (0.1%, w/v, 0.5 mL). Reducing power was determined by measuring the formation of

Pearl's Prussian blue at 700 nm. Ascorbic acid was used as the positive standard.

Evaluation of analgesic activity

Randomly screened animals were divided into four groups – Groups I, II, III, and IV – consisting of six mice in each. Groups I and II were considered as control and positive control, respectively, whereas groups III and IV were test groups.

Acetic acid-induced writhing test

Mice ($n = 6$) were administered orally with diclofenac sodium 25 mg/kg body weight as standard drug, root extract at the doses of 250 and 500 mg/kg and control as 1% Tween-80 in distilled water. Thirty minutes after the treatment, the mice were subjected to intraperitoneal injection of 0.7% acetic acid at the dose of 10 mL/kg to induce writhing. After 5 min of acetic acid administration, the writhes were counted for 10 min. The numbers of writhes of standard and extract were compared to controls.^[35]

Hot-plate test

Animals were screened based on their reaction time at 3-5 s when subjected to pain stimulus. Mice were placed on a hot plate at $55 \pm 0.5^\circ\text{C}$. Reaction time was recorded when mice licked their fore and hind paws and jumped at before (0 min) and 30, 60, 90, and 120 min after oral administration of extract at the doses of 250 and 500 mg/kg, standard drug morphine, at the dose of 5 mg/kg (i.p.) and control as 1% Tween-80 in distilled water (10 mL/kg, p.o.). Cutoff point was considered 15 s to avoid accidental damage of the paws of mice.^[36]

Evaluation of in vivo antidiarrheal activity

In vivo antidiarrheal activity was investigated by castor oil-induced diarrheal episode in mice.^[37] Test animals were divided into control, positive control, and test groups containing six mice in each group. The control group received 1% Tween-80 in distilled water (10 mL/kg) whereas the positive control group received standard loperamide (2 mg/kg). The test groups received root extract at doses of 250 and 500 mg/kg. All the treatments were administered orally, 1 h prior to oral administration of castor oil at the dose of 0.5 mL per mouse. Each mouse was placed in an individual cage and the floor was lined with blotting paper. The blotting paper was changed every hour. The observation period was 4 h. The latent period and total stool count after 4 h were recorded. Antidiarrheal activity was expressed as the prolongation of latent period and percentage of inhibition of defecation.

Brine shrimp lethality assay for general toxicity

General toxicity of the extract was tested by the brine shrimp lethality assay.^[38] Artificial sea water was prepared by dissolving 20 g of NaCl and 18 g of table salt in 1 L of distilled water and was filtered off to obtain a clear solution. A rectangular tank was divided into two

unequal compartments by a porous separator. The larger compartment was darkened while the smaller one was kept illuminated. Eggs of *Artemia salina* were hatched at $25\text{--}30^\circ\text{C}$ for 24-48 h. The larvae (nauplii) were attracted by the light and moved to the smaller compartment through the holes. They were then collected by a pipette. Extract was dissolved in distilled water with DMSO and transferred to test tubes to obtain concentrations of 320, 160, 80, 40, 20, 10, and 5 $\mu\text{g}/\text{mL}$ in 5 mL artificial sea water with ten nauplii in each test tube. Anticancer drug vincristine sulphate was used as positive control at concentrations of 5, 2.5, 1.25, 0.625, and 0.312 $\mu\text{g}/\text{mL}$. The concentration of DMSO did not exceed 0.01% in any of the test tubes. Control test tubes were subjected to DMSO in artificial seawater at the same concentration as in test tubes for samples. After 24 h incubation at $25\text{--}30^\circ\text{C}$, the number of viable nauplii was counted using a magnifying glass.

Evaluation of anthelmintic activity

Anthelmintic activity of the root extract was investigated on live parasites *P. cervi* and *H. contortus* of cattle.^[39,40] The parasites were divided into different groups consisting of six parasites in each group. Extracts at concentrations of 25, 50, 100, and 200 mg/mL and reference standard albendazole at the concentration of 15 mg/mL of 10 mL in PBS were prepared and transferred to Petri dishes. The control group was treated with 0.1% Tween-80 in PBS. Six parasites were placed in each Petri dish and observed. The time of paralysis was recorded when no movement was observed unless shaken vigorously. The death time was recorded after ensuring that the parasites did not move when shaken vigorously, dipped in warm water (50°C), or subjected to external stimuli. Anthelmintic activity was expressed as the time required for paralysis and death of parasites as compared to control.

Statistical analysis

All the values were expressed as mean \pm Standard error for mean (SEM). Student's *t*-test was used to estimate significant differences between the test and control groups. Statistical analysis was performed with Prism 5.0 (GraphPad software Inc., San Diego, CA). Results were considered as statistically significant when the *P* value was less than the alpha factor (0.05).

RESULTS

DPPH scavenging activity

Both the extract and ascorbic acid showed a linear increase in DPPH scavenging activity at lower concentrations until they reached a plateau at higher concentrations [Figure 1]. IC_{50} values for both test and standard were within the linear region. Using the values of the linear region, IC_{50} of the extract and ascorbic acid was determined as 44.86 and 13.25 $\mu\text{g}/\text{mL}$, respectively.

TPC determination

The TPC of *M. seminifera* root extract was 537.89 mg GAE/100 g of dried plant material [Figure 2].

Reducing power assay

The root extract of *M. seminifera* displayed concentration-dependent increase in the reducing power. At the concentrations of 15.62, 31.25, 62.5, 125, and 250 µg/mL, the extract showed absorbance of 0.482, 0.528, 0.657, 0.687, and 0.758, whereas at the same concentrations, standard ascorbic acid showed absorbance of 0.725, 0.855, 1.236, 1.688, and 2.523, respectively [Figure 3].




Figure 1: 2,2-diphenyl-1-picrylhydrazyl scavenging activity of *Musa seminifera* root




Figure 2: Standard calibration curve of gallic acid




Figure 3: Reducing power of *Musa seminifera* root

The acetic acid-induced writhing test

The extract revealed significant and dose-dependent pain inhibition at both dose levels tested. The extract showed 42.11% and 69.32% writhing inhibition at doses of 250 and 500 mg/kg body weight, respectively, whereas standard diclofenac sodium (25 mg/kg) showed 78.95% writhing inhibition [Table 1].

The hot-plate test

The extract significantly and dose-dependently increased the pain threshold. The extract showed maximum reaction times of 5.77 and 6.48 s at doses of 250 and 500 mg/kg body weight, respectively, whereas morphine (5 mg/kg) showed a reaction time of 9.30 s [Table 2].

In vivo antidiarrheal activity

The extract showed significant and dose-dependent prolongation of latent period as well as decrease in number of stools throughout the whole observation period compared to control. The extract demonstrated 57.89% and 77.19% inhibition of defecation at doses of 250 and 500 mg/kg body weight, respectively, whereas loperamide (3 mg/kg) showed 88.58% inhibition of defecation [Table 3].

The brine shrimp lethality assay

LC₅₀ was calculated using LdP line Probit analysis software, USA. The LC₅₀ for *M. seminifera* root extract was found to be 36.21 µg/mL and that of vincristine sulphate was 0.43 µg/mL.

Anthelmintic activity

The root extract revealed significant anthelmintic activity in a concentration-dependent manner. The extract, at concentrations of 25, 50, 100, and 200 mg/mL exhibited paralysis at 24.17, 11.17, 6.16, and 2.25 min and death at 33.33, 21.67, 9.50, and 5.16 min, respectively, for *P. cervi*, and paralysis at 39.33, 23.00, 13.50, and 7.16 min and death at 51.33, 32.00, 22.83, and 13.00 min, respectively, for *H. contortus*. Standard drug albendazole at the concentration of 15 mg/mL showed paralysis at 12.50 min and death at 20.00 min for *P. cervi* and paralysis at 24.17 min and death at 35.50 min for *H. contortus* [Table 4].

Table 1: Effect of *Musa seminifera* root on acetic acid induced writhing in mice

Treatment (n=6)	Dose (mg/kg)	No. of writhes	% Inhibition
Control	1% tween-80 in water	19±0.73	-
Diclofenac sodium	25	4±0.51*	78.95
Extract 250	250	11±0.77**	42.11
Extract 500	500	5.83±0.65*	69.32

Values are expressed as mean±SEM, SEM: Standard error for mean, n: Number of mice (6), *P<0.001 and **P<0.01 versus control, Student's t-test

Table 2: Effect of *Musa seminifera* root in hot-plate test in mice

Treatment (n=6)	Dose (mg/kg)	Reaction time(s)				
		0 min	30 min	60 min	90 min	120 min
Morphine	5	3.52±0.12*	6.24±0.13**	7.32±0.13**	9.30±0.12**	8.20±0.08**
Extract 250	250	3.50±0.10*	4.37±0.07**	5.77±0.08**	5.05±0.04**	4.02±0.07**
Control	1% tween-80 in water	3.15±0.06	3.26±0.04	3.27±0.08	3.23±0.07	3.23±0.06
Extract 500	500	3.47±0.11*	5.15±0.09**	6.48±0.11**	5.90±0.09**	4.90±0.08**

Values are expressed as mean±SEM, SEM: Standard error for mean, n: Number of mice (6), *P<0.05 and **P<0.001 versus control, Student's t-test

Table 3: Effect of *Musa seminifera* root on castor oil induced diarrhea in mice

Treatment (n=6)	Dose (mg/kg)	Onset of diarrhea (min)	No. of stools after 4 h	% Inhibition of defecation
Control	1% tween-80 in water	33±3.59*	22.80±1.63*	-
Loperamide	3	197.40±0.77*	2.60±0.50*	88.59
Extract 250	250	101±3.64*	9.60±0.74*	57.89
Extract 500	500	178.80±5.00*	5.20±0.58*	77.19

Values are expressed as mean±SEM, SEM: Standard error for mean, n: Number of mice (6), *P<0.001 versus control, Student's t-test

Table 4: Anthelmintic activity of *Musa seminifera* root

Treatment (n=6)	Conc.(mg/ml)	Time (min) taken for paralysis (P) and death (D)			
		Paramphistomum cervi		Haemonchus contortus	
		P	D	P	D
Control	0.1% Tween-80 in PBS	-	-	-	-
Albendazole	15	12.50±0.42*	20.00±0.57*	24.17±0.47*	35.50±0.42*
Extract 25	25	24.17±0.65*	33.33±0.80*	39.33±0.66*	51.33±0.49*
Extract 50	50	11.17±0.47*	21.67±0.71*	23.00±0.63*	32.00±0.57*
Extract 100	100	6.16±0.40*	9.50±0.42*	13.50±0.56*	22.83±0.70*
Extract 200	200	2.25±0.25*	5.16±0.40*	7.16±0.30*	13.00±0.51*

Values are expressed as mean±SEM, SEM: Standard error for mean, n: Number of parasites (6), * P<0.001 versus control, Student's t-test, PBS: Phosphate-buffered saline

DISCUSSION

Free-radicals are responsible for many disease conditions including rheumatoid arthritis, cardiovascular disorders, cysticfibrosis, inflammation, neurodegenerative diseases (e.g., Parkinsonism, Alzheimer's disease), AIDS, and cancer.^[41] Antioxidants can prevent these disease conditions by scavenging harmful free-radicals. *M. seminifera* showed potential antioxidant activity in the DPPH radical-scavenging assay. The antioxidant activity of plant extracts is generally due to phenolic components such as flavonoids, chalcones, and polyhydroxy benzoic acid derivatives.^[42] The TPC of the extract was much higher than some of the phenol-rich fruits such as strawberry and plum (244.1 and 303 mg GAE/100 g, respectively);this supports the observed strong antioxidant activity.^[33] Besides neutralizing free radicals, reducing power may serve as a significant indicator to assess the antioxidant activity of natural

extracts.^[43] The reducing power assay is based on the fact that the substances having reaction potential react with potassium ferricyanide (Fe_3^+) to reduce into potassium ferrocyanide (Fe_2^+), which reacts with ferric chloride to form ferric ferrous complex (Prussian's blue) that has an absorption maximum at 700 nm. Reducing ability was increased with the increase in concentration of the extract, which indicated that some compounds in the extract are able to react with free radicals to convert them into stable forms as well as to terminate chain reactions.

The extract showed analgesic activity in both of the methods. The abdominal constriction response, i.e., 'writhing' induced by acetic acid, is a model to assess peripherally acting analgesic activity.^[44,45] The extract significantly and dose dependently reduced writhing count compared to control, which indicated potential peripheral analgesic activity through prostaglandin

inhibition. The hot-plate test is a model to assess centrally acting analgesic activity, which focuses on changes only in the spinal cord level.^[46-48] The extract significantly increased the pain threshold, probably through central mechanisms involving these receptor systems.

Despite the use of conventional antidiarrheal therapies, many plants are used in Bangladesh to treat diarrhea. Castor oil liberates ricinolic acid, which results in irritation and inflammation of intestinal mucosa; subsequently, intestinal motility and secretion are stimulated due to the release of prostaglandins.^[49] The *M. seminifera* root extract significantly and dose dependently reduced the quantity of feces as well as prolonged the onset of diarrhea. The result suggested that the extract might have potential antidiarrheal activity via the antisecretory mechanism.

The brine shrimp lethality bioassay is an easy and sensitive bench-top assay for predicting important pharmacological activities such as enzyme inhibition, ion channel interference, and cytotoxic activity.^[37,50,51] The correlation between the brine shrimp lethality bioassay and *in vitro* growth inhibition of rapidly growing human tumor cell lines was established by National Cancer Institute (NCI, USA).^[52] In the present study, both the extract and vincristine sulphate showed a gradual increase in mortality rate with the increase in concentration. The LC₅₀ value for the crude extract was found to be very low, signifying that the extract may contain potent pharmacologically active compound(s).

The anthelmintic activity of plant materials is usually judged on the basis of loss of movement or paralysis and complete destruction or death of live parasites in *in vitro* studies.^[53-55] The extract showed significant and concentration-dependent decrease in both paralysis and death time compared with standard albendazole. Although the exact mechanism involved with the observed anthelmintic activity of the extract is not known, polyphenolics present in the extract may play a role in it. Polyphenolic compounds such as tannins have been found to be responsible for the strong anthelmintic activity by interfering in energy generation in helminths by uncoupling oxidative phosphorylation.^[56]

CONCLUSION

The results of the present pharmacological investigation support the uses of this plant in folk medicine. The extract showed significant and dose-dependent antioxidant, analgesic, antidiarrheal and anthelmintic activities, and general toxicity.

ACKNOWLEDGMENT

We thank the International Centre for Diarrhoeal Disease and Research, Bangladesh (ICDDR, B) for providing experimental mice, and Beximco Pharmaceuticals Ltd., Bangladesh, for the standard diclofenac sodium and albendazole.

REFERENCES

1. Partha P, Hossain AB. Ethanobotanical investigation into the Mandi ethnic community in Bangladesh. *Bangladesh J Plant Taxon* 2007;14:129-45.
2. Ghani A. Medicinal plants of Bangladesh: Chemical constituents and uses. 2nd ed. Dhaka, Bangladesh: The Asiatic Society of Bangladesh; 2003. p. 315.
3. Hossain MS, Alam MB, Asadujjaman M, Zahan R, Islam MM, Mazumder ME, et al. Antidiarrhoeal, antioxidant and antimicrobial activities of the *Musa sapientum* seed. *Avicenna J Med Biotech* 2011;3:95-105.
4. Pari L, Maheswari JU. Hypoglycaemic effect of *Musa sapientum* L. in alloxan-induced diabetic rats. *J Ethnopharmacol* 1999;68:321-5.
5. Fagbemi JF, Ugoji E, Adenipekun T, Adelowotan O. Evaluation of the antimicrobial properties of unripe banana (*Musa sapientum* L.), lemon grass (*Cymbopogon citratus* S.) and turmeric (*Curcuma longa* L.) on pathogens. *Afr J Biotechnol* 2009;8:1176-82.
6. Agarwal PK, Singh A, Gaurav K, Goel S, Khanna HD, Goel RK. Evaluation of wound healing activity of extracts of plantain banana (*Musa sapientum* var. *paradisiaca*) in rats. *Indian J Exp Biol* 2009;47:32-40.
7. Tewtrakul S, Itharat A, Thammaratwasik P, Oraikul B. Anti-allergic and anti-microbial activities of some Thai crops. *Songklanakarin J Sci Technol* 2008;30:467-73.
8. Zeid AH, Abou S. Chemical and biological study of the leaves of some *Musa* species. *Egyptian J Pharma Sci* 1999;39:379-98.
9. Akihisa T, Kimura Y, Kokke WCM, Takase S, Yasukawa K, Jin-Nai A, et al. 4-Epicycloeucaenone and 4-epicyclomusalenone: Two 3-oxo-28-norcycloartanes from the fruit peel of *Musa sapientum* L. *Chem Pharm Bull* 1997;45:744-6.
10. Mallick C, Chatterjee K, Guhabiswas M, Ghosh D. Antihyperglycemic effects of separate and composite extract of root of *Musa paradisiaca* and leaf of *Coccinia indica* in streptozotocin-induced diabetic male albino rat. *Afr J Tradit Complement Altern Med* 2007;4:362-71.
11. Alisi CS, Nwyanwu CE, Akujobi CO, Ibegbulem CO. Inhibition of dehydrogenase activity in pathogenic bacteria isolates by aqueous extracts of *Musa paradisiaca* (var *Sapientum*). *Afr J Biotechnol* 2008;7:1821-5.
12. Otálvaro F, Görls H, Hölscher D, Schmitt B, Echeverri F, Quiñones W, et al. Dimeric phenylphenalenones from *Musa acuminata* and various Haemodoraceae species. Crystal structure of anigorootin. *Phytochemistry* 2002;60:61-6.
13. Abad T, McNaughton-Smith G, Fletcher WQ, Echeverri F, Diaz-Peñate R, Tabraue C, et al. Isolation of (S)-(+)-naproxene from *Musa acuminata*. Inhibitory effect of naproxene and its 7-methoxy isomer on constitutive COX-1 and inducible COX-2. *Planta Med* 2000;66:471-3.
14. Opitz S, Otálvaro F, Echeverri F, Quiñones W, Schneider B. Isomeric oxabenzochrysenones from *Musa acuminata* and *Wachendorfia thyrsiflora*. *Nat Prod Lett* 2002;16:335-8.

15. Miah MN, Bachar SC, Nahar L, Rahman MS, Rashid MA, Hadiuzzaman S, *et al.* Composition of the volatiles of *Citrus macroptera* var *annamensis* and evaluation of bioactivity. *J Essential Oil Bearing Plants* 2010;13:211-8.
16. Alam MA, Subhan N, Chowdhury SA, Awal MA, Mostofa M, Rashid MA, *et al.* *Anthocephalus cadamba* extract shows hypoglycemic effect and eases oxidative stress in alloxan-induced diabetic rats. *Braz J Pharmacog Rev Bras Farmacog* 2011;21:154-64.
17. Mazid MA, Nahar L, Datta BK, Bashar SA, Sarker SD. Potential antitumour activity of two *Polygonum* species. *Arch Biol Sci* 2011;63:465-8.
18. Ara A, Arifuzzaman M, Ghosh CK, Hashem MA, Ahmad MU, Bachar SC, *et al.* Anti-inflammatory activity of *Adenanthera pavonina* in experimental animals. *Braz J Pharmacog Rev Bras Farmacog* 2010;20:929-32.
19. Mazid MA, Datta BK, Bachar SC, Bashar SMA, Nahar L, Sarker SD. Analgesic and anti-inflammatory activities of *Polygonum stagninum* (Polygonaceae). *Pharm Biol* 2010;48:770-4.
20. Mazid MA, Datta BK, Nahar L, Bashar SM, Bachar SC, Sarker SD. Analgesic, anti-inflammatory and diuretic properties of *Polygonum barbatum* var *barbata*. *Braz J Pharmacog Rev Bras Farmacog* 2009;19:749-54.
21. Rahman AK, Chowdhury AK, Ali HA, Raihan SZ, Ali MS, Nahar L, *et al.* Antibacterial activity of two limonoids from *Swietenia mahagoni* against multiple-drug-resistant (MDR) bacterial strains. *J Nat Med* 2009;63:41-5.
22. Alam MA, Subhan N, Awal MA, Alam MS, Sarder M, Nahar L, *et al.* Antinociceptive and anti-inflammatory properties of *Ruellia tuberosa*. *Pharm Biol* 2009;47:209-14.
23. Alam MA, Akter R, Subhan N, Rahman MM, Majumder MM, Nahar L, *et al.* Antidiarrhoeal property of the hydroethanolic extract of the flowering tops of *Anthocephalus cadamba*. *Braz J Pharmacog Rev Bras Farmacog* 2008;18:155-9.
24. Subhan N, Alam MA, Ahmed F, Shahid IJ, Nahar L, Sarker SD. Bioactivity of *Excoecaria agallocha*. *Braz J Pharmacog Rev Bras Farmacog* 2008;18:521-6.
25. Mondal S, Paul SK, Uddin SJ, Nahar L, Auzi AA, Sarker SD. A comparative study on the *in vitro* antibacterial activity of the pneumatophores of *Heritiera fomes* and *Xylocarpus moluccensis*. *ARS Pharmaceutica* 2008;49:51-6.
26. Uddin SJ, Nahar L, Shilpi JA, Shoeb M, Borkowski T, Gibbons S, *et al.* Gedunin, a limonoid from *Xylocarpus granatum*, inhibits the growth of CaCo-2 colon cancer cell line *in vitro*. *Phytother Res* 2007;21:757-61.
27. Sarker SD, Uddin SJ, Shilpi JA, Rouf R, Ferdous ME, Nahar L. Neuropharmacological properties of *Xylocarpus moluccensis*. *Fitoterapia* 2007;78:107-11.
28. Datta BK, Nahar L, Rahman MM, Gray AI, Auzi AA, Sarker SD. Polygosomic acid, a new cadinane sesquiterpene, from *Polygonum viscosum* inhibits the growth of drug-resistant *Escherichia coli* and *Staphylococcus aureus* (MRSA) *in vitro*. *J Nat Med* 2007;61:391-6.
29. Uddin SJ, Shilpi JA, Middleton M, Byres M, Shoeb M, Nahar L, *et al.* Swarnalin and *cis*-swarnalin, two new tetrahydrofuran derivatives with free radical scavenging activity, from the aerial parts of *Cuscuta reflexa*. *Nat Prod Res* 2007;21:663-8.
30. Uddin SJ, Shilpi JA, Rahman MT, Ferdous M, Rouf R, Sarker SD. Assessment of neuropharmacological activities of *Pandanus foetidus* (Pandanaaceae) in mice. *Pharmazie* 2006;61:362-4.
31. Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* 1983;16:109-10.
32. Ebrahimzadeh MA, Nabavi SM, Nabavi SF, Eslami B. Free radical scavenging ability of methanolic extract of *Hyoscyamus squarrosus* leaves. *Pharmacologyonline* 2009;2:796-802.
33. Marinova D, Ribarova F, Atanassova M. Total phenolics and total flavonoids in Bulgarian fruits and vegetables. *J Univ Chem Technol Metallurgy* 2005;40:255-60.
34. Oyaizu M. Studies on product of browning reaction prepared from glucosamine. *Jpn J Nutri* 1986;44:307-15.
35. Koster R, Anderson M, De Beer EJ. Acetic acid for analgesics screening. *Fed Proc* 1959;18:412-7.
36. Eddy NB, Leimbach D. Synthetic analgesics. II. Diethylnbutenyl- and diethylnbutylamines. *J Pharmacol Exp Ther* 1953;107:385-93.
37. Abdullahi AL, Agho MO, Amos S, Gamaniel KS, Wambebe C. Antidiarrhoeal activity of the aqueous extract of *Terminalia avicennoides* roots. *Phytother Res* 2001;15:431-4.
38. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp: A convenient general bioassay for active plant constituents. *Planta Med* 1982;45:31-4.
39. Tandon V, Pal P, Roy B, Rao HS, Reddy KS. *In vitro* anthelmintic activity of root-tuber extract of *Flemingia vestita*, an indigenous plant in Shillong, India. *Parasitol Res* 1997;83:492-8.
40. Hossain E, Chandra G, Nandy AP, Mandal SC, Gupta JK. Anthelmintic effect of a methanol extract of *Bombax malabaricum* leaves on *Paramphistomum explanatum*. *Parasitol Res* 2012;110:1097-102.
41. Halliwell H. Free radicals, antioxidants and human disease: Curiosity, cause or consequence? *Lancet* 1994;344:721-4.
42. Wang K, Yang C, Zhang Y. Phenolic antioxidants from Chinese toon (fresh young leaves and shoots of *Toona sinensis*). *Food Chem* 2007;101:365-71.
43. Meir S, Kanner J, Akiri B, Hadas SP. Determination and involvement of aqueous reducing compounds in oxidative defense systems of various senescing leaves. *J Agric Food Chem* 1995;43:1813-7.
44. Gené RM, Segura L, Adzet T, Marin E, Iglesias J. *Heterotheca inuloides*: Anti-inflammatory and analgesic effect. *J Ethnopharmacol* 1998;60:157-62.
45. Roberts LJ, Morrow JD. Analgesic-antipyretic and anti-inflammatory agents and drugs employed in the treatment of gout. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. 10th ed. New York: McGraw-Hill; 2001. p. 687-732.
46. Vongtau HO, Abbah J, Mosugu O, Chindo BA, Ngazal IE, Salawu AO, *et al.* Antinociceptive profile of the methanolic extract of *Neorautanenia mitis* root in rats and mice. *J Ethnopharmacol* 2004;92:317-24.
47. Headley PM, O'Shaughnessy CT. Evidence for opiate and dopamine interaction in striatum. *Br J Pharmacol* 1985;86:700.
48. Wigdor S, Wilcox GL. Central and systemic morphine-induced antinociception in mice: Contribution of descending serotonergic and noradrenergic pathways. *J Pharmacol Exp Ther* 1987;242: 90-5.
49. Pierce NF, Carpenter CC Jr, Elliott HL, Greenough WB 3rd. Effects of prostaglandins, theophylline, and cholera exotoxin upon transmucosal water and electrolyte movement in the canine jejunum. *Gastroenterology* 1971;60:22-32.
50. Anderson JE, Goetz CM, McLaughlin JL, Suffness M. A blind comparison of simple bench-top bioassays and human tumor cell cytotoxicities as antitumor prescreens. *Phytochem Anal* 1991;2:107-11.
51. Borowitz JL, McLaughlin JL. Evidence for calcium channels in brine shrimp: Diltiazem protects shrimp against cadmium. *Bull Environ Contam Toxicol* 1992;48:435-40.

52. Silva TM, Nascimento RJ, Batista MM, Agra MF, Camara CA. Brine shrimp bioassay of some species of *Solanum* from Northeastern Brazil. *Brazil J Pharmacol* 2007;17:35-8.
53. Goto C, Kasuya S, Koga K, Ohtomo H, Kagei N. Lethal efficacy of extract from *Zingiber officinale* (traditional Chinese medicine) or 6-shogaol and 6-gingerol in *Anisakis* larvae *in vitro*. *Parasitol Res* 1990;76:653-6.
54. Robinson RD, Williams LA, Lindo JF, Terry SI, Mansingh A. Inactivation of *strongyloides stercoralis* filariform larvae *in vitro* by six Jamaican plant extracts and three commercial anthelmintics. *West Indian Med J* 1990;39:213-7.
55. Togo J, Santamarina MT, Peris D, Ubeira FM, Leiro SL, Sanmartin ML. *In vitro* effect of anthelmintics on *Anisakis simplex* survival. *Jpn J Parasitol* 1992;41:473-80.
56. Martin RJ. Modes of action of anthelmintic drugs. *Vet J* 1997;154:11-34.

Cite this article as: Saha S, Shilpi JA, Mondal H, Gofur R, Billah M, Nahar L, *et al.* Bioactivity studies on *Musa seminifera* Lour. *Phcog Mag* 2013;9:315-22.

Source of Support: Nil. **Conflict of Interest:** None declared.