



Published in final edited form as:

Diabetologia. 2011 February ; 54(2): 440–450. doi:10.1007/s00125-010-1956-9.

Tissue-specific dysregulation of hexose-6-phosphate dehydrogenase and glucose-6-phosphate transporter production in *db/db* mice as a model of type 2 diabetes

Y. Wang,

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

Y. Nakagawa,

Department of Pediatrics, Hamamatsu University School of Medicine, Hamamatsu, Japan

L. Liu,

Department of Endocrinology & Metabolism, Shanghai Jiaotong University Affiliated Sixth People's Hospital, Shanghai Diabetes Institute, Shanghai, People's Republic of China

W. Wang,

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

X. Ren,

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

A. Anghel,

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

K. Lutfy,

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

T. C. Friedman, and

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA

Y. Liu

Division of Endocrinology, Metabolism & Molecular Medicine, Charles Drew University of Medicine & Sciences, UCLA School of Medicine, 1731 E. 120th St, Los Angeles, CA 90059, USA
dryanjunliu@hotmail.com

Abstract

Aims/hypothesis—Tissue-specific amplification of glucocorticoid action through 11 - hydroxysteroid dehydrogenase type 1 (11 -HSD1) affects the development of the metabolic

© Springer-Verlag 2010

Correspondence to: Y. Liu.

Y. Nakagawa and Y. Wang contributed equally to this study.

Electronic supplementary material The online version of this article (doi:10.1007/s00125-010-1956-9) contains supplementary material, which is available to authorised users.

Duality of interest The authors declare that there is no duality of interest associated with this manuscript.

syndrome. Hexose-6-phosphate dehydrogenase (H6PDH) mediates intracellular NADPH availability for 11 β -HSD1 and depends on the glucose-6-phosphate transporter (G6PT). Little is known about the tissue-specific alterations of H6PDH and G6PT and their contributions to local glucocorticoid action in *db/db* mice.

Methods—We characterised the role of H6PDH and G6PT in pre-receptor metabolism of glucocorticoids by examining the production of the hepatic 11 β -HSD1-H6PDH-G6PT system in *db/db* mice.

Results—We observed that increased production of hepatic H6PDH in *db/db* mice was paralleled by upregulation of hepatic G6PT production and responded to elevated circulating levels of corticosterone. Treatment of *db/db* mice with the glucocorticoid antagonist RU486 markedly reduced production of both H6PDH and 11 β -HSD1 and improved hyperglycaemia and insulin resistance. The reduction of H6PDH and 11 β -HSD1 production by RU486 was accompanied by RU486-induced suppression of hepatic *G6pt* (also known as *Slc37a4*) mRNA. Incubation of mouse primary hepatocytes with corticosterone enhanced G6PT and H6PDH production with corresponding activation of 11 β -HSD1 and PEPCK: effects that were blocked by RU486. Knockdown of *H6pd* by small interfering RNA showed effects comparable with those of RU486 for attenuating the corticosterone-induced H6PDH production and 11 β -HSD1 reductase activity in these intact cells. Addition of the G6PT inhibitor chlorogenic acid to primary hepatocytes suppressed H6PDH production.

Conclusions/interpretation—These findings suggest that increased hepatic H6PDH and G6PT production contribute to 11 β -HSD1 upregulation of local glucocorticoid action that may be related to the development of type 2 diabetes.

Keywords

11 β -HSD1; G6PT; G6PT inhibitor; H6PDH; H6PDH siRNA; Insulin resistance; NADPH; Type 2 diabetes

Introduction

Excess glucocorticoid (GC) production strongly affects the development of type 2 diabetes and obesity via activation of intracellular GC receptor (GR) [1–5]. In rodents, increased GC production promotes hepatic gluconeogenesis and adipocyte differentiation and induces insulin resistance and obesity in genetically obese *db/db* and *ob/ob* mice [5–8]. However, tissue GC action can also be regulated at the pre-receptor level by the 11 β -hydroxysteroid dehydrogenases. Metabolic tissues such as liver and adipose tissue abundantly produces 11 β -hydroxysteroid dehydrogenase (11 β -HSD1), a resident enzyme of the endoplasmic reticulum (ER) lumen that acts *in vivo* as an NADPH-dependent reductase to generate active cortisol (corticosterone in rodents) from inactive 11-keto cortisone (11-dehydrocorticosterone) [9–11]. 11 β -HSD1 plays an important role in the regulation of tissue GC action [12–15]. Indeed, enhanced 11 β -HSD1 could result in the production of excess tissue GCs and induction of GC-receptor-mediated local GC action related to glucose homeostasis, insulin action and adiposity, all of which are associated with the development of type 2 diabetes and visceral obesity [16–18]. Pharmacological evidence further validates the role of 11 β -HSD1 in the control of insulin resistance and obesity as 11 β -HSD1 inhibitors reduced hepatic glucose output and improved insulin sensitivity in obese animal models, as well as in patients with type 2 diabetes [19–23]. These studies implicate the importance of 11 β -HSD1 in the pathogenesis of metabolic syndrome and obesity.

The role of 11 β -HSD1 in the pre-receptor activation of GCs is entirely dependent on the production of its cofactor NADPH in the ER lumen [24, 25]. NADPH is regenerated by

hexose-6-phosphate dehydrogenase (H6PDH) [26, 27], a microsomal enzyme located in the lumen of the ER and principally produced in hepatocytes and adipocytes, sites of 11 β -HSD1 and GR [28, 29]. In these target tissues, H6PDH uses glucose 6-phosphate (G6P) and NADP to produce NADPH. The supply of G6P to H6PDH is ensured by a functional membrane ER protein, the G6P transporter (G6PT), which specifically transports cytosolic G6P into the ER and therefore regulates H6PDH activity via modulation of G6P level within the ER [30, 31]. The generation of NADPH by H6PDH can be used by NADPH-dependent 11 β -HSD1 reductase activity [32, 33]. H6PDH is thus a potential candidate supplying NADPH for 11 β -HSD1-induced amplification of tissue GC production [34–36].

In the present study, we characterised the metabolic phenotype of H6PDH and G6PT in relation to pre-receptor metabolism of GCs by examining the effects of the GC antagonist RU486 on the production of the 11 β -HSD1–H6PDH–G6PT system in the liver and adipose tissues of *db/db* mice. We also assessed the direct interaction of G6PT and H6PDH in the control of 11 β -HSD1 driving tissue GC action in primary cultures of hepatocytes from *db/db* mice by using *H6pdh* small interfering (si)RNA and the G6PT inhibitor chlorogenic acid (CA). Finally, we examined the hormonal regulation of H6PDH and G6PT in these intact cells.

Methods

Animals

Male C57BL/KsJ-obese (*db/db*) mice and their lean littermates (*db/+*) were purchased at 10 weeks of age from Taconic Farms (Hudson, NY, USA) and housed in a room illuminated daily from 07:00 to 19:00 hours (12 h light/dark cycle) with free access to water and standard laboratory chow. RU486 (25 mg/kg) or vehicle was given by intraperitoneal injection twice each day (at 07:00 and 19:00 hours) for 3 days or 3 weeks as previously described by Friedman et al. and Liu et al. [5, 15]. Body weight and food intake were recorded daily. All animal experiments were approved by the Charles Drew University Institutional Animal Care and Use Committee.

Insulin tolerance test

For the insulin tolerance test (ITT), animals were fasted for 12 h, and blood samples were drawn at different times following insulin injection (2 U/kg i. p.; Novolin R; Eli Lilly, Indianapolis, IN, USA).

Biochemical assays

Blood glucose levels were determined by the glucose oxidase method [36]. Plasma corticosterone levels were determined by RIA using mouse corticosterone as a standard (ICN Biomedicals, Costa Mesa, CA, USA). Plasma insulin levels were measured by RIA using rat insulin as a standard (Crystal Chemicals, Chicago, IL, USA).

Cell culture and treatment

Primary hepatocytes were isolated from male *db/db* mice [21] and seeded onto collagen-coated dishes in DMEM/F-12 medium with 10% FBS at 37°C for 4 h. Cells were then washed with PBS, and the medium was changed to DMEM/F-12 medium without FCS. After 12 h, cells were treated with corticosterone (1×10^{-7} – 1×10^{-5} mol/l) or insulin (1×10^{-6} mol/l) in the presence or absence of RU486 (1×10^{-6} mol/l) for 48 or 72 h. The G6PT inhibitor CA (0–200 μ mol/l) was added to primary cultures of hepatocytes for 48 h. This pharmacological dose of CA is similar to doses previously used in mouse 3T3-L1 pre-adipocytes and glioma cells [37, 38].

siRNA experiments in primary cultures of hepatocytes

Cells were transfected with the *H6pd* siRNA (SABiosciences siRNA ID KM26265; SABiosciences, Frederick, MD, USA), or negative control siRNA (SABiosciences) using lipofectamine 2000 transfection agent according to the manufacturer's protocol (Invitrogen, Carlsbad, CA, USA). After 4 h, cells were treated with vehicle or corticosterone (1×10^{-6} mol/l) for an additional 48 h.

Microsomal enzymatic activity assays

The liver microsomal pellet was obtained and 11 β -HSD1 activity was evaluated by addition of 0.4–1 mmol/l NADPH and 900 nmol/l unlabelled 11-dehydrocorticosterone (11-DHC) with 100 nmol/l 11- 3 H]DHC as tracer to microsomes in Krebs–Ringer buffer solution at 37°C for 10–30 min [36, 39]. The liver microsomes (1 mg/ml protein) were permeabilised with 1% Triton-100 to allow the free access of the cofactor to the intraluminal enzyme. Steroids were separated by thin-layer chromatography (TLC) and analysed by scintillation counting. The percentage of interconversion of 3 H]corticosterone and 11- 3 H]DHC was calculated from the radioactivity in each fraction. Reductase activity of 11 β -HSD1 was also evaluated by immunoassay of the corticosterone produced from 11-DHC using a sensitive corticosterone ELISA kit.

H6PDH enzyme activity was determined by spectrophotometric measurement of NADPH production in the presence of G6P and NADP using absorbance at 340 nm with a spectrophotometer (Ultrospec 2100, Amersham Biosciences) [36, 40]. Protein, 50 μ g, from liver microsomes was incubated with 0.5–5 mmol/l G6P, 1–5 mmol/l NADP and 100 mmol/l glycine buffer solution at 22°C for 0–5 min. Specific activities were calculated and expressed as micromoles NADPH production per minute per milligram protein. The protein concentration was measured by the Bradford assay (Bio-Rad Protein Assay Kit, Bio-Rad, Hercules, CA, USA).

11 β -HSD1 and H6PDH activity in primary hepatocytes

Cells were incubated with 2 nmol/l 11- 3 H]DHC with 18 nmol/l unlabelled 11-DHC for 10–30 min [41]. Steroids were extracted from 1.0 ml of culture medium with ethyl acetate and separated by TLC. Enzyme activity levels were determined by counting the radioactivity. For the H6PDH activity in vitro, protein extracts, 20 μ g, from primary hepatocytes were incubated with 2 mmol/l G6P as substrate in 100 μ l total volume of glycine buffer (pH 10.0) with 0.5 mmol/l NADP as a cofactor. The changes in absorbance at 340 nm were measured during 25 min at 5 min intervals [36].

RNA extraction and real-time quantitative PCR analysis

Total RNA was extracted using a single-step extraction method (RNAzol B, Invitrogen). cDNA synthesis from 2.0 μ g RNA was performed using high capacity RNA-to-cDNA Kit (Applied Biosystems, Carlsbad, CA, USA). Real-time primers were designed with Primer express software 2.0 (Applied Biosystems) and are listed in the Electronic supplementary material (ESM) Table 1. Amplification of each target cDNA was then performed with SYBR Green I Master Kits in the ABI Prism 7700 Sequence Detection System (Applied Biosystems) according to the protocols recommended by the manufacturer. All reactions were carried out using the following cycling parameters: 55°C for 2 min and 95°C for 10 min, following by 40 cycles of 95°C for 15 s and 60°C for 1 min. Threshold cycle (C_t) readings for each of the unknown samples were then used to calculate the amount of target genes and were normalised to the signal of 18S rRNA. Data analysis is based on the C_t method.

G6P uptake measurement

Microsomal G6P uptake measurements were performed according to protocols previously published by Hiraiwa et al. [42]. Briefly, 50–100 µg of microsomes were incubated in 50 mmol/l sodium cacodylate buffer containing 250 mmol/l sucrose (pH 6.5) in the presence of 0.2–1 mmol/l G6P plus [U-¹⁴C]G6P (American Radiolabeled Chemicals, St Louis, MO, USA) at 22°C for 3 min. The reaction was stopped by filtering through a nitrocellulose filter and samples were quickly washed with an ice-cold buffer containing 50 mmol/l TRIS-HCl, pH 7.4 and 250 mmol/l sucrose. Microsomes were subsequently permeabilised with 0.2% deoxycholate to abolish intra-vesicular G6P uptake. The radioactivity associated with microsomes retained by filters was measured by liquid-scintillation counting.

Western blot analysis

Protein, 25 µg samples from liver microsomes or 50 µg total cellular proteins from primary hepatocytes, was analysed by SDS-PAGE as described by Liu et al. [36]. Membranes were incubated with a polyclonal anti-11 β -HSD1 antibody (1:1,500; Affinity Bioreagents, Rockford, IL, USA) and mouse anti-human H6PDH antibody (1:2,000; Novus Biological, Littleton, CO, USA). The 11 β -HSD1 and H6PDH protein signal was quantified with the use of the Eagle Eye II Quantitation System (Stratagene, La Jolla, CA, USA).

Statistical analyses

All data are expressed as the mean \pm SEM. The normality of the distribution of data was established using the Wilks–Shapiro test, and outcome measures between groups were compared by Student's *t* test. To compare multiple groups, one-way ANOVA used. If ANOVA reveals significant differences, then individual group comparisons were performed by using the Newman–Keul's post hoc test. The differences among groups were considered significant at $p < 0.05$.

Results

Characterisation of H6PDH and G6pt levels in db/db mice db/db

mice had higher body weight and plasma levels of insulin, corticosterone and blood glucose than matched littermates (Table 1). Plasma levels of cholesterol and triacylglycerols in *db/db* mice were significantly higher than those in lean mice (Table 1). 11 β -HSD1 reductase activity in liver microsomes of *db/db* mice was significantly increased to 47% over that of *db/+* controls ($p < 0.01$; Fig. 1a). Western blot analysis revealed the 11 β -HSD1 protein production in the liver of *db/db* mice was increased 1.7-fold over that of *db/+* controls ($p < 0.01$; Fig. 1b). H6PDH activity and protein production in the liver of *db/db* mice were increased 1.57- and 1.9-fold, respectively, compared with *db/+* control mice ($p < 0.001$; Fig. 1c, d). Real-time RT-PCR analysis revealed that hepatic 11 β -*Hsd1* (also known as *Hsd11b1*) mRNA levels were increased 2.8-fold in *db/db* mice and level was positively correlated with *H6pd* mRNA levels ($p < 0.01$), which increased to 1.8-fold in liver over that of *db/+* animals ($p < 0.01$; Fig. 2a). Similarly, the hepatic *G6pt* (also known as *Slc37a4*) mRNA levels in *db/db* mice were significantly increased by 1.9-fold over that of *db/+* controls ($p < 0.01$; Fig. 2a). Moreover, the [U-¹⁴C]G6P uptake in hepatic microsomes of *db/db* mice was higher than that of *db/+* controls ($p < 0.01$; Fig. 2b). The levels of hepatic *Pepck* and *G6Pase* mRNA in *db/db* mice were higher than the respective in *db/+* mice ($p < 0.01$; Fig. 2c). AUC analysis showed that the glucose levels in *db/db* mice were significantly higher than those in vehicle-treated *db/+* animals (Fig. 2d). In addition, subcutaneous and epididymal adipose 11 β -*Hsd1* mRNA levels were higher than those of lean controls (Fig. 3a, b). Moreover, *H6pd* mRNA levels were also markedly increased in subcutaneous fat and in epididymal fat in *db/db* mice compared with lean levels ($p < 0.05$), respectively. However, there was no difference in *G6pt*

mRNA levels in subcutaneous and epididymal fat between *db/db* mice and lean controls (Fig. 3a, b).

The effects of RU486 treatment on the phenotypes of type 2 diabetes in *db/db* mice

As shown in Table 1, RU486 compared with vehicle treatment for 3 weeks lowered blood glucose levels with no significant changes in body weight and levels of plasma insulin and lipids in *db/db* mice. However, plasma corticosterone levels were increased in *db/db* mice after RU486 treatment ($p < 0.001$). In contrast, hepatic 11 β -HSD1 reductase activity was reduced 1.6-fold in RU486-treated *db/db* mice (Fig. 1a). Western blot analysis revealed that 11 β -HSD1 protein production was decreased 2.5-fold in RU486-treated *db/db* mice ($p < 0.01$ vs *db/db* controls; Fig. 1b). Similarly, treatment of *db/db* mice with RU486 for 3 weeks significantly reduced hepatic H6PDH activity to 45% that of vehicle-treated *db/db* mice ($p < 0.01$; Fig. 1d); this reduction in enzyme activity occurred in parallel with decreased hepatic H6PDH protein production in RU486-treated *db/db* mice (Fig. 1c). Real-time RT-PCR analysis showed that in *db/db* mice, RU486 reduced hepatic 11 β -*Hsd1* and *H6pd* mRNA expression, respectively, to the levels of vehicle-treated *db/+* mice (Fig. 2a). In parallel with the decrease in *H6pd* mRNA levels, hepatic *G6pt* mRNA expression was decreased by twofold in *db/db* mice after RU486 treatment (Fig. 2a). Consistent with decreased *G6pt* expression, RU486 reduced [U - ^{14}C]G6P uptake in the liver microsomes, although it did not restore G6P uptake levels to those of *db/+* controls (Fig. 2b). Moreover, RU486 also significantly reduced hepatic *Pepck* mRNA levels, but did not exert significant effects on the hepatic *G6Pase* (also known as *G6pc*) mRNA expression in *db/db* mice (Fig. 2c). The AUC glucose levels were reduced in *db/db* mice after RU486 treatment (Fig. 2d). RU486 treatment for 3 weeks also reduced the *G6pt* mRNA level with simultaneous reduction in H6PDH and 11 β -HSD1 expression in subcutaneous fat of *db/db* mice as compared with controls (Fig. 3a), but did not alter epididymal fat G6PT, H6PDH and 11 β -HSD1 expression (Fig. 3b). However, RU486 did not exert significant effects on 11 β -*Hsd1*, *H6pd* and *G6pt* mRNA expression in adipose tissues from *db/+* mice (Fig. 3a, b). In addition, a shorter treatment of *db/db* mice with RU486 for 3 days also significantly reduced the expression of 11 β -*Hsd1*, *H6pd* and *G6pt* in the liver and improved glucose levels, but did not change the slope of insulin tolerance curve (data not shown), indicating that blocking the action of GC with RU486 leads to changes in gene expression that are independent of improvement in insulin sensitivity.

H6PDH and G6PT abundance in primary cultures of hepatocytes

To confirm our *in vivo* observations, we investigated the effects of exogenous corticosterone and insulin on G6PT and H6PDH in primary hepatocytes from *db/db* mice. Western blot analysis showed that treatment of hepatocytes with increasing doses of corticosterone led to a concentration-dependent induction of H6PDH protein production (Fig. 4a). A concentration of corticosterone (1×10^{-6} mol/l), similar to that occurring in *db/db* mice *in vivo*, significantly increased NADPH production in the presence of G6P as a substrate (Fig. 4b). The increase in H6PDH activity was consistent with real-time RT-PCR analyses, which showed that corticosterone increased *H6pd* mRNA levels 1.6-fold in primary hepatocytes compared with vehicle ($p < 0.001$; Fig. 4c). In contrast, treatment of hepatocytes with both corticosterone and RU486 (10^{-6} mol/l) for 48 h failed to increase H6PDH activity (Fig. 4b) or *H6pdh* mRNA levels (Fig. 4c). Moreover, co-treatment with corticosterone and RU486 also blocked the corticosterone-induced changes in 11 β -*Hsd1*, *Pepck* and *G6Pase* mRNA expression (Fig. 4d) in these intact cells. However, no changes in *H6pdh* mRNA expression were observed after 48 h in cells exposed to a relatively high concentration of insulin (1×10^{-6} mol/l; Fig. 4c). Extending our findings on H6PDH activity, corticosterone-induced expression of *G6pt* mRNA in primary hepatocytes was reduced significantly by RU486 (Fig. 5a, b).

The effects of H6PDH siRNA and G6PT inhibitor (CA) on H6PDH and 11 β -HSD1

As shown in Fig.6, H6PDH protein and *H6pd* mRNA levels were markedly reduced in intact hepatocytes transfected with *H6pd* siRNA in comparison to cells transfected with siRNA control, respectively ($p < 0.001$, Fig. 6a, b). Similarly, in comparison with control levels, 11 -HSD1 reductase activity in primary hepatocytes after transfection with *H6pdh* siRNA was decreased by 44% ($p < 0.01$; Fig. 6c). Furthermore, *H6pd* siRNA attenuated the corticosterone-mediated increase in H6PDH protein and 11 -HSD1 reductase activity in these intact cells in comparison with control levels (Fig. 6), indicating that suppression of H6PDH by siRNA exerted effects comparable with those of RU486 on reducing the corticosterone-induced H6PDH levels and 11 -HSD1 activity.

The effects of the G6PT inhibitor CA on H6PDH and 11 -HSD1 are shown in Fig. 7. CA significantly reduced H6PDH protein production in primary hepatocytes compared with vehicle treatment for 48 h (Fig. 7a). Consequently, the mRNA levels of *H6pd* were decreased by 1.6-fold in these intact cells after CA treatment (Fig. 7b). In agreement with reduction of H6PDH, CA treatment also reduced 11 -*Hsd1* mRNA expression as compared with that of controls ($p < 0.01$; Fig. 7b).

Discussion

We found that the phenotype of type 2 diabetes in *db/db* mice was associated with the induction of hepatic H6PDH activity and gene expression. H6PDH activity and protein expression were elevated in liver microsomes from *db/db* mice, and this was accompanied by the induction of hepatic 11 -HSD1 reductase activity. This elevated production of hepatic H6PDH corresponded with increased hepatic *Pepck* and *G6pase* mRNA levels, high blood glucose and insulin resistance, suggesting that elevated hepatic H6PDH levels may contribute to the development of type 2 diabetes in our animal model. Induction of hepatic *H6pd* expression could increase the generation of the crucial co-factor NADPH and thus enhance 11 -HSD1 upregulation of local GC production, which would lead to gluconeogenesis linked to the development of hyperglycaemia in diabetic animals [15, 21]. Moreover, *H6pd* and 11 -*Hsd1* mRNA levels were also increased in the adipose fat of *db/db* mice. Our present results are consistent with recent reports that adipose *H6pd* and 11 -*Hsd1* mRNA were increased in patients with type 2 diabetes [43]. To our knowledge, hepatic H6PDH has not been measured in patients with type 2 diabetes. Our findings are also supported by earlier reports that increased hepatic 11 -HSD1 production may contribute to the pathogenesis of type 2 diabetes in *db/db* mice [21, 44].

In *db/db* mice, excess corticosterone production has been shown to be crucial for the development of hyperglycaemia, insulin resistance and obesity, and adrenalectomy reverses these changes [6–8]. In the present study, we observed that the induction of hepatic H6PDH activity and *H6pd* gene expression was associated with the elevated circulating levels of corticosterone. This was validated using isolated hepatocytes from *db/db* mice in which corticosterone increased H6PDH activity at the transcriptional level. Increased H6PDH activity could then provide the cofactor NADPH to 11 -HSD1, suggesting that GCs could positively promote their own pre-receptor metabolism through activation of NADPH availability to 11 -HSD1 reductase activity. Our results are consistent with those of earlier reports that cortisone increased H6PDH activity leading to 11 -HSD1 amplifying cortisol production in rat liver microsomal fractions [34]. These data support our suggestion that the induction of hepatic H6PDH production may, in part, result from excess circulating corticosterone action.

It is well known that pharmacological blockade of GR by RU486 antagonises hypercortisolaemia-related hyper-glycaemia and insulin resistance in *db/db* mice as well as

in patients with Cushing's syndrome [5, 8]. In the present study, we found that treatment of *db/db* mice with RU486 markedly reduced production of hepatic H6PDH and 11 β -HSD1 and improved hyperglycaemia and insulin resistance. Moreover, we observed that RU486 treatment reversed corticosterone-induced production of H6PDH activity and NADPH and prevented activation of 11 β -HSD1 and PEPCK production, consistent with a recent study reporting that the dexamethasone-induced upregulation of *H6pd* mRNA can be blocked by RU486 in mouse 3T3-L1 adipocytes [45]. Moreover, we also observed that suppression of *H6pd* by siRNA decreased 11 β -HSD1 reductase activity and showed effects comparable with those of RU486 on blocking the corticosterone-induced H6PDH and 11 β -HSD1 in intact hepatocytes. Reduction of hepatic H6PDH expression by RU486 could decrease the ability of H6PDH utilising NADP to regenerate NADPH, thereby limiting the intraluminal NADPH exposure to 11 β -HSD1 linked to the reduction of 11 β -HSD1 and the improvement of glycaemic variables. This is in agreement with a recent study reporting that the role of RU486 in glycaemic homeostasis is associated with reduction of hepatic 11 β -HSD1 levels in type 2 diabetic mice [46].

In addition, the current study also observed a positive relationship between hepatic H6PDH activity and G6PT abundance in *db/db* mice and *db/+* controls. Indeed, earlier studies showed that H6PDH requires G6P transport to maintain its metabolic substrate G6P availability within the ER lumen [30, 47, 48]. We observed that the induction of H6PDH production is associated with an elevation of *G6pt* mRNA levels with increased hepatic intraluminal G6P uptake. In contrast, decreased hepatic G6PT abundance after RU486 treatment reduced the intraluminal hepatic G6P uptake with corresponding suppression of H6PDH activity. In agreement with the reduction of H6PDH levels, the current study also showed that the G6PT inhibitor CA, at doses used in vitro [37, 38], markedly reduced H6PDH levels and led to the suppression of 11 β -HSD1 in intact mouse hepatocytes, indicating that G6PT is required to maintain H6PDH levels. Our findings are consistent with recent studies reporting that G6P availability in the ER lumen was reduced, leading to decreased H6PDH and 11 β -HSD1 activity in mice with global deletion of G6PT [48]. This is also indirectly supported by a recent study demonstrating that the G6PT inhibitor S3483 (derived from CA) decreased G6P-induced H6PDH levels and resulted in the inhibition of reductase activity of 11 β -HSD1 in rat liver microsomes [33, 49]. These data support our hypothesis that suppression of G6PT production in hepatocytes may be an additional mechanism of the RU486-mediated reduction of H6PDH production. Moreover, we also observed that RU486 prevented corticosterone-mediated induction of G6PT production in mouse hepatocytes, suggesting that endogenous GCs could exert positive effects on G6PT production in liver that is likely to occur through a GR-mediated mechanism. These findings are supported by a recent study which showed that there are three GC response elements present in the promoter region of the *G6pt* gene [50]. This is also in agreement with previous studies showing that the GR signalling pathway was required for the regulation of 11 β -HSD1 and PEPCK production by GCs [15, 51].

In summary, we showed that increased hepatic H6PDH activity and G6PT expression may contribute to 11 β -HSD1 upregulation of local GC action linked to the development of type 2 diabetes. We also found that the induction of hepatic G6PT and H6PDH production may be mediated, at least in part, through the action of elevated circulating GCs. The benefits of RU486 in *db/db* mice may be associated with the endogenous inactivation of 11 β -HSD1 amplifying tissue GC action through reducing production of H6PDH and G6PT. These findings raise the possibility that tissue-specific modulation of H6PDH and G6PT production may be a new strategy to be coupled with the investigation of 11 β -HSD1 as a therapeutic target in the metabolic syndrome.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Y. Liu is supported by NIH grants KOI DK073272 and SC1DK087655. T. C. Friedman is supported by NIH grants R01 DA14659, K21 DA00276 and an endowment grant S21MD000103-NIH/NCMHD.

Abbreviations

11 -HSD1	11 -Hydroxysteroid dehydrogenase type 1
CA	Chlorogenic acid
11-DHC	11-Dehydrocorticosterone
ER	Endoplasmic reticulum
GC	Glucocorticoid
G6P	Glucose 6-phosphate
G6Pase	Glucose-6-phosphatase
G6PT	Glucose-6-phosphate transporter
GR	Glucocorticoid receptor
H6PDH	Hexose-6-phosphate dehydrogenase
ITT	Insulin tolerance test
TLC	Thin layer chromatography

References

1. Rizza RA, Mandarino LJ, Gerich JE. Cortisol-induced insulin resistance in man: impaired suppression of glucose production and stimulation of glucose utilization due to a postreceptor defect of insulin action. *J Clin Endocrinol Metab.* 1982; 54:131–138. [PubMed: 7033265]
2. Bjorntorp P, Rosmond R. Obesity and Cortisol. *Nutrition.* 2000; 16:924–936. [PubMed: 11054598]
3. Kahn BB, Flier JS. Obesity and insulin resistance. *J Clin Invest.* 2000; 106:473–481. [PubMed: 10953022]
4. Havel PJ, Busch BL, Curry DL, et al. Predominately glucocorticoid agonist actions of RU-486 in young specific-pathogen-free Zucker rats. *Am J Physiol.* 1996; 271:R710–R717. [PubMed: 8853395]
5. Friedman JE, Sun Y, Yun JS. Phosphoenolpyruvate carboxykinase (GTP) gene transcription and hyperglycemia are regulated by glucocorticoids in genetically obese *db/db* mice transgenic mice. *J Biol Chem.* 1997; 272:31475–31481. [PubMed: 9395482]
6. Bamberger CM, Schulte HM, Chrousos GP. Molecular determinants of glucocorticoid receptor function and tissue sensitivity to glucocorticoids. *Endocr Rev.* 1996; 17:245–261. [PubMed: 8771358]
7. Shimomura Y, Bray GA, Lee M. Adrenalectomy and steroid treatment in obese (*ob/ob*) and diabetic (*db/db*) mice. *Horm Metab Res.* 1987; 19:295–299. [PubMed: 2957297]
8. Nieman LK, Chrousos GP, Kellner C, et al. Successful treatment of Cushing's syndrome with the glucocorticoid antagonist RU486. *J Clin Endocrinol Metab.* 1985; 61:536–540. [PubMed: 2991327]
9. Seckl JR, Walker BR. Minireview: 11 beta-hydroxysteroid dehydrogenase type: a tissue-specific amplifier of glucocorticoid action. *Endocrinology.* 2001; 142:1371–1376. [PubMed: 11250914]

10. Jamieson PM, Chapman KE, Edwards CR, et al. 11 beta-hydroxysteroid dehydrogenase is an exclusive 11 beta-reductase in primary cultures of rat hepatocytes: effect of physicochemical and hormonal manipulations. *Endocrinology*. 1995; 136:4754–4761. [PubMed: 7588203]
11. Bujalska IJ, Kumar S, Stewart PM. Does central obesity reflect Cushing's disease of the omentum? *Lancet*. 1997; 349:1210–1213. [PubMed: 9130942]
12. Stewart PM, Boulton A, Kumar S, Clark PM, Shackleton CH. Cortisol metabolism in human obesity: impaired cortisone Cortisol conversion in subjects with central adiposity. *J Clin Endocrinol Metab*. 1999; 84:1022–1027. [PubMed: 10084590]
13. Napolitano A, Voice MW, Chapman KE, et al. 11 -hydroxysteroid dehydrogenase 1 in adipocytes: expression is differentiation-dependent and hormonally regulated. *J Steroid Biochem Mol Biol*. 1998; 164:251–260. [PubMed: 9618026]
14. Rajan V, Edwards CR, Seckl JR. 11 -Hydroxysteroid dehydrogenase in cultured hippocampal cells reactivates inert 11-dehydrocorticosterone, potentiating neurotoxicity. *J Neurosci*. 1996; 16:65–70. [PubMed: 8613810]
15. Liu Y, Nakagawa Y, Friedman TC, et al. Increased glucocorticoid receptor and 11 -hydroxysteroid dehydrogenase type 1 expression in hepatocytes may contribute to the phenotype of type 2 diabetes in *db/db* mice. *Diabetes*. 2005; 54:32–40. [PubMed: 15616008]
16. Tiosano D, Eisentein I, Hochberg Z, et al. 11beta-Hydroxysteroid dehydrogenase activity in hypothalamic obesity. *J Clin Endocrinol Metab*. 2003; 88:379–384. [PubMed: 12519880]
17. Masuzaki H, Paterson J, Flier JS, et al. A transgenic model of visceral obesity and the metabolic syndrome. *Science*. 2001; 294:2166–2170. [PubMed: 11739957]
18. Paterson JM, Morton NM, Fievet C, et al. Metabolic syndrome without obesity: hepatic overexpression of 11 -hydroxysteroid dehydrogenase type 1 in transgenic mice. *Proc Natl Acad Sci USA*. 2010; 101:7088–7093. [PubMed: 15118095]
19. Kotelevtsev Y, Holmes MC, Burchell A, et al. 11beta-Hydroxysteroid dehydrogenase type 1 knockout mice show attenuated glucocorticoid-inducible responses and resist hyperglycemia on obesity or stress. *Proc Natl Acad Sci USA*. 1997; 94:14924–14929. [PubMed: 9405715]
20. Alberts P, Engblom L, Edling N, et al. Selective inhibition of 11beta-hydroxysteroid dehydrogenase type 1 decrease blood glucose concentrations in hyperglycaemic mice. *Diabetologia*. 2002; 45:1528–1532. [PubMed: 12436336]
21. Liu Y, Yan C, Wang Y, et al. Liver X receptor agonist T0901317 inhibition of glucocorticoid receptor expression in hepatocytes may contribute to the amelioration of diabetic syndrome in *db/db* mice. *Endocrinology*. 2006; 147:5061–5068. [PubMed: 16873540]
22. Wang SJ, Birtles S, de Schoolmeester J, et al. Inhibition of 11beta-hydroxysteroid dehydrogenase type 1 reduces food intake and weight gain but maintains energy expenditure in diet-induced obese mice. *Diabetologia*. 2006; 49:1333–1337. [PubMed: 16612591]
23. Andrews RC, Rooyackers O, Walker BR. Effects of the 11beta-hydroxysteroid dehydrogenase inhibitor carbenoxolone on insulin sensitivity in men with type 2 diabetes. *J Clin Endocrinol Metab*. 2003; 88:285–291. [PubMed: 12519867]
24. Odermatt A, Arnold P, Stauffer A, et al. The N-terminal anchor sequences of 11beta-hydroxysteroid dehydrogenases determine their orientation in the endoplasmic reticulum membrane. *J Biol Chem*. 1999; 274:28762–28770. [PubMed: 10497248]
25. Mziaut H, Korza G, Hand AR, et al. Targeting proteins to the lumen of endoplasmic reticulum using N-terminal domains of 11beta-hydroxysteroid dehydrogenase and the 50-kDa esterase. *J Biol Chem*. 1999; 274:14122–14129. [PubMed: 10318829]
26. Mason PJ, Stevens D, Diez A, et al. Human hexose-6-phosphate dehydrogenase (glucose 1-dehydrogenase) encoded at lp36: coding sequence and expression. *Blood Cells Mol Dis*. 1999; 25:30–37. [PubMed: 10349511]
27. Ozols J. Isolation and the complete amino acid sequence of lumenal endoplasmic reticulum glucose-6-phosphate dehydrogenase. *Proc Natl Acad Sci USA*. 1993; 90:5302–5306. [PubMed: 8506377]
28. Hewitt KN, Walker EA, Stewart PM. Minireview: hexose-6-phosphate dehydrogenase and redox control of 11beta-hydroxysteroid dehydrogenase type 1 activity. *Endocrinology*. 2005; 146:2539–2543. [PubMed: 15774558]

29. McCormick KL, Wang X, Mick GJ. Evidence that the 11 beta-hydroxysteroid dehydrogenase (11beta-HSD1) is regulated by pentose pathway flux. Studies in rat adipocytes and microsomes. *J Biol Chem.* 2006; 281:341–347. [PubMed: 16234247]
30. van Schaftingen E, Gerin I. The glucose-6-phosphatase system. *Biochem J.* 2002; 362:513–532. [PubMed: 11879177]
31. Chou JY, Matern D, Mansfield BC, et al. Type I glycogen storage diseases: disorders of the glucose-6-phosphatase complex. *Curr Mol Med.* 2002; 2:121–143. [PubMed: 11949931]
32. Atanasov AG, Nashev LG, Schweizer RA, et al. Hexose-6-phosphate dehydrogenase determines the reaction direction of 11beta-hydroxysteroid dehydrogenase type 1 as an oxoreductase. *FEBS Lett.* 2004; 571:129–133. [PubMed: 15280030]
33. Banhegyi G, Benedetti A, Fulceri R, et al. Cooperativity between 11 beta-hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase in the lumen of the endoplasmic reticulum. *J Biol Chem.* 2004; 279:27017–27021. [PubMed: 15090536]
34. Draper N, Walker EA, Bujalska IJ, et al. Mutations in the genes encoding 11beta-hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase interact to cause cortisone reductase deficiency. *Nat Genet.* 2003; 34:434–439. [PubMed: 12858176]
35. Lavery GG, Walker EA, Araper N, et al. Hexose-6-phosphate dehydrogenase knock-out mice lack 11beta-hydroxysteroid dehydrogenase type 1-mediated glucocorticoid generation. *J Biol Chem.* 2006; 281:6546–6551. [PubMed: 16356929]
36. Liu YJ, Wang Y, Nakagawa Y, et al. Inhibition of hepatic glucocorticoid receptor and hexose-6-phosphate dehydrogenase expression ameliorates diet-induced insulin resistance and obesity in mice. *J Mol Endocrinol.* 2008; 41:53–64. [PubMed: 18524870]
37. Hsu CL, Huang SL, Yen GC. Inhibitory effect of phenolic acids on the proliferation of 3T3-L1 preadipocytes in relation to their antioxidant activity. *J Agric Food Chem.* 2006; 54:4191–4197. [PubMed: 16756346]
38. Belkaid A, Currie JC, Desganes J, et al. The chemo-preventive properties of chlorogenic acid reveal a potential new role for the microsomal glucose-6-phosphate translocase in brain tumor progression. *Cancer Cell Int.* 2006; 6:7. [PubMed: 16566826]
39. Marcolongo P, Senesi S, Gava B, et al. Metyrapone prevents cortisone-induced preadipocyte differentiation by depleting luminal NADPH of the endoplasmic reticulum. *Biochem Pharmacol.* 2008; 76:382–390. [PubMed: 18599022]
40. Nammi S, Dembele K, Nyomba BLG. Increased 11beta-hydroxysteroid dehydrogenase type-1 and hexose-6-phosphate dehydrogenase in liver and adipose tissue of rat offspring exposed to alcohol in utero. *Am J Physiol Regul Integr Comp Physiol.* 2007; 292:R1101–R1109. [PubMed: 17122334]
41. Liu Y, Nakagawa Y, Wang Y, et al. Leptin activation of corticosterone production in hepatocytes may contribute to the reversal of obesity and hyperglycemia in leptin-deficient *ob/ob* mice. *Diabetes.* 2003; 52:1409–1416. [PubMed: 12765951]
42. Hiraiwa H, Pan CJ, Lin B, et al. Inactivation of the glucose 6 phosphate transporter causes glycogen storage disease type 1b. *J Biol Chem.* 1999; 274:5532–5536. [PubMed: 10026167]
43. Uckaya G, Karadurmus N, Kutlu O, et al. Adipose tissue 11-beta-hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase gene expressions are increased in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract.* 2008; 2:S135–S140. [PubMed: 18963204]
44. Nakano S, Inada Y, Masuzaki H, et al. Bezafibrate regulates the expression and enzyme activity of 11beta-hydroxysteroid dehydrogenase type 1 in murine adipose tissue and 3T3-L1 adipocytes. *Am J Physiol Endocrinol Metab.* 2007; 292:E1213–E1222. [PubMed: 17190905]
45. Balachandran A, Guan H, Sellan M, et al. Insulin and dexamethasone dynamically regulate adipocyte 11beta-hydroxysteroid dehydrogenase type 1. *Endocrinology.* 2008; 149:4069–4407. [PubMed: 18467433]
46. Taylor, Al; Frizzell, N.; McKillop, AM., et al. Effect of RU486 on hepatic and adipocyte gene expression improves diabetes control in obesity-type 2 diabetes. *Horm Metab Res.* 2009; 41:899–904. [PubMed: 19670152]

47. Piccirella S, Czegle I, Lizak B, et al. Uncoupled redox systems in the lumen of the endoplasmic reticulum. Pyridine nucleotides stay reduced in an oxidative environment. *J Biol Chem.* 2006; 281:4671–4677. [PubMed: 16373343]
48. Warlker EA, Ahmed A, Lavery GG, et al. 11beta-Hydroxysteroid dehydrogenase type 1 regulation by intracellular glucose 6-phosphate provides evidence for a novel link between glucose metabolism and hypothalamo-pituitary- adrenal axis function. *J Biol Chem.* 2007; 282:27030–270306. [PubMed: 17588937]
49. Marcolongo P, Piccirella S, Schesi S, et al. The glucose-6-phosphate transporter-hexose-6-phosphate dehydrogenase-11beta- hydroxysteroid dehydrogenase type 1 system of the adipose tissue. *Endocrinology.* 2007; 148:2487–2495. [PubMed: 17303657]
50. Hiraiwa H, Chou YJ. Glucocorticoids activate transcription of the gene for the glucose-6-phosphate transporter, deficient in glycogen storage disease type 1b. *DNA Cell Biol.* 2001; 20:447–453. [PubMed: 11560776]
51. Opherck C, Tranche F, Kellendonk C, et al. Inactivation of the glucocorticoid receptor in hepatocytes leads to fasting hypoglycemia and ameliorates hyperglycemia in streptozotocin-induced diabetes mellitus. *Mol Endocrinol.* 2004; 18:1346–1353. [PubMed: 15031319]





Fig. 1. 11 -HSD1 and H6PDH activity and protein levels in liver microsomes of lean *db/+* and obese *db/db* mice treated with vehicle or RU486. **a** 11 -HSD1 reductase activity was measured in mouse liver microsomes using 11-DHC as substrate in the presence of NADPH. Production and relative quantification of 11 -HSD1 (**b**) and H6PDH (**c**) protein was done relative to the amount of GAPDH. **d** H6PDH activity was measured in liver microsomes on the basis of NADPH formation using 2 mmol/l G6P as substrate in the presence of NAD. Data are mean ± SEM from seven to eight mice (per group). ***p*<0.01, ****p*<0.001 vs *db/+* controls; †*p*<0.05, ††*p*<0.01 vs *db/db* controls. White bars, vehicle; black bars, RU486

**Fig. 2.**

a Quantitative real-time RT-PCR analysis demonstrating the relative alterations of *Hsd1*, *H6pdh* and *G6pt* mRNA expression in the livers of *db/+* and *db/db* mice treated with vehicle or RU486. White bars, *db/+*; black bars, *db/+*+RU486; hatched bars, *db/db*; grey bars, *db/db*+RU486. **b** Uptake of [^{14}C]G6P into the liver microsomes of lean *db/+* and *db/db* mice treated with vehicle or RU486. White squares, *db/+*; black squares, *db/+*+RU486; white circles, *db/db*; black circles, *db/db*+RU486. **c** Expression levels of *Pepck* and *G6pase* mRNA in the liver of *db/+* and *db/db* mice treated with vehicle or RU486. White bars, *db/+*; black bars, *db/+*+RU486; hatched bars, *db/db*; grey bars, *db/db*+RU486. **d** The effects of RU486 on AUC of plasma glucose level in ITT. Data are mean \pm SEM from seven to eight mice per group. ** $p < 0.01$ vs *db/+* controls; $\dagger\dagger p < 0.01$ vs *db/db* controls




Fig. 3. Adipose 11 -*Hsd1*, *H6pd* and *G6pt* mRNA expression in (a) subcutaneous and (b) epididymal fat of lean *db/+* and obese *db/db* mice treated with vehicle or RU486. Relative expression of mRNA levels was measured by RT-PCR and normalised to 18S rRNA expression. * $p < 0.05$, ** $p < 0.01$ vs *db/+* control; †† $p < 0.01$ vs *db/db* controls. White bars, *db/+*; hatched bars, *db/+* + RU486; black bars, *db/db*; grey bars, *db/db*+RU486




Fig. 4. Effects of corticosterone and insulin on primary cultures of *db/db* mouse hepatocytes. Levels of H6PDH production (a) and activity (b) in cells treated with corticosterone for 48 h. In (b), white circles, control; black squares, corticosterone; black triangles, corticosterone + RU486. The levels of *H6pdh*, *11-Hsd1*, *G6Pase* and *Pepck* mRNA expression (c, d) in cells treated with corticosterone (1×10^{-6} mol/l) or insulin (1×10^{-6} mol/l) in the presence or absence of RU486 (1×10^{-6} mol/l) for 48 h. In (d), white bars, *11-Hsd1*; black bars, *G6Pase*; grey bars, *Pepck*. Production and relative quantification of H6PDH protein levels are expressed relative to the amount of GAPDH. Relative H6PDH activity was measured on the basis of NADPH production at 5 min intervals for 20 min. Values are the mean \pm SEM from three separate culture preparations. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs controls; †† $p < 0.01$, ††† $p < 0.001$ vs corticosterone-treated hepatocytes. C, control; B, corticosterone; I, insulin; RU, RU486




Fig. 5.
a Effect of different doses of corticosterone on *G6pt* mRNA expression in primary cultures of hepatocytes from *db/db* mice. **b** Hepatocytes were incubated with corticosterone (10^{-6} mol/l) or insulin (10^{-6} mol/l) for 48 h in the absence or present of RU486. Expression and relative quantification of *G6pt* mRNA levels expressed relative to the amount of 18S in vehicle-treated cells. Values are the mean±SEM from three separate culture preparations. ** $p < 0.01$, *** $p < 0.001$ vs vehicle-treated cells. C, control; B, corticosterone; I, insulin; RU, RU486




Fig. 6. Suppression of *H6pd* expression by *H6pd* siRNA decreased 11 -HSD1 reductase activity and attenuated the effects of corticosterone on H6PDH and 11 -HSD1 in primary culture of hepatocytes. Cells were transfected with either *H6pd* siRNA (si) or negative control siRNA (N) and cultured with or without corticosterone (**b**; 1×10^{-6} mol/l) for 48 h. **a** Western blots showing effect of *H6pd* siRNA inhibition on H6PDH protein in cells. **b** *H6pd* and *11 -Hsd1c* mRNAs were measured by real- time RT-PCR. White bars, *H6pd*; black bars, *11 -Hsd1c*. 11 -HSD1 reductase activity was expressed as the percentage of [³H]DHC converted to [³H]corticosterone (b) in medium from hepatocytes. Values are the mean±SEM from three separate culture preparations. ** $p < 0.01$, *** $p < 0.001$ vs negative control; ††† $p < 0.001$ vs corticosterone-treated hepatocytes




Fig. 7. The effects of CA on H6PDH and 11 -HSD1 in primary cultures of hepatocytes from *db/db* mice. **a** Primary hepatocytes were incubated with increased concentrations of the G6PT inhibitor CA (0– 200 μmol/l) for 48 h and H6PDH protein levels were determined by western blot analysis. **b** Hepatocytes were incubated with CA (150 μmol/l) for 48 h. *H6pdh* and *11 -Hsd1* mRNA were measured by real-time RT-PCR. Values are the mean±SEM from three separate culture preparations. ** $p < 0.01$ vs vehicle-treated hepatocytes. White bars, vehicle; black bars, CA

Table 1

Body weight, blood glucose, plasma insulin, lipid and corticosterone

Variable	<i>db/+ + vehicle</i>	<i>db/+ + RU486</i>	<i>db/db + vehicle</i>	<i>db/db + RU486</i>
<i>n</i>	7	7	8	8
Body weight (g)	31±1.5	31.5±1.8	55±4.3 ^{***}	48.8±3.6
Glucose (mmol/l)	8.2±0.5	7.3±0.6	30±2.3 ^{***}	11.7±1.2 ^{†††}
Insulin (pmol/l)	2.8±0.5	2.4±0.4	17.2±2.9 ^{***}	15±3.1
Cholesterol (mmol/l)	1.6±0.1	57±5.7	110±7 ^{**}	94±12
Triacylglycerols (mmol/l)	1.5±0.2	1.4±0.13	2.5±0.25 ^{**}	2.1 ±0.2
Corticosterone (nmol/l)	122±25	295±34 ^{**}	469±110 ^{**}	870±121 ^{†††}

Data are mean ± SEM from seven to eight mice per group

^{**}
 $p < 0.01$ ^{***}
 $p < 0.001$ vs *db/+* vehicle^{†††}
 $p < 0.001$ vs *db/db* + vehicle