



Published in final edited form as:

J Org Chem. 2013 September 6; 78(17): 8909–8914. doi:10.1021/jo401498w.


One-Pot Anti-Markovnikov Hydroamination of Unactivated Alkenes by Hydrozirconation and Amination

Alexandra E. Strom and John F. Hartwig

Department of Chemistry, University of California, Berkeley, Berkeley, CA 94720

John F. Hartwig: jhartwig@berkeley.edu

Abstract



A one-pot hydroamination of alkenes is reported. The synthesis of primary and secondary amines from unactivated olefins was accomplished in the presence of a variety of functional groups. Hydrozirconation, followed by amination with nitrogen electrophiles, provides exclusive anti-Markovnikov selectivity, and most products are isolated in high yields without the use of column chromatography.

The conversion of alkenes to alkylamines is a challenging, but synthetically important transformation.^{1, 2} This transformation traditionally has been conducted by hydroboration, oxidation, and conversion of the alcohol or aldehyde to the amine. A more efficient method would be the combination of hydrometallation and direct conversion of the alkylmetal intermediate to the amine. The combination of hydroboration and reaction of the alkylboron intermediate with nitrogen electrophiles to form the corresponding amine products was described as early as 1964³ (Scheme 1a), but these reactions are limited to the formation of primary amines and occurred in modest yields. A more general method to perform anti-Markovnikov amination of alkenes in high yields with readily available reagents and wide functional group tolerance has not been reported.^{1, 4}

The combination of hydroboration and amination of the resulting alkylboron compound has been revisited since the initial report.⁵⁻⁸ Recently, amination of alkyl boranes to form tertiary amines has been accomplished with copper catalysts.⁹⁻¹² In addition, three equivalents of *n*-butyl lithium and MeONH_2 have been shown to react with pinacol boronate esters to form primary amines and anilines (Scheme 1c).¹³

Correspondence to: John F. Hartwig, jhartwig@berkeley.edu.

Supporting Information **Available**. Spectroscopic data for all compounds. The Material is available free of charge via the Internet at <http://pubs.acs.org>

Despite these recent advances in the synthesis of amines from alkenes, there are a number of limitations. Particularly relevant to the work we report, methods for the synthesis of secondary amines from alkenes are limited to isolated examples, and these reactions proceed in low yield (Scheme 1b).¹⁴⁻¹⁸ In addition, the utility of reactions to form primary amines is limited by the properties of the reagent used for amination. Hydrazoic acid has been used as the reagent to provide primary amines from alkyl boron reagents in high yields,⁶ but the explosion hazard of hydrazoic acid limits the utility of this method. Moreover, reactions conducted with the reagent formed from *n*-butyl lithium and a solution of MeONH₂ require long times with boronic esters at elevated temperatures, and 3 equivalents of each are required for acceptable yields (Scheme 1).

To address these limitations, we followed an alternative strategy involving alkylzirconium intermediates. Srebnik and Zheng reported the stepwise addition of Schwartz's reagent (bis(cyclopentadienyl)zirconium chloride hydride), followed by *O*-mesitylsulfonyl hydroxylamine (MSH), to form primary alkylamine products from alkenes.¹⁹ Although this reaction provides a route to terminal primary amines, MSH decomposes over time and requires a multi-step synthesis from hydroxylamine.²⁰ Thus, a method is needed to convert alkylzirconium compounds to primary and secondary alkylamines in high yield with an accessible reagent for the sequence of hydrozirconation and amination to be practical.

Herein we report a method for the synthesis of primary and secondary amines from alkenes by a one-pot procedure based on hydrozirconation that occurs in good to high yields without isolation of intermediates. This protocol is based on preliminary data for the conversion of *N*-allyl indoles to the corresponding amines²¹ and is sufficiently mild to tolerate a range of aprotic functional groups. This process includes commercially available and readily synthesized reagents and forms products that were isolated, in most cases, without column chromatography. These attributes make the reaction suitable for the one-pot, anti-Markovnikov hydroamination of complex molecules.

Our studies on the hydroamination of alkenes via hydrozirconation began with an evaluation of the conditions for the conversion of the alkylzirconium compounds to the corresponding amines. The reaction of 1-octylzirconocene chloride (generated from 1-octene and Schwartz reagent) with *N*-methylhydroxylamine-*O*-sulfonic acid was complete in 30 minutes at 50 °C, but the yields of the *N*-methyloctylamine product were variable. Higher yields were obtained consistently if the alkylzirconocene formed in the first step was used immediately.²² Simple evaporation of the solvent from the final solution and aqueous extraction provided pure amine products. Acid-sensitive substrates were subjected to a modified aqueous extraction and purified by column chromatography.

With these conditions in hand, the reactions of a variety of alkenes were evaluated (Table 1). The amination reaction was conducted on a 4 mmol scale without reduction in yield (entry 2), and, at 10 mmol scale, the yield remained synthetically useful (entry 3). The reaction in entry 14 shows that a Boc-protected secondary amine is stable to Schwartz reagent, and the reactions in entries 13, 7, and 8 show that the reaction occurs with strained and 1,1-disubstituted alkenes to form the corresponding amines cleanly. Amination of the sterically hindered vinyl cyclohexane (entry 11) and *tert*-butyl ethylene (entry 12) proceeded in good

yield, although 1.1 equivalents of *tert*-butyl ethylene were necessary to ensure complete conversion of Schwartz's reagent. Vinylarenes (entries 9 and 10) were converted to substituted phenethylamines without competing amination at the benzylic position. Yields were lower for the reaction of 3-trifluoromethyl styrene than that of 3,4-dimethoxy styrene, and this lower yield is consistent with a higher equilibrium concentration of the benzylic zirconocene complex due to stabilization from the electron deficient arene.

The synthesis of primary amines from commercially available hydroxylamine-*O*-sulfonic acid was also studied (Table 2). The amination step was complete in minutes at room temperature, and the products were isolated in high yields. In one case, the combination of hydrozirconation and conversion of the alkylzirconium intermediate to the primary amine (entry 4) gave a mixture of benzylic and terminal amines; the same sequence conducted with *N*-methylhydroxylamine-*O*-sulfonic acid formed only the terminal amine (Table 1, entry 6). The absence of the benzylic amine product from the sequence that generates the secondary amine suggests that the amination of the branched alkylzirconocene does not occur with the more hindered *N*-methyl aminating reagent and the remaining starting material is converted to the protonolysis product, propylbenzene.

It is well established that the hydrozirconation of internal alkenes forms terminal alkylzirconium products. To test if this feature of hydrozirconation would lead to a method for the one-pot conversion of internal alkenes to terminal amines, the hydrozirconation-amination process was conducted with a mixture of internal alkenes (Table 2, entry 5). 1-Octylamine was the only observed product, indicating complete conversion to the linear alkylzirconocene and amination of the linear alkylzirconocene intermediate.

The alkyl substituent on nitrogen was varied to investigate the potential of this route to form additional terminal secondary amines from alkenes (Table 3). Under the standard conditions, the combination of hydrozirconation of 1-octene, followed by amination of the alkylzirconium compound with *N*-(3-phenylpropyl)hydroxylamine-*O*-sulfonic acid, consistently provided 40-50% yields of alkylamine product (entry 1). However, when this sequence was conducted with the amination step at 80 °C, the *N*-(3-phenylpropyl)amine product was isolated in 72% yield (entry 2). *N*-heptyl hydroxylamine-*O*-sulfonic acid was synthesized from heptanal in 3 steps, and the combination of hydrozirconation of octene and reaction with this *N*-alkyl hydroxylamine-*O*-sulfonic acid afforded 79% yield of *N*-heptyl octylamine (entry 3) following the conditions from entry 2. The presence of additional substituents *alpha* to nitrogen was detrimental to the yield of the desired amine (entries 4 and 5); no product was observed in the case of *N,N*-dimethyl hydroxylamine-*O*-sulfonic acid to form tertiary amines at any temperature (entry 6).

To investigate the ability of this method to be used with complex molecules containing basic nitrogens and oxygenated frameworks, the hydroamination of the *cinchona* alkaloid 9-*O*-benzyl quinine **1a** was conducted. This substrate might be expected to quench the zirconium hydride by binding of the nitrogen in the quinoline or highly basic quinuclidine. This substrate also contains a doubly benzylic ether, which could be cleaved under reducing conditions. Despite these possible side reactions, the transformation of the alkene to the secondary alkylamine proceeded cleanly to afford the product **1b** in 92% yield.

The reaction of mannose-derived **2a** was conducted to assess the ability of the hydrozirconation-amination sequence to be conducted with a highly oxygenated substrate. Given the oxophilicity and Lewis acidity of zirconium, cleavage or epimerization at the anomeric position of **2a** might be expected to occur. However, this reaction of the alkene formed the secondary alkylamine product **2b** in 66% isolated yield. The sole byproduct observed from this reaction was formed by reduction of the alkene, presumably due to protonolysis of the intermediate alkyl zirconocene.

In summary, a one-pot hydroamination of unactivated alkenes has been developed. The synthesis of primary and secondary amines from alkenes containing a variety of functional groups occurs under the reported conditions, and this method was found to be applicable to the derivatization of complex molecules.

Experimental Section

General

Unless otherwise stated, reactions were performed using standard Schlenk and drybox techniques. Liquid alkene substrates were stored over 4 Å molecular sieves in the drybox. Bis(cyclopentadienyl) zirconium chloride hydride was purchased from Strem Chemicals and stored at room temperature in a drybox. Hydroxylamine-*O*-sulfonic acid was purchased from Sigma Aldrich and stored at 0 °C or in the drybox freezer at -30 °C. *N*-Boc-(2,2-diphenyl-hept-6-enyl)-methyl-amine,²³ 9*O*-benzyl quinine,²⁴ 2-methyl-1-octene, and 2-allylphenyl 2,3,4,6-tetra-*O*-benzyl- α -D-mannopyranoside²⁵ were prepared by literature procedures. Thin-layer chromatography (TLC) plates were visualized by UV fluorescence quenching, potassium permanganate, or *p*-anisaldehyde staining. Chemical shifts are reported in ppm relative to the residual solvent peak (CDCl₃ = 7.26 ppm for ¹H and 77.0 ppm for ¹³C; (CD₃)₂SO = 2.50 ppm for ¹H and 39.52 ppm for ¹³C) and are reported relative to Me₄Si (δ 0.00). Mass spectrometry analyses (ESI-MS) were performed using an ion trap analyzer.

N-methylhydroxylamine-*O*-sulfonic acid

A 250 mL Schlenk flask was charged with *N*-methylhydroxylamine hydrochloride (5.00 g, 59.9 mmol) and a stir bar, dried under high vacuum for one hour, and backfilled with nitrogen. Under a constant flow of nitrogen, 10 mL of dry, degassed DCM was added, and the flask was cooled to -78 °C. Aliquots of ClSO₃H⁴ (1 mL) were added every 15 min to total 8 portions (8.00 mL, 120. mmol). The reaction mixture was stirred at -78 °C for an additional 30 min and warmed to room temperature. Dry, degassed Et₂O (20 mL) was added slowly, and the solid was filtered under nitrogen, washing with three 10 mL portions of Et₂O to yield 5.7 g of white, fluffy solid (74% yield), which could be stored indefinitely in the drybox at -30 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 2.88 (s, 3H) ¹³C NMR (101 MHz, DMSO) δ 36.4. HRMS-ESI (*m/z*): [M - H]⁻¹ calcd for CH₄NO₄S, 125.9867; found 125.9866.

N-isopropylhydroxylamine-O-sulfonic acid

A 250 mL Schlenk flask was charged with *N*-isopropylhydroxylamine hydrochloride (623 mg, 5.62 mmol) and a stir bar, dried under high vacuum for one hour, and backfilled with nitrogen. Under a constant flow of nitrogen, 20 mL of dry, degassed DCM was added, and the flask was cooled to -78 °C. Aliquots of ClSO₃H were added every 15 min over 2 hours, to total 1.50 equiv (563 μL, 8.43 mmol). The reaction mixture was stirred at -78 °C for an additional 15 min and warmed to room temperature over 30 min. Dry, degassed Et₂O (10 mL) was added slowly, and the solid was filtered under nitrogen and washed with three 5 mL portions of Et₂O to yield 640 mg of white, fluffy solid (74% yield), which could be stored indefinitely in the drybox at -30 °C. ¹H NMR (500 MHz, DMSO) δ 3.54 (h, J = 6.5 Hz, 1H), 1.21 (d, J = 6.5 Hz, 6H). ¹³C NMR (126 MHz, DMSO) δ 52.6, 16.6. HRMS-ESI (m/z): [M - H]⁻¹ calcd for C₃H₈NO₄S, 154.0180; found 154.0180.

N-(3-phenylpropyl)hydroxylamine-O-sulfonic acid

3-phenylpropionaldehyde oxime

The title compound was prepared via a modified literature procedure.²⁷ In a 50 mL flask, a solution of 3-phenylpropionaldehyde (1.98 mL, 15.0 mmol) in 10 mL of H₂O was stirred, open to air. A solution of hydroxylamine hydrochloride (5.21 g, 75 mmol) in EtOH (10 mL) was added at room temperature, and the solution was cooled to 0 °C. NaOH (1.0 g, excess) in 5 mL of H₂O was added slowly, and the solution was warmed to room temperature. The reaction was heated at 70 °C and stirred for 2 hours. The reaction was allowed to cool to room temperature, and the precipitate was filtered, washed with water, dried under high vacuum, and used without further purification.

N-(3-phenylpropyl)hydroxylamine

In a 100 mL flask under air, 3-phenylpropionaldehyde oxime (745 mg, 5.00 mmol) was stirred at 0 °C in MeOH (10 mL). A mixture of 6N HCl(aq) and MeOH (1:1 v/v) and a solution of NaBH₃CN (408 mg, 6.50 mmol) in MeOH were alternately added dropwise over 30 min, maintaining pH <3. The solution was warmed to room temperature, maintaining pH <3, and stirred until no oxime remained, as determined by thin-layer chromatography (about 30 min). The reaction was quenched with brine (10 mL), basified with aqueous 6N NaOH to pH >9, and transferred to a separatory funnel. The solution was extracted with 2 × 20 mL Et₂O, and the combined organic layers were washed with brine, dried with MgSO₄, and concentrated in vacuo. The crude material was taken up in benzene (10 mL) and treated with conc HCl (1 equiv). The solvent was removed under high vacuum and the crude HCl salt was used in the next step.

N-(3-phenylpropyl)hydroxylamine-O-sulfonic acid

A 50 mL Schlenk flask was charged with solid *N*-(3-phenylpropyl)hydroxylamine hydrochloride (915 mg, 4.87 mmol) and a stir bar, dried under high vacuum for one hour, and backfilled with nitrogen. Under a constant flow of nitrogen, 8 mL of dry, degassed DCM was added, and the flask was cooled to -78 °C. Aliquots of ClSO₃H were added every 15 min over 2 hours to total 1.50 equiv (488 μL, 7.31 mmol). The reaction mixture was stirred at -78 °C for an additional 15 min and warmed to room temperature over 30 min.

Dry, degassed Et₂O (10 mL) was added slowly, and the precipitate was filtered under nitrogen and washed with three 5 mL portions of Et₂O to yield 890 mg of white, fluffy solid (79%), which could be stored indefinitely in the drybox at -30 °C. ¹H NMR (600 MHz, DMSO) δ 11.24 – 11.00 (br s, 1H), 7.30 (m, 2H), 7.21 (m, 3H), 6.06 – 3.94 (br s, 1H), 3.18 (m, 2H), 2.65 (m, 2H), 1.90 (m, 2H). ¹³C NMR (151 MHz, DMSO) δ 140.7, 128.5, 128.3, 126.1, 49.2, 31.7, 25.0. HRMS–ESI (*m/z*): [M – H]⁻¹ calcd for C₉H₁₂NO₄S, 230.0493; found 230.0494.

***N*-(heptyl)hydroxylamine-*O*-sulfonic acid**

1-heptanal oxime

The title compound was prepared via a modified literature procedure.²⁷ In a 100 mL round bottom flask, a solution of 3-phenylpropionaldehyde (1.41 mL, 10.0 mmol) in 15 mL of H₂O was stirred, open to the air. A solution of hydroxylamine hydrochloride (5.21 g, 75 mmol) in EtOH (5 mL) was added at room temperature, and the solution was cooled to 0 °C. NaOH (0.6 g, 15 mmol) in 5 mL of H₂O was added slowly, and the solution was warmed to room temperature. The reaction was heated at 70 °C and stirred for 1 hour. The reaction was allowed to cool to room temperature, and the product was allowed to precipitate over night. The precipitate was filtered, washed with water, dried under high vacuum, and used without further purification (800 mg).

***N*-(heptyl)hydroxylamine**

In a 50 mL round bottom flask under air, heptanal oxime (794 mg, 6.14 mmol) was stirred at -60 °C in MeOH (10 mL). A mixture of 6N HCl(aq) and MeOH (1:1 v/v) and a solution of NaBH₃CN (502 mg, 8.00 mmol) in MeOH (5 mL) were alternately added dropwise over 30 min, maintaining pH <3. The solution was warmed to room temperature, maintaining pH <3, and stirred until no oxime remained, as determined by thin-layer chromatography (about 30 min). The reaction was quenched with brine (10 mL), basified with aqueous 6N NaOH to pH >8, and transferred to a separatory funnel. The solution was extracted with 2 × 20 mL Et₂O, and the combined organic layers were washed with brine, dried with MgSO₄, and concentrated in vacuo. The crude material was taken up in benzene (10 mL) and treated with conc HCl (1 equiv). The solvent was removed under high vacuum and the crude HCl salt was used in the next step (736 mg).

***N*-(heptyl)hydroxylamine-*O*-sulfonic acid**

A 50 mL Schlenk flask was charged with solid *N*-(3-phenylpropyl)hydroxylamine hydrochloride (736 mg, 5.61 mmol) and a stir bar, dried under high vacuum for one hour, and backfilled with nitrogen. Under a constant flow of nitrogen, 5 mL of dry, degassed DCM was added, and the flask was cooled to -78 °C. Aliquots of ClSO₃H were added every 15 min over 1.5 hours to total 1.50 equiv (560 μL, 7.31 mmol). The reaction mixture was stirred at -78 °C for an additional 15 min and warmed to room temperature over 30 min. The reaction was cooled to -78 °C and dry, degassed Et₂O (10 mL) was added slowly. The precipitate was filtered under nitrogen and washed with three 5 mL portions of Et₂O to yield 1.04 g of white, fluffy solid (88%), which could be stored indefinitely in the drybox at -30 °C. ¹H NMR (600 MHz, DMSO) δ 5.00 – 3.40 (br s, 2H), 3.25 – 2.99 (m, 2H), 1.73 – 1.46

(m, 2H), 1.38 – 1.15 (m, 8H), 0.86 (t, $J = 5.1$ Hz, 3H). ^{13}C NMR (151 MHz, DMSO) δ 49.6, 31.0, 28.2, 25.7, 23.1, 22.0, 134.0. $[\text{M} - \text{H}]^{-1}$ calcd for $\text{C}_7\text{H}_{16}\text{NO}_4\text{S}$, 210.0806; found 210.0804.

***N*-(*tert*-butoxy carbonyl)-(2,2-diphenyl-hept-6-enyl)-7-methylamine**

A 20 mL vial was loaded with a stir bar, (2,2-diphenyl-hept-6-enyl)-methylamine (279 mg, 1.00 mmol), 3 mL THF, and triethylamine (420 mL, 3.00 mmol). Di-*tert*-butyl dicarbonate (275 mg, 1.10 mmol) was added, and the mixture was stirred at room temperature for one hour. The reaction was diluted with H_2O (20 mL) and EtOAc (20 mL) in a separatory funnel, and the aqueous phase was extracted with 2×20 mL EtOAc. The combined organic layers were washed with brine, dried with MgSO_4 , and concentrated by rotary evaporation. The product was lyophilized in benzene under high vacuum to yield 379 mg of white solid (quant.) as a mixture of rotamers, and used without further purification. NMR analysis was performed at 60 °C. ^1H NMR (600 MHz, CDCl_3) δ 7.30 – 7.24 (m, 4H), 7.24 – 7.17 (m, 6H), 5.78 – 5.63 (m, 1H), 4.93 (d, $J = 17.1$ Hz, 1H), 4.89 (d, $J = 10.1$ Hz, 1H), 4.05 (s, 2H), 2.22 – 2.04 (m, 5H), 2.04 – 1.90 (m, 2H), 1.45 (s, 9H), 1.24 – 1.11 (m, 2H). ^{13}C NMR (151 MHz, CDCl_3) δ 156.8, 147.1, 138.9, 128.6, 127.9, 126.0, 114.4, 79.4, 57.4, 51.5, 36.5, 36.3, 34.4, 28.4, 23.9.

General Procedure A for the Hydrozircation/Amination of Alkenes

In a drybox, an oven-dried, screw-top, 1 dram vial was charged with a stir bar, and bis(cyclopentadienyl) zirconium chloride hydride (103 mg, 0.400 mmol, 1.00 equiv).⁶ THF (1.5 mL, 3.75 M) and alkene (0.400 mmol, 1.00 equiv) were added. The vial was sealed with a cap equipped with a PTFE-lined septum and stirred at room temperature. After 0.5 to 4 h the reaction mixture became homogeneous,⁷ and *N*-alkylhydroxylamine-*O*-sulfonic acid (0.600 mmol, 1.5 equiv) was added. The vial was re-sealed, removed from the drybox, and heated at 50 °C for 30 min with stirring. The reaction was cooled to room temperature, diluted with wet THF (5 mL), and concentrated in vacuo. The crude solids were dissolved in aqueous 1 M HCl (5 mL), transferred to a separatory funnel, and washed twice with Et_2O . The combined washes were extracted once more with 5 mL of 1 M HCl, and the combined aqueous phases were basified to pH >8 with 1 M NaOH and extracted twice with Et_2O . The organic phases from the second extraction were washed with brine, dried with Na_2SO_4 , and concentrated in vacuo to yield the desired product.

General Procedure B for the Hydrozirconation/Amination of Alkenes

In a drybox, an oven dried, 1 dram vial was charged with a stir bar, and bis(cyclopentadienyl) zirconium chloride hydride (103 mg, 0.400 mmol, 1.00 equiv).⁶ THF (1.5 mL) and alkene (0.400 mmol, 1.00 equiv) were added. The vial was sealed with a cap equipped with a PTFE-lined septum and stirred at room temperature. After 0.5 to 4 h the reaction mixture became homogeneous, and hydroxylamine-*O*-sulfonic acid (67.9 mg, 0.600 mmol, 1.5 equiv) was added. The vial was re-sealed, stirred at room temperature for 30 min, and removed from the drybox. The reaction was diluted with wet THF (5 mL), and concentrated in vacuo. The crude solids were dissolved in aqueous 1 M HCl (5 mL), transferred to a separatory funnel, and washed twice with Et_2O . The combined washes were

extracted once more with 5 mL of 1 M HCl, and the combined aqueous phases were basified to pH >8 with 1 M NaOH and extracted twice with Et₂O. The organic phases from the second extraction were washed with brine, dried with Na₂SO₄, and concentrated in vacuo to yield the desired product.

General Procedure C for the Hydrozirconation/Amination of Alkenes

General Procedure A was followed until the workup step. The crude solids were dissolved in aqueous 1 M NaOH (5 mL), transferred to a separatory funnel, diluted with H₂O, and extracted twice with Et₂O. The combined organic phases were washed with brine, dried with Na₂SO₄, and concentrated in vacuo. The crude residue was purified via column chromatography, eluting with 0.5% NH₄OH, 1-10% MeOH in DCM (v/v).

N-(3-phenylpropyl)octylamine

In a drybox, an oven-dried, 1 dram vial was charged with a stir bar and bis(cyclopentadienyl) zirconium chloride hydride (51.6 mg, 0.200 mmol, 1.00 equiv),⁶ THF (0.75 mL) and 1-octene (31.4 mL, 0.200 mmol, 1.00 equiv) were added. The vial was sealed with a cap equipped with a PTFE-lined septum and stirred at room temperature. After 1 h the reaction mixture became homogeneous, and *N*-(3-phenylpropyl)hydroxylamine-*O*-sulfonic acid (69.6 mg, 0.300 mmol, 1.5 equiv) was added. The vial was re-sealed, removed from the drybox, and heated at 80 °C for 30 min with stirring. The reaction was cooled to room temperature, diluted with wet THF (2 mL), and concentrated in vacuo. The crude solids were dissolved in aqueous 1 M NaOH (5 mL), transferred to a separatory funnel, diluted with H₂O, and extracted twice with Et₂O. The combined organic phases were washed with brine, dried with Na₂SO₄, and concentrated in vacuo. The crude residue was purified via column chromatography eluting with 0.5% NH₄OH, 1-10% MeOH in DCM (v/v) to yield 37 mg of *N*-(3-phenylpropyl)octylamine (clear oil, 74%). ¹H NMR (600 MHz, CDCl₃) δ 7.34 – 7.27 (m, 2H), 7.25 – 7.14 (m, 3H), 2.74 – 2.61 (m, 4H), 2.59 (t, *J* = 7.3 Hz, 2H), 1.89 – 1.75 (m, 2H), 1.47 (m, 2H), 1.36 – 1.18 (m, 10H), 0.88 (t, *J* = 6.9 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 142.2, 128.3, 128.3, 125.7, 50.0, 49.5, 33.7, 31.8, 31.7, 30.1, 29.5, 29.2, 27.4, 22.6, 14.1.

N-(*tert*-butoxy carbonyl)-(2,2-diphenyl-heptyl)-7-methylamino)-methylamine

General procedure C. 120 mg (mixture of rotamers, clear oil, 74%). ¹H NMR (600 MHz, CDCl₃) δ 7.34 – 7.20 (m, 5H), 7.18 – 7.13 (m, 5H), 4.09 – 3.88 (m, 2H), 2.54 – 2.42 (m, 2H), 2.38 (s, 3H), 2.18 – 1.87 (m, 6H), 1.41 (m, 9H), 1.34 – 1.17 (m, 2H), 1.11 – 0.94 (m, 2H). ¹³C NMR (151 MHz, CDCl₃) δ 186.5, (186.1), 147.5, (146.5), 128.5, (128.3), 127.8, 126.0, (125.8), 79.8, (79.1), 57.9, (56.6), 52.1, (51.8), 51.3, (51.0), 36.5, 36.4, (36.3), 29.8, (29.3), 28.4, 28.1, (27.8), 24.4, (24.1). HRMS–ESI (*m/z*): [M + H]⁺ calcd for C₂₆H₃₉N₂O₂, 411.3007; found 411.2999.

9-*O*-benzyl-10,11-dihydro-11-methylamino-quinine

In a drybox, an oven-dried, screw-top, 1 dram vial was charged with a stir bar, and bis(cyclopentadienyl) zirconium chloride hydride (51.6 mg, 0.200 mmol, 1.00 equiv). THF-*d*₈ (1.0 mL) and 9-*O*-benzyl quinine (82.9 mg, 0.200 mmol, 1.00 equiv) were added, the vial

was sealed with a cap equipped with a PTFE-lined septum, and the reaction was stirred at room temperature. After 30 min the reaction mixture became homogeneous, and the solution was analyzed by NMR spectroscopy. The amount of remaining olefin was determined to be approximately 0.2 equiv, and a further portion of bis(cyclopentadienyl) zirconium chloride hydride (16.6 mg, 0.0640 mmol, 0.322 equiv) was added. The reaction was stirred for 1 h, at which point the reaction mixture became homogeneous, and *N*-methylhydroxylamine-*O*-sulfonic acid (38.1 mg, 0.300 mmol, 1.50 equiv) was added. The vial was re-sealed, removed from the drybox, and heated at 50 °C for 30 min with stirring. The reaction was cooled to room temperature, diluted with wet THF (2 mL), and concentrated in vacuo. The crude solids were dissolved in aqueous 1 M HCl (5 mL), transferred to a separatory funnel, diluted with H₂O, and washed twice with Et₂O. The combined washes were extracted once more with 5 mL of 1 M HCl, and the combined aqueous phases were basified to pH >8 with aqueous 1 M NaOH and extracted twice with Et₂O. The organic phases from the second extraction were washed with brine, dried with Na₂SO₄, and concentrated in vacuo to yield 82 mg of the desired product as a white solid (92%). NMR analysis was performed at 60 °C to minimize peak broadening, due to constrained rotation at C9. ¹H NMR (600 MHz, CDCl₃) δ 8.74 (d, *J* = 4.4 Hz, 1H), 8.03 (d, *J* = 9.2 Hz, 1H), 7.55 (s, 1H), 7.47 (d, *J* = 4.2 Hz, 1H), 7.41 – 7.28 (m, 6H), 5.82 – 5.64 (m, 1H), 4.55 – 4.43 (m, 2H), 4.01 (s, 3H), 3.70 – 3.56 (m, 1H), 3.35 – 3.21 (m, 2H), 2.90 – 2.78 (m, 1H), 2.60 (m, 3H), 2.39 (s, 3H), 2.08 – 1.97 (m, 1H), 1.91 – 1.75 (m, 3H), 1.65 – 1.47 (m, 4H), 1.27 (br s, 1H). ¹³C NMR (151 MHz, CDCl₃) δ 158.4, 150.4, 147.3(×2), 144.7, 137.4, 131.8, 128.5, 127.9, 127.7, 127.2, 122.4, 118.6, 101.2, 77.2, 71.3, 59.9, 57.5, 56.5, 48.6, 43.3, 34.8, 32.8, 32.4, 29.7, 25.7. HRMS: (*m/z* + 1) Calc'd 446.2802 Found 446.2799. HRMS–ESI (*m/z*): [M + H]⁺ calcd for C₂₈H₃₆N₃O₂, 446.2802; found 446.2799.

2-(3-methylaminopropyl)phenyl-2,3,4,6-tetra-*O*-benzyl- α -D-mannopyranoside

General Procedure C. 26 mg (colorless solid, 66%). ¹H NMR (500 MHz, CDCl₃) δ 7.45 – 7.26 (m, 18H), 7.24 – 7.08 (m, 5H), 6.94 (td, *J* = 7.3 Hz, 1.2 Hz, 1H), 5.58 (d, *J* = 1.8 Hz, 1H), 4.93 (d, *J* = 10.8 Hz, 1H), 4.80 (s, 1H), 4.80 (s, 1H), 4.71 (m, 2H), 4.67 (d, *J* = 12.0 Hz, 1H), 4.56 (d, *J* = 10.8 Hz, 1H), 4.48 (d, *J* = 11.9 Hz, 1H), 4.17 (t, *J* = 9.4 Hz, 1H), 4.07 (dd, *J* = 9.4 Hz, 3.0 Hz, 1H), 3.98 – 3.92 (m, 1H), 3.88 – 3.76 (m, 2H), 3.70 (d, *J* = 9.5 Hz, 1H), 2.56 – 2.47 (m, 4H), 2.35 (s, 3H), 1.75 – 1.67 (m, 2H). ¹³C NMR (151 MHz, CDCl₃) δ 154.3, 138.5, 138.38, 138.34, 138.2, 131.0, 129.9, 128.4, 128.4, 128.3, 128.2, 127.9, 127.8, 127.8, 127.74, 127.71, 127.68, 127.6, 127.4, 127.2, 122.1, 114.5, 96.5, 79.9, 75.1, 74.9, 74.8, 73.3, 72.73, 72.67, 72.4, 69.1, 51.8, 36.5, 30.3, 28.0. HRMS–ESI (*m/z*): [M + H]⁺ calcd for C₄₄H₅₀N₁O₆, 688.3633; found 688.3616.

(3,3-dimethyl)butylmethylamine

General procedure A was followed, using 1.10 equivalents of 3,3-dimethyl-1-butene (56.7 μ L, 0.400 mmol) to yield 42 mg (clear oil, 91%). ¹H NMR (400 MHz, CDCl₃) δ 2.63 – 2.52 (m, 2H), 2.44 (s, 3H), 2.21 (s, 2H), 1.45 – 1.32 (br s, 1H), 0.89 (s, 9H). ¹³C NMR (101 MHz, CDCl₃) δ 48.2, 43.7, 36.4, 29.8, 29.5.

(6-chloro)hexylmethylamine

General procedure A. 50 mg (clear oil, 83%). ^1H NMR (600 MHz, CDCl_3) δ 3.53 (t, $J=6.7$ Hz, 2H), 2.59 (t, $J=7.2$ Hz, 2H), 2.44 (s, 3H), 2.07 (br s, 1H), 1.81 – 1.71 (m, 2H), 1.58 – 1.47 (m, 2H), 1.47 – 1.40 (m, 2H), 1.40 – 1.28 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 51.8, 45.0, 36.3, 32.4, 29.5, 26.7, 26.5. HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_7\text{H}_{17}\text{ClN}$, 150.1045; found 150.1044.

N-methyl octylamine

General procedure A. 51 mg (clear oil, 89%). ^1H NMR (400 MHz, CDCl_3) δ 2.54 (t, $J=7.2$ Hz, 2H), 2.41 (s, 3H), 1.62 – 1.53 (br s, 1H), 1.52 – 1.41 (m, 2H), 1.26 (m, 10H), 0.86 (t, $J=6.7$ Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 52.1, 36.4, 31.8, 29.8, 29.5, 29.2, 27.3, 22.6, 14.0.³⁰

N-methyl(3-trimethylsilyl)propylamine

General procedure A. 48 mg (yellow oil, 83%). ^1H NMR (500 MHz, CDCl_3) δ 2.54 (t, $J=7.1$ Hz, 2H), 2.41 (s, 3H), 2.13 (br s, 1H), 1.52 – 1.37 (m, 2H), 0.49 – 0.42 (m, 2H), -0.04 (s, 9H). ^{13}C NMR (75 MHz, CDCl_3) δ 55.5, 36.5, 24.3, 14.3, -1.5

N-methyl(3-phenyl)propylamine

General procedure A. 50 mg (yellow oil, 83%). ^1H NMR (400 MHz, CDCl_3) δ 7.37 – 7.29 (m, 2H), 7.28 – 7.20 (m, 3H), 2.78 – 2.60 (m, 4H), 2.49 (s, 3H), 1.88 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 142.1, 128.3, 128.3, 125.7, 51.5, 36.3, 33.6, 31.4.³¹

N-(2-cyclohexyl)ethyl methylamine

General procedure A. 50 mg (yellow oil, 89%). ^1H NMR (400 MHz, CDCl_3) δ 2.62 – 2.49 (m, 2H), 2.40 (s, 3H), 1.78 – 1.52 (m, 4H), 1.35 (m, 2H), 1.28 – 1.00 (m, 5H), 0.88 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 49.7, 37.5, 36.5, 35.6, 33.4, 26.6, 26.3.³²

N-methyl(2-phenyl)propylamine

General procedure A. 41 mg (yellow oil, 69%). ^1H NMR (500 MHz, CDCl_3) δ 7.31 (t, $J=7.6$ Hz, 1H), 7.25 – 7.17 (m, 3H), 2.96 (tq, $J=7.0$ Hz, 7.0 Hz, 1H), 2.74 (d, $J=7.2$ Hz, 2H), 2.38 (s, 3H), 1.87 – 1.42 (m, 1H), 1.26 (d, $J=7.0$ Hz, 4H). ^{13}C NMR (75 MHz, CDCl_3) δ 145.2, 128.5, 127.1, 126.3, 59.2, 39.8, 36.3, 20.0.³¹

1-octylamine

General procedure B. 48 mg (clear oil, 92%). ^1H NMR (500 MHz, CDCl_3) δ 2.65 (t, $J=6.8$ Hz, 2H), 1.69 (br s, 2H), 1.50 – 1.34 (m, 2H), 1.34 – 1.14 (m, 12H), 0.85 (t, $J=6.8$ Hz, 3H). ^{13}C NMR (75 MHz, CDCl_3) δ 42.1, 33.7, 31.8, 29.4, 29.3, 26.8, 22.6, 14.1.³³

(2-phenyl)propylamine

General procedure B. 48 mg (yellow, 88%). ^1H NMR (600 MHz, CDCl_3) δ 7.31 (t, $J=7.4$ Hz, 2H), 7.24 – 7.17 (m, 3H), 2.90 – 2.80 (m, 2H), 2.79 – 2.70 (m, 1H), 1.56 (br s, 2H), 1.25 (d, $J=6.9$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 144.9, 128.5, 127.3, 126.3, 49.4, 43.4, 19.2.³⁴

N-methyl-exo-2-norbornylamine

General procedure A. 41 mg (yellow, 81%). ^1H NMR (500 MHz, CDCl_3) δ 2.44 (dd, $J=7.4$, 3.5 Hz, 1H), 2.36 (s, 3H), 2.16 (d, $J=3.9$ Hz, 1H), 1.55 (ddd, $J=12.4$, 7.5, 2.3 Hz, 1H), 1.50 – 1.36 (m, 3H), 1.13 – 1.00 (m, 4H), 1.23 (br s, 1H), 1.04 (d, $J=2.1$ Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 63.8, 40.2, 39.6, 35.6, 34.8, 34.3, 28.5, 26.9.³⁵

3,4-dimethoxy-N-methylphenethylamine

General procedure A. 55 mg (yellow oil, 71%). ^1H NMR (400 MHz, CDCl_3) δ 6.86 – 6.59 (m, 3H), 3.86 (s, 3H), 3.84 (s, 3H), 2.81 (t, $J=6$ Hz, 2H), 2.75 (t, $J=6$ Hz, 1H), 2.43 (s, 1H), 2.10 (br s, 1H). ^{13}C NMR (151 MHz, CDCl_3) δ 148.84, 148.82, 147.4, 132.3, 120.5, 111.86, 111.85, 111.84, 111.2, 111.2, 55.9, 55.8, 53.1, 36.1, 35.5.³⁶

3-trifluoromethyl-N-methylphenethylamine

General procedure A. 63 mg (yellow oil, 78%). ^1H NMR (600 MHz, CDCl_3) δ 7.62 – 7.28 (m, 4H), 2.97 – 2.84 (m, 4H), 2.75 (br s, 1H), 2.46 (s, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 140.6, 132.1, 130.8 (q, $J=32.0$ Hz), 128.9, 125.3 (q, $J=3.7$ Hz), 124.1 (q, $J=272.1$ Hz), 123.1 (q, $J=7.1$, 3.3 Hz), 52.6, 35.9, 35.6. ^{19}F NMR (376 MHz, CDCl_3) δ -61.8.

(2-cyclohexyl)ethylamine

General procedure B. 36 mg (yellow oil, 70%). ^1H NMR (500 MHz, CDCl_3) δ 2.69 (t, $J=7.0$ Hz, 2H), 1.86 (br s, 2H), 1.74 – 1.56 (m, 5H), 1.43 – 1.29 (m, 3H), 1.27 – 1.01 (m, 3H), 0.99 – 0.76 (m, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 41.4, 39.6, 35.2, 33.3, 26.5, 26.3.³⁷

N,2-dimethyloctan-1-amine

General procedure C. 45 mg (yellow oil, 72%). ^1H NMR (300 MHz, CDCl_3) δ 2.47 (dd, $J=11.5$, 5.9 Hz, 1H), 2.41 (s, 3H), 2.33 (dd, $J=11.5$, 7.3 Hz, 1H), 1.58 (m, 1H), 1.48 (br s, 1H), 1.40 – 1.15 (m, 9H), 1.15 – 0.99 (m, 1H), 0.97 – 0.77 (m, 6H). ^{13}C NMR (126 MHz, CDCl_3) δ 58.8, 36.7, 35.0, 33.0, 31.9, 29.6, 26.9, 22.6, 18.1, 14.1. HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{10}\text{H}_{24}\text{N}$, 158.1903; found 158.1902.

N-heptyloctan-1-amine

General procedure C, heating to 80 °C. 74 mg (yellow oil, 81%). ^1H NMR (500 MHz, CDCl_3) δ 2.78 – 2.42 (m, 4H), 1.52 – 1.34 (m, 4H), 1.34 – 1.00 (m, 19H), 0.86 (t, $J=6.8$ Hz, 6H). ^{13}C NMR (75 MHz, CDCl_3) δ 50.24 (2), 32.07 (2), 30.17 (2), 29.8, 29.52, 29.48, 27.7, 27.6, 22.90, 22.86, 14.3 (2).

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


We thank the NIH-NIGMS (GM-055382) for support of this work.

References

1. Muller TE, Hultzsich KC, Yus M, Foubelo F, Tada M. *Chem Rev.* 2008; 108:3795–3892. [PubMed: 18729420]
2. Muller TE, Beller M. *Chem Rev.* 1998; 98:675–703. [PubMed: 11848912]
3. Brown HC, Heydkamp WR, Breuer E, Murphy WS. *J Am Chem Soc.* 1964; 86:3565–3566.
4. Barker TJ, Jarvo ER. *Synthesis.* 2011; 24:3954–3964.
5. Jigajinni VB, Pelter A, Smith K. *Tetrahedron Lett.* 1978; 19:181–182.
6. Kabalka GW, Henderson DA, Varma RS. *Organometallics.* 1987; 6:1369–1370.
7. Kabalka GW, Wang Z, Goudgaon NH. *Synthetic Comm.* 1989; 19:2409–2414.
8. Genêt JP, Hajicek J, Bischoff L, Greck C. *Tetrahedron Lett.* 1992; 33:2677–2680.
9. Matsuda N, Hirano K, Satoh T, Miura M. *Angew Chem Int Ed.* 2012; 51:3642–3645.
10. Rucker RP, Whittaker AM, Dang H, Lalic G. *J Am Chem Soc.* 2012; 134:6571–6574. [PubMed: 22469028]
11. Rucker RP, Whittaker AM, Dang H, Lalic G. *Angew Chem Int Ed.* 2012; 51:3953–3956.
12. Zhu C, Li G, Ess DH, Falck JR, Kürti L. *J Am Chem Soc.* 2012; 134:18253–18256. [PubMed: 23082853]
13. Mlynarski SN, Karns AS, Morken JP. *J Am Chem Soc.* 2012; 134:16449–16451. [PubMed: 23002712]
14. Suzuki A, Sono S, Itoh M, Brown HC, Midland MM. *J Am Chem Soc.* 1971; 93:4329–4330.
15. Brown HC, Midland MM, Levy AB. *J Am Chem Soc.* 1972; 94:2114–2115.
16. Brown HC, Midland MM, Levy AB. *J Am Chem Soc.* 1973; 95:2394–2396.
17. Akimoto I, Suzuki A. *Synthetic Comm.* 1981; 11:475–480.
18. Kabalka GW, Wang Z. *Synthetic Comm.* 1990; 20:231–237.
19. Zheng B, Srebniak M. *J Org Chem.* 1995; 60:1912–1913.
20. Mendiola J, Rincón JA, Mateos C, Soriano JF, de Frutos Os, Niemeier JK, Davis EM. *Org Process Res Dev.* 2009; 13:263–267.
21. Stanley LM, Hartwig JF. *Angew Chem Int Ed.* 2009; 48:7841–7844.
22. See Supporting Information for details
23. Bexrud JA, Eisenberger P, Leitch DC, Payne PR, Schafer LL. *J Am Chem Soc.* 2009; 131:2116–2118. [PubMed: 19173650]
24. Li H, Wang Y, Tang L, Deng L. *J Am Chem Soc.* 2004; 126:9906–9907. [PubMed: 15303849]
25. Luo SY, Tripathi A, Zulueta MML, Hung SC. *Carbohydrate Research*; Elsevier Ltd. 2012; 352:197–201.
26. Plastic pipette tips affixed to an adjustable volume pipette were used to transfer chlorosulfonic acid
27. Biraboneye AC, Madonna S, Maher P, Kraus JL. *ChemMedChem.* 2010; 5:79–85. [PubMed: 19943277]
28. Care was taken not to exceed 1 equivalent of Schwartz reagent
29. Allowing the hydrozirconation reaction to stir overnight before addition of aminating reagent resulted in reduced yields (61% for amination of 1-octene). Consumption of Schwartz reagent (insoluble in THF) was monitored visually
30. Stearns JF, Rapoport H. *J Org Chem.* 1977; 42:3609–3614.
31. Hanada S, Ishida T, Motoyama Y, Nagashima H. *J Org Chem.* 2007; 72:7551–7559. [PubMed: 17760460]
32. Furstoss R, Tadayoni R, Esposito G, Lacrampe J, Heumann A, Waegell B. *Can J Chem.* 1976; 54:3569–3577.
33. Card RJ, Schmitt JL. *J Org Chem.* 1981; 46:754–757.
34. Bocchinfuso R, Robinson JB. *Eur J Med Chem.* 1999; 34:293–300.
35. Pienemann T, Schaefer HJ. *Synthesis.* 1987; 11:1005–1007.
36. Theodore LJ, Nelson WL. *J Org Chem.* 1987; 52:1309–1315.
37. Kukula P, Koprivova K. *J Catal.* 2005; 234:161–171.

**Scheme 1.**

Anti-Markovnikov amination of alkenes (yields based on 1 equiv of alkene).



Scheme 2. Hydroamination of complex substrates

Table 1
Reactions to form *N*-methylamines

entry ^a	alkene	product	yield ^b
1			89
2			92 ^c
3			79 ^d
4			83
5			89
6			84
7			67
8			61 ^e
9			53
10			69
11			67
12			71 ^f

entry ^a	alkene	product	yield ^b
13			80
14			72 ^e

^a conditions: alkene (0.4 mmol), THF (1.5 mL), Cp₂ZrHCl (0.4 mmol), MeNHOSO₃H (0.6 mmol).

^b Isolated yield (%) without column chromatography, average of two runs.

^c 4 mmol scale.

^d 10 mmol scale.

^e isolated yield (%) after column chromatography.

^f 1.1 equiv alkene, yield based on Cp₂ZrHCl.

Table 2

Reactions to form primary amines.

entry ^a	alkene	product	yield ^b
1			92
2			94
3			71
4			88 ^c
5		86 : 14 linear : branched 	78

^a conditions: alkene (0.4 mmol), THF (1.5 mL), Cp₂ZrHCl (0.4 mmol); H₂NOSO₃H (0.6 mmol).

^b isolated yield (%) without column chromatography.

^c combined yield of linear and branched products.

Table 3

Reactions to form secondary amines.

entry ^a	aminating reagent	product	yield ^b
1			46
2			72 ^c
3			79 ^c
4			44
5			<5
6			-

^a conditions: alkene (0.2 mmol), THF (0.75 mL), Cp₂ZrHCl (0.2 mmol); aminating reagent (0.3 mmol), 50 °C, 30 min.

^b yield determined by NMR spectroscopy with CH₂Br₂ as an internal standard.

^c reaction performed at 80 °C, isolated yield (%) after column chromatography, average of two runs.