

Review Article

Listeriosis during Pregnancy: A Public Health Concern

Teresa Mateus,^{1,2} Joana Silva,¹ Rui L. Maia,³ and Paula Teixeira¹

¹ Centro de Biotecnologia e Química Fina (CBQF) Laboratório Associado, Escola Superior de Biotecnologia, Universidade Católica Portuguesa/Porto, Rua Dr. António Bernardino Almeida, 4200-072 Porto, Portugal

² Departamento de Medicina Veterinária (EUVG), Escola Universitária Vasco da Gama, Coimbra, Portugal

³ Faculdade de Ciências Humanas e Sociais—Universidade Fernando Pessoa, Porto, Portugal

Correspondence should be addressed to Paula Teixeira; pcteixeira@porto.ucp.pt

Received 2 July 2013; Accepted 29 August 2013

Academic Editors: A. Canellada, A. Malek, and C. M. Peterson

Copyright © 2013 Teresa Mateus et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Listeria was first described in 1926 by Murray, Webb, and Swann, who discovered it while investigating an epidemic infection among laboratory rabbits and guinea pigs. The role of *Listeria monocytogenes* as a foodborne pathogen was definitively recognized during the 1980s. This recognition was the consequence of a number of epidemic human outbreaks due to the consumption of contaminated foods, in Canada, in the USA and in Europe. Listeriosis is especially severe in immunocompromised individuals such as pregnant women. The disease has a low incidence of infection, although this is undeniably increasing, with a high fatality rate amongst those infected. In pregnant women listeriosis may cause abortion, fetal death, or neonatal morbidity in the form of septicemia and meningitis. Improved education concerning the disease, its transmission, and prevention measures for immunocompromised individuals and pregnant women has been identified as a pressing need.

1. Introduction

Listeria monocytogenes is a Gram positive, facultative intracellular, foodborne pathogen responsible for cases and outbreaks of listeriosis. *Listeria* was first described in 1926 by Murray et al. who discovered it while investigating an epidemic infection among laboratory rabbits and guinea pigs [1]. At that time, it was given the name *Bacterium monocytogenes* because infection in the animals was characterized by monocytosis. The following year, Pirie isolated an identical bacterium from the liver of several gerbils and proposed the name *Listerella* for the genus in honor of Lord Lister a prominent surgeon of the time [2]. Despite considerable confusion in the nomenclature of the pathogen until 1940, the official name *Listeria monocytogenes* was adopted in the Sixth Edition of Bergey's Manual of Determinative Bacteriology [3].

The first cases of human listeriosis were reported by Nyfeldt in 1929 [4]. The increased number of reported cases during the 1980s in several countries, and the evidence of foodborne transmission, turned listeriosis into a recognized foodborne disease [5, 6]. Listeriosis has emerged as an atypical foodborne illness of major public health concern because

of the severity of the disease (meningitis, septicemia, and abortion), the high case fatality rate (20–30% of cases), the long incubation period, and the predilection for individuals who have an underlying condition which leads to impairment of T-cell-mediated immunity [7]. In addition, the ubiquitous nature of *L. monocytogenes* and its ability to survive and grow in harsh conditions (wide pH range, high salt concentration, refrigeration temperatures) makes this pathogen of high concern to the food industry.

It has been reported that the annual incidence of listeriosis ranges between 1 and 10 per million population [8]. Although listeriosis is less common than other foodborne diseases, 19% and 17% of the known causes of foodborne disease-related deaths occurring in the USA and France, respectively, are caused by *L. monocytogenes* [9, 10].

Most authors comment that 99% of the human *Listeria* infections are from foodborne origin [11–14]. *Listeria monocytogenes* has been recovered from several foods such as meat, milk, and fish products; ice cream; vegetables; and several ready-to-eat foods [15–17]. There are recent surveys of *L. monocytogenes* in foods that demonstrate the presence of this bacterium in several of these foods and ready-to-eat

products [12, 17–23]. Also, there have been several listeriosis outbreaks worldwide [24–28], and certain foods have been described as “high risk” for listeriosis. In recent years the annual number of reported listeriosis cases has increased in several European countries [29–33]. Goulet et al. [30] and Shetty et al. [6] suggest that this increase is particularly due to the increasing population aged more than 60 years, or less than 60 years, although with an immunocompromised predisposing condition.

Listeria monocytogenes is responsible for cases and outbreaks of febrile gastrointestinal disease in otherwise healthy people and invasive listeriosis, which more usually affects pregnant, newborns, the elderly, and immunocompromised individuals. According to Warriner and Namvar [25] non-invasive listeriosis is associated with a huge intake of *L. monocytogenes*.

Listeria monocytogenes has been differentiated into 13 serotypes; serotypes 1/2a, 1/2b, and 4b have been involved in the majority of reported human listeriosis cases [33].

Subtype characterization of *L. monocytogenes* isolates from listeriosis outbreaks has suggested that many outbreaks were caused by a small number of *L. monocytogenes* epidemic clones [34], that is, by a closely related group of isolates that evolved clonally. Epidemic clones of *L. monocytogenes* have been implicated in several outbreaks and sporadic cases of listeriosis worldwide [35–37], and for this reason, although not yet proven, they have been considered more virulent than other strains.

2. High Risk Foods for *Listeria* Infection

High-risk foods have all of the following properties: (1) have the potential for contamination with *L. monocytogenes* (2) support the growth of *L. monocytogenes* to high numbers, (3) do not require further cooking at home (4) require refrigeration, (5) have an extended shelf life, and (6) are contaminated with high levels of the pathogen at the time of consumption [13, 38, 39]. Ready-to-eat seafood such as smoked fish or mussels, raw seafood including sushi and sashimi, premixed raw vegetables including coleslaw, meat products such as sausages, pâté, and rillettes, hot dogs, ham, salami, chicken wraps and deli turkey breast, unpasteurized milk, soft-serve ice creams, and soft cheeses are examples of foods that have been associated with transmission of listeriosis.

The minimum number of cells capable of causing disease is not known although epidemiological data suggest that it is high (ca. 10^6 CFU/g) [40]; the infectious dose depends on the strain virulence and the host susceptibility [40, 41].

The characteristics of the disease (the hugely variable and potentially very long incubation periods, the low attack rates, and the rarity of identification of specific food vehicles), provide limited data for calculation of dose responses [42].

3. Mechanisms of Infection

Listeria monocytogenes has the ability to induce its own entry into host cells, usually mammalian cells (human and animal), such as macrophages, epithelial cells, and endothelial cells

of the gastrointestinal tract [43]. After ingestion, those cells surviving the low pH in the stomach pass through to the small intestine—the first site where invasion occurs—disseminate from the mesenteric lymph nodes to the spleen and the liver and from there *L. monocytogenes* can reach the brain or the placenta [44], causing, respectively, infections of the central nervous system (CNS) mostly in immunocompromised patients and intrauterine/cervical infections in pregnant women [40].

Listeria monocytogenes uses various proteins, including some internalins to adhere and to invade the host cells. Once in the intracellular phagocytic vacuole, bacteria secrete listeriolysins and phospholipases that allow it to lyse the vacuolar membrane and avoid intracellular killing. Upon being released into the cytoplasm, *L. monocytogenes* can multiply and induce the formation of actin filaments which will allow it to move in the cytoplasm until it reaches the plasma membrane. Subsequently, adjacent cells are invaded through plasma membrane protrusions and cell-to-cell spread. Through this cycle *L. monocytogenes* can move from one host cell to another cell, without being in the extracellular environment, thus escaping from the human T-cell immune system, and invading other tissues and organs [45, 46].

4. Incubation Period for Listeriosis

Unlike other foodborne diseases, the incubation period for listeriosis can be long. Goulet et al. [47] recently reported an overall median incubation period of invasive listeriosis of 8 days, ranging from 1 to 67 days. For pregnancy-associated cases the authors reported a longer incubation period (median 27.5 days, ranging from 17 to 67 days) than for CNS cases (median 9 days, ranging from 1 to 14 days) and for bacteraemia cases (median 2 days, ranging 1 to 12 days). For febrile gastrointestinal disease, the median incubation period was reported as 24 hours, ranging from 6 hours to 10 days [47].

5. Individuals at Higher Risk of Listeriosis

Although exposure and colonization may occur in any person, those patients without predisposing factors represent less than 20% of the cases [48]. Silk et al. [49] reported an incidence rate of listeriosis of 0.27 cases per 100,000 general US population. There are well-defined risk groups for human invasive listeriosis: pregnant women and their fetuses or neonates, the elderly, and cellular immunocompromised individuals such as those who are receiving corticosteroids or chemotherapy, hemodialysis, transplants, diabetic, HIV carriers, and drug dependents [40, 50]. Ogunmodede et al. [51] found that pregnant women have an infection risk twenty times higher than other healthy adults. Incidence rates among pregnant women and adults aged more than 65 years were reported in the USA as 3.42 and 1.21 cases per 100,000 population, respectively [49]. Pregnant Hispanic women are also recognized as having a much higher infection risk than pregnant women of other ethnicities [52, 53], probably due to the habit of the consuming soft cheese [13, 54]. In a recent study by Pouillot et al. [53] it was demonstrated

that a significant increase in listeriosis incidence begins among persons as young as 45 years and that incidence rates subsequently increase steadily with age.

6. Invasive Listeriosis

In adults, the most common clinical form of listeriosis is meningitis, due to the bacterial tropism to the central nervous system [44, 55]. Salamano et al. [54] reported that meningitis caused by *Listeria* represents 5% to 11% of all acute bacterial meningitis in adults, reaching 14% in patients aged more than 45 years. Apparently, individuals with predisposing factors seem to suffer more from septicemia rather than meningitis, but this situation may be due to the followup of these individuals by the health system, making them more available for blood collection and culture in the case of a febrile episode [41]. A survey of 782 listeriosis cases reported in 20 countries, showed that 43% of the infections were related with pregnancy, 29% with septicemia, 24% with central nervous system infection, and 4% were atypical forms of the infection [56]. *Listeria* infections may produce sequelae, but the incidence is rarely determined. Up to 11% of neonates and 30% of patients that survive central nervous system infections, suffer from residual symptoms and psychiatric sequelae [52, 56, 57]. Salamano et al. [54] reported a high level of recurrences (7%).

6.1. Listeriosis during Pregnancy. During pregnancy, cellular immunity is minimal due to the increased progesterone [58, 59], making pregnant women particularly susceptible to intracellular microorganisms like *L. monocytogenes* [51, 60]. The vertical cell-to-cell transmission is frequent since *L. monocytogenes* shows uterus and placenta tropism [48, 51]. Immunological aspects of *Listeria* infection during pregnancy were reviewed by Poulsen and Czuprynski [61].

Goulet et al. [30] reported that the incidence in pregnant women and neonates appears to be decreasing in Europe. On the contrary Sisó et al. [62] reported an increase in the incidence of listeriosis between 2000 and 2010 in a tertiary referral hospital in Barcelona. It is important to point out that listeriosis is not often diagnosed in pregnancy, due to the microbiological diagnostic techniques being difficult, and also due to the histological alterations in the placenta being similar to other diseases, hindering, therefore, a definitive assessment of the importance of listeriosis in pregnancy health [63]. Moreover, pregnant women may be asymptomatic or present nonspecific clinical symptoms (e.g., flu-like symptoms, backache, headache, vomiting/diarrhea, muscle pains, and sore throat) [64–66]. Gupta et al. [67] stated that microbiological examinations are not usually made on spontaneous or premature abortions; thus, it is difficult to infer the actual incidence of listeriosis [68].

Infection by *L. monocytogenes* during pregnancy may result in serious outcomes including miscarriage, stillbirth, chorioamnionitis, preterm delivery, and maternal and neonatal sepsis [65, 66].

It is agreed that there is an increased susceptibility for listeriosis in late pregnancy. A third of the listeriosis cases affect pregnant women mostly in the third trimester

of pregnancy [48, 58, 59], rarely in the second and only exceptionally occurs in the first trimester [63]. Studies by Sisó et al. [62] reported an incidence of the infection two times higher after 28 weeks of gestation than in the first trimesters. Listeriosis during early gestations generally has a poorer prognosis for fetuses as opposed to later gestations and commonly results in miscarriage or stillbirth [62, 65, 67]. DiMaio [48] commented that iron supplement, commonly prescribed during pregnancy, favors the development of listeriosis.

A transplacental fetal infection may result in abortion, as already stated, premature birth, or an infected neonate [54, 68]. Laciari et al. [68] Doyle [58] and Schwab and Edelweiss [69] agree that most cases of the intrauterine infection are a consequence of the transplacental dissemination after bacteremia in pregnant women and less frequently due to an ascending contamination from the pregnant woman's lower genital tract colonized by *L. monocytogenes*, as the vagina and cervix are also potentially carrier sites of this microorganism [48]. More commonly, infection during the passage of the fetus through the birth canal may result in the nosocomial transmission in the neonatal nursery [70]. Two distinct infection forms are described in the newborns.

6.1.1. Early-Onset Neonatal Listeriosis. This is acquired *in utero*, by transplacental transmission, and is also known as granulomatosis infantiseptica. Disease in pregnancy precedes the fetal infection, although symptoms may not be specific [48, 70]. This infection, generally occurring at a mean of 36 h after birth [71], and probably due to aspiration of infected amniotic liquid, is characterized by clinical features like septicemia (81–88%), respiratory distress or pneumonia (38%), and meningitis (24%) [72, 73]. The formation and dissemination of abscesses and granulomas in multiple organs may occur [55]. The mortality rate for infants born alive approaches 20%, and the frequency of abortion and stillbirth increases the overall mortality rate to more than 50% [74].

6.1.2. Late-Onset Neonatal Listeriosis. Infection occurs particularly during delivery, by contamination, with symptoms such as meningitis or meningoencephalitis together with septicemia, occurring 2 or 3 weeks after delivery [70]. In these cases deliveries are not complicated and the transmission may be in the passage through the birth canal; the more frequent occurrence during deliveries by caesarean section being relevant, suggesting nosocomial transmission [48, 52].

The mortality rate associated with late-onset disease is ca. 10% [69], but as commented by DeWaal et al. [75] and Buzby [76] a high rate of surviving babies develop severe and chronic neurological complications, for example, delayed mental development and blindness.

6.2. Diagnosis of Listeriosis. The diagnosis of listeriosis depends upon detecting the growth of the microorganisms in corporal fluids normally considered as sterile—blood, amniotic, and cerebrospinal fluids [48, 57]. A rapid diagnosis would be important for preventing recurrences in pregnancies and to treat neonatal infections that may cause meningitis with sequelae such as hydrocephaly, delayed mental

development, convulsions, and other neurological symptoms [58, 75, 76].

The diagnosis of listeriosis in pregnancy and early treatment would be possible if all the febrile episodes during pregnancy were evaluated through a blood culture [67, 68]. For this purpose, pregnant women should report any disease symptom, because listeriosis can be apparently harmless for the pregnant woman but severely infect the fetus [52, 75, 77].

6.3. Treatment of Listeriosis. Listeriosis often requires antimicrobial therapy. The treatment of choice consists of a β -lactam antibiotic, normally ampicillin, alone or in combination with an aminoglycoside, classically gentamicin. Second-line agents in case of allergy to β -lactams or in certain disease states include trimethoprim/sulfamethoxazole, erythromycin, vancomycin, and the fluoroquinolones [78]. Resistance of clinical isolates of *L. monocytogenes* to these antibiotics is low [79]. Early diagnosis and antibiotic administration increases the probability of a favorable outcome [62, 78].

7. How to Reduce the Incidence of Listeriosis

Due to the high mortality rate of listeriosis, it is necessary to reduce its incidence. For that purpose, it is urgent to identify the most efficacious strategies based on risk assessment from food production to consumption [13, 48].

According to the International Life Sciences Institute [13], the use of scientifically-based messages, to high-risk groups and in health professionals, education is one of the main strategies to adopt concerning the reduction of listeriosis incidence, because it is generally accepted that in some foods, *L. monocytogenes* absence may not be possible. These strategies should minimize food contamination and educate the population in general and risk groups in particular, as well as health professionals and care providers [13, 41]. Cates et al. [80] concluded from their study about listeriosis information transmission to pregnant women that if we want the message to be well delivered to those at risk, it should be targeted and appealing to the pregnant woman; provide detailed information about sources, prevention measures, morbidity, and mortality rates (but as simple as “1 in 4”, rather than 25%); give emphasis to words like “abortion”, “premature birth”; and be transmitted by the obstetrician or other doctor. “Preventative” public alerts about concerning listeriosis prevention are already available (e.g., <http://www.cdc.gov/pregnancy/infections-listeria.html>; <http://ideas.health.vic.gov.au/diseases/listeria-facts.asp>; <http://americanpregnancy.org/pregnancycomplications/listeria.html>; <http://www.listeriosisprevention.com/facts.html>; <http://www.health.state.mn.us/divs/idepc/diseases/listeriosis/index.html>). Wong et al. [81] commented that in the USA, doctors are considered by the population as the most credible source of information about health. Doctors are important food safety educators as during the year, they are in contact with 80% of the population, and also because people are more likely to change behaviors if they have been recently ill. Hence, campaigns about food safety education targeted for doctors are recommended [81], especially for obstetricians, as

pregnant women will not change their attitudes if they do not recognize the source of the information or if they do not believe it [82].

Acknowledgments

This work was supported by the National Funds from FCT—Fundação para a Ciência e a Tecnologia through Projects PTDC/AGR-ALI/64662/2006 and PEst-OE/EQB/LA0016/2011.

References

- [1] E. G. D. Murray, R. A. Webb, and M. B. R. Swann, “A disease of rabbits characterised by a large mononuclear leucocytosis, caused by a hitherto undescribed bacillus *Bacterium monocytogenes* (n.Sp.),” *Journal of Pathology and Bacteriology*, vol. 29, no. 4, pp. 407–439, 1926.
- [2] J. H. H. Pirie, “A new disease of veld rodents ‘tiger river disease,’” *Publication of the South African Institute of Medical Research*, vol. 3, pp. 163–186, 1927.
- [3] M. L. Gray and A. H. Killinger, “*Listeria monocytogenes* and listeric infections,” *Bacteriological Reviews*, vol. 30, no. 2, pp. 309–382, 1966.
- [4] A. Nyfeldt, “Etiologie de la mononucleose infectieuse,” *Comptes Rendus des Seances de la Societe de Biologie*, vol. 101, pp. 590–591, 1929.
- [5] F. Ueda, R. Anahara, F. Yamada, M. Mochizuki, Y. Ochiai, and R. Hondo, “Discrimination of *Listeria monocytogenes* contaminated commercial Japanese meats,” *International Journal of Food Microbiology*, vol. 105, no. 3, pp. 455–462, 2005.
- [6] A. Shetty, J. McLauchlin, K. Grant, D. O’Brien, T. Howard, and E. M. Davies, “Outbreak of *Listeria monocytogenes* in an oncology unit associated with sandwiches consumed in hospital,” *Journal of Hospital Infection*, vol. 72, no. 4, pp. 332–336, 2009.
- [7] F. Allerberger, “*Listeria*: growth, phenotypic differentiation and molecular microbiology,” *FEMS Immunology and Medical Microbiology*, vol. 35, no. 3, pp. 183–189, 2003.
- [8] V. Goulet, M. Hebert, C. Hedberg et al., “Incidence of listeriosis and related mortality among groups at risk of acquiring listeriosis,” *Clinical Infectious Diseases*, vol. 54, no. 5, pp. 652–660, 2012.
- [9] V. Vaillant, H. de Valk, E. Baron et al., “Foodborne infections in France,” *Foodborne Pathogens and Disease*, vol. 2, no. 3, pp. 221–232, 2005.
- [10] E. Scallan, R. M. Hoekstra, F. J. Angulo et al., “Foodborne illness acquired in the United States—major pathogens,” *Emerging Infectious Diseases*, vol. 17, no. 1, pp. 7–15, 2011.
- [11] M. Wiedmann, “An integrated science-based approach to dairy food safety: *Listeria monocytogenes* as a model system,” *Journal of Dairy Science*, vol. 86, no. 6, pp. 1865–1875, 2003.
- [12] C. Mena, G. Almeida, L. Carneiro, P. Teixeira, T. Hogg, and P. A. Gibbs, “Incidence of *Listeria monocytogenes* in different food products commercialized in Portugal,” *Food Microbiology*, vol. 21, no. 2, pp. 213–216, 2004.
- [13] ILSI, “Achieving continuous improvement in reductions in foodborne listeriosis—a risk-based approach,” *Journal of Food Protection*, vol. 68, no. 9, pp. 1932–1994, 2005.
- [14] E. Hofer, C. M. F. Reis, and C. B. Hofer, “Sorovares de *Listeria monocytogenes* e espécies relacionadas, isoladas de material

- clínico humano,” *Revista da Sociedade Brasileira de Medicina Tropical*, vol. 39, pp. 32–37, 2006.
- [15] P. C. C. Araújo, R. M. Franco, L. A. T. Oliveira, and J. C. A. P. Carvalho, “Ocorrência de *Listeria monocytogenes* em produtos de carne de peru comercializados na cidade de Niterói,” *Acta Scientiae Veterinariae*, vol. 30, pp. 19–25, 2002.
- [16] K. Kwiatek, “Occurrence of *Listeria monocytogenes* in selected food of animal origin,” *Bulletin of the Veterinary Institute in Pulawy*, vol. 48, pp. 269–272, 2004.
- [17] V. Garrido, A. I. Vitas, and I. García-Jalón, “Survey of *Listeria monocytogenes* in ready-to-eat products: prevalence by brands and retail establishments for exposure assessment of listeriosis in Northern Spain,” *Food Control*, vol. 20, no. 11, pp. 986–991, 2009.
- [18] V. Garrido, I. García-Jalón, A. I. Vitas, and M. Sanaa, “Listeriosis risk assessment: simulation modelling and “what if” scenarios applied to consumption of ready-to-eat products in a Spanish population,” *Food Control*, vol. 21, no. 3, pp. 231–239, 2010.
- [19] A. A. Fallah, S. S. Saei-Dehkordi, M. Rahnama, H. Tahmasby, and M. Mahzounieh, “Prevalence and antimicrobial resistance patterns of *Listeria* species isolated from poultry products marketed in Iran,” *Food Control*, vol. 28, no. 2, pp. 327–332, 2012.
- [20] S. T. Lambertz, C. Nilsson, A. Bråndenmark et al., “Prevalence and level of *Listeria monocytogenes* in ready-to-eat foods in Sweden 2010,” *International Journal of Food Microbiology*, vol. 160, no. 1, pp. 24–31, 2012.
- [21] Y. Moreno, J. Sánchez-Contreras, R. M. Montes, J. García-Hernández, L. Ballesteros, and M. A. Ferrús, “Detection and enumeration of viable *Listeria monocytogenes* cells from ready-to-eat and processed vegetable foods by culture and DVC-FISH,” *Food Control*, vol. 27, no. 2, pp. 374–379, 2012.
- [22] H. Jamali, L. C. Chai, and K. L. Thong, “Detection and isolation of *Listeria* spp. and *Listeria monocytogenes* in ready-to-eat foods with various selective culture media,” *Food Control*, vol. 32, no. 1, pp. 19–24, 2013.
- [23] T. Kramarenko, M. Roasto, K. Meremäe, M. Kuningas, P. Pölsama, and T. Elias, “*Listeria monocytogenes* prevalence and serotype diversity in various foods,” *Food Control*, vol. 30, no. 1, pp. 24–29, 2013.
- [24] J. Lundén, R. Tolvanen, and H. Korkeala, “Human listeriosis outbreaks linked to dairy products in Europe,” *Journal of Dairy Science*, vol. 87, no. 1, pp. E6–E11, 2004.
- [25] K. Warriner and A. Namvar, “What is the hysteria with *Listeria*?” *Trends in Food Science and Technology*, vol. 20, no. 6-7, pp. 245–254, 2009.
- [26] C. L. Little, C. F. L. Amar, A. Awofisayo, and K. A. Grant, “Hospital-acquired listeriosis associated with sandwiches in the UK: a cause for concern,” *Journal of Hospital Infection*, vol. 82, no. 1, pp. 13–18, 2012.
- [27] M. Yde, M. Naranjo, W. Mattheus et al., “Usefulness of the European epidemic intelligence information system in the management of an outbreak of listeriosis, Belgium, 2011,” *Eurosurveillance*, vol. 17, no. 38, article 1, 2012.
- [28] E. J. Cartwright, K. A. Jackson, S. D. Johnson, L. M. Graves, B. J. Silk, and B. E. Mahon, “Listeriosis outbreaks and associated food vehicles, United States, 1998–2008,” *Emerging Infectious Diseases*, vol. 19, no. 1, pp. 1–9, 2013.
- [29] J. G. Denny and J. McLauchlin, “Human *Listeria monocytogenes* infections in European opportunity for improved European surveillance,” *Eurosurveillance*, vol. 13, no. 13, article 5, 2008.
- [30] V. Goulet, C. Hedberg, A. Le Monnier, and H. de Valk, “Increasing incidence of listeriosis in France and other European countries,” *Emerging Infectious Diseases*, vol. 14, no. 5, pp. 734–740, 2008.
- [31] A. K. Jensen, S. Ethelberg, B. Smith et al., “Substantial increase in listeriosis, Denmark 2009,” *Eurosurveillance*, vol. 15, no. 12, article 4, 2010.
- [32] F. Allerberger and M. Wagner, “Listeriosis: a resurgent foodborne infection,” *Clinical Microbiology and Infection*, vol. 16, no. 1, pp. 16–23, 2010.
- [33] G. Almeida, A. Morvan, R. Magalhães et al., “Distribution and characterization of *Listeria monocytogenes* clinical isolates in Portugal, 1994–2007,” *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 29, no. 10, pp. 1219–1227, 2010.
- [34] S. Kathariou, “*Listeria monocytogenes* virulence and pathogenicity, a food safety perspective,” *Journal of Food Protection*, vol. 65, no. 11, pp. 1811–1829, 2002.
- [35] V. Chenal-Francisque, J. Lopez, T. Cantinelli et al., “World wide distribution of major clones of *Listeria monocytogenes*,” *Emerging Infectious Diseases*, vol. 17, no. 6, pp. 1110–1112, 2011.
- [36] H. C. den Bakker, E. D. Fortes, and M. Wiedmann, “Multilocus sequence typing of outbreak-associated *Listeria monocytogenes* isolates to identify epidemic clones,” *Foodborne Pathogens and Disease*, vol. 7, no. 3, pp. 257–265, 2010.
- [37] S. J. Knabel, A. Reimer, B. Verghese et al., “Sequence typing confirms that a predominant *Listeria monocytogenes* clone caused human listeriosis cases and outbreaks in Canada from 1988 to 2010,” *Journal of Clinical Microbiology*, vol. 50, no. 5, pp. 1748–1751, 2012.
- [38] FAO/WHO (Food and Agriculture Organisation/World Health Organisation), “Risk assessment of *Listeria monocytogenes* in ready-to-eat foods,” *Microbiological Risk Assessment Series*, vol. 5, pp. 1–98, 2004, <ftp://ftp.fao.org/docrep/fao/010/y5394e/y5394e.pdf>.
- [39] M. M. Guerra and F. A. Bernardo, “O risco da listeriose e a identificação do perigo—Revisão,” *Revista Portuguesa de Ciências Veterinárias*, vol. 99, pp. 69–76, 2004.
- [40] J. A. Vázquez-Boland, M. Kuhn, P. Berche et al., “*Listeria* pathogenesis and molecular virulence determinants,” *Clinical Microbiology Reviews*, vol. 14, no. 3, pp. 584–640, 2001.
- [41] B. Swaminathan and P. Gerner-Smidt, “The epidemiology of human listeriosis,” *Microbes and Infection*, vol. 9, no. 10, pp. 1236–1243, 2007.
- [42] J. McLauchlin, R. T. Mitchell, W. J. Smerdon, and K. Jewell, “*Listeria monocytogenes* and listeriosis: a review of hazard characterisation for use in microbiological risk assessment of foods,” *International Journal of Food Microbiology*, vol. 92, no. 1, pp. 15–33, 2004.
- [43] P. Cossart, “Molecular and cellular basis of the infection by *Listeria monocytogenes*: an overview,” *International Journal of Medical Microbiology*, vol. 291, no. 6-7, pp. 401–409, 2001.
- [44] M. Lecuit, “Human listeriosis and animal models,” *Microbes and Infection*, vol. 9, no. 10, pp. 1216–1225, 2007.
- [45] J. Pizarro-Cerdá and P. Cossart, “Subversion of cellular functions by *Listeria monocytogenes*,” *Journal of Pathology*, vol. 208, no. 2, pp. 215–223, 2006.
- [46] N. E. Freitag, G. C. Port, and M. D. Miner, “*Listeria monocytogenes*—from saprophyte to intracellular pathogen,” *Nature Reviews Microbiology*, vol. 7, no. 9, pp. 623–628, 2009.
- [47] V. Goulet, L. A. King, V. Vaillant, and H. de Valk, “What is the incubation period for listeriosis?” *BMC Infectious Diseases*, vol. 13, article 11, 2013.

- [48] H. DiMaio, "Listeria infection in women," *Primary Care Update for Ob/Gyns*, vol. 7, no. 1, pp. 40–45, 2000.
- [49] B. J. Silk, K. A. Date, K. A. Jackson et al., "Invasive listeriosis in foodborne diseases active surveillance network, 2004–2009: further targeted prevention needed for higher-risk groups," *Clinical Infectious Diseases*, vol. 54, supplement 5, pp. S396–S404, 2012.
- [50] P. Mook, S. J. O'Brien, and I. A. Gillespie, "Concurrent conditions and human listeriosis, England, 1999–2009," *Emerging Infectious Diseases*, vol. 17, no. 1, pp. 38–43, 2011.
- [51] F. Ogunmodede, J. L. Jones, J. Scheffel, E. Kirkland, J. Schulkin, and R. Lynfield, "Listeriosis prevention knowledge among pregnant women in the USA," *Infectious Disease in Obstetrics and Gynecology*, vol. 13, no. 1, pp. 11–15, 2005.
- [52] A. R. Delgado, "Listeriosis in pregnancy," *Journal of Midwifery and Women's Health*, vol. 53, no. 3, pp. 255–259, 2008.
- [53] R. Pouillot, K. Hoelzer, K. A. Jackson, O. L. Henao, and B. J. Silk, "Relative risk of listeriosis in foodborne diseases active surveillance network (FoodNet) sites according to age, pregnancy, and ethnicity," *Clinical Infectious Diseases*, vol. 54, supplement 5, pp. S405–S410, 2012.
- [54] R. Salamano, A. Braselli, A. Hoppe, R. Monteghirfo, and T. Silva, "Neurolisterioses en adultos—seis casos clínicos," *Arquivos de Neuro-Psiquiatria*, vol. 63, no. 4, pp. 1063–1069, 2005.
- [55] E. Hofer, R. S. do Nascimento, and M. A. de Oliveira, "Meningitis due to *Listeria monocytogenes*. Case reports in patients of Distrito Federal, Brazil," *Revista da Sociedade Brasileira de Medicina Tropical*, vol. 31, no. 2, pp. 173–178, 1998.
- [56] FAO/WHO (Food and Agriculture Organisation/World Health Organisation), "Risk assessment of *Listeria monocytogenes* in ready-to-eat foods," *Microbiological Risk Assessment Series*, vol. 4, pp. 1–78, 2004, <http://www.who.int/foodsafety/publications/micro/en/mra4.pdf>.
- [57] A. Platnaris, A. Hatzimichael, S. Ktenidou-Kartali et al., "A case of *Listeria* meningoencephalitis complicated by hydrocephalus in an immunocompetent infant," *European Journal of Pediatrics*, vol. 168, no. 3, pp. 343–346, 2009.
- [58] M. E. Doyle, "Virulence characteristics of *Listeria monocytogenes*," *Fri Briefings*, Food Research Institute, University of Wisconsin, pp. 1–13, 2001.
- [59] B. Scheule, "Food safety education: health professionals' knowledge and assessment of WIC client needs," *Journal of the American Dietetic Association*, vol. 104, no. 5, pp. 799–803, 2004.
- [60] A. Leber, M. L. Zenclussen, A. Teles et al., "Pregnancy: tolerance and suppression of immune responses," *Methods in Molecular Biology*, vol. 677, pp. 397–417, 2011.
- [61] K. P. Poulsen and C. J. Czuprynski, "Pathogenesis of listeriosis during pregnancy," *Animal Health Research Reviews*, vol. 14, no. 1, pp. 30–39, 2013.
- [62] C. Sisó, A. Goncé, J. Bosch, M. D. Salvia, S. Hernández, and F. Figueras, "Listeriosis in pregnancy: a secular trend in a tertiary referral hospital in Barcelona," *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 31, no. 9, pp. 2125–2132, 2012.
- [63] J. P. Schwab and M. I. A. Edelweiss, "Identificação imunohistoquímica de *Listeria monocytogenes* em placentas fixadas em formol e embebidas em parafina," *Revista Brasileira de Ginecologia e Obstetrícia*, vol. 25, pp. 501–505, 2003.
- [64] K. A. Jackson, M. Iwamoto, and D. Swerdlow, "Pregnancy-associated listeriosis," *Epidemiology and Infection*, vol. 138, no. 10, pp. 1503–1509, 2010.
- [65] E. Mylonakis, M. Paliou, E. L. Hohmann, S. B. Calderwood, and E. J. Wing, "Listeriosis during pregnancy: a case series and review of 222 cases," *Medicine*, vol. 81, no. 4, pp. 260–269, 2002.
- [66] R. F. Lamont, J. Sobel, S. Mazaki-Tovi et al., "Listeriosis in human pregnancy: a systematic review," *Journal of Perinatal Medicine*, vol. 39, no. 3, pp. 227–236, 2011.
- [67] V. Gupta, V. Gautam, N. Mehta, I. Kumari, and R. M. Joshi, "Listeriosis in second trimester of pregnancy: case report from India," *Japanese Journal of Infectious Diseases*, vol. 56, no. 2, pp. 60–61, 2003.
- [68] A. L. Laciari, R. P. Hasuoka, S. M. Correa, A. M. Miranda, and O. N. P. Centorbi, "Symptomatic hydrocephalus in a newborn infected with *Listeria monocytogenes*," *Brazilian Journal of Microbiology*, vol. 31, no. 1, pp. 9–11, 2000.
- [69] J. P. Schwab and M. I. A. Edelweiss, "Identificação de *Listeria monocytogenes* em placentas humanas e espécimes de aborto pela técnica de imunohistoquímica," *Jornal Brasileiro de Patologia e Medicina Laboratorial*, vol. 39, pp. 111–114, 2003.
- [70] K. M. Posfay-Barbe and E. R. Wald, "Listeriosis," *Seminars in Fetal and Neonatal Medicine*, vol. 14, no. 4, pp. 228–233, 2009.
- [71] J. P. Relier, "Listeriosis," *Journal of Antimicrobial Chemotherapy*, vol. 5, pp. 51–57, 1979.
- [72] J. McLauchlin, "Human listeriosis in Britain, 1967–85, a summary of 722 cases: 1. Listeriosis during pregnancy and in the newborn," *Epidemiology and Infection*, vol. 104, no. 2, pp. 181–189, 1990.
- [73] M. J. Linnan, L. Mascola, X. Lou et al., "Epidemic listeriosis associated with Mexican-style cheese," *The New England Journal of Medicine*, vol. 319, no. 13, pp. 823–828, 1988.
- [74] W. F. Schlech, "Foodborne listeriosis," *Clinical Infectious Diseases*, vol. 31, no. 3, pp. 770–775, 2000.
- [75] S. DeWaal, D. Birkett, and Z. Enga, "Unexpected consequences: miscarriage and birth defects from tainted food," Center for Science in the Public Interest Washington, DC, USA, 2000, <http://www.cspinet.org/foodsafety/conseqco.pdf>.
- [76] J. C. Buzby, "Children and microbial foodborne illness," *Food Review*, vol. 24, pp. 32–37, 2001.
- [77] R. Bortolussi, "*Listeria monocytogenes* infections in neonates," *Seminars in Pediatric Infectious Diseases*, vol. 10, no. 2, pp. 111–118, 1999.
- [78] M. E. Temple and M. C. Nahata, "Treatment of listeriosis," *Annals of Pharmacotherapy*, vol. 34, no. 5, pp. 656–661, 2000.
- [79] A. Morvan, C. Moubareck, A. Leclercq et al., "Antimicrobial resistance of *Listeria monocytogenes* strains isolated from humans in France," *Antimicrobial Agents and Chemotherapy*, vol. 54, no. 6, pp. 2728–2731, 2010.
- [80] S. C. Cates, H. L. Carter-Young, S. Conley, and B. O'Brien, "Pregnant women and listeriosis: preferred educational messages and delivery mechanisms," *Journal of Nutrition Education and Behavior*, vol. 36, no. 3, pp. 121–127, 2004.
- [81] S. Wong, R. Marcus, M. Hawkins et al., "Physicians as food-safety educators: a practices and perceptions survey," *Clinical Infectious Diseases*, vol. 38, supplement 3, pp. S212–S218, 2004.
- [82] E. C. Redmond and C. J. Griffith, "Consumer food handling in the home: a review of food safety studies," *Journal of Food Protection*, vol. 66, no. 1, pp. 130–161, 2003.