

The Ferric Enterobactin Transporter Fep Is Required for Persistent *Salmonella enterica* Serovar Typhimurium Infection

Toni A. Nagy,^a Sarah M. Moreland,^{a*} Helene Andrews-Polymenis,^b Corrella S. Detweiler^a

Department of Molecular, Cellular and Developmental Biology, University of Colorado—Boulder, Boulder, Colorado, USA^a; Texas A&M University System Health Science Center, College Station, Texas, USA^b

Most bacterial pathogens require iron to grow and colonize host tissues. The Gram-negative bacterium *Salmonella enterica* serovar Typhimurium causes a natural systemic infection of mice that models acute and chronic human typhoid fever. *S. Typhimurium* resides in tissues within cells of the monocyte lineage, which limit pathogen access to iron, a mechanism of nutritional immunity. The primary ferric iron import system encoded by *Salmonella* is the siderophore ABC transporter FepBDGC. The Fep system has a known role in acute infection, but it is unclear whether ferric iron uptake or the ferric iron binding siderophores enterobactin and salmochelin are required for persistent infection. We defined the role of the Fep iron transporter and siderophores in the replication of *Salmonella* in macrophages and in mice that develop acute followed by persistent infections. Replication of wild-type and iron transporter mutant *Salmonella* strains was quantified in cultured macrophages, fecal pellets, and host tissues in mixed- and single-infection experiments. We show that deletion of *fepB* attenuated *Salmonella* replication and colonization within macrophages and mice. Additionally, the genes required to produce and transport enterobactin and salmochelin across the outer membrane receptors, *fepA* and *iroN*, are needed for colonization of all tissues examined. However, salmochelin appears to be more important than enterobactin in the colonization of the spleen and liver, both sites of dissemination. Thus, the FepBDGC ferric iron transporter and the siderophores enterobactin and salmochelin are required by *Salmonella* to evade nutritional immunity in macrophages and cause persistent infection in mice.

In humans, *Salmonella enterica* serovars Typhi and Paratyphi A, B, and C cause typhoid fever, a systemic infection with mortality rates exceeding 15% if left untreated (1, 2). *S. enterica* serovar Typhimurium causes murine typhoid in mice, a disease that models both the acute and chronic stages of human typhoidal infection. Upon ingestion, *Salmonella* bacteria pass through the stomach and traverse the intestinal epithelial barrier to colonize Peyer's patches, mesenteric lymph nodes, spleen, and liver (3, 4). *Salmonella* bacteria reside within cells of the monocyte lineage, typically macrophages that contain the bacteria within a specialized membrane-bound compartment called a *Salmonella*-containing vesicle (5, 6).

Iron is an essential nutrient for animals. However, iron must be tightly regulated, as it can both accept and donate electrons, leading to the generation of damaging free radicals (7). Dietary ferric iron (Fe^{3+}) is reduced to ferrous iron (Fe^{2+}) by a ferrireductase (DcytB) on the luminal side of intestinal epithelial cells (8, 9). Ferrous iron is then taken up by enterocytes via DMT1/Nramp2/Slc11A2, exported into the blood via ferroportin 1, and oxidized to ferric iron by hephaestin on the basolateral membrane (10). Blood ferric iron is captured by transferrin, a high-affinity iron binding protein (K_d [dissociation constant] = 10^{22} M^{-1}) (11). In tissues, cells endocytose transferrin, and ferric iron is released upon endosome acidification. The released ferric iron is then reduced (12) and transported to the cytosol (13). Cytosolic iron is sequestered by ferritin, which accounts for about 16% of iron stores in the body (14).

A major host defense against infection is nutritional immunity, the sequestration of essential molecules, including metals, to prevent pathogen outgrowth (15). Sequestration of iron is an effective antimicrobial defense because iron is a cofactor required for crucial processes, including energy production and DNA replication. For instance, during acute or chronic immune activation, ferritin-

bound iron accumulates in macrophages, a process that withholds iron from extracellular pathogens and is characteristic of the anemia of chronic disease (14, 16). In contrast, macrophages infected with *Salmonella* increase the export of iron, and the spleen and liver of *Salmonella*-infected mice do not accumulate iron (17–19). Decreased iron concentrations in macrophages limit *Salmonella* replication (20, 21), highlighting the importance of delineating how *Salmonella* acquires iron under such starved conditions.

Salmonella acquires ferric iron by secreting the catecholate siderophores enterobactin and salmochelin. Once bound to iron, enterobactin and salmochelin transit the bacterial outer membrane receptors FepA and IroN, respectively, in a TonB-dependent manner (22). Enterobactin and salmochelin then bind to FepB in the periplasm for import through FepDGC (23). During infection, macrophages respond to gamma interferon ($\text{IFN-}\gamma$) by increasing the secretion of lipocalin-2, a siderophore-capturing protein (20, 24–27). Lipocalin-2 binds enterobactin but not salmochelin, a glycosylated derivative of enterobactin that gives *Salmonella* a selective growth advantage in the intestine (26).

Whether *Salmonella* requires access to ferric iron has been examined in mice that lack functional Nramp1/Slc11a1, a ferrous

Received 1 April 2013 Returned for modification 8 May 2013

Accepted 10 August 2013

Published ahead of print 19 August 2013

Editor: S. M. Payne

Address correspondence to Toni A. Nagy, toni.nagy@colorado.edu.

* Present address: Sarah M. Moreland, School of Medicine, Temple University, Philadelphia, Pennsylvania, USA.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/IAI.00412-13

iron transporter in the membrane of the phagosome. Nramp1-deficient mice have dysregulated iron metabolism and are exquisitely sensitive to microbial pathogens that reside within vesicles, including *Salmonella*, *Mycobacterium tuberculosis*, and *Leishmania* species. *Salmonella* strains lacking *fepA* and *iroN* have no apparent phenotype in Nramp1^{-/-} mice in single and mixed infections (23, 28) or in survival assays upon intragastric or intraperitoneal inoculation compared with the wild type (29). Together, these studies indicate that enterobactin and salmochelin uptake via FepA and/or IroN is not essential for *Salmonella* infection of Nramp1^{-/-} mice.

Nramp1^{+/+} mice, such as the Sv129S6 strain, survive acute infection and develop chronic infection. In these animals, the synthesis and secretion of salmochelin are required for *Salmonella* virulence upon intraperitoneal inoculation, as measured by survival assays (26, 30). Intraperitoneal inoculation allows bacteria to bypass the gastrointestinal tract and directly access the spleen (31). However, Bearson et al. demonstrated that upon intranasal inoculation of piglets, there was no colonization defect of a triple mutant lacking *fepA*, *iroN*, and *cirA* compared to the wild-type strain (32). CirA is a third outer membrane receptor for siderophore-mediated iron uptake via FepBDGC (27, 28, 33). However, CirA does not directly bind enterobactin or salmochelin but rather captures catecholate breakdown products containing ferric iron. In the work presented here, we establish the requirement of *Salmonella* for FepB and the enterobactin and salmochelin binding proteins FepA and IroN for gastric and deep-tissue colonization of Nramp1^{+/+} mice upon orogastric inoculation.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *Salmonella enterica* serovar Typhimurium wild-type strain SL1344 (34) and mutant derivatives were grown overnight at 37°C with aeration in Luria-Bertani (LB) medium prior to infections. Antibiotics were used at the following concentrations: streptomycin at 30 µg/ml and kanamycin at 30 µg/ml.

Deletion mutants in genes involved in iron uptake and metabolism were constructed by using lambda Red methodology (35, 36). *ΔiroN*, *ΔfepA*, and *ΔentC* mutants of background strain 14028 were generated previously by using a similar methodology (36), and the *ΔfepB* mutant was generated in this study by using primers *fepB* fwd (5'-GCGCTAACCTAAGAGTAAAACGTCGCTCTGTCAACTGTGTAGGCTGGAGCTGCTTC-3') and *fepB* rev (5'-AATCGGTCTGGTCAGTCGGATAAGACTCGATAAGGCATATGAATATCCTCCTTAG-3'). PCR confirmed the insertion of the cassette by using site-specific and internal kanamycin primers. P22 phage lysates were prepared from each mutant strain and used to transduce each mutation into wild-type strain SL1344.

Restoration of *fepB* was achieved by P22 phage transduction of a wild-type copy into the *fepB*-deleted strain. To enrich for colonies in which *fepB* was restored to the wild type, P22-exposed bacteria were first incubated in M9 minimal medium for 6 h. During this time, the wild-type strain reaches an optical density at 600 nm (OD₆₀₀) of ~0.5, while the *ΔfepB* strain is still in the lag phase. Cells were diluted and plated onto streptomycin-containing LB plates. Colonies were replica plated onto kanamycin-containing LB plates, and those that grew on streptomycin but not kanamycin were tested and confirmed for restoration of a wild-type *fepB* copy by PCR.

Measurement of growth in vitro. Cultures grown overnight were diluted to an OD₆₀₀ of 0.01 in 200 µl of LB or M9 minimal medium (M9) with the vehicle control, ferric chloride (Sigma), or 2,2'-dipyridyl (Sigma). Bacteria were grown in 96-well plates with shaking in a Synergy2 plate reader (BioTek) at 37°C for 17 h. At 20-min intervals, the OD₆₀₀ was recorded. Addition of 25 to 400 µM deferasirox in broth is not practical

because the presence of deferasirox enhances *Salmonella* growth under these conditions for unknown reasons (data not shown).

Murine infection. This study was carried out in strict accordance with the recommendations in the *Guide for the Care and Use of Laboratory Animals* of the National Institutes of Health (37), and all protocols were approved by the University of Colorado Institutional Committees for Biosafety and for Animal Care and Use. For competitive infection studies, 7-week-old male and female 129SvEvTac (Nramp1^{+/+}) mice (Taconic Laboratories) bred in-house were fasted for 2 h prior to orogastric inoculation with a total of 2 × 10⁹ CFU of a 1:1 mixture of streptomycin-resistant wild-type (1 × 10⁹ CFU) and kanamycin-resistant mutant (1 × 10⁹ CFU) bacteria in 100 µl phosphate-buffered saline (PBS), as verified by plating for CFU on selective LB agar. For single-infection experiments, mice were fasted for 2 h and orogastrically inoculated with 1 × 10⁹ bacteria in 100 µl PBS. The infectious dose was verified by plating for CFU on selective LB agar. During the period in which poor grooming may be observed, mice were monitored twice daily. Cages containing “scruffy” mice were supplied with food and water in small dishes on the floor of the cage to alleviate suffering from dehydration and malnourishment. Two weeks after inoculation, infected animals were euthanized by CO₂ asphyxiation, followed by cervical dislocation. Spleen, liver, mesenteric lymph nodes, Peyer’s patches, and cecum were collected; homogenized in 1 ml PBS; and then serially diluted for plating to enumerate CFU. Competitive indexes (CIs) were calculated as follows: (CFU_{wild type}/CFU_{mutant}) output/(CFU_{wild type}/CFU_{mutant}) input.

Gentamicin protection assays. RAW264.7 cells stably transfected with the pHbA-1-*neo* expression plasmid containing the full-length *Scl11a1* cDNA (25) were seeded at 1.5 × 10⁵ cells per well in poly-L-lysine-coated 24-well tissue culture plates. Cells were activated with 20 ng/ml lipopolysaccharide (LPS) (*S. enterica* serovar Typhimurium LPS; Sigma-Aldrich) and 20 U/ml IFN-γ (PeproTech) for 18 h. Bacteria opsonized with normal mouse serum (Sigma) were added to macrophages at a multiplicity of infection of 10:1 (bacteria to macrophage). After 30 min, free and loosely adherent bacteria were removed by washing with PBS twice, and cells were incubated for a further 1.5 h at 37°C in fresh medium supplemented with gentamicin (100 µg/ml) to kill extracellular bacteria. Medium was then exchanged for medium supplemented with 10 µg/ml gentamicin to inhibit extracellular bacterial growth. At 2, 18, and 24 h, wells were washed twice with prewarmed PBS, incubated with 1% Triton X-100 for 5 min, and lysed, and serial dilutions were plated to enumerate CFU.

Statistics. *P* values were calculated with GraphPad Prism 5 (GraphPad Software Inc.) and considered significant if the *P* value was <0.05. For nonparametric data, Wilcoxon signed-rank or Mann-Whitney tests were used. Otherwise, one-sample *t* tests or Student’s *t* tests were used.

RESULTS

***Salmonella fepB* ferric iron transporter mutants colonize mice poorly in competitive infections.** We constructed a mutant *Salmonella* Typhimurium SL1344 strain with a deletion of *fepB*, encoding a protein crucial for ferric iron uptake. To establish whether FepB is needed for *Salmonella* survival and growth in tissue, mixed-infection experiments were performed in Sv129S6 (Nramp1^{+/+}) mice. These mice become chronically infected with *Salmonella* and can be used to monitor infection long term (38). We evaluated the number of wild-type organisms and the number of *fepB* mutant organisms in feces, a close proxy for the number of organisms in the cecum (39), at 1 and 3 days postinfection. At 1 day postinfection, equivalent numbers of wild-type and *ΔfepB* mutant bacteria were recovered from fecal pellets, indicating that the *ΔfepB* mutant colonizes the intestine well initially (Fig. 1A). However, by 3 days postinfection, the *ΔfepB* mutant had 100- to 10,000-fold-decreased colonization relative to the wild-type organism (Fig. 1A).




FIG 1 *Salmonella* requires the FepB ferric iron transporter for persistent infection of Sv129S6 (*Nramp1*^{+/+}) mice. (A and B) Mice were orogastrically inoculated with a 1:1 mixture of wild-type (WT) and $\Delta fepB$ strains. (A) One and 3 days postinfection (P.I.), fecal samples were collected and homogenized to determine bacterial loads. *, $P = 0.003$ compared with the null hypothesis. (B) At 2 weeks, mice were sacrificed, and the spleen, liver, mesenteric lymph nodes (MLN), Peyer's patches (PP), and cecum were immediately homogenized and plated to determine bacterial loads. Dashed lines indicate the limit of detection based on average tissue weight. Each symbol represents one mouse. *, $P < 0.05$; **, $P < 0.00001$ compared with the null hypothesis. (C and D) Mice were orogastrically inoculated with the wild-type or $\Delta fepB$ strain. (C) One and 3 days postinfection, fecal samples were collected and homogenized to determine bacterial loads in the intestine. *, $P = 0.03$ versus the wild type. (D) After 2 weeks, mice were sacrificed, and the spleen, liver, mesenteric lymph nodes, Peyer's patches, and cecum were immediately homogenized and plated to determine bacterial loads. Each symbol represents one mouse ($n = 5$). *, $P < 0.05$; **, $P < 0.001$ versus the wild type for each organ. Dashed lines indicate the limit of detection based on average tissue weight.

At 2 weeks postinfection, mice were euthanized, and tissues were collected to enumerate CFU. In gut-associated tissues, including the cecum, Peyer's patches, and mesenteric lymph nodes, the numbers of $\Delta fepB$ strain bacteria were 100-fold reduced relative to the numbers of wild-type organisms (Fig. 1B). In the spleen and liver, the phenotype was more severe, as $\Delta fepB$ colonies were recovered 1,000 times less frequently than wild-type colonies. These data suggest that ferric iron transport by the Fep system is required for tissue colonization in the context of coinfection with wild-type bacteria.

***Salmonella* requires the FepB ferric iron transporter for persistent infection of mice.** We used single-infection experiments in Sv129S6 (*Nramp1*^{+/+}) mice to establish whether the ferric iron transporter mutant colonizes tissues normally in the absence of the wild-type organism. At 1 day postinfection, the $\Delta fepB$ mutant was recovered at slightly but significantly lower numbers from fecal pellets of mice infected with the $\Delta fepB$ mutant than from mice infected with the wild-type organism (Fig. 1C). Within 3 days postinfection, the difference in recovery from fecal pellets was >10,000-fold, with mice infected with the $\Delta fepB$ mutant exhibiting a strong colonization defect. The large difference in relative strain recovery at 1 and 3 days suggests that the $\Delta fepB$ mutant initially colonizes the intestine well. At 2 weeks postinfection, there were 10^3 to 10^6 CFU per gram of wild-type *Salmonella* bacteria in each tissue examined, while tissues of mice infected with the $\Delta fepB$ mutant lacked detectable colonization (Fig. 1D). These results, in combination with the mixed-infection experiments,

clearly demonstrate that *Salmonella* infection of *Nramp1*^{+/+} mice requires the FepB ferric iron transporter in all tissues examined.

***Salmonella* requires the FepB ferric iron transporter for replication in macrophages.** During persistent infection, *S. Typhimurium* resides within macrophages (5, 38). To establish whether FepB is required for *Salmonella* replication in macrophages, macrophage-like *Nramp1*^{+/+} tissue culture cells were individually infected with equivalent numbers of wild-type or $\Delta fepB$ mutant *Salmonella* bacteria. Infected cells were lysed, and CFU were enumerated over a time course (Fig. 2A). The wild-type strain replicated during the first 18 to 24 h, while CFU recovered for the $\Delta fepB$ mutant strain remained similar throughout the experiment. Expression of *fepB* on a plasmid is toxic to *Escherichia coli* (40) and *Salmonella* (data not shown). Therefore, to confirm that *fepB* is needed for replication in macrophages, wild-type *fepB* was restored in the *fepB*-deleted strain by phage transduction. Replication of the restored strain in macrophages was comparable to that of the wild-type strain. Macrophages were also treated with 50 μ M deferasirox, a ferric iron-specific chelator, over a time course of infection. At each time point examined, replication of the wild type in the presence of deferasirox was significantly reduced compared to the replication of bacteria in the presence of the vehicle alone (Fig. 2B), similar to results obtained by Nairz et al. (41). These results demonstrate that ferric iron transport via FepB is required for replication of *Salmonella* in macrophages.

***Salmonella* can utilize ferric iron via FepB as its sole iron source.** Next, we determined whether ferric iron is sufficient to




FIG 2 *Salmonella* requires ferric iron (*fepB*) for growth in macrophages. (A) Cell culture macrophages (Nramp1⁺) were infected with the wild-type, $\Delta fepB$, or *fepB*-restored (*fepB*⁺) strain at a multiplicity of infection of 10. At 2, 18, and 24 h postinfection, macrophages were lysed, serially diluted, and plated to determine numbers of CFU/ml. Error bars indicate standard errors of the means ($n \geq 3$ experiments). *, $P < 0.05$ versus the wild type. (B) Cell culture macrophages (Nramp1⁺) were infected with the wild type in the absence or presence of 50 μ M deferasirox at a multiplicity of infection of 10. At 2, 18, and 24 h postinfection, macrophages were lysed, serially diluted, and plated to determine numbers of CFU/ml. Error bars indicate standard errors of the means ($n \geq 3$ experiments). *, $P < 0.05$ versus the wild type.

support *Salmonella* growth in broth. Control experiments demonstrated no significant growth differences between the $\Delta fepB$ mutant and the wild-type strain in nutrient-rich medium (LB medium) at 37°C over 16 h (Fig. 3A). The addition of the iron chelator 2,2'-dipyridyl, which has a similar binding affinity for ferrous and ferric iron (42), prevented the growth of both strains similarly. In M9 minimal medium without added iron, the wild-type strain grew significantly better than the $\Delta fepB$ mutant strain (Fig. 3B). Treatment with 100 but not 50 μ M ferric chloride significantly increased growth of the wild-type strain. Growth of the $\Delta fepB$ mutant strain was not improved by the addition of ferric chloride. Chelation of iron with 2,2'-dipyridyl in M9 medium reduced the growth of both wild-type *Salmonella* and the $\Delta fepB$ strain, as observed in LB medium. Collectively, these observations suggest that *Salmonella* can utilize ferric iron acquired via the catecholate transporter FepB as its sole iron source.

***Salmonella* requires Fe³⁺ binding siderophore uptake via FepA and IroN in mice.** *Salmonella* acquires ferric iron by producing and secreting the siderophores enterobactin and salmochelin, which, once bound to iron, are transported through the bacterial outer membrane receptor FepA (enterobactin) or IroN (enterobactin or salmochelin) and bind FepB in the periplasm. To determine whether ferric iron uptake via one or both of these siderophores is necessary for persistent infection, mice were orogastrically inoculated with the individual wild-type, $\Delta fepA$, or $\Delta iroN$ strain or with a $\Delta iroN \Delta fepA$ double mutant strain. After 2 weeks, mice were sacrificed, and tissue colonization was determined. Tissue colonization by $\Delta fepA$ and $\Delta iroN$ single and double mutants was significantly reduced

compared to that by the wild-type strain in the cecum (Fig. 4A). The $\Delta fepA$ mutant also had a significant colonization defect in Peyer's patches compared to wild-type *Salmonella* (Fig. 4B). In contrast, the $\Delta iroN$ mutant strain colonized the mesenteric lymph nodes, spleen, and liver poorly (Fig. 4C and D). Deletion of both *iroN* and *fepA* resulted in a severe colonization defect in the mesenteric lymph nodes, spleen, and liver compared to wild-type bacteria. These results suggest that both FepA and IroN are utilized in all tissues but that there may be a greater requirement for IroN in deep tissues during *Salmonella* infection of Nramp1^{+/+} mice.

***fepA* or *iroN* is required for growth in macrophages.** To establish whether *fepA* and/or *iroN* is required for *Salmonella* replication in macrophages, macrophage-like Nramp1^{+/+} tissue culture cells were singly infected with the *Salmonella* wild-type, $\Delta fepA$, $\Delta iroN$, or $\Delta iroN \Delta fepA$ mutant strain. Over a time course of infection, CFU enumeration indicated that deletion of $\Delta fepA$ or $\Delta iroN$ did not significantly affect the ability of *Salmonella* to replicate in macrophages compared to wild-type bacteria. However, deletion of both $\Delta iroN$ and $\Delta fepA$ prevented *Salmonella* replication within macrophages at between 2 and 24 h postinfection (Fig. 5). These results demonstrate that *Salmonella* requires either *iroN* (salmochelin or enterobactin outer membrane uptake receptor) or *fepA* (enterobactin outer membrane uptake receptor) for replication in macrophages.

***Salmonella* requires siderophore synthesis in mice.** Bacteria overproduce siderophores in strain backgrounds lacking siderophore import genes such as *iroN* and *fepA* (28). Excessive siderophore production could result in local iron starvation and thus




FIG 3 *Salmonella* can grow with ferric iron as the sole iron source. (A) The indicated strains were grown in LB medium supplemented with 200 μ M iron chelator 2,2'-dipyridyl (dipi). The optical density at 600 nm was monitored for 16 h. Error bars indicate standard deviations ($n \geq 3$ experiments). *, $P < 0.05$ versus the wild type; **, $P < 0.05$ versus the $\Delta fepB$ mutant. (B) The indicated strains were grown in M9 minimal medium supplemented with ferric chloride (Fe³⁺) and/or 200 μ M dipi. The optical density at 600 nm was monitored for 16 h. Error bars indicate standard deviations ($n \geq 3$ experiments). *, $P < 0.05$ versus the wild type; **, $P < 0.05$ versus the $\Delta fepB$ mutant.




FIG 4 *Salmonella* requires enterobactin and salmochelin during persistent infection of mice. In separate experiments (left and right), mice were orogastrically inoculated with the wild-type, $\Delta iroN$, $\Delta fepA$, or $\Delta iroN \Delta fepA$ strain. After 2 weeks, mice were sacrificed, and the cecum (A), Peyer's patches (B), mesenteric lymph nodes (C), spleen (D), and liver (E) were immediately homogenized and plated to determine bacterial loads. Each symbol represents one mouse ($n = 5$ to 10). *, $P < 0.05$ versus the wild type; **, $P < 0.003$ versus the wild type.

poor tissue colonization *in vivo*. To determine whether the *in vivo* growth defects observed for the $\Delta iroN$ and $\Delta fepA$ strains reflected enhanced siderophore production, a strain unable to import or produce enterobactin and salmochelin ($\Delta iroN \Delta fepA \Delta entC$) was generated. Mice orogastrically inoculated with individual wild-type or $\Delta iroN \Delta fepA \Delta entC$ bacteria were sacrificed at 2 weeks postinfection, and tissue colonization was determined (Fig. 6).

Tissue colonization by the triple mutant strain was significantly reduced compared to that of the wild-type strain in the cecum, Peyer's patches, mesenteric lymph nodes, and spleen, although the difference in liver colonization was not significant. These observations indicate that the reduced tissue colonization observed for $\Delta fepA$ and $\Delta iroN$ mutants is not dependent upon siderophore overproduction.




FIG 5 *Salmonella* requires either *fepA* or *iroN* for growth in macrophages. Cell culture macrophages (Nramp1⁺) were infected with the wild-type, $\Delta iroN$, $\Delta fepA$, or $\Delta iroN \Delta fepA$ strain at a multiplicity of infection of 10. At 2, 18, and 24 h postinfection, macrophages were lysed, serially diluted, and plated to determine numbers of CFU/ml. In panel B, the CFU index is defined as the number of CFU at 18 h relative to the number of CFU at 2 h (18 h) and the number of CFU at 24 h relative to the number of CFU at 18 h (24 h). Error bars indicate standard deviations ($n \geq 3$ experiments). *, $P < 0.05$ versus the wild type.




FIG 6 *Salmonella* requires siderophore synthesis in mice. Mice were orogastrically inoculated with the wild-type or Δ iroN Δ fepA Δ entC strain. After 2 weeks, mice were sacrificed, and the spleen, liver, mesenteric lymph nodes, Peyer's patches, and cecum were immediately homogenized and plated to determine bacterial loads. Each symbol represents one mouse ($n \geq 7$). *, $P < 0.006$ versus the wild type for each organ; **, $P < 0.0006$ versus the wild type for each organ.

DISCUSSION

Salmonella encodes transporters for both ferric and ferrous iron. Ferrous iron is taken up by FeoB (43) and by divalent cation transporters (MntH and SitA) (44, 45). FeoB and SitA have known roles in BALB/c Nramp1^{-/-} mice during acute infection (43, 44). Ferric iron is acquired upon the import of iron-bound siderophores through FepA and IroN and siderophore capture in the periplasm by FepB (23). Previous work established that *fepA* and *iroN* are dispensable for infection of BALB/c Nramp1^{-/-} mice (23, 28, 29). We demonstrate that siderophore capture and import via *fepB* are required for *Salmonella* survival and replication within macrophages as well as for colonization of mice upon orogastric inoculation. In addition, the ability to produce and/or take up either the enterobactin or salmochelin siderophores is needed for *Salmonella* colonization of both gastric and deep tissues in Sv129S6 Nramp1^{+/+} mice. Our observations are consistent with those of Gorbacheva et al., who found that an *S. Typhi* mutant strain lacking the ability to synthesize enterobactin (and salmochelin) failed to replicate in human macrophages (46). The growth defect observed in strains lacking *fepB* was more severe than that in strains unable to produce or import either siderophore. This may reflect that FepBDGC also takes up derivatives of catecholates that are imported by CirA (27, 28, 33). Overall, the data suggest that *S. Typhimurium* requires ferric iron during persistent infection of mice. Moreover, the results from infection of mice with single mutant strains indicate that enterobactin acquisition through FepA is important for colonization of the cecum and Peyer's patches and that enterobactin or salmochelin import via IroN is important for colonization of all tissues examined. Our studies thus provide an experimental foundation to make hypotheses about the temporal-spatial requirements of siderophores during different stages of infection.

Mice that lack hemochromatosis gene alleles (*Hfe*) have increased resistance to *Salmonella* infection. This is due to decreased levels of macrophage iron via an enhanced production of ferroportin (47) as well as increased iron sequestration via augmented lipocalin-2 production, which scavenges enterobactin (25). These results are in concordance with our current studies indicating a preferential uptake of ferric iron via IroN and FepA during infection. Because hemochromatosis studies have been performed in Nramp1^{-/-} mice, it will be of interest to determine if similar

mechanisms occur in Nramp1^{+/+} mice, which tightly regulate iron stores and mimic the acute-to-chronic progression of human typhoid fever.

Patients with β -thalassemia and thus ferric iron overload, including within macrophages, have increased susceptibility to bacterial infection. Mice with experimentally induced iron overload have significantly enhanced *Salmonella* growth rates in all tissues examined (48, 49). Dysregulation of hepcidin, a peptide that normally limits excess iron absorption, leads to iron overload and increased susceptibility to *Salmonella* infection (50). The mechanism by which iron overload increases susceptibility to *Salmonella* infection remains to be fully resolved, but the results of previous studies and the current work indicate that iron overload may provide abundant stores of ferric iron for bacterial utilization.

There are at least two pathways by which *Salmonella* may access ferric iron in macrophages. First, ferric iron is transported to different tissues by the host protein transferrin, which binds cognate receptors on cells, including macrophages (51). Host transferrin receptor expression is increased in response to many intracellular pathogens, such as *Francisella tularensis*, *Mycobacterium tuberculosis*, and *Ehrlichia* (52–54). Upon acidification of transferrin-containing endosomes, ferric iron is released from transferrin and may be accessed by these pathogens. In Nramp1^{-/-} macrophages, *Salmonella* does not require transferrin expression for successful intracellular survival (52). However, Nramp1⁺ macrophages limit *Salmonella* access to iron (20, 25). Therefore, *Salmonella* may, like other pathogens that live within vesicles, intercept transferrin-containing vesicles to obtain ferric iron. Ferric iron released from transferrin can alternatively be transported to the cytosol and stored within ferritin. Some cytosolic microbes, such as *Neisseria meningitidis* and *Listeria monocytogenes*, utilize ferritin as an iron source through mechanisms that directly degrade ferritin pools in the cytosol (55, 56). Since *Salmonella* is typically an intravesicular pathogen in macrophages, it may not be able to directly access cytosolic ferritin. However, iron starvation and cellular stress induce autophagy of ferritin, followed by ferritin degradation in vesicles (57–59). *Salmonella* has been demonstrated to induce autophagy in macrophages (60, 61) and can also direct vesicular trafficking events (62). Thus, one hypothesis is that *Salmonella* facilitates fusion of the vesicle where it resides with ferritin-containing vesicles and then uses enterobactin or salmochelin to remove Fe³⁺ from the cytosolic storage protein ferritin. The addition of exogenous iron-loaded ferritin to Nramp1^{-/-} macrophage-like cells increases *Salmonella* survival, consistent with this hypothesis (63).

Together, our studies combined with others highlight the particular importance of ferric iron in both nutritional immunity and *Salmonella* survival in the host. *Salmonella* has successfully evolved ferric iron-specific siderophores, enterobactin and salmochelin, that may be especially suited to take advantage of intracellular ferric iron storage pools. The current work also indicates that *Salmonella* may utilize siderophores differentially based upon tissue localization. Modulation of host iron regulators or the preferentially utilized *Salmonella* iron uptake systems has the potential to become pivotal in the development of new pharmacological approaches to control and/or prevent disease.

ACKNOWLEDGMENTS

We thank all the members of the Detweiler laboratory, M. Carolina Pilonieta, Erin McDonald, Eugenia Silva-Herzog, Christopher English, Garth

Kreitz, and Angelika Krivenko, for helpful discussions and technical help over the course of this project.

This work was supported by NIH grants 1F32AI094766 (T.A.N.), AI072492, and AI095395 (C.S.D.).

We declare that there are no conflicts of interest.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

- Miller SI, Pegues DA. 2009. *Salmonella* species, including *Salmonella typhi*, p 2887–2904. In Mandell GL, Bennett JE, Dolin R(ed), Mandell, Douglas, and Bennett's principles and practice of infectious diseases, 7th ed. Churchill Livingstone, Philadelphia, PA.
- Ohl ME, Miller SI. 2001. *Salmonella*: a model for bacterial pathogenesis. *Annu. Rev. Med.* 52:259–274.
- Baumler AJ, Tsolis RM, Heffron F. 1996. Contribution of fimbrial operons to attachment to and invasion of epithelial cell lines by *Salmonella typhimurium*. *Infect. Immun.* 64:1862–1865.
- Jones BD, Ghori N, Falkow S. 1994. *Salmonella typhimurium* initiates murine infection by penetrating and destroying the specialized epithelial M cells of the Peyer's patches. *J. Exp. Med.* 180:15–23.
- Nix RN, Altschuler SE, Henson PM, Detweiler CS. 2007. Hemophagocytic macrophages harbor *Salmonella enterica* during persistent infection. *PLoS Pathog.* 3:e193. doi:10.1371/journal.ppat.0030193.
- Silva-Herzog E, Detweiler CS. 2010. *Salmonella enterica* replication in hemophagocytic macrophages requires two type three secretion systems. *Infect. Immun.* 78:3369–3377.
- Fenton HJH. 1894. Oxidation of tartaric acid in presence of iron. *J. Chem. Soc. Trans.* 65:899–911.
- Latunde-Dada GO, Van der Westhuizen J, Vulpe CD, Anderson GJ, Simpson RJ, McKie AT. 2002. Molecular and functional roles of duodenal cytochrome B (Dcytb) in iron metabolism. *Blood Cells Mol. Dis.* 29:356–360.
- McKie AT, Barrow D, Latunde-Dada GO, Rolfs A, Sager G, Mudaly E, Mudaly M, Richardson C, Barlow D, Bomford A, Peters TJ, Raja KB, Shirali S, Hediger MA, Farzaneh F, Simpson RJ. 2001. An iron-regulated ferric reductase associated with the absorption of dietary iron. *Science* 291:1755–1759.
- Vulpe CD, Kuo YM, Murphy TL, Cowley L, Askwith C, Libina N, Gitschier J, Anderson GJ. 1999. Hephaestin, a ceruloplasmin homologue implicated in intestinal iron transport, is defective in the sla mouse. *Nat. Genet.* 21:195–199.
- Aisen P, Listowsky I. 1980. Iron transport and storage proteins. *Annu. Rev. Biochem.* 49:357–393.
- Ohgami RS, Campagna DR, Greer EL, Antiochos B, McDonald A, Chen J, Sharp JJ, Fujiwara Y, Barker JE, Fleming MD. 2005. Identification of a ferrireductase required for efficient transferrin-dependent iron uptake in erythroid cells. *Nat. Genet.* 37:1264–1269.
- Su MA, Trenor CC, Fleming JC, Fleming MD, Andrews NC. 1998. The G185R mutation disrupts function of the iron transporter Nramp2. *Blood* 92:2157–2163.
- Gkouvatsos K, Papanikolaou G, Pantopoulos K. 2012. Regulation of iron transport and the role of transferrin. *Biochim. Biophys. Acta* 1820:188–202.
- Kehl-Fie TE, Skaar EP. 2010. Nutritional immunity beyond iron: a role for manganese and zinc. *Curr. Opin. Chem. Biol.* 14:218–224.
- Weiss G, Goodnough LT. 2005. Anemia of chronic disease. *N. Engl. J. Med.* 352:1011–1023.
- Brown DE, Libby S, Moreland SM, McCoy MW, Brabb T, Stepanek A, Fang FC, Detweiler CS. 27 February 2013. *Salmonella enterica* causes more severe inflammatory disease in C57/BL6 Nramp1G169 mice than Sv129S6 mice. *Vet. Pathol.* [Epub ahead of print.] doi:10.1177/0300985813478213.
- Nairz M, Fritsche G, Crouch ML, Barton HC, Fang FC, Weiss G. 2009. Slc11a1 limits intracellular growth of *Salmonella enterica* sv. Typhimurium by promoting macrophage immune effector functions and impairing bacterial iron acquisition. *Cell. Microbiol.* 11:1365–1381.
- Brown DE, McCoy MW, Pilonieta MC, Nix RN, Detweiler CS. 2010. Chronic murine typhoid fever is a natural model of secondary hemophagocytic lymphohistiocytosis. *PLoS One* 5:e9441. doi:10.1371/journal.pone.0009441.
- Nairz M, Fritsche G, Brunner P, Talasz H, Hantke K, Weiss G. 2008. Interferon-gamma limits the availability of iron for intramacrophage *Salmonella typhimurium*. *Eur. J. Immunol.* 38:1923–1936.
- Chlosta S, Fishman DS, Harrington L, Johnson EE, Knutson MD, Wessling-Resnick M, Cherayil BJ. 2006. The iron efflux protein ferroportin regulates the intracellular growth of *Salmonella enterica*. *Infect. Immun.* 74:3065–3067.
- Skare JT, Ahmer BMM, Seachord CL, Darveau RP, Postle K. 1993. Energy transduction between membranes: TonB, a cytoplasmic membrane protein, can be chemically cross-linked in vivo to the outer membrane receptor FepA. *J. Biol. Chem.* 268:16302–16308.
- Rabsch W, Voigt W, Reissbrodt R, Tsolis RM, Bäumler AJ. 1999. *Salmonella typhimurium* IroN and FepA proteins mediate uptake of enterobactin but differ in their specificity for other siderophores. *J. Bacteriol.* 181:3610–3612.
- Flo TH, Smith KD, Sato S, Rodriguez DJ, Holmes MA, Strong RK, Akira S, Aderem A. 2004. Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 432:917–921.
- Fritsche G, Nairz M, Libby SJ, Fang FC, Weiss G. 2012. Slc11a1 (Nramp1) impairs growth of *Salmonella enterica* serovar Typhimurium in macrophages via stimulation of lipocalin-2 expression. *J. Leukoc. Biol.* 92:353–359.
- Raffatelli M, George MD, Akiyama Y, Hornsby MJ, Nuccio SP, Paixao TA, Butler BP, Chu H, Santos RL, Berger T, Mak TW, Tsolis RM, Bevins CL, Solnick JV, Dandekar S, Bäumler AJ. 2009. Lipocalin-2 resistance confers an advantage to *Salmonella enterica* serotype Typhimurium for growth and survival in the inflamed intestine. *Cell Host Microbe* 5:476–486.
- Hantke K, Nicholson G, Rabsch W, Winkelmann G. 2003. Salmochelins, siderophores of *Salmonella enterica* and uropathogenic *Escherichia coli* strains, are recognized by the outer membrane receptor IroN. *Proc. Natl. Acad. Sci. U. S. A.* 100:3677–3682.
- Rabsch W, Methner U, Voigt W, Tschäpe H, Reissbrodt R, Williams PH. 2003. Role of receptor proteins for enterobactin and 2,3-dihydroxybenzoylserine in virulence of *Salmonella enterica*. *Infect. Immun.* 71:6953–6961.
- Williams PH, Rabsch W, Methner U, Voigt W, Tschäpe H, Reissbrodt R. 2006. Catecholate receptor proteins in *Salmonella enterica*: role in virulence and implications for vaccine development. *Vaccine* 24:3840–3844.
- Crouch ML, Castor M, Karlinsey JE, Kalthorn T, Fang FC. 2008. Biosynthesis and IroC-dependent export of the siderophore salmochelin are essential for virulence of *Salmonella enterica* serovar Typhimurium. *Mol. Microbiol.* 67:971–983.
- Hensel M, Shea JE, Gleeson C, Jones MD, Dalton E, Holden DW. 1995. Simultaneous identification of bacterial virulence genes by negative selection. *Science* 269:400–403.
- Bearson BL, Bearson SM, Uthe JJ, Dowd SE, Houghton JO, Lee I, Toscano MJ, Lay DC, Jr. 2008. Iron regulated genes of *Salmonella enterica* serovar Typhimurium in response to norepinephrine and the requirement of fepDGC for norepinephrine-enhanced growth. *Microbes Infect.* 10:807–816.
- Hantke K. 1990. Dihydroxybenzoylserine—a siderophore for *E. coli*. *FEMS Microbiol. Lett.* 67:5–8.
- Merritt FF, Smith BP, Reina-Guerra M, Habasha F, Johnson E. 1984. Relationship of cutaneous delayed hypersensitivity to protection from challenge exposure with *Salmonella typhimurium* in calves. *Am. J. Vet. Res.* 45:1081–1085.
- Datsenko KA, Wanner B. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* 97:6640–6645.
- Santiviago CA, Reynolds MM, Porwollik S, Choi SH, Long F, Andrews-Polymenis HL, McClelland M. 2009. Analysis of pools of targeted *Salmonella* deletion mutants identifies novel genes affecting fitness during competitive infection in mice. *PLoS Pathog.* 5:e1000477. doi:10.1371/journal.ppat.1000477.
- National Research Council. 2011. Guide for the care and use of laboratory animals, 8th ed. National Academies Press, Washington, DC.
- Monack DM, Bouley DM, Falkow S. 2004. *Salmonella typhimurium* persists within macrophages in the mesenteric lymph nodes of chronically infected Nramp1^{+/+} mice and can be reactivated by IFN γ neutralization. *J. Exp. Med.* 199:231–241.
- Sivula CP, Bogomolnaya LM, Andrews-Polymenis HL. 2008. A comparison of central colonization of *Salmonella enterica* serotype Typhimurium in

- white leghorn chicks and *Salmonella*-resistant mice. *BMC Microbiol.* 8:182–195.
40. Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H, Mori H. 2005. Complete set of ORF clones of *Escherichia coli* ASKA library (a complete set of *E. coli* K-12 ORF archive): unique resources for biological research. *DNA Res.* 12:291–299.
 41. Nairz M, Schleicher U, Schroll A, Sonnweber T, Theurl I, Ludwiczek S, Talasz H, Brandacher G, Moser PL, Muckenthaler MU, Fang FC, Bogdan C, Weiss G. 2013. Nitric oxide-mediated regulation of ferroportin-1 controls macrophage iron homeostasis and immune function in *Salmonella* infection. *J. Exp. Med.* 210:855–873.
 42. Zhou T, Ma Y, Kong X, Hider RC. 2002. Design of iron chelators with therapeutic application. *Coord. Chem. Rev.* 232:151–171.
 43. Tsolis RM, Bäumlér AJ, Heffron F, Stojiljkovic I. 1996. Contribution of TonB- and Feo-mediated iron uptake to growth of *Salmonella* Typhimurium in the mouse. *Infect. Immun.* 64:4549–4556.
 44. Boyer E, Bergevin I, Malo D, Gros P, Cellier MF. 2002. Acquisition of Mn(II) in addition to Fe(II) is required for full virulence of *Salmonella enterica* serovar Typhimurium. *Infect. Immun.* 70:6032–6042.
 45. Janakiraman A, Slauch JM. 2000. The putative iron transport system SitABCD encoded on SPI1 is required for full virulence of *Salmonella typhimurium*. *Mol. Microbiol.* 35:1146–1155.
 46. Gorbacheva VY, Faundez G, Godfrey HP, Cabello FC. 2001. Restricted growth of ent(–) and tonB mutants of *Salmonella enterica* serovar Typhi in human Mono Mac 6 monocytic cells. *FEMS Microbiol. Lett.* 196:7–11.
 47. Wang L, Johnson EE, Shi HN, Walker WA, Wessling-Resnick M, Cherayil BJ. 2008. Attenuated inflammatory responses in hemochromatosis reveal a role for iron in the regulation of macrophage cytokine translation. *J. Immunol.* 181:2723–2731.
 48. Caroline L, Kozinn PJ. 1969. Infection and iron overload in thalassemia. *Ann. N. Y. Acad. Sci.* 165:148–155.
 49. Sawatzki G, Hoffmann FA, Kubanek B. 1983. Acute iron overload in mice: pathogenesis of *Salmonella typhimurium* infection. *Infect. Immun.* 39:659–665.
 50. Yuki KE, Eva MM, Richer E, Chung D, Paquet M, Cellier M, Canonne-Hergaux F, Vaulont S, Vidal SM, Malo D. 2013. Suppression of hepcidin expression and iron overload mediate *Salmonella* susceptibility in ankyrin 1 ENU-induced mutant. *PLoS One* 8:e55331. doi:10.1371/journal.pone.0055331.
 51. Theurl I, Fritsche G, Ludwiczek S, Garimorth K, Bellmann-Weiler R, Weiss G. 2005. The macrophage: a cellular factory at the interphase between iron and immunity for the control of infections. *Biometals.* 18:359–367.
 52. Pan X, Tamilselvam B, Hansen EJ, Daefler S. 2010. Modulation of iron homeostasis in macrophages by bacterial intracellular pathogens. *BMC Microbiol.* 10:64. doi:10.1186/1471-2180-10-64.
 53. Barnewall RE, Ohashi N, Rikihisa Y. 1999. *Ehrlichia chaffeensis* and *E. senetsu*, but not the human granulocytic ehrlichiosis agent, colocalize with transferrin receptor and up-regulate transferrin receptor mRNA by activating iron-responsive protein 1. *Infect. Immun.* 67:2258–2265.
 54. Clemens DL, Horwitz M. 1996. The *Mycobacterium tuberculosis* phagosome interacts with early endosomes and is accessible to exogenously administered transferrin. *J. Exp. Med.* 184:1349–1355.
 55. Larson JA, Howie HL, So M. 2004. *Neisseria meningitidis* accelerates ferritin degradation in host epithelial cells to yield an essential iron source. *Mol. Microbiol.* 53:807–820.
 56. Schaible UE, Kaufmann SH. 2004. Iron and microbial infection. *Nat. Rev. Microbiol.* 2:946–953.
 57. Konijn AM, Glickstein H, Vaisman B, Meyron-Holtz EG, Slotki IN, Cabantchik ZI. 1999. The cellular labile iron pool and intracellular ferritin in K562 cells. *Blood* 94:2128–2134.
 58. Roberts S, Bomford A. 1988. Ferritin iron kinetics and protein turnover in K562 cells. *J. Biol. Chem.* 263:19181–19187.
 59. Ollinger K, Roberg K. 1997. Nutrient deprivation of cultured rat hepatocytes increases the desferrioxamine-available iron pool and augments the sensitivity to hydrogen peroxide. *J. Biol. Chem.* 272:23707–23711.
 60. Mesquita FS, Thomas M, Sachse M, Santos AJ, Figueira R, Holden DW. 2012. The *Salmonella* deubiquitinase SseL inhibits selective autophagy of cytosolic aggregates. *PLoS Pathog.* 8:e1002743. doi:10.1371/journal.ppat.1002743.
 61. Hernandez LD, Pypaert M, Flavell RA, Galán JE. 2003. A *Salmonella* protein causes macrophage cell death by inducing autophagy. *J. Cell Biol.* 163:1123–1131.
 62. Knodler LA, Steele-Mortimer O. 2003. Taking possession: biogenesis of the *Salmonella*-containing vacuole. *Traffic* 4:587–599.
 63. Nairz M, Theurl I, Ludwiczek S, Theurl M, Mair SM, Fritsche G, Weiss G. 2007. The co-ordinated regulation of iron homeostasis in murine macrophages limits the availability of iron for intracellular *Salmonella typhimurium*. *Cell. Microbiol.* 9:2126–2140.