

Enzyme Assay Guided Isolation of an α -Amylase Inhibitor Flavonoid from *Vaccinium arctostaphylos* Leaves

Bahman Nickavar^{a*} and Gholamreza Amin^b

^aDepartment of Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^bDepartment of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

Abstract

The management of postprandial hyperglycemia is an important strategy in the control of diabetes mellitus and complications associated with the disease, especially in the diabetes type 2. Therefore, inhibitors of carbohydrate hydrolyzing enzymes can be useful in the treatment of diabetes and medicinal plants can offer an attractive strategy for the purpose. *Vaccinium arctostaphylos* leaves are considered useful for the treatment of diabetes mellitus in some countries. In our research for antidiabetic compounds from natural sources, we found that the methanol extract of the leaves of *V. arctostaphylos* displayed a potent inhibitory activity on pancreatic α -amylase activity ($IC_{50} = 0.53$ (0.53 – 0.54) mg/mL). The bioassay-guided fractionation of the extract resulted in the isolation of quercetin as an active α -amylase inhibitor. Quercetin showed a dose-dependent inhibitory effect with IC_{50} value 0.17 (0.16 – 0.17) mM.

Keywords: α -Amylase inhibition; Diabetes mellitus; Quercetin; *Vaccinium arctostaphylos*.

Introduction

Diabetes mellitus is a chronic metabolic disease characterized by elevated blood glucose levels (1, 2). The control of hyperglycemia is critical in the management of diabetes because acute and chronic complications can occur if the blood glucose concentration is not kept in normal levels (3, 4). One therapeutic approach for diabetic patients, especially type 2 diabetes, is to retard the absorption glucose by inhibition of carbohydrate hydrolyzing enzymes, such as α -amylase and α -glucosidases. The inhibition of the enzymes leads to decrease in the digestion and absorption of carbohydrates, thereby decreasing the postprandial hyperglycemia (1, 5). The use of natural products as complementary

approaches to existing medications for the treatment of diabetes mellitus is growing in the world and many plant species in different countries are known to have antidiabetic effects (6, 7). Various members of *Vaccinium* genus are one of the most frequently used natural antidiabetic remedies from plant origin (6, 8, 9). It comprises nearly 200 species, most of them are found in the northern hemisphere (10). The genus is represented in Iran only by the species *V. arctostaphylos* L (11). The plant is a resource of traditional Iranian herbal medicine, which is highly recommended for the treatment of diabetes (12). The decoction and/or infusion of the berries and leaves of *V. arctostaphylos* are effectively used for the treatment of diabetes in Iran, Republic of Georgia and Russia (12, 13).

To the best of our knowledge, there are few studies on antidiabetic properties of *V. arctostaphylos*. Feshani *et al.*, have shown that

* Corresponding author:

E-mail: bnickavar@yahoo.com

Table 1. α -Amylase inhibitory activities and IC₅₀ values of the leaf extract of *V. arctostaphylos* and its active compound quercetin.

Concentration	Inhibition (%) ^a	IC ₅₀ ^b
<i>Leaf extract (mg/mL)</i>		
1.00	85.13 ± 0.84	
0.80	75.28 ± 1.07	
0.64	60.62 ± 0.90	0.53 (0.53 – 0.54) mg/mL
0.51	36.87 ± 0.91	
0.41	19.12 ± 0.92	
<i>Quercetin (mM)</i>		
0.464	72.23 ± 0.86	
0.297	55.30 ± 0.73	
0.190	43.58 ± 1.02	0.17 (0.16 – 0.17) mM
0.122	22.02 ± 0.82	
0.078	10.06 ± 0.90	

^a The data are expressed as means ± SEM for five experiments in each group.

^b The IC₅₀ values were established by logarithmic regression curves with normalized data (using the computer software GraphPad Prism 3.02 for Windows) and presented as their respective 95% confidence limits.

the extract of *V. arctostaphylos* berries was an effective antihyperglycemic, antioxidant and antihyperlipidemic agent in alloxan-diabetic rats (9). Also, based on the study of Nickavar and Amin, the berries of *V. arctostaphylos* had a potent inhibitory activity on porcine pancreatic α -amylase enzyme (14). However, no study has so far been conducted concerning the antidiabetic activity of the plant leaves. This investigation is a continuation of our previous work to study the antidiabetic and α -amylase inhibitory activities of *Vaccinium arctostaphylos* and to identify the active constituents of the plant.

Experimental

Plant material

The leaves of *V. arctostaphylos* were collected from the forest region of Asalem in the north of Iran in August 2002. Voucher specimens were deposited in the herbarium of the Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran (no. 6520 THE).

Chemicals

All of the chemicals used in this study were purchased from Sigma-Aldrich Chemical Co. (France) and/or Merck Company (Germany).

Extraction, chromatography and spectroscopy

The dried and ground leaves (500 g) were extracted with 90% methanol three times by maceration method and then, the extract was concentrated *in vacuo*. The crude extract (27 g) was diluted with water and partitioned with *n*-C₆H₁₂, CHCl₃ and EtOAc, successively. The inhibitory effects of the crude extract and all fractions were studied on α -amylase activity. The ethyl acetate fraction displayed the highest inhibitory activity. The ethyl acetate fraction was subjected to more fractionation by column chromatography on silica gel (70–230 mesh) using petroleum-ether / ethyl acetate system as the eluent and the eluent polarity was increased by increasing the ratio of EtOAc during the process. Fraction F5 which showed a high inhibitory activity, was purified by repeated preparative layer chromatography on coated plates with silica gel (230–400 mesh) using EtOAc/CH₃COOH/HCOOH/H₂O (100:11:11:26, v/v/v/v) as the best developing solvent system. Finally, fraction F₅ yielded the pure active compound 1 (64 mg). The UV light (366 nm) was used for the visualization of bands before and after spraying with AlCl₃.

The pure compound was identified on the basis of spectral and chromatographical studies. The UV-Vis spectrum was recorded on a Shimadzu UV-160A spectrophotometer in




Figure 1. Chemical structure of quercetin.

methanol. The $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were taken on a Bruker FT 500 in DMSO-d_6 and chemical shifts were recorded as δ -values. The EI-MS was obtained on an Agilent Technologies 5973 spectrometer.

α -Amylase inhibition test

The α -amylase inhibitory activity was determined using the method described previously by Nickavar and Amin (15, 16). Briefly, 1 mL of the porcine pancreatic α -amylase enzyme solution (0.5 IU/mL) in 20 mM phosphate buffer (pH 6.9) was incubated with 1 mL of each test (at various concentrations) for 30 min. The reaction was initiated by adding 1 mL of 0.5% soluble potato starch solution and the mixture was incubated for 3 min at 25°C . Then, 1 mL of the color reagent (96 mM 3, 5-dinitrosalicylic acid and 5.31 M sodium potassium tartrate in 2 M sodium hydroxide) was added and the mixture was placed in a water bath at 85°C . After 15 min, the reaction mixture was diluted with distilled water and the absorbance value was determined at 540 nm. Individual blanks were prepared for correcting the background absorbance. In this case, the color reagent solution was added prior to the addition of starch solution and the mixture then placed in the water bath immediately. Controls were representative of the 100% enzyme activity. They were conducted in an identical fashion replacing tests with 1 mL of the solvent. Acarbose, a well-known α -amylase inhibitor, was used as positive control. The inhibition percentage of α -amylase was assessed by the following formulae:

$$I_{\alpha\text{-amylase}}(\%) = 100 \cdot \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right)$$

where A_{control} is the absorbance of each control and A_{sample} is the net absorbance of each sample. The net absorbance of each sample was calculated by following the equation:

$$A_{\text{sample}} = A_{\text{test}} - A_{\text{blank}}$$

where A_{test} is the absorbance of each test and A_{blank} is the absorbance of each blank.

The $I_{\alpha\text{-amylase}}(\%)$ for each sample was plotted against the logarithm of the sample concentration, and a logarithmic regression curve was established in order to calculate the IC_{50} value.

Results and Discussion

The methanol extract of *V. arctostaphylos* leaves showed a dose-dependent inhibitory effect on the α -amylase activity [$\text{IC}_{50} = 0.53$ (0.53–0.54) mg/mL] (Table 1).

In order to identify the active components, solvent-solvent partition performed with $n\text{-C}_6\text{H}_{12}$, CHCl_3 and EtOAc, successively. The ethyl acetate fraction revealed the highest activity therefore; it was selected for further separation. The chromatographical analysis of the ethyl acetate fraction showed flavonoid compounds. The most active flavonoid compound was isolated as the pale yellow amorphous powder (64 mg). It had $R_f = 0.48$

on TLC (silica gel 60) with EtOAc/CH₃COOH/HCOOH/H₂O (100:11:11:26, v/v/v/v). The spectroscopic data for the compound were as follows:

UV-Vis: λ_{\max} (in CH₃OH) = 260, 275 (shoulder), 380 nm; + AlCl₃ = 265, 455 nm; + AlCl₃ + HCl = 265, 425; + NaOAc = 275, 380 nm (degradation); + NaOAc + H₃BO₃: 260, 395 nm; + NaOMe = rapid degradation.

¹H-NMR (500 MHz, in DMSO-d₆), δ : 6.15 (1H, d, J = 1.8 Hz, H-6), 6.39 (1H, d, J = 1.8 Hz, H-8), 6.88 (1H, d, J = 8.4 Hz, H-5'), 7.53 (1H, dd, J = 8.4, 1.9 Hz, H-6'), 7.65 (1H, d, J = 1.9 Hz, H-2').

¹³C-NMR (125 MHz, in DMSO-d₆), δ : 94.2 (C-8), 99.0 (C-6), 103.8 (C-10), 115.9 (C-2'), 116.5 (C-5'), 120.8 (C-6'), 122.8 (C-1'), 136.5 (C-3), 145.9 (C-3'), 147.6 (C-2'), 148.5 (C-4'), 157.0 (C-5), 161.5 (C-9), 164.8 (C-7), 176.0 (C-4).

EI-MS (70 eV), m/z (I%): 302 (100%) (M⁺).

The spectral data of the compound showed that it was quercetin (Figure 1) and all of its data were matched with those reported in the literature (17).

In this study, quercetin inhibited α -amylase activity in a dose-dependent manner. The IC₅₀ values for α -amylase inhibition by quercetin and acarbose (as the positive control) were 0.17 (0.16 – 0.17) mM and 0.033 (0.031–0.036) mM, respectively (Figure 1 and Table 1).

The genus of *Vaccinium* generally produces a variety of phenolic metabolites especially anthocyanins, flavonols, phenolic acids, procyanidines, etc (18). Quercetin has already been isolated from leaves of some species belonging to the genus *Vaccinium* such as *V. reticulatum* and *V. calycinum*, *V. myrtillus*, *V. angustifolium*, *V. vitis-idaea*, etc (19-22). Phytochemical studies on the different parts of *V. arctostaphylos* show the occurrence of flavonoids and coumarins in leaves, phenolic acids and their derivatives in leaves and unripe berries and anthocyanins in ripe berries (23-30). However, the study is the first report on the inhibitory effect of *V. arctostaphylos* leaves on α -amylase and isolation of quercetin as their active component.

Based on these experimental results, it can be concluded that a part of antidiabetic effects

observed from *V. arctostaphylos* leaves might be due to the inhibition of the α -amylase by the flavonoid quercetin.

References

- (1) Ali H, Houghton PJ and Soumyanath A. α -amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *J. Ethnopharmacol.* (2006) 107: 449-455.
- (2) Li YH, Wen SP, Kota BP, Peng G, Li GQ, Yamahara J and Roufogalis BD. *Punica granatum* flower extract, a potent α -glucosidase inhibitor, improves postprandial hyperglycemia in Zucker diabetic fatty rats. *J. Ethnopharmacol.* (2005) 99: 239-244.
- (3) Ye F, Shen Z and Xie M. α -glucosidase inhibition from a Chinese medical herb (*Ramulus mori*) in normal and diabetic rats and mice. *Phytomedicine* (2002) 9: 161-166.
- (4) Funke I and Melzig MF. Traditionally used plants in diabetes therapy: phytotherapeutics as inhibitors of α -amylase activity. *Braz. J. Pharmacogn.* (2006) 16: 1-5.
- (5) Kwon YI, Jang HD and Shetty K. Evaluation of *Rhodiola crenulata* and *Rhodiola rosea* for management of Type II diabetes and hypertension. *Asia Pac. J. Clin. Nutr.* (2006) 15: 425-432.
- (6) Martineau LC, Couture A, Spoor D, Benhaddou-Andaloussi A, Harris C, Meddah B, Leduc C, Burt A, Vuong T, Mai Le P, Prentki M, Bennett SA, Arnason JT and Haddad PS. Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. *Phytomedicine* (2006) 13: 612-623.
- (7) Wang L, Zhang XT, Zhang HY, Yao HY and Zhang H. Effect of *Vaccinium bracteatum* Thunb. leaves extract on blood glucose and plasma lipid levels in streptozotocin-induced diabetic mice. *J. Ethnopharmacol.* (2010) 130: 465-469.
- (8) Helmstädter A and Schuster N. *Vaccinium myrtillus* as an antidiabetic medicinal plant - Research through the ages. *Pharmazie* (2010) 65: 315-321.
- (9) Feshani AM, Kouhsari SM and Mohammadi S. *Vaccinium arctostaphylos*, a common herbal medicine in Iran: Molecular and biochemical study of its antidiabetic effects on alloxan-diabetic Wistar rats. *J. Ethnopharmacol.* (2011) 133: 67-74.
- (10) Morazzoni P and Bombardelli E. *Vaccinium myrtillus* L. *Fitoterapia* (1996) 67: 3-29.
- (11) Mozaffarian V. *A Dictionary of Iranian Plants Names*. Farhang Moaser Publishers, Tehran (1998) 570.
- (12) Amin G. *Popular Medicinal Plants of Iran*. Research Deputy of Health Ministry, Tehran (1991) 126.
- (13) Abidov M, Ramazanov A, Jimenez DRM and Chkhikvishvili I. Effect of Blueberin on fasting glucose, C-reactive protein and plasma aminotransferases, in female volunteers with diabetes type 2: double-blind, placebo controlled clinical study. *Georgian Med. News* (2006) 66-71.

- (14) Nickavar B and Amin G. Bioassay-guided separation of an alpha-amylase inhibitor anthocyanin from *Vaccinium arctostaphylos* berries. *Z. Naturforsch.* (2010) 65c: 567-570.
- (15) Nickavar B and Yousefian N. Inhibitory effects of six *Allium* species on α -Amylase enzyme activity. *Iranian J. Pharm. Res.* (2009) 8: 53-57.
- (16) Nickavar B, Abolhasani L and Izadpanah H. α -Amylase inhibitory activities of six *Salvia* species. *Iranian J. Pharm. Res.* (2008) 7: 297-303.
- (17) David JM, Cruz FG, Guedes MLS and Cháve JP. Flavonol glycosides from *Davilla flexuosa*. *J. Braz. Chem. Soc.* (1996) 7: 115-118.
- (18) Chen C, Li Y and Xu Z. Chemical principles and bioactivities of blueberry. *Yaoxue Xuebao* (2010) 45: 422-429.
- (19) Bohm BA and Koupai-Abyazani MR. Flavonoids and condensed tannins from leaves of Hawaiian *Vaccinium reticulatum* and *Vaccinium calycinum* (Ericaceae). *Pac. Sci.* (1994) 48: 458-463.
- (20) Fraisse D, Carnat A and Lamaison JL. Polyphenolic composition of the bilberry leaf. *Ann. Pharmaceut. Franc.* (1996) 54: 280-283.
- (21) McIntyre KL, Harris CS, Saleem A, Beaulieu LP, Ta CA, Haddad PS and Amason JT. Seasonal phytochemical variation of anti-glycation principles in lowbush blueberry (*Vaccinium angustifolium*). *Planta Med.* (2009) 75: 286-292.
- (22) Smolarz HD, Matysik G and Wojciak-Kosior M. High-performance thin-layer chromatographic and densitometric determination of flavonoids in *Vaccinium myrtillus* L. and *Vaccinium vitis-idaea* L. *J. Planar Chromatogr.* (2000) 13: 101-105.
- (23) Latti AK, Kainulainen PS, Hayirlioglu-Ayaz S, Ayaz FA and Riihinen KR. Characterization of anthocyanins in Caucasian blueberries (*Vaccinium arctostaphylos* L.) native to Turkey. *J. Agric. Food Chem.* (2009) 57: 5244-5249.
- (24) Mzhavanadze VV. Chlorogenic acid from the leaves of blueberry (*Vaccinium arctostaphylos*). *Soobshch. Akad. Nauk Gruz. SSR.* (1971) 62: 189-192.
- (25) Mzhavanadze VV. Glycosides of kaempferol from leaves of bilberry (*Vaccinium arctostaphylos*). *Soobshch. Akad. Nauk Gruz. SSR.* (1971) 62: 445-447.
- (26) Mzhavanadze VV, Targamadze IL and Dranik LI. Polyphenols of leaves of *Vaccinium arctostaphylos*. *Khim. Prir. Soedin.* (1971) 7: 546.
- (27) Mzhavanadze VV, Targamadze IL and Dranik LI. Phenol compounds from the leaves of *Vaccinium arctostaphylos*. *Khim. Prir. Soedin.* (1972) 8: 124.
- (28) Mzhavanadze VV, Targamadze IL and Dranik LI. Phenolic compounds of unripe blueberry fruits *Vaccinium arctostaphylos*. *Soobshch. Akad. Nauk Gruz. SSR.* (1972) 68: 205-208.
- (29) Mzhavanadze VV, Targamadze IL and Dranik LI. Dynamics of the content of chlorogenic and p-coumaric acids in the leaves and fruits of blueberry *Vaccinium arctostaphylos*. *Soobshch. Akad. Nauk Gruz. SSR.* (1972) 66: 449-452.
- (30) Nickavar B and Amin G. Anthocyanins from *Vaccinium arctostaphylos* berries. *Pharm. Biol.* (2004) 42: 289-291.

This article is available online at <http://www.ijpr.ir>