



CrossMark  
click for updates

## Correction


**Cite this article:** Ruta M, Angielczyk KD, Fröbisch J, Benton MJ. 2013 Decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids. *Proc R Soc B* 280: 20132414. <http://dx.doi.org/10.1098/rspb.2013.2414>

# Decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids

Marcello Ruta, Kenneth D. Angielczyk, Jörg Fröbisch and Michael J. Benton

*Proc. R. Soc. B* 280, 20131071 (7 October 2013; Published online 14 August 2013)  
(doi:10.1098/rspb.2013.1071)

Figure 3*h* in the original paper contains an inaccuracy. The bottom right panel (3*h*) is identical to panel 3*g*. The correct panel 3*h*, describing the plot of the rarefied distance from founder, is reproduced here together with all other panels of figure 3. The caption of that figure is slightly altered.



**Figure 3.** (a–d) Comparisons between anomodont disparity (rarefied median values, grey circles) and diversity (white squares) through time intervals t1–t8; the disparity values are for the sum of ranges (a,c) and the sum of variances (b,d); the error bars around the diversity values are calculated as  $\pm\sqrt{N}$ , where  $N$  is the number of taxa in any given interval; (a,b) comparisons based on the number of taxa present in the phylogeny; (c,d) comparisons based on the total number of known anomodont taxa. (e–h) Distance from the founder (white rhombs), expressed as the average generalized Euclidean distance of taxa from *Biseridens* ('founder' taxon); (e,f) average distance of taxa binned by time intervals; (g,h) average distance of major groups. Vertical grey bars mark the Permian–Triassic boundary.