

Interleukin-1 and Tumor Necrosis Factor- α Trigger Restriction of Hepatitis B Virus Infection via a Cytidine Deaminase Activation-induced Cytidine Deaminase (AID)*

Received for publication, July 12, 2013, and in revised form, September 8, 2013. Published, JBC Papers in Press, September 11, 2013, DOI 10.1074/jbc.M113.501122

Koichi Watashi^{†1}, Guoxin Liang[§], Masashi Iwamoto[‡], Hiroyuki Marusawa[¶], Nanako Uchida[‡], Takuji Daito[‡], Kouichi Kitamura[§], Masamichi Muramatsu[§], Hirofumi Ohashi[‡], Tomoko Kiyohara[‡], Ryosuke Suzuki[‡], Jisu Li^{||}, Shuping Tong^{||}, Yasuhito Tanaka^{**}, Kazumoto Murata^{**}, Hideki Aizaki^{†2}, and Takaji Wakita[‡]

From the [†]Department of Virology II, National Institute of Infectious Diseases, Tokyo 162-8640, Japan, the [§]Department of Molecular Genetics, Kanazawa University Graduate School of Medical Science, Kanazawa 920-8640, Japan, the [¶]Department of Gastroenterology and Hepatology, Kyoto University Graduate School of Medicine, Kyoto 606-8507, Japan, the ^{||}Liver Research Center Rhode Island Hospital, Warren Alpert School of Medicine, Brown University, Providence, Rhode Island 02903, the ^{**}Department of Virology and Liver Unit, Nagoya City University Graduate School of Medicinal Sciences, Nagoya 467-8601, Japan, and the [‡]Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine, Ichikawa 272-8516, Japan

Background: Cytokines and host factors triggering innate immunity against hepatitis B virus (HBV) are not well understood.

Results: IL-1 and TNF α induced cytidine deaminase AID, an anti-HBV host factor, and reduced HBV infection into hepatocytes.

Conclusion: IL-1/TNF α reduced host susceptibility to HBV infection through AID up-regulation.

Significance: Proinflammatory cytokines modulate HBV infection through a novel innate immune pathway involving AID.

Virus infection is restricted by intracellular immune responses in host cells, and this is typically modulated by stimulation of cytokines. The cytokines and host factors that determine the host cell restriction against hepatitis B virus (HBV) infection are not well understood. We screened 36 cytokines and chemokines to determine which were able to reduce the susceptibility of HepaRG cells to HBV infection. Here, we found that pretreatment with IL-1 β and TNF α remarkably reduced the host cell susceptibility to HBV infection. This effect was mediated by activation of the NF- κ B signaling pathway. A cytidine deaminase, activation-induced cytidine deaminase (AID), was up-regulated by both IL-1 β and TNF α in a variety of hepatocyte cell lines and primary human hepatocytes. Another deaminase APOBEC3G was not induced by these proinflammatory cytokines. Knockdown of AID expression impaired the anti-HBV effect of IL-1 β , and overexpression of AID antagonized HBV infection, suggesting that AID was one of the responsible factors for the anti-HBV activity of IL-1/TNF α . Although AID induced hypermutation of HBV DNA, this activity was dispensable for the anti-

HBV activity. The antiviral effect of IL-1/TNF α was also observed on different HBV genotypes but not on hepatitis C virus. These results demonstrate that proinflammatory cytokines IL-1/TNF α trigger a novel antiviral mechanism involving AID to regulate host cell permissiveness to HBV infection.

The intracellular immune response can eliminate pathogens from a host, and host cells possess different mechanisms to counteract viral infection depending on the virus type. Human immunodeficiency virus (HIV) infection is restricted by cellular proteins designated as restriction factors, including APOBEC3G (A3G),³ TRIM5 α , tetherin/BST-2, and SAMHD1 (1, 2). All of these factors can be induced by stimulation with interferon (IFN). Hepatitis C virus (HCV) is eliminated by type I and III IFNs derived from dendritic cells or infected hepatocytes (3–6). In hepatocytes, this process involves a series of antiviral factors that are downstream genes of IFN, IFN-stimulated genes (ISGs). Influenza virus spread and virulence is inhibited by cytokines such as IFNs and TNF α . Responsive genes for these mechanisms include IFN-induced cellular Mx proteins that are dynamin-like GTPases (7, 8). However, these cytokine-induced antiviral immune responses are poorly understood in hepatitis B virus (HBV) infection.

* This work was supported by grants-in-aid from the Ministry of Health, Labor, and Welfare, Japan, the Ministry of Education, Culture, Sports, Science, and Technology, Japan, and the Japan Society for the Promotion of Science and incentive support from the Viral Hepatitis Research Foundation of Japan.

✂ Author's Choice—Final version full access.

¹ To whom correspondence may be addressed: Dept. of Virology II, National Institute of Infectious Diseases, Tokyo, 1-23-1 Toyama, Shinjuku-ku, Tokyo, 162-8640, Japan. Tel.: 81-3-5285-1111; Fax: 81-3-5285-1161; E-mail: kwatashi@nih.go.jp.

² To whom correspondence may be addressed: Dept. of Virology II, National Institute of Infectious Diseases, Tokyo, 1-23-1 Toyama, Shinjuku-ku, Tokyo, 162-8640, Japan. Tel.: 81-3-5285-1111; Fax: 81-3-5285-1161; E-mail: aizaki@nih.go.jp.

³ The abbreviations used are: A3G, APOBEC3G; AID, activation-induced cytidine deaminase; HBV, hepatitis B virus; HCV, hepatitis C virus; ISG, IFN-stimulated gene; QNZ, 6-amino-4-(4-phenoxyphenylethylamino)quinazolinone; GEq, genome equivalent; PHH, primary human hepatocyte; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; ISRE, interferon sensitivity-responsive element; cccDNA, covalently closed circular DNA.

Anti-HBV Activity of IL-1 and TNF α Mediated by AID

HBV infection is a worldwide health problem affecting more than 350 million people and is a major cause of the development of liver cirrhosis and hepatocellular carcinoma (9–11). During the course of infection, a number of cytokines and chemokines are up-regulated in HBV-infected patients, including IFN α / γ / λ , TNF α , IL-1, IL-6, IL-10, IL-12, IL-15, and IL-8 (12–15). Some of these cytokines are reported to suppress HBV replication (3, 16–21). In particular, type I, II, and III IFNs suppress the replication of HBV *in vitro* and *in vivo* (19, 20, 22–26). Although one of the downstream genes of IFN, A3G, has the potential to reduce HBV replication (27–34), it is still under discussion whether this protein is responsible for the anti-HBV activity of type I IFN, because it has been previously reported by Trono and co-workers (28, 35) that the induction of A3G does not explain the IFN-induced inhibition of HBV replication. Moreover, these studies were carried out using an HBV transgene that only reproduces a portion of the whole HBV life cycle, mainly focusing on intracellular HBV replication.

Here, we screened for cytokines and chemokines that affected HBV infection in HepaRG cells, a human hepatocyte cell line susceptible to HBV infection and reproducing the whole HBV life cycle (36, 37). IL-1 and TNF α decreased the host cell permissiveness to HBV infection, and this effect was at least partly mediated by the induction of activation-induced cytidine deaminase (AID). The anti-HBV activity of IL-1/TNF α was mechanistically different from that of IFN α . This study presents the activity of IL-1/TNF α to suppress HBV infection into hepatocytes independent of the effect on immune cells and the physiological role of AID in this machinery. Moreover, as far as we know, this is the first report to show the AID function to inhibit the infection of human pathogenic virus.

EXPERIMENTAL PROCEDURES

Reagents—All cytokines were purchased from PeproTech or R & D Systems. Heparin was obtained from Mochida Pharmaceutical. Lamivudine, PD98059, SP600125, SB203580, and Bay11-7082 were obtained from Sigma. Entecavir was obtained from Santa Cruz Biotechnology. BMS-345541 and 6-amino-4-(4-phenoxyphenylethylamino)quinazoline (QNZ) were purchased from Merck.

Cell Culture—HepaRG cells (Biopredic) were cultured with Williams' medium E (Invitrogen) supplemented with 2 mM L-glutamine, 200 units/ml penicillin, 200 μ g/ml streptomycin, 10% FBS, 5 μ g/ml insulin (Wako), 20 ng/ml EGF (PeproTech), 50 μ M hydrocortisone (Sigma), and 2% DMSO (Sigma). HepG2, HepAD38 (kindly provided by Dr. Seeger at Fox Chase Cancer Center) (38), and HepG2.2.15 cells (a kind gift from Dr. Urban at Heidelberg University) (39) were cultured with DMEM/F-12 + GlutaMAX (Invitrogen) supplemented with 10 mM HEPES (Invitrogen), 200 units/ml penicillin, 200 μ g/ml streptomycin, 10% FBS, 50 μ M hydrocortisone, and 5 μ g/ml insulin in the presence (HepAD38 and HepG2.2.15) or absence (HepG2) of 400 μ g/ml G418 (Nacalai Tesque). HepAD38 cells were cultured with 0.3 μ g/ml tetracycline when terminating HBV induction. Huh-7.5.1 cells (kindly provided from Dr. Chisari at Scripps Research Institute) were cultured as described previously (40). Primary human hepatocytes (PHH) isolated from urokinase-type plasminogen activator transgen-

ic/SCID mice inoculated with PHH (PhoenixBio) or purchased from Lonza were cultured with DMEM supplemented with 20 mM HEPES, 100 units/ml penicillin, 100 μ g/ml streptomycin, 10% FBS, and 44 mM NaHCO₃ or with 1 mM pyruvate, nonessential amino acids, 20 mM HEPES, 200 units/ml penicillin, 200 μ g/ml streptomycin, 10% FBS, 0.25 μ g/ml insulin (Wako), 5 ng/ml EGF, and 50 nM dexamethasone.

HBV Preparation and Infection—HBV used in this study was mainly derived from HepAD38 cells, which is classified as genotype D (38). Media from HepAD38 cells at days 7–31 post-induction of HBV by depletion of tetracycline were recovered every 3 days. Media were cleared through a 0.45- μ m filter and precipitated with 10% PEG8000 and 2.3% NaCl. The precipitates were washed and resuspended with medium at \sim 200-fold concentration. The HBV DNA was quantified by real time PCR. HBV genotype A and C in Fig. 7B was recovered from the media of HepG2 cells transfected with the plasmid pHBV/Aeus and pHBV/C-AT (41).

HepaRG cells were infected with HBV at 2000 (Fig. 7B) or 6000 (other figures) genome equivalent (GEq)/cell in the presence of 4% PEG8000 for 16 h as described previously (36). Urban and co-workers (42) reported that more than 10³ GEq/cell amount of HBV derived from HepAD38 or HepG2.2.15 cells (*i.e.* 1.25–40 \times 10⁴ GEq/cell) as inoculum was required for efficient infection into HepaRG cells. The anti-HBV effect of IL-1/TNF α shown in this study was also observed when inoculated with HBV at 300 GEq/cell (data not shown).

Extraction of DNA and RNA—HBV DNA was extracted from the cells or from the medium using a DNA kit (Qiagen) according to the manufacturer's protocol. Total RNA was recovered with RNeasy mini kit (Qiagen) according to the manufacturer's protocol.

Real Time PCR and RT-PCR—HBV DNA was quantified by real time PCR analysis using the primer set 5'-ACTCACC-AACCTCCTGTCCT-3' and 5'-GACAAACGGGCAACAT-ACCT-3' and probe 5'-carboxyfluorescein (FAM)-TATCG-CTGGATGTGTCTGCGGCGT-carboxytetramethylrhodamine (TAMRA)-3' (43). The PCR was performed at 50 $^{\circ}$ C for 2 min, 94 $^{\circ}$ C for 10 min, and 50 cycles of 94 $^{\circ}$ C for 15 s and 60 $^{\circ}$ C for 1 min. Detection of cccDNA was achieved using 5'-CGTCTGTGCCTTCTCATCTGC-3' and 5'-GCACAG-CTTGAGGCTTGAA-3' as primers and 5'-CTGTAGGC-ATAAATTGGT (MGB)-3' as a probe (44). This primer-probe set theoretically detected neither relaxed circular DNA nor HBV DNA integrated into host genome but can capture cccDNA as described previously (44). For quantification of cellular mRNA, cDNA was synthesized from extracted RNA using SuperScriptIII (Invitrogen), followed by PCR with TaqMan Gene Expression Master Mix (Applied Biosystems) and primer-probe set (TaqMan Gene Expression Assay, Applied Biosystems) or with Power SYBR Green PCR Master Mix (Applied Biosystems) and 5'-AAATGTC-CGCTGGGCTAAGG-3' and 5'-GGAGGAAGAGCAATT-CCACGT-3' as primers for AID.

RT-PCR was performed as described previously (45) using a one-step RNA PCR kit (Takara). Primers for amplifying each gene were as follows: 5'-CTCTGAGGTTTAGCATTTCA-3' and 5'-CTCCAGGTCCAAAATGAATA-3' for *cIAP*; 5'-GCA-

GATTTATCAACGGCTTT-3' and 5'-CAGTTTTCCACCA-CAACAAA-3' for XIAP; 5'-TAGCCAACATGCTCCTCACAGAC-3' and 5'-TCTTCTACCACTGGTTTCATGC-3' for ISG56; 5'-GCCTTTTCATCCAAATGGAATTC-3' and 5'-GAAATCTGTTCTGGGCTCATG-3' for PKR; and 5'-CCATGGAGAAGGCTGGGG-3' and 5'-CAAAGTTGTCATGGATGACC-3' for GAPDH, respectively.

ELISA—HBs protein was quantified by ELISA using plates incubated at 4 °C overnight with a sheep anti-HBs antibody at 1:5000 dilution (Maxisorp nunc-immuno plate, Nunc catalog no. 439454) followed by coating with 0.2% BSA, 0.02% NaN₃, 1× PBS at 4 °C until use. Samples were incubated with the plates for 2 h and after washing with TBST four times, horseradish peroxidase-labeled rabbit anti-HBs antibody was added for 2 h. The substrate solution (HCV core ELISA kit: Ortho) was reacted for 30 min before the A₄₅₀ values were measured.

Indirect Immunofluorescence Analysis—Indirect immunofluorescence analysis was performed essentially as described previously (45). After fixation with 4% paraformaldehyde and permeabilization with 0.3% Triton X-100, an anti-HBc antibody (DAKO, catalog no. B0586) was used as the primary antibody.

MTT Assay—The MTT assay was performed as described previously (46).

Immunoblot Analysis—Immunoblot analysis was performed as described previously (47). The polyclonal antibody against AID was generated using a peptide derived from AID protein as an immunogen as described previously for preparation of the anti-AID antibody 1 (48). The specificity of the antibody was described previously (48, 49).

Lentiviral Vector-mediated Gene Transduction—Lentivirus carrying shRNAs was prepared with 293T cells transfected with expression plasmids for HIV-1 Gag-Pol, VSV G, and shRNAs (sh-control, sh-cyclophilin A, sh-AID(1), sh-AID(2); Mission shRNA) (Sigma) with Lipofectamine 2000 (Invitrogen). Recovered lentiviral vector was transduced into HepaRG cells followed by selection with 1.5 μ g/ml puromycin. Lentivirus overexpressing AID, AID mutant, A3G, or the control lentivirus was recovered using expression plasmids for HIV-1 Gag-Pol, Rev, VSV G, and the corresponding expression vector as described previously (50).

Southern Blot Analysis—Southern blot was performed as described previously (41). After digestion of free nucleic acids with DNase I and RNase A, cell lysates were digested with proteinase K, and HBV DNA in the core particles was extracted with phenol/chloroform, followed by isopropyl alcohol precipitation. Probe was prepared by cutting pHBV/D-IND60 (41) with SacII and BspHI to generate a full-length HBV DNA probe and labeled with AlkPhos direct labeling reagents (GE Healthcare). Labeled bands were visualized with CDP-star detection reagent (GE Healthcare).

Quantification of Nucleocapsid-associated HBV RNA—After digestion of free nucleic acids with DNase I and RNase A, nucleocapsid was precipitated with PEG8000 (41). Total RNA was then extracted from the resuspended precipitates. HBV RNA was quantified by real time RT-PCR with 5'-TCCCTCGCCTCGCAGACG-3' and 5'-GTTTCCCACCTTAT-

GAGTC-3' as primers with Power SYBR Green PCR Master Mix (Applied Biosystems).

Co-immunoprecipitation Assay—Co-immunoprecipitation assay was essentially performed as described (45).

Differential DNA Denaturation PCR—Differential DNA denaturation PCR was performed as described previously (51).

Reporter Assay—DNA transfection was performed with pNF- κ B-luc or pISRE-TA-luc (Stratagene) and pRL-TK (Promega), which express firefly luciferase driven by NF- κ B or ISRE and *Renilla* luciferase by herpes simplex virus thymidine kinase promoter, respectively, and Polyethylenimine Max (Polysciences Inc., catalog no. 24765). After compound or cytokine treatment, cells were lysed, and luciferase activities were measured as described previously (52). A reporter carrying HBV core promoter was constructed by inserting the DNA fragment (1413–1788 nucleotide number) of HBV DNA (D-IND60) into pGL4.28 vector (Promega) (41). In the reporter assay using this construct (Fig. 1H), HX531, a retinoid X receptor antagonist was used as a positive control as retinoid X receptor was involved in the transcription from the core promoter (53).

RESULTS

IL-1 Reduced Host Cell Susceptibility to HBV Infection—To evaluate the effect of cytokines and chemokines on susceptibility to HBV infection, we treated HepaRG cells (36) with cytokines for 3 h prior to and 16 h during HBV infection, followed by culture without stimuli for an additional 12 days (Fig. 1A, lower scheme). Heparin, a competitive inhibitor of HBV attachment (54), was used as a positive control and decreased secretion of the viral envelope surface protein (HBs) from HBV-infected cells (Fig. 1A, upper graph, lane 38), which suggests a successful HBV infection in this experiment. Examination of 36 cytokines and chemokines revealed that IL-1 β drastically decreased protein secretion from HBs (Fig. 1A, upper graph, lane 8). Although IFNs had a strong anti-HBV effect by a continuous treatment after HBV infection (Fig. 3C, panel b, and data not shown), they had only a limited effect in this screening where cytokines were only pretreated and cotreated with HBV (Fig. 1A, lanes 2–7). HBc protein expression (Fig. 1B) and HBV DNA (Fig. 1C) in the cells and medium (Fig. 1D) were significantly decreased by treatment with IL-1 β without cytotoxicity (Fig. 1G). HBV cccDNA and HBV RNA was also decreased in infected cells treated with IL-1 β (Fig. 1, E and F). IL-1 β did not decrease HBV core promoter activity at least in HepG2 cells (Fig. 1H). These results suggest that IL-1 β suppressed HBV infection to HepaRG cells. IL-1 β did not decrease the expression of sodium taurocholate cotransporting polypeptide (NTCP), a recently reported HBV entry receptor (data not shown) (55). Similar results were obtained using primary human hepatocytes (Fig. 1I).

NF- κ B Signaling Was Critical for Anti-HBV Activity—As shown in Fig. 2A, IL-1 β suppressed HBV infection in a dose-dependent manner. This anti-HBV effect was reversed by cotreatment with a neutralizing antibody for the IL-1 receptor, IL-1RI (Fig. 2B), suggesting that receptor engagement was required for anti-HBV activity. IL-1Ra is a natural antagonist that associates with IL-1RI but does not trigger downstream signal transduc-

Anti-HBV Activity of IL-1 and TNF α Mediated by AID




FIGURE 1. Suppression of HBV infection by IL-1 β . *A*, upper graph, HepaRG cells were pretreated with cytokines at 100 ng/ml (except for IFN α and IFN β at 100 IU/ml) or heparin at 25 units/ml as a positive control or were left untreated (*control*) for 3 h and then infected with HBV in the presence of each stimuli for 16 h. After washing, cells were cultured in normal growth medium for 12 days. HBs protein secreted into the medium was quantified by ELISA. Lower scheme indicates the treatment procedure for HepaRG cells. Black and dashed line boxes indicate the periods with and without treatment, respectively. *B–G* and *I*, HepaRG cells (*B–G*) or PHH (*I*) were treated as shown in *A* with or without 100 ng/ml IL-1 β or 25 units/ml heparin as a positive control. HBc protein in the cells (red) was detected by indirect immunofluorescence analysis, and the nucleus was stained with DAPI (blue) at 12 days post-infection (*B*). HBV DNA (*C* and *I*), cccDNA (*E*), and HBV RNA (*F*) in the cells as well as HBV DNA in the medium (*D*) were detected. Cell viability was quantified by MTT assay (*G*). *HBV*(–) in *I* indicates uninfected cells. All of the data, except in *I*, are based on the average of three independent experiments. *I* shows the average results from one representative experiment, but the reproducibility of the data were confirmed in three independent experiments. *H*, reporter plasmid carrying the HBV core promoter was transfected with HepG2 cells and then treated with or without IL-1 β (1, 10, and 100 ng/ml) and an retinoid X receptor antagonist HX531 as a positive control for 6 h. Luciferase activity was measured.

tion (56). Treatment with IL-1Ra did not decrease HBV infectivity (Fig. 2C), suggesting that signal transduction triggered by IL-1 was required for anti-HBV activity.

To identify the signal transduction pathway essential for anti-HBV activity, we treated HepaRG cells with PD98059, SP600125, SB203580, and Bay11-7082, which are inhibitors for MEK, JNK, p38, and NF- κ B, respectively (57). As shown in Fig. 2D, only cotreatment with Bay11-7082 significantly removed the anti-HBV effect of IL-1 β . Luciferase assay and RT-PCR analysis indicated that Bay11-7082, but not other inhibitors, blocked the transactivation of NF- κ B (Fig. 2E, upper panels) and NF- κ B downstream genes, *cIAP* and *XIAP* (Fig. 2E, lower

panels). Additional NF- κ B inhibitors, BMS-345541 and QNZ (Fig. 2G), also reversed the anti-HBV effect of IL-1 β (Fig. 2F). These data suggest a critical role for NF- κ B activation in the anti-HBV activity. Additionally, IL-1 β did not augment the activity of interferon sensitivity-responsive element (ISRE) and mRNAs for ISGs, *ISG56*, and double-stranded RNA-dependent protein kinase (*PKR*) in HepaRG cells (Fig. 2H), suggesting that the anti-HBV activity is independent of ISG up-regulation. TNF α , another cytokine that activates NF- κ B signaling (Fig. 2E, lower panels), also inhibited HBV infection (Fig. 2I). Thus, NF- κ B activation in host hepatocytes was critical for the anti-HBV activity of proinflammatory cytokines.




FIGURE 2. **NF- κ B activation triggered by IL-1 and TNF α was critical for anti-HBV activity.** A–D, F, and I, HepaRG cells were left untreated (*control*) or treated with varying concentrations of IL-1 β (1, 10, 30, and 100 ng/ml) or 25 units/ml heparin (A), with 30 ng/ml IL-1 β together with or without a neutralizing anti-IL-1RI antibody at 20 μ g/ml (B), with 10 ng/ml IL-1 β or varying concentrations of IL-1Ra (10, 30, and 100 ng/ml) (C), with 3 ng/ml IL-1 β together with or without PD98059, SP600125, SB203580, or Bay11-7082 (D), or QNZ or BMS-345541 (F), or with TNF α (10, 100, and 300 ng/ml) (I) according to the treatment schedule shown in Fig. 1A. HBV infection was monitored by HBs protein secretion into the medium in A, C, D, F, and I and with Hbc protein in the cells in B. E, G, and H, NF- κ B (E and G) and ISRE activity (H) were measured by reporter assay in the cells transfected with the reporter plasmid expressing luciferase driven from five tandem repeats of NF- κ B elements (E, upper graph, and G) or ISRE (H, upper graph) or by RT-PCR in the cells (E and H, lower panels) upon signaling inhibitors used in D and F together with or without IL-1 β (E and G), or upon IL-1 β (10, 30, and 100 ng/ml) or IFN α 100 IU/ml as a positive control (H) for 6 h. The white and black bars in the upper graph of E and G show the data in the absence or presence of IL-1 β , respectively. Bands for mRNA for *clAP*, *XIAP*, and *GAPDH* (E) or *ISG56*, *PKR*, and *GAPDH* (H) are presented in the lower panels. All of the data are based on averages of three independent experiments.

Early Phase of HBV Infection as Well as HBV Replication Were Impaired by IL-1 Treatment—Although heparin, an attachment inhibitor, could block HBV infection only if added together with the HBV inoculum, pretreatment with IL-1 β before HBV infection was sufficient to show anti-HBV activity (Fig. 3A, panel b). This activity was amplified by a prolonged

treatment time of up to 12 h (Fig. 3B). Intriguingly, HBV cellular DNA was also reduced by IL-1 β treatment following HBV infection (Fig. 3C, panel b). In contrast, IFN α was not effective by pretreatment (Figs. 3C, panel a, and 1A), although it did decrease HBV DNA by treatment after HBV infection (Fig. 3C, panel b), consistent with previous reports that IFN α can sup-

Anti-HBV Activity of IL-1 and TNF α Mediated by AID




FIGURE 3. Defining the steps of the HBV life cycle targeted by IL-1 β . A, HepaRG cells were pretreated with IL-1 β or heparin for 3 h and then infected with HBV in the presence (A, panel a) or absence (A, panel b) of IL-1 β or heparin for 16 h. HBV infection was monitored with HBs protein secretion from the infected cells. Only pretreatment with IL-1 β and not heparin could inhibit HBV infectivity. *d*, day. B, HepaRG cells were pretreated with IL-1 β or left untreated (–) for the indicated time (h) and infected with HBV without IL-1 β . Anti-HBV activity was amplified by a prolonged treatment time. C, panel a, HepaRG cells were pretreated with 10 ng/ml IL-1 β , 100 IU/ml IFN α , or 1 μ M lamivudine for 3 h, followed by infection with HBV for 16 h in the absence of cytokines (pretreatment). C, panel b, HepaRG cells were infected with HBV for 16 h without pretreatment. After washing out the input virus, cells were cultured in normal medium for the first 8 days and then cultured with IL-1 β , IFN α , or lamivudine for the following 4 days (post-treatment). HBV DNA in the cells was measured by real time PCR. IL-1 β showed an anti-HBV activity in both pretreatment and post-treatment, although an anti-HBV effect of IFN α was seen only with post-treatment. D, HepAD38 cells were treated with 100 ng/ml IL-1 β or 1 μ M lamivudine, or left untreated for 6 days in the absence of tetracycline. HBV replication was evaluated by measurement of HBV DNA in the medium. E, HepaRG cells were pretreated with IL-1 β , lamivudine, or heparin for 3 h or left untreated and infected with HBV for 16 h in the presence or absence of each compound. After trypsinization and extensive washing of the cells, cellular DNA was immediately recovered to detect HBV DNA. HBV DNA at 16 h post-infection was decreased by treatment with IL-1 β but not lamivudine.

press HBV replication (19, 20, 26). Thus, the anti-HBV activity of IL-1 β is likely to be mechanistically different from that of IFN α .

The HBV life cycle can be divided into at least two phases as follows: 1) the early phase of infection that includes attachment, entry, nuclear import, and cccDNA formation; and 2) the late phase representing HBV replication, including transcription, assembly, reverse transcription, DNA synthesis, and viral release (58). The early phase of HBV infection is not supported, but HBV DNAs persistently replicate in HepAD38 cells in the presence of tetracycline (38). IL-1 β decreased the HBV DNA levels in HepAD38 cells (Fig. 3D), suggesting suppression of HBV replication. In addition, to examine the early phase preceding HBV replication, we infected HepaRG cells with HBV in the presence of IL-1 β for 16 h and then immediately recovered cellular DNA in the trypsinized cells for quantification of HBV DNA (Fig. 3E). This procedure likely detected HBV DNA that had been internalized and evaded the host restriction before initiation of HBV replication because lamivudine showed no effect on the amount of DNA detected (Fig. 3E). In this experiment, IL-1 β significantly decreased HBV DNA (Fig. 3E). cccDNA was also decreased by IL-1 β , suggesting that the early phase of HBV infection before cccDNA formation was also interrupted by IL-1 β .

IL-1 and TNF α Induced the Expression of AID—The innate immune pathway against HBV infection remains largely unknown. Recently, accumulating evidence suggested that several APOBEC family proteins, especially A3G, suppressed HBV replication when overexpressed (27–33). In contrast, there was no report available suggesting the anti-HBV function of other restriction factors against HIV, TRIM5 α , tetherin/BST-2, and SAMHD1. We then investigated APOBEC family proteins as a candidate for an anti-HBV effector. The APOBEC family includes APOBEC1 (A1), A2, A3s, A4, and AID (59). Because some of these proteins are reported to be up-regulated in cytokine-stimulated hepatocytes (27, 28, 60, 61), we examined the expression of these genes in cells treated with IL-1 β , TNF α , and IFN α as a control for 12 h. The mRNA levels of A1, A2, and A3A were below the detection threshold. A3G and A3F mRNA were significantly expressed in HepaRG cells, and their expression levels were remarkably increased by IFN α treatment (Fig. 4A), as observed in other reports (27, 28, 61). IL-1 β and TNF α did not significantly up-regulate A3s, and only AID was up-regulated 6–10-fold by both cytokines (Fig. 4A). Induction of A3s by both IL-1 β and TNF α was not observed at any time point examined until 12 h (data not shown). In contrast, induction of AID mRNA by IL-1 β and TNF α was conserved in human hepatocyte cell lines, such as HepG2 and FLC4 cells, and in primary human hepatocytes (Fig. 4B). AID protein production was also increased in primary human hepatocytes by treatment with IL-1 β and TNF α (Fig. 4C). This AID induction by IL-1 β was suggested to be NF- κ B-dependent, because the up-regulation of AID mRNA was canceled by addition of NF- κ B inhibitors, Bay11-7082 or QNZ (Fig. 4D).

AID Played a Significant Role in the IL-1-mediated restriction of HBV—To examine the function of AID during HBV infection, we transduced AID ectopically into HepaRG cells using a lentiviral vector (Fig. 5A, left panel). The susceptibility of these

AID-overexpressing cells to HBV was decreased by approximately one-third compared with the parental or empty vector-transduced HepaRG cells (Fig. 5A, right panel), suggesting that AID can restrict HBV infection. An AID mutant AID(M139V), with reported diminished activity to support class switching (48), also decreased the susceptibility to HBV infection, although the reduction in HBV susceptibility was moderate compared with the case of the wild type AID (Fig. 5B).

To examine the relevance of endogenous AID in the anti-HBV activity of IL-1, we transduced a lentiviral vector carrying a short hairpin RNA (shRNA) against AID (sh-AID) or a non-relevant protein cyclophilin A (Fig. 5C), and we observed the anti-HBV activity of IL-1 β in these cells. IL-1 β decreased HBV infection in the control and sh-cyclophilin A-transduced cells by \sim 3.0-fold as determined by HBs secretion (Fig. 5D, lanes 1 and 2, black bars). In contrast, anti-HBV activity of IL-1 β was limited to only 1.6–1.7-fold in the cells transduced with sh-AIDs (Fig. 5D, lanes 3 and 4, black bars). Such relieved anti-HBV activity following AID knockdown was not observed in the case for heparin treatment (Fig. 5D, lanes 1–4, gray bars). Similar results were obtained by monitoring intracellular HBV DNA after infection (data not shown). Although the anti-HBV effect of IL-1 β was not completely blunted, these data suggest that AID plays a significant role in mediating the anti-HBV effect of IL-1 β .

Similar observations were obtained in HBV-replicating cells overexpressing AID (Fig. 5, E and F). Core particle-associated HBV DNA in HepG2 cells transfected with an HBV-encoding plasmid was decreased by overexpression with AID as well as with A3G (Fig. 5E, lanes 1 and 3). Intriguingly, HBV DNA in core particles was also decreased by expression of an AID mutant AID(H56Y), which contains a mutation in the cytidine deaminase motif and is derived from a class switch deficiency patient (Fig. 5E, lane 2) (48). Southern blot also showed that the HBV rcDNA level in HepG2.2.15 cells was reduced by transduction with AID and another mutant AID(M139V), with diminished activity to support class switching (Fig. 5F) (48). These data suggest that AID could suppress HBV replication, and this restriction activity can be still observed with reduced enzymatic activity. In addition, AID was shown to interact with HBV core protein by coimmunoprecipitation assay (Fig. 5G). Moreover, overexpression of AID reduced the levels for nucleocapsid-associated HBV RNA (Fig. 5H). These results further suggest an antiviral activity of AID against HBV replication.

AID Could Induce Hypermutation of HBV DNA—Major enzymatic activity for APOBEC family proteins is the introduction of hypermutation in target DNA/RNA, and hypermutation accounts for antiviral activity for A3G against HIV-1 to some extent (2). Several groups reported that APOBEC family proteins could induce hypermutation in HBV DNA (27, 30, 32, 34). Next we asked whether AID could induce hypermutations in HBV DNA. In differential DNA denaturation PCR analysis, a high content of A/T bases introduced by hypermutation decreased denaturation temperatures (51). As shown in Fig. 6A, ectopic expression of AID decreased the denaturation temperature of HBV DNA as shown by that of A3G. Sequence analyses of the HBV DNA X region amplified at 83 °C by differential DNA denaturation PCR indicated a massive accumulation of

Anti-HBV Activity of IL-1 and TNF α Mediated by AID




FIGURE 4. AID expression was induced by IL-1 β and TNF α . *A*, mRNAs for A3B, -C, -D, -F, -G, -H and AID were quantified by real time RT-PCR analysis in HepaRG cells treated with 100 ng/ml IL-1 β , 100 ng/ml TNF α , or 100 IU/ml IFN α for 12 h or left untreated. *Graphs* show the relative expression levels compared with the controls set at 1. *B*, AID mRNA was detected in HepG2, FLC4 cells, and PHH treated with IL-1 β , TNF α , or IFN α or left untreated. Induction of AID by IL-1 β and TNF α was observed in HepG2 and FLC4 cells and primary human hepatocytes. *C*, AID protein (upper panel) and actin levels as an internal control (lower panel) were examined by immunoblot of primary human hepatocytes treated with IL-1 β or TNF α or left untreated. *D*, AID mRNA was detected in PHH treated with 100 ng/ml IL-1 β in the presence or absence of NF- κ B inhibitors, Bay11-7082, or QNZ for 12 h.

G-to-A mutations by AID (Fig. 6B). The frequency of G-to-A mutations was augmented by AID expression (Fig. 6C). In this experiment, AID(IP8Bdel), a hyper-active mutant of AID (62), further promoted the accumulation of the G-to-A and C-to-T mutations, although AID(H56Y) showed mutations in HBV DNA equivalent with mock GFP control sample (Fig. 6C). Thus, AID had the potential to introduce hypermutation in nucleocapsid-associated HBV DNA.

IL-1 Suppressed the Infection of Different HBV Genotypes but Not That of HCV—We examined whether the antiviral activity of IL-1 β and TNF α could be generalized to other viruses or was specific to HBV. As shown in Fig. 7A, the production of infec-

tious HCV and HCV core proteins in the medium was not significantly altered by treatment with these cytokines in HCV-infected cells, compared to when IFN α was used as a positive control (Fig. 7A). In contrast, IL-1 suppressed the infection of HBV genotype A and C into HepaRG cells (Fig. 7B) as well as genotype D (Fig. 1C). These data suggest that the antiviral activity of proinflammatory cytokines IL-1 and TNF α is specific to HBV.

DISCUSSION

In this study, cytokine screening revealed that IL-1 and TNF α decreased the host cell susceptibility to HBV infection.




FIGURE 5. AID played a significant role in IL-1-mediated anti-HBV activity. *A and B, left panels*, HepaRG cells were transduced with a lentiviral vector carrying the expression plasmid for AID (RG-AID), AID(M139V) mutant (RG-AID(M139V)) (B), or the control vector (RG-EV). Protein expression for AID (upper panel) and actin (lower panel) in these cells, the parental HepaRG cells (HepaRG), and those transiently transfected with AID expression plasmid (AID overexpression) (A) was examined by immunoblot. *Right panels*, these cells were infected with HBV followed by detection of secreted HBs protein as Fig. 1A. AID-transduced cells were less susceptible to HBV infection. *C*, HepaRG cells were transduced with lentiviral vector carrying shRNAs for AID (RG-shAID#1 and RG-shAID#2) or for cyclophilin A (RG-shCyPA) as a control. AID mRNA (left panel) and protein (right panel) were quantified by real time RT-PCR and immunoblot analysis. *D*, cells produced in *C* were infected with HBV in the absence or presence of IL-1 β or heparin, and HBs was detected in the medium as in Fig. 1A to examine the anti-HBV effect of IL-1 β and heparin. The fold reduction of HBV infection by IL-1 β treatment is shown as IL-1 β anti-HBV above the graph. The white, gray, and black bars indicate HBs value of the cells without treatment and with heparin and IL-1 β treatment, respectively. The anti-HBV activity of IL-1 β but not heparin was reduced in the AID-transduced cells. *E*, AID and its mutant suppressed HBV replication. HepG2 cells were cotransfected with GFP-tagged AID, AID(H56Y), A3G, and GFP itself along with an HBV-encoding plasmid. Following 3 days, cytoplasmic nucleocapsid HBV DNA was quantified (upper graph), and the overexpressed proteins as well as actin were detected (lower panels). *F*, lentiviral vectors carrying AID, AID(M139V) mutant, A3G, or an empty vector (empty vector) were transduced or left untransduced (no transduction) into HepG2.2.15 cells. Nucleocapsid associated HBV DNA in these cells or in HepG2 cells (HBV-) was detected by Southern blot (upper panel). AID (middle panel) and A3G protein (lower panel) were also detected by immunoblot. *G*, HBV core interacted with AID. HepAD38 cells transduced without (no transduction) or with AID-expressing vector or the empty vector (empty vector) were lysed and treated with anti-core antibody (1st panel) or control normal IgG (2nd panel) for immunoprecipitation (IP). Total fraction without immunoprecipitation (3rd to 5th panels) was also recovered to detect AID (1st to 3rd panels), HBV core (5th panel), and actin (5th panel) by immunoblot. WB, Western blot. *H*, HBV RNA in core particles was extracted as shown under "Experimental Procedures" in HepG2 cells overexpressing HBV DNA together with or without AID or A3G.

Anti-HBV Activity of IL-1 and TNF α Mediated by AID




FIGURE 6. AID could induce hypermutation of HBV DNA. *A* and *B*, HepG2 cells were cotransfected with an expression vector for GFP-tagged AID, HA-tagged A3G, or GFP along with an HBV-encoding plasmid. 3 days after transfection, nucleocapsid-associated HBV DNA was extracted, and differential DNA denaturation PCR was performed to amplify the X gene segments. The numbers above the panels in *A* show denaturing temperatures. The X gene fragment amplified at 83 °C in the AID sample was cloned in to a T vector and sequenced in *B*. Alignment of independent five clones with reference sequence (X02763) is indicated. *C*, AID and its mutant (JP8Bdel) induced G-to-A and C-to-T hypermutations in HBV DNA. HepG2 cells were transfected with expression vectors of GFP-tagged AID, AID(H56Y), AID(JP8Bdel), or GFP itself together with HBV encoding plasmid. Three days after transfection, cells were harvested, and nucleocapsid-associated HBV DNA was extracted. X gene fragments were amplified at 94 °C and cloned in T vector. 55 clones were sequenced as described under "Experimental Procedures." The numbers indicate the clone numbers carrying the mutation. *D*, expression of GFP, GFP-tagged AID, AID(H56Y), and AID(JP8Bdel) is shown by immunoblot.

This antiviral mechanism is rather unique, given that the intracellular immune response against viruses is typically triggered by IFNs. So far, type I, II, and III IFNs are reported to suppress the replication step of the HBV life cycle (19, 20, 25, 26). In contrast, we suggest that IL-1 and TNF α inhibit the early phase of HBV infection as well as the replication. This is consistent with cumulative clinical evidence suggesting that these proinflammatory cytokines contribute to HBV elimination (63–65).

IL-1 and TNF α are generally produced mainly in macrophages and also in other cell types, including T cells and endothelial cells (66). Although the main producer cells of these cytokines in hepatitis B patients are not defined, it has been reported that the secretion of IL-1 and TNF α in nonparenchymal cells were increased by HBV infection into hepatocytes (67). TNF α production in macrophages was augmented by addition of recombinant Hbc (68). A number of clinical studies cumulatively




FIGURE 7. **Antiviral activity of AID was specific to HBV.** A, Huh-7.5.1 cells were pretreated with IL-1 β , TNF α , or IFN α for 3 h or left untreated and then cocultured with HCV for 4 h. After washing HCV and cytokines and culturing the cells with normal medium for 72 h, the infectivity of HCV (left panel) as well as HCV core protein (right panel) in the medium was quantified. B, HepaRG cells were treated with IL-1 β or heparin or left untreated for 3 h prior to and 16 h during infection of HBV genotype A (left graph) or C (right graph) as shown in Fig. 1A. HBV infection was monitored with cellular HBV DNA at 12 days after the infection as Fig. 1C.

show that serum levels of IL-1 and TNF α are increased in hepatitis B patients (12). Recently, it has been a significant clinical problem that HBV reactivates during the course of treatment with immunosuppressants such as anti-TNF α agents (64, 65). Taken together, it is proposed that acute or chronic HBV infection induces IL-1/TNF α from macrophages or other cells in the liver of infected patients, which can directly suppress HBV infection in hepatocytes, in addition to their immunomodulatory effects to the host immune cells. Although IL-1 level in HBV-infected patients varies between papers, Daniels *et al.* (63) reported that the peak IL-1 β level in HBV-infected patients was 9–36 ng/ml under Toll-like receptor stimulation, at which concentration IL-1 β showed significant anti-HBV effects in this study. In general, downstream genes of NF- κ B include a number of antiviral factors such as *viperin*, *iNOS*, and *RANTES* (69). Although some of these genes may function cooperatively for IL-1- and TNF α -induced anti-HBV machinery, our data suggest that AID, at least in part, plays a role in the elimination of HBV that was potentiated by proinflammatory cytokines IL-1 and TNF α .

AID belongs to APOBEC family proteins that share enzyme activity to convert cytidine to uracil in mainly DNA, and occasionally RNA (51, 70, 71). Although AID was initially identified in B cells, chronic inflammation can trigger its expression in hepatocytes (60). The induction of AID was reportedly mediated by NF- κ B (60), consistent with the results in this study. Although AID in B cells is essential for class switch recombination and somatic hypermutation of immunoglobulin genes (70, 72), the physiological role of AID in hepatocytes is unknown.

Although expression of AID in hepatocytes is still lower than in B cells, AID is reportedly expressed in the liver both in cell culture and *in vivo* settings (34, 60). Our results raise the idea that AID plays a role in innate antiviral immunity. AID also has a role in virus-induced pathogenesis as it was reported to counteract oncogenesis induced by Abelson-murine leukemia virus (73). In addition, AID was reported to restrict L1 retrotransposition, which can predict the role of AID in innate immunity (74). This study is significant in that it revealed a biological function of AID in viral infection itself, linking it to the restriction of a pathogenic human virus. It will be interesting to analyze the role of AID in the infection process of other viruses in the future.

Although the mechanism for AID suppression of the HBV life cycle is the subject of future study, AID possibly targets the early phase of HBV infection, including entry as well as the replication stage, including assembly and reverse transcription (Fig. 3). It has been recently reported that chicken AID reduced cccDNA of duck HBV possibly through targeting cccDNA as well as nucleocapsid-associated HBV DNA (75). This study is likely to support the idea that AID may target cccDNA formed after HBV entry into hepatocytes, and also associates with nucleocapsid-associated HBV DNA during HBV replication, although it is not clear whether the innate immune machinery against HBV/duck HBV is conserved in human and chicken cells. A3G blocked HBV replication through the inhibition of reverse transcriptase (29), packaging of pregenomic RNA (33), and the destabilization of packaged pregenomic RNA (31) independently of its deaminase activity, and it also induced hypermutation of HBV DNA (27, 30, 32, 34). It was recently reported that AID was packaged into the HBV nucleocapsid (51). Moreover, AID induced C-to-T and G-to-A hypermutations in HBV DNA/RNA, although the anti-HBV activity has not been demonstrated so far (51). The hypermutation activity of AID was likely to be dispensable for its anti-HBV replication function (Figs. 5 and 6), as reported for APOBEC3G by several groups (29, 30, 33). Further analysis is required to elucidate the precise mechanisms for AID-mediated suppression of the HBV life cycle.

In conclusion, we have identified that host cell susceptibility to HBV infection is modulated by IL-1 and TNF α , and AID is involved in this machinery. This sheds new light on the link between proinflammatory cytokines and the development of the innate antiviral defense.

Acknowledgments—HepAD38, HepG2.2.15, and Huh-7.5.1 cells were kindly provided by Dr. Seeger at Fox Chase Cancer Center, Dr. Urban at Heidelberg University, and Dr. Chisari at Scripps Research Institute. We are grateful to M. Matsuda, T. Date, T. Mizoguchi, Y. Hirama, M. Sasaki, H. Aoyagi, and S. Nakajima for technical and secretarial assistance. We also thank Dr. Ishida at PhoenixBio, Dr. Sugiyama at National Center for Global Health and Medicine, and all of the members of the Department of Virology II, National Institute of Infectious Diseases, for their helpful discussions.

REFERENCES

1. St Gelais, C., and Wu, L. (2011) SAMHD1: a new insight into HIV-1 restriction in myeloid cells. *Retrovirology* 8, 55

2. Strebel, K., Luban, J., and Jeang, K. T. (2009) Human cellular restriction factors that target HIV-1 replication. *BMC Med.* **7**, 48
3. Bertoletti, A., and Ferrari, C. (2003) Kinetics of the immune response during HBV and HCV infection. *Hepatology* **38**, 4–13
4. Lemon, S. M. (2010) Induction and evasion of innate antiviral responses by hepatitis C virus. *J. Biol. Chem.* **285**, 22741–22747
5. Saito, T., and Gale, M., Jr. (2007) Principles of intracellular viral recognition. *Curr. Opin. Immunol.* **19**, 17–23
6. Saito, T., and Gale, M., Jr. (2008) Regulation of innate immunity against hepatitis C virus infection. *Hepatology Res.* **38**, 115–122
7. Haller, O., Stertz, S., and Kochs, G. (2007) The Mx GTPase family of interferon-induced antiviral proteins. *Microbes Infect.* **9**, 1636–1643
8. Seo, S. H., and Webster, R. G. (2002) Tumor necrosis factor α exerts powerful anti-influenza virus effects in lung epithelial cells. *J. Virol.* **76**, 1071–1076
9. Loomba, R., and Liang, T. J. (2007) Treatment of chronic hepatitis B. *Antivir. Ther.* **12**, Suppl. 3, H33–H41
10. Pawlowsky, J. M., Dusheiko, G., Hatzakis, A., Lau, D., Lau, G., Liang, T. J., Locarnini, S., Martin, P., Richman, D. D., and Zoulim, F. (2008) Virologic monitoring of hepatitis B virus therapy in clinical trials and practice: recommendations for a standardized approach. *Gastroenterology* **134**, 405–415
11. Waris, G., and Siddiqui, A. (2003) Regulatory mechanisms of viral hepatitis B and C. *J. Biosci.* **28**, 311–321
12. Ait-Goughoulte, M., Lucifora, J., Zoulim, F., and Durantel, D. (2010) Innate antiviral immune responses to hepatitis B virus. *Viruses* **2**, 1394–1410
13. Kakumu, S., Fuji, A., Yoshioka, K., and Tahara, H. (1989) Serum levels of α -interferon and γ -interferon in patients with acute and chronic viral hepatitis. *Hepatogastroenterology* **36**, 97–102
14. Protzer, U., Maini, M. K., and Knolle, P. A. (2012) Living in the liver: hepatic infections. *Nat. Rev. Immunol.* **12**, 201–213
15. Rossol, S., Marinos, G., Carucci, P., Singer, M. V., Williams, R., and Naoumov, N. V. (1997) Interleukin-12 induction of Th1 cytokines is important for viral clearance in chronic hepatitis B. *J. Clin. Invest.* **99**, 3025–3033
16. Cavanaugh, V. J., Guidotti, L. G., and Chisari, F. V. (1997) Interleukin-12 inhibits hepatitis B virus replication in transgenic mice. *J. Virol.* **71**, 3236–3243
17. Guo, J. T., Zhou, H., Liu, C., Aldrich, C., Saputelli, J., Whitaker, T., Barrasa, M. I., Mason, W. S., and Seeger, C. (2000) Apoptosis and regeneration of hepatocytes during recovery from transient hepadnavirus infections. *J. Virol.* **74**, 1495–1505
18. McClary, H., Koch, R., Chisari, F. V., and Guidotti, L. G. (2000) Relative sensitivity of hepatitis B virus and other hepatotropic viruses to the antiviral effects of cytokines. *J. Virol.* **74**, 2255–2264
19. Pagliaccetti, N. E., Chu, E. N., Bolen, C. R., Kleinstein, S. H., and Robek, M. D. (2010) λ and α interferons inhibit hepatitis B virus replication through a common molecular mechanism but with different *in vivo* activities. *Virology* **401**, 197–206
20. Robek, M. D., Boyd, B. S., and Chisari, F. V. (2005) λ interferon inhibits hepatitis B and C virus replication. *J. Virol.* **79**, 3851–3854
21. Thompson, A. J., Colledge, D., Rodgers, S., Wilson, R., Revill, P., Desmond, P., Mansell, A., Visvanathan, K., and Locarnini, S. (2009) Stimulation of the interleukin-1 receptor and Toll-like receptor 2 inhibits hepatitis B virus replication in hepatoma cell lines *in vitro*. *Antivir. Ther.* **14**, 797–808
22. Bertoletti, A., Maini, M. K., and Ferrari, C. (2010) The host-pathogen interaction during HBV infection: immunological controversies. *Antivir. Ther.* **15**, Suppl. 3, 15–24
23. Chang, J., Block, T. M., and Guo, J. T. (2012) The innate immune response to hepatitis B virus infection: Implications for pathogenesis and therapy. *Antivir. Res.* **96**, 405–413
24. Guidotti, L. G., Ando, K., Hobbs, M. V., Ishikawa, T., Runkel, L., Schreiber, R. D., and Chisari, F. V. (1994) Cytotoxic T lymphocytes inhibit hepatitis B virus gene expression by a noncytolytic mechanism in transgenic mice. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 3764–3768
25. Guidotti, L. G., Ishikawa, T., Hobbs, M. V., Matzke, B., Schreiber, R., and Chisari, F. V. (1996) Intracellular inactivation of the hepatitis B virus by cytotoxic T lymphocytes. *Immunity* **4**, 25–36
26. Xu, C., Guo, H., Pan, X. B., Mao, R., Yu, W., Xu, X., Wei, L., Chang, J., Block, T. M., and Guo, J. T. (2010) Interferons accelerate decay of replication-competent nucleocapsids of hepatitis B virus. *J. Virol.* **84**, 9332–9340
27. Bonvin, M., Achermann, F., Greeve, I., Stroka, D., Keogh, A., Inderbitzin, D., Candinas, D., Sommer, P., Wain-Hobson, S., Vartanian, J. P., and Greeve, J. (2006) Interferon-inducible expression of APOBEC3 editing enzymes in human hepatocytes and inhibition of hepatitis B virus replication. *Hepatology* **43**, 1364–1374
28. Jost, S., Turelli, P., Mangeat, B., Protzer, U., and Trono, D. (2007) Induction of antiviral cytidine deaminases does not explain the inhibition of hepatitis B virus replication by interferons. *J. Virol.* **81**, 10588–10596
29. Nguyen, D. H., Gummuluru, S., and Hu, J. (2007) Deamination-independent inhibition of hepatitis B virus reverse transcription by APOBEC3G. *J. Virol.* **81**, 4465–4472
30. Noguchi, C., Hiraga, N., Mori, N., Tsuge, M., Imamura, M., Takahashi, S., Fujimoto, Y., Ochi, H., Abe, H., Maekawa, T., Yatsuji, H., Shirakawa, K., Takaori-Kondo, A., and Chayama, K. (2007) Dual effect of APOBEC3G on hepatitis B virus. *J. Gen. Virol.* **88**, 432–440
31. Rösler, C., Köck, J., Kann, M., Malim, M. H., Blum, H. E., Baumert, T. F., and von Weizsäcker, F. (2005) APOBEC-mediated interference with hepadnavirus production. *Hepatology* **42**, 301–309
32. Suspène, R., Guétard, D., Henry, M., Sommer, P., Wain-Hobson, S., and Vartanian, J. P. (2005) Extensive editing of both hepatitis B virus DNA strands by APOBEC3 cytidine deaminases *in vitro* and *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 8321–8326
33. Turelli, P., Mangeat, B., Jost, S., Vianin, S., and Trono, D. (2004) Inhibition of hepatitis B virus replication by APOBEC3G. *Science* **303**, 1829
34. Vartanian, J. P., Henry, M., Marchio, A., Suspène, R., Aynaud, M. M., Guétard, D., Cervantes-Gonzalez, M., Battiston, C., Mazzaferro, V., Pineau, P., Dejean, A., and Wain-Hobson, S. (2010) Massive APOBEC3 editing of hepatitis B viral DNA in cirrhosis. *PLoS Pathog.* **6**, e1000928
35. Turelli, P., Liagre-Quazzola, A., Mangeat, B., Verp, S., Jost, S., and Trono, D. (2008) APOBEC3-independent interferon-induced viral clearance in hepatitis B virus transgenic mice. *J. Virol.* **82**, 6585–6590
36. Gripon, P., Rumin, S., Urban, S., Le Seyec, J., Glaise, D., Cannie, I., Guyomard, C., Lucas, J., Trepo, C., and Guguen-Guillouzo, C. (2002) Infection of a human hepatoma cell line by hepatitis B virus. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 15655–15660
37. Hantz, O., Parent, R., Durantel, D., Gripon, P., Guguen-Guillouzo, C., and Zoulim, F. (2009) Persistence of the hepatitis B virus covalently closed circular DNA in HepaRG human hepatocyte-like cells. *J. Gen. Virol.* **90**, 127–135
38. Ladner, S. K., Otto, M. J., Barker, C. S., Zaifert, K., Wang, G. H., Guo, J. T., Seeger, C., and King, R. W. (1997) Inducible expression of human hepatitis B virus (HBV) in stably transfected hepatoblastoma cells: a novel system for screening potential inhibitors of HBV replication. *Antimicrob. Agents Chemother.* **41**, 1715–1720
39. Sells, M. A., Zelent, A. Z., Shvartsman, M., and Acs, G. (1988) Replicative intermediates of hepatitis B virus in HepG2 cells that produce infectious virions. *J. Virol.* **62**, 2836–2844
40. Zhong, J., Gastaminza, P., Cheng, G., Kapadia, S., Kato, T., Burton, D. R., Wieland, S. F., Uprichard, S. L., Wakita, T., and Chisari, F. V. (2005) Robust hepatitis C virus infection *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 9294–9299
41. Sugiyama, M., Tanaka, Y., Kato, T., Orito, E., Ito, K., Acharya, S. K., Gish, R. G., Kramvis, A., Shimada, T., Izumi, N., Kaito, M., Miyakawa, Y., and Mizokami, M. (2006) Influence of hepatitis B virus genotypes on the intra- and extracellular expression of viral DNA and antigens. *Hepatology* **44**, 915–924
42. Schulze, A., Mills, K., Weiss, T. S., and Urban, S. (2012) Hepatocyte polarization is essential for the productive entry of the hepatitis B virus. *Hepatology* **55**, 373–383
43. Liu, Y., Hussain, M., Wong, S., Fung, S. K., Yim, H. J., and Lok, A. S. (2007) A genotype-independent real-time PCR assay for quantification of hepatitis B virus DNA. *J. Clin. Microbiol.* **45**, 553–558
44. Mason, A. L., Xu, L., Guo, L., Kuhns, M., and Perrillo, R. P. (1998) Molecular basis for persistent hepatitis B virus infection in the liver after clearance of serum hepatitis B surface antigen. *Hepatology* **27**, 1736–1742

45. Koyanagi, M., Hijikata, M., Watashi, K., Masui, O., and Shimotohno, K. (2005) Centrosomal P4.1-associated protein is a new member of transcriptional coactivators for nuclear factor- κ B. *J. Biol. Chem.* **280**, 12430–12437
46. Watashi, K., Yeung, M. L., Starost, M. F., Hosmane, R. S., and Jeang, K. T. (2010) Identification of small molecules that suppress microRNA function and reverse tumorigenesis. *J. Biol. Chem.* **285**, 24707–24716
47. Watashi, K., Khan, M., Yedavalli, V. R., Yeung, M. L., Strelbel, K., and Jeang, K. T. (2008) Human immunodeficiency virus type 1 replication and regulation of APOBEC3G by peptidyl prolyl isomerase Pin1. *J. Virol.* **82**, 9928–9936
48. Ta, V. T., Nagaoka, H., Catalan, N., Durandy, A., Fischer, A., Imai, K., Nonoyama, S., Tashiro, J., Ikegawa, M., Ito, S., Kinoshita, K., Muramatsu, M., and Honjo, T. (2003) AID mutant analyses indicate requirement for class-switch-specific cofactors. *Nat. Immunol.* **4**, 843–848
49. Matsumoto, Y., Marusawa, H., Kinoshita, K., Endo, Y., Kou, T., Morisawa, T., Azuma, T., Okazaki, I. M., Honjo, T., and Chiba, T. (2007) *Helicobacter pylori* infection triggers aberrant expression of activation-induced cytidine deaminase in gastric epithelium. *Nat. Med.* **13**, 470–476
50. Endo, Y., Marusawa, H., Kou, T., Nakase, H., Fujii, S., Fujimori, T., Kinoshita, K., Honjo, T., and Chiba, T. (2008) Activation-induced cytidine deaminase links between inflammation and the development of colitis-associated colorectal cancers. *Gastroenterology* **135**, 889–898
51. Liang, G., Kitamura, K., Wang, Z., Liu, G., Chowdhury, S., Fu, W., Koura, M., Wakae, K., Honjo, T., and Muramatsu, M. (2013) RNA editing of hepatitis B virus transcripts by activation-induced cytidine deaminase. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 2246–2251
52. Marusawa, H., Hijikata, M., Watashi, K., Chiba, T., and Shimotohno, K. (2001) Regulation of Fas-mediated apoptosis by NF- κ B activity in human hepatocyte derived cell lines. *Microbiol. Immunol.* **45**, 483–489
53. Raney, A. K., Johnson, J. L., Palmer, C. N., and McLachlan, A. (1997) Members of the nuclear receptor superfamily regulate transcription from the hepatitis B virus nucleocapsid promoter. *J. Virol.* **71**, 1058–1071
54. Schulze, A., Gripon, P., and Urban, S. (2007) Hepatitis B virus infection initiates with a large surface protein-dependent binding to heparan sulfate proteoglycans. *Hepatology* **46**, 1759–1768
55. Yan, H., Zhong, G., Xu, G., He, W., Jing, Z., Gao, Z., Huang, Y., Qi, Y., Peng, B., Wang, H., Fu, L., Song, M., Chen, P., Gao, W., Ren, B., Sun, Y., Cai, T., Feng, X., Sui, J., and Li, W. (2012) Sodium taurocholate cotransporting polypeptide is a functional receptor for human hepatitis B and D virus. *Elife* **1**, e00049
56. Arend, W. P. (1991) Interleukin 1 receptor antagonist. A new member of the interleukin 1 family. *J. Clin. Invest.* **88**, 1445–1451
57. Requena, P., Daddaoua, A., Guadix, E., Zarzuelo, A., Suárez, M. D., Sánchez de Medina, F., and Martínez-Augustín, O. (2009) Bovine glycomacropptide induces cytokine production in human monocytes through the stimulation of the MAPK and the NF- κ B signal transduction pathways. *Br. J. Pharmacol.* **157**, 1232–1240
58. Locarnini, S., and Zoulim, F. (2010) Molecular genetics of HBV infection. *Antivir. Ther.* **15**, Suppl. 3, 3–14
59. Goila-Gaur, R., and Strelbel, K. (2008) HIV-1 Vif, APOBEC, and intrinsic immunity. *Retrovirology* **5**, 51
60. Endo, Y., Marusawa, H., Kinoshita, K., Morisawa, T., Sakurai, T., Okazaki, I. M., Watashi, K., Shimotohno, K., Honjo, T., and Chiba, T. (2007) Expression of activation-induced cytidine deaminase in human hepatocytes via NF- κ B signaling. *Oncogene* **26**, 5587–5595
61. Tanaka, Y., Marusawa, H., Seno, H., Matsumoto, Y., Ueda, Y., Kodama, Y., Endo, Y., Yamauchi, J., Matsumoto, T., Takaori-Kondo, A., Ikai, I., and Chiba, T. (2006) Anti-viral protein APOBEC3G is induced by interferon- α stimulation in human hepatocytes. *Biochem. Biophys. Res. Commun.* **341**, 314–319
62. Ito, S., Nagaoka, H., Shinkura, R., Begum, N., Muramatsu, M., Nakata, M., and Honjo, T. (2004) Activation-induced cytidine deaminase shuttles between nucleus and cytoplasm like apolipoprotein B mRNA editing catalytic polypeptide 1. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 1975–1980
63. Daniels, H. M., Meager, A., Eddleston, A. L., Alexander, G. J., and Williams, R. (1990) Spontaneous production of tumour necrosis factor α and interleukin-1 β during interferon- α treatment of chronic HBV infection. *Lancet* **335**, 875–877
64. Esteve, M., Saro, C., González-Huix, F., Suarez, F., Forné, M., and Viver, J. M. (2004) Chronic hepatitis B reactivation following infliximab therapy in Crohn's disease patients: need for primary prophylaxis. *Gut* **53**, 1363–1365
65. Manzano-Alonso, M. L., and Castellano-Tortajada, G. (2011) Reactivation of hepatitis B virus infection after cytotoxic chemotherapy or immunosuppressive therapy. *World J. Gastroenterol.* **17**, 1531–1537
66. Hacham, M., Argov, S., White, R. M., Segal, S., and Apte, R. N. (2000) Distinct patterns of IL-1 α and IL-1 β organ distribution—a possible basis for organ mechanisms of innate immunity. *Adv. Exp. Med. Biol.* **479**, 185–202
67. Hösel, M., Quasdorff, M., Wiegmann, K., Webb, D., Zedler, U., Broxtermann, M., Tedjokusumo, R., Esser, K., Arzberger, S., Kirschning, C. J., Langenkamp, A., Falk, C., Büning, H., Rose-John, S., and Protzer, U. (2009) Not interferon, but interleukin-6 controls early gene expression in hepatitis B virus infection. *Hepatology* **50**, 1773–1782
68. Cooper, A., Tal, G., Lider, O., and Shaul, Y. (2005) Cytokine induction by the hepatitis B virus capsid in macrophages is facilitated by membrane heparan sulfate and involves TLR2. *J. Immunol.* **175**, 3165–3176
69. Dev, A., Iyer, S., Razani, B., and Cheng, G. (2011) NF- κ B and innate immunity. *Curr. Top. Microbiol. Immunol.* **349**, 115–143
70. Delker, R. K., Fugmann, S. D., and Papavasiliou, F. N. (2009) A coming-of-age story: activation-induced cytidine deaminase turns 10. *Nat. Immunol.* **10**, 1147–1153
71. Muramatsu, M., Sankaranand, V. S., Anant, S., Sugai, M., Kinoshita, K., Davidson, N. O., and Honjo, T. (1999) Specific expression of activation-induced cytidine deaminase (AID), a novel member of the RNA-editing deaminase family in germinal center B cells. *J. Biol. Chem.* **274**, 18470–18476
72. Muramatsu, M., Kinoshita, K., Fagarasan, S., Yamada, S., Shinkai, Y., and Honjo, T. (2000) Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* **102**, 553–563
73. Gourzi, P., Leonova, T., and Papavasiliou, F. N. (2006) A role for activation-induced cytidine deaminase in the host response against a transforming retrovirus. *Immunity* **24**, 779–786
74. MacDuff, D. A., Demorest, Z. L., and Harris, R. S. (2009) AID can restrict L1 retrotransposition suggesting a dual role in innate and adaptive immunity. *Nucleic Acids Res.* **37**, 1854–1867
75. Chowdhury, S., Kitamura, K., Simadu, M., Koura, M., and Muramatsu, M. (2013) Concerted action of activation-induced cytidine deaminase and uracil-DNA glycosylase reduces covalently closed circular DNA of duck hepatitis B virus. *FEBS Lett.* (2013) **587**, 3148–3152