

Genomics update

Single-cell genomics: unravelling the genomes of unculturable microorganisms

Victor de Jager^{1,2,3} and Roland J. Siezen^{1,2,4,5,6*}

¹Netherlands Bioinformatics Centre, Nijmegen, The Netherlands.

²Centre for Molecular and Biomolecular Informatics, NCMLS, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands.

³Laboratory of Microbiology, Wageningen University, Wageningen, The Netherlands.

⁴TI Food and Nutrition, Wageningen, The Netherlands.

⁵NIZO food research, Ede, The Netherlands.

⁶Kluyver Centre for Genomics of Industrial Fermentation, Delft, The Netherlands.

Microbial genomics and related transcriptomics methods rely on culturing techniques to obtain enough DNA suitable for high-throughput sequencing without resorting to DNA amplification techniques. A few microgram of DNA is needed for most common next-generation sequencing methods. For transcriptome analysis, sufficient cDNA is needed to measure low abundance mRNA copies in the cell. However, the large majority of microbes on earth resist cultivation, hampering research into their relevant gene pool, ecological niche or industrial relevance. For example, many environmental or gut-related species cannot be grown outside their natural habitat. Even if we isolate the metagenome or the metatranscriptome from these environments, this reveals only a fragmented sequence landscape that is difficult to assign to individual species. Although enrichment techniques or metatranscriptome analysis of previously unculturable species have been shown to assist in directed culturing, e.g. of a *Rikenella*-like bacterium (Bomar *et al.*, 2011), the unravelling of a complex metagenome into its individual genomes and their organization is impossible using current technologies.

A major challenge is the analysis of bacteria and other organisms living inside a complex matrix, like biofilms. Metagenome or transcriptome analysis of microorganisms has been described for biofilms consisting of a single species by scraping of the biofilm to obtain

enough material (Holmes *et al.*, 2006), but for multi-species biofilms this method results in a metagenome or metatranscriptome dataset. The solution to these challenges may be the isolation and genomic analysis of unculturable single cells isolated from such environments. Here we describe in brief the state-of-the-art in single-cell microbial genomics.

Single-cell isolation

Several methods exist to extract and investigate single microbial cells from their environment. Flow cytometry or fluorescence-activated cell sorting (FACS) has been used since the 1970s and its applications in microbiology were recognized early (Fouchet *et al.*, 1993); recent advances are described by Müller and Nebe-von-Caron (2010), Wang and Bodovitz (2010), and Wang *et al.* (2010). Micromanipulation has been described by Kvist *et al.* (2007) and more recently by Woyke *et al.* (2010). Microfluidic device techniques are shown to be effective by combining the separation of cells and subsequently performing biochemical reactions on the device itself, thereby maximizing reaction yield (Marcy *et al.*, 2007a) (Fig. 1).

Single-cell genome sequencing and data analysis

Whereas classical next-generation sequencing to determine an organism's genome sequence relies on pooling DNA from 10^6 – 10^8 cells, single-cell genomics relies on whole-genome amplification from a single cell. Most studies rely on Multiple displacement amplification (MDA), a biochemical amplification technique using random primers and ϕ 29 DNA polymerase (Dean *et al.*, 2001; Raghunathan *et al.*, 2005; Zhang *et al.*, 2006; Marcy *et al.*, 2007a). Other amplification techniques like random-primed PCR result in a more over- and under-representation of different regions of the template DNA and generate very short fragments (Dean *et al.*, 2001; Hosono *et al.*, 2003). MDA, however, results in fragments of 12–100 kb rendering them suitable for sequencing. Although the complete microbial genome from a single cell can be amplified to amounts required for current

*For correspondence. E-mail r.siezen@cmbi.ru.nl; Tel. (+31) 2436 19559; Fax (+31) 2436 19395.




Fig. 1. Photograph of a single-cell isolation and genome amplification chip capable of processing nine samples in parallel (eight cells, one positive control).
 A. To visualize the architecture, the channels and chambers have been filled with blue food colouring and the control lines to actuate the valves have been filled with red food colouring (scale bar 5 mm).
 B. Schematic diagram of the automated sorting procedure. Closed valves are shown in red, open valves are transparent. Cells are drawn in green.
 C. Typical result of cell sorting showing for each unit (seven with a single cell and one negative control without a cell) a colour combination of a phase contrast image (gray) and a fluorescence image (green). A green overlaid square has been placed around the cell to ease visualization, whereas a red crossed square indicates the absence of cell. Scale bar is 100 μm . Reprinted from Marcy *et al.* (2007a).

sequencing methods without *a priori* sequence knowledge, early studies suggested that up to 40% of the genomic sequence was missed (Podar *et al.*, 2007; Marcy *et al.*, 2007b; Woyke *et al.*, 2009) (Table 1).

An overview of an MDA set-up using a microfluidic device is shown in Fig. 2, although FACS-based methods are also often reported in literature (Rodrigue *et al.*, 2009; Siegl and Hentschel, 2010). All DNA in the

Table 1. Examples of single-cell genome sequencing.

Microorganism	Assembled bases (Mb)	Estimated % genome recovery	Scaffolds	Contigs	GC%	Single cell separation	Isolation source	Reference
TM7a (new phylum)	2.865	?	1825		34.3	Microfluidics	Human mouth biofilm	Marcy <i>et al.</i> (2007b)
TM7_GTL1 (new phylum)	0.679	?		132	48.5	FISH/FACS	Soil	Podar <i>et al.</i> (2007)
<i>Prochlorococcus</i> MED4		95		755		FACS	Sea water; lab culture ^a	Rodrigue <i>et al.</i> (2009)
<i>Flavobacterium</i> MS024-2A	1.905	91		17	36	Flow cytometer	Coastal water, Maine, USA	Woyke <i>et al.</i> (2009)
<i>Flavobacterium</i> MS024-3C	1.505	78		21	39	Flow cytometer	Coastal water, Maine, USA	Woyke <i>et al.</i> (2009)
Cand. <i>Sulcia muelleri</i> DMIN	0.244	100	1	1	22.5	Micromanipulator	Symbiont from insect bacteriome (green sharpshooter)	Woyke <i>et al.</i> (2010)
<i>Poribacteria</i>	1.885	66		1597	53.4	FACS	Symbiont from marine sponge	Siegl <i>et al.</i> (2011)
Cand. <i>Nitrosoarchaeum limnia</i> SFB1	1.690 ^b	95	26	136	32.4	Microfluidics, laser tweezer	Ammonia-oxidizing enrichment culture; sediment water, San Francisco bay, USA	Blainey <i>et al.</i> (2011)

a. Method validation using strain with known genome sequence.

b. Pooled sequence data from five individual cells; see Table 2.




Fig. 2. A mixture of cells sampled from a complex microbial ecosystem is introduced into the chip. Single cells are selected using an optical trap, and are sorted into chambers for cell lysis and genome amplification. Genomes are amplified in nanolitre MDA reactions to produce larger quantities of DNA (shown are SYBR Green–stained products in microfluidic reaction chambers). Sequencing libraries are created from the amplified genomic DNA for sequencing on a high-throughput DNA sequencer. The sequence reads are assembled to recover the genome sequence, which is annotated to identify genes and pathways present in the original cell. Reprinted by permission from Macmillan Publishers Ltd: Nature Methods (Kalisky and Quake, 2011), copyright 2011. The microfluidics image was reprinted from Leslie (2011).

initial sample will be amplified, which renders the method very prone to DNA contamination. Another disadvantage of the initial method is uneven amplification of the genome, which results in high-coverage sequencing of the amplified genomic regions while remaining sequences may not be sufficiently covered (Zhang *et al.*, 2006). Marcy *et al.* (2007a) demonstrated that reducing MDA reaction volumes lowers non-specific synthesis from contaminant DNA templates and unfavourable interactions between primers. The work of Rodrigue *et al.* (2009) demonstrated a biochemical method to normalize the products obtained in MDA reactions. They also discussed the problem of chimera formation linking non-contiguous chromosomal regions in MDA (Dean *et al.*, 2001; Zhang *et al.*, 2006), which may hamper sequence assembly and render mate-pair data less efficient in contig positioning. Several other single-cell techniques are described in recent reviews by Wang and Bodovitz (2010), Kalisky and Quake (2011), and Pan *et al.* (2011). As data analysis from single-cell amplified genomes is equally challenging, the software framework SmashCell has been developed to automate the main steps in sequence assembly, gene prediction, annotation and visualization (Harrington *et al.*, 2010).

Single-cell genome sequences of uncultured microorganisms

Examples of sequencing of single amplified genomes (SAGs) are listed in Table 1. Woyke *et al.* (2010) describe using a micro-displacement technique to sequence a genome from an uncultured single cell of *Candidatus Sulcia muelleri* DMIN, a symbiont isolated from the bacteriome of the green sharpshooter *Draeculacephala minerva*. This polyploid bacterium has an estimated 200–900 genome copies per cell. Of the 57 Mb of sequence generated, approximately 90% was of contaminant origin, as estimated by mapping to a previously sequenced genome of *Sulcia* and phylogenetic analysis with blastx and MEGAN (Mitra *et al.*, 2009). The remaining reads were assembled into a draft genome, misassemblies due to chimeras were corrected manually, and subsequent application of primer walking, sequencing PCR products and Illumina sequencing resulted in a final finished genome (Fig. 3).

Siegl *et al.* (2011) used FACS to isolate cells from the candidate phylum *Poribacteria* and subsequently MDA to obtain a SAG. These bacteria are almost exclusively found in marine sponges as symbionts and resist cultivation efforts. The SAG of 1.88 Mb was contained in 1597




Fig. 3. *Sulcia* cell isolation and sequence coverage, closure and polishing locations along the *Sulcia* DMIN single cell genome. A. Micromanipulation of the single *Sulcia* cell from the sharpshooter bacteriome metasample. B. Sequence coverage including closure and polishing locations along the finished, circular *Sulcia* DMIN. Reprinted from Woyke *et al.* (2010). For figure details see the original article.

contigs, which covered an estimated two-thirds of the total genomic DNA based on the distribution of tRNA genes and their specificities found in the contigs. Nevertheless, a comprehensive overview of poribacterial metabolism could be deduced (Fig. 4). The extensive Sup-type polyketide synthases found in the SAG of *Poribacteria* confirmed the previously proposed assignment of Sup-PKS to this species. With the finding of a second putative PKS system showing high similarity to the lipopolysaccharide type I PKS WcbR from *Nitrosomonas* and *Burkholderia*, as well as RkpA from *Sinorhizobium fredii*, they suggested that *Poribacteria* contain at least two different types of PKS systems and their products may be involved in sponge–microbe interactions. This study showed that single-cell genomics is highly capable of dissecting the

genomic information from unculturable bacteria, shedding light on genomic organization, metabolic functions and possibly new insight in the debate on the origin of sponge bioactive compounds.

Ammonia-oxidizing archaea (AOA) are among the most abundant microbes on Earth, and may significantly impact global nitrogen and carbon cycles. Five single cells were isolated from a low-salinity sediment AOA-enrichment culture using a microfluidic device and laser tweezers, and DNA was amplified and sequenced separately from each cell (Blainey *et al.*, 2011) (Tables 1 and 2). Individually, three single-cell datasets gave assemblies of more than 1 Mb at sequencing depths of 10× to 30×, and an estimated 60% genomic coverage each; the low coverage is considered typical due to MDA amplification bias. Surprisingly,




Fig. 4. A schematic overview of poribacterial metabolism as deduced from SAG sequencing. Reprinted by permission from Macmillan Publishers Ltd: The ISME Journal (Siegl *et al.*, 2011), copyright 2011.

each of the single-cell assemblies represented a different 60% of the target genome, and combining the five datasets led to a single-cell assembly representing > 95% of the *Nitrosoarchaeum limnia* genome. Based on nucleotide identity comparisons, this AOA is proposed to represent a new genus of Crenarchaeota. In contrast to other described AOA, this low-salinity archaeum appears to be motile, based on the presence of numerous motility and

chemotaxis-associated genes in the genome (Blainey *et al.*, 2011).

Single-cell transcriptomics, metabolomics and proteomics

Recent reports on single-cell transcriptomics discuss mainly the analysis of polyadenylated mRNA of eukary-

Table 2. Assembly statistics for sequencing of three single cells of *Nitrosoarchaeum limnia* SFB1, and consensus genome (reads from metagenome and five single cells).

Assembly statistics	Cell 23	Cell 21	Cell 3	Five single cells co-assembly	Consensus single cells and metagenome
Raw read bases	17 107 411	52 341 561	29 999 202	118 796 782	150 994 537
Assembly bases	1 094 113	1 039 820	1 041 604	1 690 404	1 769 573
Scaffolds	68	76	83	26	2
Unscaffolded contigs	287	177	265	110	29
Estimated % genome coverage	62	59	59	95	99

Adapted from Table 1 of Blainey *et al.* (2011).

© 2011 The Authors

Journal compilation © 2011 Society for Applied Microbiology and Blackwell Publishing Ltd, *Microbial Biotechnology*, 4, 431–437

otes. A comprehensive overview of the technologies involved is given by Tang *et al.* (2011). In short, the single-cell methods exploit reverse transcription using oligo(dT) primers to convert mRNAs with poly(A) tails into cDNAs, followed by uniform amplification and sequencing (RNA-seq). However, currently no single-cell analysis reports are known that exploit protocols for mRNA extraction from bacterial cells, for instance using the MessageAmp II-Bacteria Kit (Ambion) as described by Frias-Lopez *et al.* (2008). Single-cell metabolome and proteome/peptidome analyses are still in their infancy, as these compounds cannot be amplified and their analysis requires technological breakthroughs in pushing the limits of detection (Rubakhin *et al.*, 2011).

Future

Since the introduction of single-cell genomics (Raghunathan *et al.*, 2005), there have been surprisingly few reports of successful reconstruction of whole genomes from single unculturable bacterial cells (Table 1). This undoubtedly reflects the extreme difficulties in the various steps of single-cell isolation, miniaturization, DNA amplification, avoidance of contamination and data analysis. Nevertheless, the pioneering examples show that it is definitely feasible to sequence genomes of single unculturable cells isolated from complex consortia, and we expect this approach to become more widespread as miniaturization technologies improve.

Recently, it has also been recognized that isogenic microbial populations (pure cultures) contain substantial cell-to-cell differences in physiological parameters such as growth rate, resistance to stress and regulatory circuit output (Ingham *et al.*, 2008; Lidstrom and Konopka, 2010). In this light, adaptation of single-cell genome sequencing using microfluidic approaches towards RNA-seq transcriptome analysis of single cells using next-generation mRNA sequencing should become increasingly important (Siezen *et al.*, 2010).

Acknowledgements

This project was carried out within the research programmes of the Kluyver Centre for Genomics of Industrial Fermentation and the Netherlands Bioinformatics Centre, which are part of the Netherlands Genomics Initiative/Netherlands Organization for Scientific Research.

References

Blainey, P.C., Mosier, A.C., Potanina, A., Francis, C.A., and Quake, S.R. (2011) Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS ONE* **6**: e16626–e16626.

Bomar, L., Maltz, M., Colston, S., and Graf, J. (2011) Directed culturing of microorganisms using metatranscriptomics. *mBio* **2**: e00012–e00011.

Dean, F.B., Nelson, J.R., Giesler, T.L., and Lasken, R.S. (2001) Rapid amplification of plasmid and phage DNA using Phi 29 DNA polymerase and multiply-primed rolling circle amplification. *Genome Res* **11**: 1095–1099.

Fouchet, P., Jayat, C., Héchard, Y., Ratinaud, M.H., and Frelat, G. (1993) Recent advances of flow cytometry in fundamental and applied microbiology. *Biol Cell* **78**: 95–109.

Frias-Lopez, J., Shi, Y., Tyson, G.W., Coleman, M.L., Schuster, S.C., Chisholm, S.W., and DeLong, E.F. (2008) Microbial community gene expression in ocean surface waters. *Proc Natl Acad Sci USA* **105**: 3805–3810.

Harrington, E.D., Arumugam, M., Raes, J., Bork, P., and Relman, D.A. (2010) SmashCell: a software framework for the analysis of single-cell amplified genome sequences. *Bioinformatics* **26**: 2979–2980.

Holmes, D.E., Chaudhuri, S.K., Nevin, K.P., Mehta, T., Methé, B.A., Liu, A., *et al.* (2006) Microarray and genetic analysis of electron transfer to electrodes in *Geobacter sulfurreducens*. *Environ Microbiol* **8**: 1805–1815.

Hosono, S., Faruqi, A.F., Dean, F.B., Du, Y., Sun, Z., Wu, X., *et al.* (2003) Unbiased whole-genome amplification directly from clinical samples. *Genome Res* **13**: 954–964.

Ingham, C.J., Beerthuyzen, M., and van Hylckama Vlieg, J. (2008) Population heterogeneity of *Lactobacillus plantarum* WCFS1 microcolonies in response to and recovery from acid stress. *Appl Environ Microbiol* **74**: 7750–7758.

Kalisky, T., and Quake, S.R. (2011) Single-cell genomics. *Nature Methods* **8**: 311–314.

Kvist, T., Ahring, B.K., Lasken, R.S., and Westermann, P. (2007) Specific single-cell isolation and genomic amplification of uncultured microorganisms. *Appl Microbiol Biotechnol* **74**: 926–935.

Leslie, M. (2011) Single-cell tech primer. *Science (New York, NY)* **331**: 24–25.

Lidstrom, M.E., and Konopka, M.C. (2010) The role of physiological heterogeneity in microbial population behavior. *Nat Chem Biol* **6**: 705–712.

Marcy, Y., Ishoey, T., Lasken, R.S., Stockwell, T.B., Walenz, B.P., Halpern, A.L., *et al.* (2007a) Nanoliter reactors improve multiple displacement amplification of genomes from single cells. *PLoS Genet* **3**: 1702–1708.

Marcy, Y., Ouverney, C., Bik, E.M., Lösekann, T., Ivanova, N., Martin, H.G., *et al.* (2007b) Dissecting biological ‘dark matter’ with single-cell genetic analysis of rare and uncultivated TM7 microbes from the human mouth. *Proc Natl Acad Sci USA* **104**: 11889–11894.

Mitra, S., Klar, B., and Huson, D.H. (2009) Visual and statistical comparison of metagenomes. *Bioinformatics (Oxford, England)* **25**: 1849–1855.

Müller, S., and Nebe-von-Caron, G. (2010) Functional single-cell analyses: flow cytometry and cell sorting of microbial populations and communities. *FEMS Microbiol Rev* **34**: 554–587.

Pan, X.-H., Zhu, H.-Y., and Marjani, S.L. (2011) Technological advances in single-cell genomic analyses. *Yi Chuan* **33**: 17–24 (article in Chinese).

Podar, M., Abulencia, C.B., Walcher, M., Hutchison, D., Zengler, K., Garcia, J.A., *et al.* (2007) Targeted access to

- the genomes of low-abundance organisms in complex microbial communities. *Appl Environ Microbiol* **73**: 3205–3214.
- Raghunathan, A., Ferguson, H.R., Jr, Bornarth, C.J., Song, W., Driscoll, M., and Lasken, R.S. (2005) Genomic DNA amplification from a single bacterium. *Appl Environ Microbiol* **71**: 3342–3347.
- Rodrigue, S., Malmstrom, R.R., Berlin, A.M., Birren, B.W., Henn, M.R., and Chisholm, S.W. (2009) Whole genome amplification and de novo assembly of single bacterial cells. *PLoS ONE* **4**: e6864–e6864.
- Rubakhin, S.S., Romanova, E.V., Nemes, P., and Sweedler, J.V. (2011) Profiling metabolites and peptides in single cells. *Nat Methods* **8**: S20–S29.
- Siegl, A., and Hentschel, U. (2010) PKS and NRPS gene clusters from microbial symbiont cells of marine sponges by whole genome amplification. *Environ Microbiol Rep* **2**: 507–513.
- Siegl, A., Kamke, J., Hochmuth, T., Piel, J., Richter, M., Liang, C., *et al.* (2011) Single-cell genomics reveals the lifestyle of Poribacteria, a candidate phylum symbiotically associated with marine sponges. *ISME J* **5**: 61–70.
- Siezen, R.J., Wilson, G., and Todt, T. (2010) Prokaryotic whole-transcriptome analysis: deep sequencing and tiling arrays. *Microb Biotechnol* **3**: 125–130.
- Tang, F., Lao, K., and Surani, M.A. (2011) Development and applications of single-cell transcriptome analysis. *Nat Methods* **8**: S6–S11.
- Wang, D., and Bodovitz, S. (2010) Single cell analysis: the new frontier in 'omics'. *Trends Biotechnol* **28**: 281–290.
- Wang, Y., Hammes, F., De Roy, K., Verstraete, W., and Boon, N. (2010) Past, present and future applications of flow cytometry in aquatic microbiology. *Trends Biotechnol* **28**: 416–424.
- Woyke, T., Xie, G., Copeland, A., Gonzalez, J.M., Han, C., Kiss, H., *et al.* (2009) Assembling the marine metagenome, one cell at a time. *PLoS ONE* **4**: e5299.
- Woyke, T., Tighe, D., Mavromatis, K., Clum, A., Copeland, A., Schackwitz, W., *et al.* (2010) One bacterial cell, one complete genome. *PLoS ONE* **5**: e10314–e10314.
- Zhang, K., Martiny, A.C., Reppas, N.B., Barry, K.W., Malek, J., Chisholm, S.W., and Church, G.M. (2006) Sequencing genomes from single cells by polymerase cloning. *Nat Biotechnol* **24**: 680–686.