

## Natural Occurrence of Toxigenic *Fusarium proliferatum* on Paddy (*Oryza sativa* L.) in Karnataka, India

<sup>1</sup>Pavagada Krishnamurthy Maheshwar and <sup>2</sup>Gottravalli Ramanayaka Janardhana \*

<sup>1</sup>Department of Microbiology, Yuvaraja's College, University of Mysore, Mysore  
570 005, Karnataka, India

<sup>2</sup>Department of Studies in Botany, University of Mysore, Manasagangotri, Mysore  
570 006, Karnataka, India

**Abstrak:** Pencemaran biji benih padi (beras bersama sekam/ sekam padi) oleh spesies *Fusarium* boleh mengakibatkan kerosakan dan diikuti dengan penghasilan mikotoksin, terutamanya fumonisin yang memberi kesan terhadap kesihatan manusia dan haiwan. Satu kajian mikologi telah dijalankan untuk menilai penghasilan fumonisin secara semulajadi oleh *Fusarium proliferatum* dalam padi yang tumbuh di kawasan-kawasan geografi berbeza di Karnataka (India). Sejumlah 65 sampel *F. proliferatum* diasingkan daripada padi dan dianalisa menggunakan *polymerase chain reaction* (PCR). Satu set primer-primer Fp3-F dan Fp4-R telah digunakan untuk mengenal pasti spesies *F. proliferatum* manakala satu lagi set primer-primer, FUM1 telah digunakan untuk mengenal pasti kebolehan menghasilkan fumonisin. Kesemua 65 sampel *F. proliferatum* memberi keputusan positif kepada kedua-dua set primer, dengan menghasilkan produk yang dibesarkan kepada saiz yang dijangka. Tambahan lagi, melalui analisa *thin layer chromatography* (TLC), fumonisin B<sub>1</sub> (FB<sub>1</sub>) telah dikesan di kesemua sampel *F. proliferatum* yang memberi keputusan PCR yang positif.

**Kata kunci:** Fp3-F/Fp4-R, FUM1, Diagnostik PCR, TLC

**Abstract:** Contamination of paddy seeds (rice with husk) by *Fusarium* species can cause spoilage and subsequent production of mycotoxins, especially fumonisins that affect human and animal health. A mycological study was conducted to evaluate the natural occurrence of fumonisin B<sub>1</sub> produced by *Fusarium proliferatum* on paddy grown in different geographic regions of Karnataka (India). A total of 65 isolates of *F. proliferatum* from paddy samples were analysed by polymerase chain reaction (PCR). One set of primers, Fp3-F and Fp4-R was employed to identify the species *F. proliferatum*, and another set of primers, FUM1 was employed to determine the fumonisin producing ability of the isolates. All 65 isolates of *F. proliferatum* scored positive with both set of primers, producing amplified products of the expected sizes. Furthermore, thin layer chromatography (TLC) analysis detected fumonisin B<sub>1</sub> (FB<sub>1</sub>) in all of the PCR positive isolates of *F. proliferatum*.

**Keywords:** Fp3-F/Fp4-R, FUM1, PCR Diagnostics, TLC

---

\* Corresponding author: grjbelur@gmail.com

## INTRODUCTION

Paddy is an important worldwide food crop and is the staple food for most of Asia. Paddy is the basic food for the Indian population, as it is a rich source of dietary energy and a good source of amino acids, such as thiamine, riboflavin, glutamic acid and niacin (FAO 2004). However, paddy is highly susceptible to fungal infections, such as *Fusarium* species (Trung *et al.* 2001), *F. proliferatum* (Matsushima) Nirenberg and *F. verticillioides* (Sacc.) Nirenberg (synonym *F. moniliforme* Sheldon), which produce mycotoxins, such as trichothecenes, zearalenones and fumonisins (Abbas *et al.* 1998; Desjardins *et al.* 2000). Fumonisin B<sub>1</sub> (FB<sub>1</sub>) and fumonisin B<sub>2</sub> (FB<sub>2</sub>) are the major toxins of the family of mycotoxins produced by *F. proliferatum* and *F. verticillioides*. While fumonisin production by *F. verticillioides* has been well studied there is limited data for *F. proliferatum* (Rheeder *et al.* 2002).

*Fusarium* contamination of paddy has been reported in many parts of the world (Tonon *et al.* 1997; Pacin *et al.* 2002), but with less frequency compared to other cereals. Despite the importance of paddy as a staple food and the reported occurrence of mycoflora and fumonisins (Park *et al.* 2005; Hinojo *et al.* 2006), there is a scarcity of information on the incidence of *Fusarium* species and fumonisin contamination in paddy produced in the Indian state of Karnataka.

The International Agency for Research on Cancer (IARC) evaluated FB<sub>1</sub> as a possible carcinogen to humans (Group 2B) (Domijan *et al.* 2005). It is important to detect fumonisins in order to prevent human and animal exposure to such toxic substances. Polymerase chain reaction (PCR) protocols based on intergenic spacer (IGS) sequences have been extensively used for the accurate detection of *Fusarium* species (Jurado *et al.* 2006). In the present investigation, species-specific primers Fp3-F and Fp4-R based on IGS sequences were used to detect *F. proliferatum* isolated from paddy. In addition, the fumonisin producing ability of these isolates was confirmed by PCR using a second set of primers, FUM1, and by thin layer chromatography (TLC) studies that detect FB<sub>1</sub>.

## MATERIALS AND METHODS

### Chemicals and Reagents

The reagents for PCR, including primers, were procured from Bangalore Genei, Bangalore (India). All other chemicals used in the study were of reagent grade (Merck Ltd., Mumbai). FB<sub>1</sub> standard was purchased from Sigma Aldrich Chemicals Pvt. Ltd., Bangalore, India.

### Collection of Paddy Seed Samples

In total, 109 lots of different paddy seed samples were collected from farmer's fields from 15 districts in the state of Karnataka during the month of June 2006. The seed samples (0.5 kg) were packed, appropriately labelled and used for further studies.

### Isolation of *Fusarium* Species

Isolation of *Fusarium* species from the 109 paddy samples was accomplished by using modified malachite green agar medium containing malachite green oxalate (2.5 mg/l) (MGA 2.5) (Bragulat *et al.* 2004). Modified Czapek-Dox agar (CZA) and Spezieller Nährstoffarmer Agar media (SNA) were used (Leslie & Summerell 2006) to maintain cultures.

For mycological analysis, 109 paddy samples were subjected to sampling by the hand-halving method (Mathur & Kongsdal 2003). A total of 200 paddy seeds from each sample were surface sterilised by treatment with 1% sodium hypochlorite solution for 1 min and rinsed twice in sterile distilled water. The surface-sterilised seeds were plated (10 per plate) on MGA 2.5 containing chloramphenicol (50 mg/l). The plates were incubated at 26°C for 5 days. The developing fungal colonies were counted and *Fusarium* species were isolated and sub-cultured on potato dextrose agar medium. The cultures were identified using fungal keys and manuals (Booth 1977; Leslie & Summerell 2006).

The isolation frequency (Fr) (Gonzalez *et al.* 1995) of the *Fusarium* isolates was calculated as below:

$$\text{Fr (\%)} = \frac{\text{Number of samples infected with } \textit{Fusarium} \text{ species} \times 100}{\text{Total number of samples analysed}}$$

### DNA Extraction

Genomic DNA was extracted from isolates of *F. proliferatum* (65), *F. verticillioides* (27), *F. semitectum* Berkeley & Ravenel (12) and *F. oxysporum* Schlechtendahl emend Snyder & Hansen (10). Each fungal species was inoculated aseptically into 500 µl of sterile potato dextrose broth in 2 ml microfuge tubes. The tubes were incubated for 5 days at 28°C and DNA was isolated from each fungal sample (Zhang *et al.* 1998). Mycelia from the microfuge tubes were centrifuged at 5000 rpm (REMI C24 Cooling Centrifuge, Chennai) for 8 min and the supernatant was discarded. The mycelium was resuspended in cell lysis buffer (2% CTAB, 1.4 M NaCl, 20 mM EDTA, 100 mM Tris-HCl, pH 8.0, pre-heated at 65°C) and heated at 65°C for 20 min in a water bath. An equal volume of phenol:chloroform (1:1) was added to each tube and centrifuged at 3000 rpm for 5 min. The supernatants were transferred to new microfuge tubes and an equal volume of isopropyl alcohol was added. The microfuge tubes were incubated at -20°C for 2 h and then centrifuged at 8000 rpm for 8 min to precipitate the DNA. The DNA pellet was air-dried and resuspended in 20 µl of nuclease free water and used directly for PCR analysis.

### Primers for PCR

A pair of primers, Fp3-F (5'CGGCCACCAGAGGATGTG 3') and Fp4-R (5' CAACACGAATCGCTTCCTGAC 3') (Jurado *et al.* 2006), specific to *F. proliferatum* was used for species identification. Another set of primers specific to the *fum1* gene (involved in fumonisin biosynthesis), FUM1 forward (5'-CCATCACAGTGGGACACAGT-3') and FUM1 reverse (5'-CGTATCGTCAG CATGATGTAGC-3') (Bluhm *et al.* 2004), was used to determine the fumonisin

producing ability of *F. proliferatum*. The expected amplicon sizes were 230 bp and 183 bp, respectively.

### **Polymerase Chain Reaction**

Genomic DNA was subjected to PCR analysis using Advanced Primus 25 Thermocycler (Peqlab, Germany). The PCR mixture for primer Fp3-F and Fp4-R contained 100 ng of genomic DNA, 1.25 µl of each primer (20 pmol), 0.5 µl of Taq DNA polymerase (3 U/µl), 2.5 µl of 10X PCR buffer, 1.0 µl of MgCl<sub>2</sub> (25 mM) and 1 µl of 2 mM dNTPs. The final volume was brought up to 25 µl with nuclease free water.

PCR conditions were set at 94°C for 4 min for the initial denaturation, followed by 35 cycles of denaturation at 94°C for 30 s, primer annealing at 60°C for 40 s, primer extension at 72°C for 1 min and the final extension at 72°C for 5 min.

The PCR mixture for FUM1 primers contained 2 µl of genomic DNA, 1 µl of each FUM1 primers (20 pmol), 0.5 µl of Taq DNA polymerase (3 U/µl), 2.5 µl of 10X PCR buffer, 2.5 µl of MgCl<sub>2</sub> (25 mM) and 1 µl of 2 mM dNTPs. The final volume was made up to 25 µl with nuclease free water.

The PCR conditions were 94°C for 4 min for the initial denaturation, followed by 35 cycles of denaturation at 94°C for 1 min, primer annealing at 60°C for 1 min, primer extension at 72°C for 1 min and the final extension at 72°C for 5 min. Amplified products were analysed on a 1.5% agarose gel in 1X TAE buffer (40 mM Tris acetate and 1.0 mM EDTA) and documented with a gel documentation system (UTP-Bio Doc, USA).

### **Sample Preparation for Thin Layer Chromatography**

PCR positive isolates of *F. proliferatum* (25) were cultured on rice medium for FB<sub>1</sub> detection. About 30 g of polished rice in 9 ml of sterile distilled water were autoclaved in a 250 ml conical flask. A conidial suspension (10<sup>6</sup>/ml) of each *F. proliferatum* isolate was prepared by adding 7 ml of sterile water to a Petri dish containing a 14-day-old fungal culture. Each flask containing the rice medium was inoculated with 5 ml of conidial suspension, mixed well and incubated at 28°C for 28 days in the dark. Rice cultures were oven-dried overnight at 60°C and finely ground in a laboratory mill. The powdered samples were stored at -20°C until further analysis (Desjardins *et al.* 2000).

### **Extraction and Cleanup**

FB<sub>1</sub> was extracted from the finely ground sample using a protocol developed by Rice *et al.* (1995). 10 g samples were placed into a 250 ml conical flask containing 50 ml of acetonitrile:water (ACN:water, 50+50, v/v) and the flask was covered and shaken for 30 min. 15 ml of the supernatant was filtered through a Whatman No. 4 filter paper. A C18 Sep Pak solid phase extraction (SPE) clean up column (Waters, USA) was preconditioned by rinsing with 2 ml ACN followed by 2 ml of 1% aqueous potassium chloride (KCl). The filtered extract (2 ml) and 1% KCl (6 ml) were mixed in a vial and applied to the column.

The solution flowed through the SPE column at a flow rate of 4 ml/min. The column was rinsed with 2 ml of 1% KCl followed by 2 ml of ACN:water (15+85, v/v). The rinse was discarded and air was forced through the column to expel all the rinse solution. FB<sub>1</sub> was eluted from the column with 2 ml of ACN:water (70+30, v/v).


Following extraction and clean up the samples were analysed by TLC (Bailey *et al.* 2005). Sample extracts (5 µl) and FB<sub>1</sub> standards (200 ng/ml and 400 ng/ml) were spotted on TLC plates (AluGram SIL G/UV 254, Machery Nagel, Germany). Separation was carried out in a mixture of 1-butanol:acetic acid:water (20+10+10, v/v/v). After drying, plates were sprayed with a solution containing methanol:0.5% p-anisaldehyde:acetic acid:sulphuric acid (85:0.5:10.5 v/v) and heated for 10 min at 110°C. The plates were observed under daylight conditions for the presence of purple FB<sub>1</sub> spots.

## RESULTS


In total, 109 paddy samples screened on MGA 2.5 containing chloramphenicol yielded 10 different *Fusarium* species namely, *F. solani* (Martius) Appel & Wollenweber emend (63.3%), *F. anthropilum* (A. Braun) Wollenweber (59.63%), *F. oxysporum* (55%), *F. semitectum* (54.12%), *F. sporotrichioides* Sherbakoff (50.45%), *F. proliferatum* (40.36%), *F. verticillioides* (24.77%), *F. graminearum* Schwabe (23.85%), *F. lateritium* Nees (16.51%) and *F. poae* (Peck) Wollenweber (14.67%). *F. proliferatum* isolates produced aerial mycelium with purple violet pigmentation on the agar medium. Many polyphialides were observed with abundant club shaped microconidia in long chains. The macroconidia were slender, thin walled, curved and 5-septate. Chlamydo spores were absent.

### PCR Amplification

The species-specific primers Fp3-F and Fp4-R used in this study amplified a 230 bp PCR products in all 65 isolates of *F. proliferatum*; such amplified products were not detected in other *Fusarium* species tested (Fig. 1). Furthermore, all 65 isolates of *F. proliferatum* and 27 isolates of *F. verticillioides* scored positive with the FUM1 set of primers by producing a 183 bp product, indicating their potential fumonisin producing ability. No such amplification was detected in other *Fusarium* species tested (Fig. 2).




**Figure 1:** Agarose gel (1.5%) showing PCR products using the Fp3-F/Fp4-R (230 bp) set of primers specific to *F. proliferatum*. Lane (M): (100 bp) DNA ladder; lane (2): *F. semitectum*; lane (3): *F. verticillioides*; lane (4–10): *F. proliferatum*.



**Figure 2:** Agarose gel (1.5%) showing the amplified products of fumonisin producing *F. proliferatum* and *F. verticillioides* with the FUM1 (183 bp) set of primers. Lane (M): (100 bp) DNA ladder; Lane (1): *F. semitectum*; lane (2): *F. oxysporum*; lane (3–4); *F. verticillioides*; lane (5–9): *F. proliferatum*.

### TLC Analysis

TLC revealed the occurrence of purple-coloured FB<sub>1</sub> spots having relative front (R<sub>f</sub>) values of 0.75, in all the 25 isolates of *F. proliferatum* (Fig. 3).



**Figure 3:** TLC plate showing purple coloured spots after spraying with a mixture of 0.5% p-anisaldehyde. Lane (1–2): FB<sub>1</sub> standards (200 ng and 400 ng, respectively); lane (3–15): culture extracts of FB<sub>1</sub> from *F. proliferatum* isolates.

### DISCUSSION

Mycotoxicological surveys of cereals are very important, especially in tropical countries, where temperature and relative humidity are favourable for fungal growth. Like all other cereals, paddy is susceptible to damage by mycotoxigenic fungi. After harvest, it is subjected to traditional sun drying for 3 to 5 days prior to being sent to the mills where it is stored for human and animal consumption. Due to this post harvest practice, paddy is susceptible to damage by fungi, especially the *Fusarium* species. *Fusariums* produce mycotoxins, such as fumonisins. Mycological analysis of 109 paddy seed samples revealed the presence of 10 different species of *Fusarium*. In terms of frequency, *F. solani* ranked the highest (63.3%). However, in terms of fumonisin producing ability, *F. proliferatum* (40.36%) ranked first. *F. proliferatum* is one of the dominant species on paddy and would probably be the main source of contamination by fumonisins. This finding is in agreement with earlier studies (Tonon *et al.* 1997; Pacin *et al.* 2002), which reported the *Fusarium* species (especially *F. proliferatum*) to frequently contaminate paddy. In a study by Abbas *et al.* (1998), paddy samples infected with *F. proliferatum* (40%) were found positive for FB<sub>1</sub> at levels of 4.3 µg/g. Infection of paddy with *F. proliferatum* in Korean polished rice revealed the presence of FB<sub>1</sub> at a concentration of 4.4–7.1 µg/g (Park *et al.* 2005).

Epidemiological studies have linked the human esophageal carcinoma to the consumption of maize with high incidences of *F. verticillioides* contamination (Marasas *et al.* 1981; Yoshizawa *et al.* 1994). Hence, there is a need for a rapid detection of the toxin producing fungi for the effective management of fumonisins in cereals. Molecular methods such as PCR have been developed for the

detection of fumonisins to replace traditional methods based on molecular probes (Jurado *et al.* 2005). Unlike methods involving culturing, PCR does not require the presence of viable organisms for detection and may be performed even with small amounts of sample.

We employed diagnostic methods based on PCR to accurately detect fumonisin producers. *F. proliferatum* were identified using species-specific primers based on IGS sequences. Additionally, the use of the FUM1 primers, designed based on the *fum1* gene, for the detection of fumonisin producing strains proved to be highly useful for identifying toxigenic *Fusarium* isolates. All 65 isolates tested scored positive with the species-specific primers. Furthermore, all 65 isolates of *F. proliferatum* were positive using the FUM1 primer set, indicating that all 65 isolates have the potential ability to produce fumonisin. Apart from the *F. proliferatum* species, 27 isolates of *F. verticillioides* also scored positive with the FUM1 set of primers.

The IGS region, which is commonly used for identification purposes in taxonomic studies, was found to be appropriate for the molecular detection of *F. proliferatum* in this study. The IGS regions contain high levels of sequence variability among the species of the same genus and allow differentiation of genetically related species (Edel *et al.* 2000; Kim *et al.* 2001; Konietzny & Greiner 2003; González-Jaén *et al.* 2004).

TLC, an economical analytical procedure, was employed for the detection of fumonisin produced by *F. proliferatum* isolates. TLC remains an important tool for mycotoxin detection in countries that often produce and export agricultural commodities but do not have expensive equipment at their disposal. As opposed to labour intensive and time consuming advanced high pressure liquid chromatography (HPLC) screening methods (Rottinghaus *et al.* 1992), TLC is a relatively easy and useful technique.

## CONCLUSION

The presence of *Fusarium* species in food and animal feed can be a threat to the food chain, as fumonisins are known to survive heat processing. The present study revealed the natural occurrence of fumonisin producing *F. proliferatum* species on paddy seeds produced in Karnataka (India). The risks associated with *F. proliferatum* contaminated paddy warrants the need for further analysis of this basic food that is intended for human and animal consumption.

## REFERENCES

- Abbas H K, Cartwright R D, Shier W T, Abouzied M M, Bird C B, Rice L G, Ross P F, Sciombato G L and Meredith F I. (1998). Natural occurrence of fumonisins in rice with *Fusarium* sheath rot disease. *Plant Disease* 82: 22–25.
- Bailly J D, Querin A, Tardieu D and Guerre P. (2005). Production and purification of fumonisins from a highly toxigenic *Fusarium verticillioides* strain. *Revue de Médecine Vétérinaire* 156 (11): 547–554.



- Bluhm B H, Cousin M A and Woloshuk C P. (2004). Multiplex real time PCR detection of fumonisin producing and trichothecenes producing groups of *Fusarium* species. *Journal of Food Protection* 67(3): 536–543.
- Booth C. (1977). *Fusarium: Laboratory guide to the identification of major species*. Kew, England: Commonwealth Mycological Institute.
- Bragulat M R, Martinez E, Castella G and Cabanes F J. (2004). Selective efficacy of culture media recommended for isolation and enumeration of *Fusarium* species. *Journal of Food Protection* 67: 207–211.
- Desjardins A E, Manandhar H K, Plattner R D, Manandhar G G, Poling S M and Maragos C M. (2000). *Fusarium* species from Nepalese rice and production of mycotoxins and gibberellic acid by selected species. *Applied and Environmental Microbiology* 66(3): 1020–1025.
- Domijan A, Peraica M, Jurjevic Z, Ivic D and Cvjetković B. (2005). Fumonisin B<sub>1</sub>, fumonisin B<sub>2</sub>, zearalenone and ochratoxin A contamination of maize in Croatia. *Food Additives and Contaminants* 22(7): 677–680.
- Edel V, Steinberg C, Gautheron N and Alabouvette C. (2000). Ribosomal DNA-targeted oligonucleotide probe and PCR assay specific for *Fusarium oxysporum*. *Mycological Research* 104(5): 518–526.
- FAO (2004). Food and Nutrition Division. Food and Agriculture Organization of the United Nations Viale delle Terme di Caracalla, Rome 00100 Italy. <http://www.fao.org/rice2004/en/f-sheet/factsheet3.pdf> (accessed on 12 February 2004).
- González-Jaén M T, Mirete S, Patiño B, Lopez-Errasquín E and Vázquez C. (2004). Genetic markers for the analysis of variability and for production of specific diagnostic sequences in fumonisin-producing strains of *Fusarium verticillioides*. *European Journal of Plant Pathology* 110: 525–532.
- Gonzalez H H, Resnik S L, Boca R T and Marasas W F O. (1995). Mycoflora of Argentinian corn harvested in the main production area in 1990. *Mycopathologia* 130: 29–36.
- Hinojo M J, Medina A, Valle-algarra F M, Gimeno-adelantado J V, Jiménez M and Mateo R. (2006). Fumonisin production in rice cultures of *Fusarium verticillioides* under different incubation conditions using an optimized analytical method. *Food Microbiology* 23: 119–127.
- Jurado M, Vázquez C, Patiño B and González-Jaén M T. (2005). PCR detection assays for the trichothecene-producing species *Fusarium graminearum*, *Fusarium culmorum*, *Fusarium poae*, *Fusarium equiseti* and *Fusarium sporotrichioides*. *Systematic and Applied Microbiology* 28(6): 562–568.
- Jurado M, Vázquez C, Marín S, Sanchis V and González-Jaén M T. (2006). PCR-based strategy to detect contamination with mycotoxigenic *Fusarium* species in maize. *Systematic and Applied Microbiology* 29(8): 681–689.

- Kim H J, Choi Y K and Min B R. (2001). Variation of the Intergenic Spacer (IGS) region of ribosomal DNA among *Fusarium oxysporum* formae speciales. *The Journal of Microbiology* 39(4): 265–272.
- Konietzny U and Greiner R. (2003). The application of PCR in the detection of mycotoxigenic fungi in foods. *Brazilian Journal of Microbiology* 34: 283–300.
- Leslie J F and Summerell B A. (2006). *The Fusarium laboratory manual* 1<sup>st</sup> ed. Ames, Iowa: Blackwell Publishing.
- Marasas W F O, Wehner F C, Van Rensburg S J and Van Schalkwyk D J. (1981). Mycoflora of corn produced in human esophageal cancer areas in Transkei, South Africa. *Phytopathology* 71: 792–796.
- Mathur S B and Kongsdal O. (2003). *Common laboratory seed health testing methods for detecting fungi*. Switzerland: International Seed Testing Association (ISTA).
- Pacin A M, Gonzalez H H L, Etcheverry M, Resnik S L, Vivas L and Espin S. (2002). Fungi associated with food and feed commodities from Ecuador. *Mycopathologia* 156: 87–92.
- Park J W, Choi S Y, Hwang H J and Kim Y B. (2005). Fungal mycoflora and mycotoxins in Korean polished rice destined for humans. *International Journal of Food Microbiology* 103: 305–314.
- Rheeder J P, Marasas W F O and Vismer H F. (2002). Production of fumonisin analogs by *Fusarium* species. *Applied and Environmental Microbiology* 68(5): 2101–2105.
- Rice L G, Ross P F, Dejong J, Plattner R D and Coats J R. (1995). Evaluation of a liquid chromatographic method for the determination of fumonisins in corn, poultry feed and *Fusarium* culture material. *Journal of AOAC International* 78: 1002–1009.
- Rottinghaus G E, Coatney C E and Minor H C. (1992). A rapid, sensitive thin layer chromatography procedure for the detection of fumonisin B<sub>1</sub> and B<sub>2</sub>. *Journal of Veterinary Diagnostic Investigation* 4: 326–329.
- Tonon S A, Marucci R S, Jerke G and Garcia A. (1997). Mycoflora of paddy and milled rice produced in the Northeastern Argentina and Southern Paraguay. *International Journal of Food Microbiology* 37: 231–235.
- Trung T S, Bailly J D, Querin A, Le Bras P and Guerre P. (2001). Fungal contamination of rice from South Vietnam, mycotoxinogenesis of selected strains and residues in rice. *Revue de Médecine Vétérinaire* 152(7): 555–560.
- Yoshizawa T, Yamashita A and Luo Y. (1994). Fumonisin occurrence in corn from high- and low-risk areas for human esophageal cancer in China. *Applied and Environmental Microbiology* 60(5): 1626–1629.
- Zhang Y P, Uyemoto J K and Kirkpatrick B C. (1998). A small-scale procedure for extracting nucleic acids from woody plants infected with various phytopathogens for PCR assay. *Journal of Virological Methods* 71: 45–50.