



Published in final edited form as:

Child Adolesc Psychiatr Clin N Am. 2014 January ; 23(1): . doi:10.1016/j.chc.2013.07.002.

Emotion Regulation: Concepts & Practice in Autism Spectrum Disorder

Carla A. Mazefsky^a and Susan W. White^b

^aAssistant Professor, Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, PA

^bAssociate Professor, Department of Psychology, Virginia Tech, Blacksburg, VA

Synopsis

The purpose of this article is to describe emotion regulation, and how emotion regulation may be compromised in autism spectrum disorder (ASD). This information may be useful for clinicians working with children with ASD who exhibit behavioral problems. Suggestions for practice are provided.

Keywords

Emotion regulation; autism spectrum disorder; therapy; behavioral problems; review

INTRODUCTION

Emotion Regulation Concepts

Imagine that you are driving to work and someone cuts you off. Your heart rate rapidly increases and you experience a wave of intense irritation, yet you manage to blare on your horn, simultaneously hit your brakes, and remain focused on safely driving. You have just engaged in effective emotion regulation (ER), which broadly encompasses the processes related to modifying your emotions to fit the context or meet one's goals^{1,2} (in this case, staying safe). Although the distinction is widely debated,³⁻⁵ emotion *regulation* differs from the *experience* of emotion itself, in that ER involves an attempt to modify the intensity or temporal features of an emotion [e.g. after the initial startle when you are cut off, you quickly experience anger when you see the driver texting (emotion); but, you attempt to keep the anger from escalating so that you can remain focused on safely driving (ER)]. ER processes can occur at the unconscious level (without realizing you are doing so, you maintain a level of fear that will keep you alert but not unable to act) and at a conscious

© 2013 Elsevier Inc. All rights reserved.

Corresponding Author: ^aCarla A. Mazefsky, Ph.D., Department of Psychiatry, University of Pittsburgh, 3811 O'Hara Street, Webster Hall, Suite 300, Pittsburgh, PA 15213, Phone: 412-246-5459, Fax: 412-246-5470, mazefskyca@upmc.edu.

Co-Author: ^bSusan W. White, PhD, Department of Psychology, Virginia Tech, 109 Williams Hall (0436), Blacksburg, VA 24061-0100, Phone: (540)231-6174, Fax (540)231-3652, sww@vt.edu

Conflicts of Interest:

Dr. Mazefsky: None

Dr. White: None

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

level (e.g. telling yourself you are okay after the incident is over). Further, ER can be response-focused as in this example, or antecedent-focused (prior to the emotion).

Most people have a characteristic and fairly stable ER style. A person's customary ER style can be generally adaptive or maladaptive, with the latter often associated with psychopathology and less appropriate behavior. Disrupted, or maladaptive, ER has been implicated as a mechanism underlying various psychiatric disorders, including depressive^{6,7} and anxiety⁸ disorders, and borderline personality disorder.⁹ Thus, poor ER is a transdiagnostic process that plays a role in many disorders in producing inappropriate emotional and behavioral reactions. The mechanisms that give rise to emotional dysregulation and how ER manifests itself, however, are more disorder-specific.⁹

Role of Emotion Regulation in ASD

Although much less studied in autism spectrum disorder (ASD) than in other psychiatric disorders, disrupted ER is likely to be a significant factor in producing aberrant behavior in ASD as well.^{10,11} One likely manifestation of ER failure in ASD is serious behavioral disturbance. Tantrums, uncontrolled outbursts, aggression, and self-injury are often interpreted as defiant or deliberate. While this interpretation is likely accurate in some circumstances, it is more often the case that these inappropriate behavioral reactions stem from ineffective management of emotional states in response to stress or overstimulation.¹²

Absent or impaired ER may be a more parsimonious explanation of serious behavioral disturbance in ASD than psychiatric comorbidity. Psychiatric diagnoses are difficult to reliably make in ASD for a variety of reasons including lack of measures validated for use with this population, difficulty assessing certain symptoms in nonverbal individuals, inadequate insight and poor temporal reporting, unique manifestations of distress in ASD, and the challenges involved in interpreting and differentiating symptoms that could be attributed to ASD or a secondary disorder (e.g. lack of positive affect as part of ASD or due to depression).^{13,14} For all of these reasons, there is growing concern that psychiatric diagnoses may be over-used in ASD.¹⁵ Many 'secondary' psychiatric problems may be more accurately conceptualized as part of the ASD itself or may stem from a fundamental problem in ER.¹⁰

ASD-Related Factors that Impede Effective Emotion Regulation

Many characteristics of ASD may interfere with effective ER (see Figure 1).¹⁶ First, alexithymia, or difficulty identifying, distinguishing, and describing emotions, has been well-documented in ASD.^{17,18,19} Although not essential for all forms of ER, recognizing and understanding your own emotions is necessary for effortful ER.²⁰ Labeling of one's emotion has been proposed as critical to successful ER,²¹ and being able to communicate to others about your emotional state is also involved in interpersonal ER aspects, such as joint problem-solving or sharing of one's emotions. Given that language competence is associated with emotional competence in typical development,²² it is also conceivable that the language and communication impairments common in ASD affect development or regulatory abilities.

As proposed by Samson, Huber, and Gross,²³ core deficits in theory of mind, or ability to take others' perspectives cognitively and affectively and to recognize one's own state of mind, may be related to poor ER. Some regulatory strategies (e.g., intentionally changing one's cognitive frame or perspective) are inherently related to perspective-taking ability. The social and cognitive deficits that define ASD also create ER challenges, particularly given that adaptive ER is context-dependent and requires one to be able to accurately identify critical aspects of the situation.¹⁰ Even among cognitively higher-functioning

individuals with ASD, there are deficits in the processing and integration of complex information.²⁴ Children with ASD may focus on the wrong information or misinterpret others' intentions due to failures in perspective-taking and lack of appreciation of others' perceptions and experiences (e.g., theory of mind deficits),²⁵ which could increase frustration. Difficulty accurately perceiving others' social and emotional cues may also interfere with the timing and implementation of effortful ER strategies. Impulsivity and impaired inhibition, which are present in as many as 50% of children with ASD,²⁶ could similarly interfere with ER. Specifically, even if the child knows what he should do when upset (e.g., take deep breaths), inability to inhibit the more potent and automatic response (e.g., hitting and yelling) could result in ineffective ER. If a child with ASD does stop to think about the situation before reacting, the tendency to be rigid and engage black-and-white thinking may still preclude a flexible and adaptive ER response.¹⁰ Even on structured neuropsychological tests in laboratory settings (which are slower and more straightforward than the fast-moving interactions that require ER in daily life), problem-solving deficits in ASD have been documented.²⁴

Finally, children with ASD may be predisposed to problems with emotional control given differences in arousal and underlying neural circuitry. Children with ASD are known to have unusual reactions to sensory information and are often sensitive to change. These differences in responses to the environment may increase reactivity and lability.²⁷ Further, although results are not entirely consistent, there is evidence to suggest that at least some of the ASD population is physiologically hyper-aroused,²⁸ particularly those who present with anxiety symptoms. Finally, although neuroimaging research on ER in ASD is limited, there is evidence that the neural structures implicated in ER in other populations differ in ASD either in function, size, or circuitry with other parts of the brain.²⁹⁻³¹ Finally, high rates of mood and anxiety disorders in the first degree family members of children with ASD,^{32,33} and conceptualization of such problems as part of the broader autism phenotype,³⁴ raise the possibility of an underlying genetic predisposition to ER problems in ASD as well. For all of these reasons, we assert that it is more appropriate to consider impaired ER as inherent to ASD itself (e.g., stemming from or directly related to having ASD).

Consistent with the heuristic of transdiagnostic processes proposed by Nolen-Hoeksema and Watkins,³⁵ ER deficits within ASD may be conceptualized as a 'proximal' risk factor for expressed psychopathology. Poor ER is proximal in that it underlies the observed psychopathology (e.g., self-injury, anxiety). Impaired ER, likewise, is influenced by 'distal' factors such as genetic predisposition, heightened baseline physiological arousal, and atypical neural circuitry.^{36,37} The influence of these distal factors on a person's ER deficits may be strengthened by problems such as alexithymia, poor perspective-taking, and inadequate response inhibition. Moderating factors then operate to influence how the impaired ER might be expressed in a given individual and at a given time (multifinality). The youngster with ASD with poor ER ability, when faced with the need to interact with others in ambiguous situations (e.g., 6th grade lunch-room), for instance, might become extremely anxious or agitated.

ILLUSTRATIVE CASE EXAMPLE

Johnny is a 12 year old boy who was diagnosed with high-functioning autism at the age of 11, based on scores above the autism cut-off on the *Autism Diagnostic Interview-Revised* and above the autism spectrum disorder cut-off on the *Autism Diagnostic Observation Schedule*. He obtained a Full Scale IQ score of 90. Johnny's family history is notable for several psychiatric disorders, with three first-degree relatives with a depression history, one with bipolar disorder, three with anxiety disorder diagnoses, and two with attention-deficit/hyperactivity disorder (ADHD). Johnny's first diagnosis, at the age of 7 years, was ADHD,

which was followed 6 months later by a diagnosis of bipolar disorder. Between the ages of 8 and 11 years old, he was on 23 different medications in an attempt to manage his emotions and behavior. Unfortunately, the medications, particularly those utilized as a treatment for bipolar disorder, were not effective, and he developed tics. During this period of time, he also had three psychiatric hospitalizations. A comprehensive assessment of his psychiatric history using the *Autism Comorbidity Interview*³⁸ at age 12 supported a past depressive disorder diagnosis but failed to confirm the presence of bipolar disorder. Specifically, there was no evidence of any mania or hypomania. It was clear, however, that problems with self-control and ER were present as a very young child. He was an irritable toddler, “easily triggered” and upset, per his mother’s report. When upset, he would bang his head or pull his hair out. Although he did “bounce between being sad or depressed and irritable or angry,” further probing revealed clear ASD-related triggers to his mood changes, such as loud, sudden noises, his schedule being changed, or a meal being late triggering a “meltdown”, and poor social-communication skills (e.g. misunderstanding sarcasm) leading to frustration. He had very poor insight into these concerns, including low scores on several self-report psychiatric screeners. His treatment plan includes a focus on increasing his self-awareness and emotion understanding, reducing the number of psychotropic medications, conducting a functional behavior assessment to identify potential antecedents to his meltdowns, addressing the identified external triggers, and teaching him ER skills.

CURRENT APPROACHES TO PRACTICE

At this time, we lack evidence-based tools with which to accurately assess ER that have been validated in clients with ASD. In light of difficulties with insight and labeling of one’s own emotions, self-report is likely to be insufficient on its own when completing the assessment and case formulation phase of treatment. Observational approaches (e.g., seeing how client responds to a mildly frustrating situation) can yield important information. It is important to assess the client’s general attention to emotional states, capacity to connect to these states and describe them, and ability to differentiate among similar but unique emotions (e.g., upset and angry). A thorough evaluation should also try to separate core impairments in receptive and expressive communication and social interaction from maladaptive ER. The types of regulatory strategies primarily relied on by the client, and those that may be under-developed, should inform treatment planning.

Although there is little controlled clinical research on the efficacy of ER-focused interventions for people with ASD, there are several plausible and potentially effective intervention approaches to consider. Drawing from the extant research on treatment of co-occurring problems, such as anxiety and mood disturbance, the predominant psychosocial approach is cognitive-behavioral therapy (CBT).^{39–41} CBT-based approaches for clients with ASD often incorporate content to address ER deficits.^{42,43} There is great variability in the degree to which ER training is included in such interventions, however, and evaluation of the relative import of various treatment components, including ER training, has not been conducted. Few treatment programs have been developed explicitly to improve ER ability in clients with ASD. The *Exploring Feelings* program⁴² is a CBT-based intervention for school-age children, with specific curricula for anxiety and for anger. It includes affect education, cognitive restructuring, and appropriate strategies to manage intense emotion. This program was recently modified for use with younger children (5–7 years) with ASD,⁴⁴ and results of an initial open trial indicated improvement in emotion lability and regulation.⁴⁵

In light of evidence that individuals with ASD tend to engage in suppression and struggle with reappraisal, targeting increased cognitive flexibility in treatment is recommended.²³ There is empirical evidence that many of the cognitive regulatory strategies (i.e.,

suppression, rumination) load onto a single latent factor and that these cognitive strategies are more strongly associated with psychopathology than adaptive cognitive strategies (e.g., reappraisal, problem-solving).⁹ Aldao and Nolen-Hoeksema⁹ proposed that clients may develop a 'default' regulatory approach that, in essence, overwhelms their ability to use newly learned, more adaptive strategies such as reappraisal. As such, treatments outside of traditional CBT, which tends to focus on restructuring of thoughts to alter feelings, may be beneficial. For example, meditation, mindfulness-training, and acceptance-based approaches may help the client to reduce attempts to suppress feelings, which might enhance his or her willingness and ability to develop stronger adaptive regulatory strategies.

Psychoeducation and acceptance-based approaches may be especially helpful, given the chronic and pervasive nature of ASD. Indeed, a premise of dialectical behavior therapy (DBT⁴⁶), a well-supported treatment for borderline personality disorder which is characterized by extreme emotion dysregulation, is balancing acceptance of self (as is) with desire for change. Although there has been consideration of how to adapt DBT for clients with ASD,⁴⁷ there have been no clinical trials and there is no published treatment outcome research. Mindfulness and acceptance-based approaches (MABIs), such as DBT, have been used extensively to treat problems with ER. MABIs differ from traditional CBT primarily in how the patient's relationship to his/her feelings and thoughts is conceptualized. Rather than identification and alteration of maladaptive or incorrect thoughts and unhelpful feelings (CBT), MABIs strive to help the patient change his relationship with (or view of) the problem, become less fused with his own thoughts (accepting a thought as just a thought), and behave in a fashion consistent with his goals and values.⁴⁸ MABIs are associated with strong and durable improvements in symptoms of anxiety and depression and increased use of adaptive ER strategies,⁴⁹ and have recently been shown to be as effective as CBT for treatment of anxiety.⁵⁰

Regardless of treatment approach, a more intensive focus on developing the client's emotional awareness and ability to recognize and report on his emotional state is often required for clients with ASD, compared to those with other (non-ASD) diagnoses. Improving emotional insight and regulatory skills can be complicated by other treatments the client is receiving, especially medication. Between one-half and three-quarters of child clients with ASD are prescribed psychoactive medications,⁵¹⁻⁵³ often for target problems that might arguably be rooted in poor ER, such as irritability.⁵⁴ If a medication is dampening an emotional response, this may further complicate efforts to work on increasing emotional insight and awareness.

Often it can be helpful to use visual strategies to work on identifying, understanding and communicating about one's own emotions. One example of a visual system that is widely used to teach these basic emotion skills is *The Incredible Five Point Scale*.⁵⁵ Although there has not been any systematic research on this program, it was developed based on clinical experience, was designed specifically for ASD and related populations, and has been applied across the range of intellectual and verbal abilities. In short, it provides a metric for helping the child to identify and communicate varying degrees of emotion from a scale of 1 (This never bothers me) to 5 (This could make me lose control). The skills are taught using specific situations identified by the child and/or a caregiver or a standard set of picture cards of different situations, which the child then learns to place into a chart with pockets corresponding to one thru five. The intent is that the child will learn to notice these situations, have a way to communicate about his or her emotions (e.g. I am at a 3, either verbally or with picture cards), and then strategies can be taught regarding what to do in those situations. Although this is just one example, and it is a fairly narrow set of skills, the concepts it embodies – e.g. incorporating visual cues, making emotion as concrete and possible, and individualizing ER interventions – are all useful approaches to consider.

Behaviors and deficits that are core to the ASD diagnosis (e.g., impaired reciprocal social behavior) must be considered and clinicians must be mindful of ASD-specific factors (both proximal and distal) that might influence ER ability. Related to this, moderators of poor ER manifestation can sometimes be managed antecedently (e.g., planning for where to sit in the lunchroom). A common approach to handling problematic behavior is to complete a functional behavior analysis and, based on results, provide individualized positive behavioral supports (see also Doehring & Hagopian, this issue).⁵⁶ This approach often focuses on modifying external factors that may trigger negative emotional reactions. This type of intervention is rooted in a long history of behavioral analysis that has a wealth of support for its effectiveness dating back to the 1960s.⁵⁷ However, a limitation is that it tends to be focused on specific problems and situations and thus the impact may not generalize to other stress-provoking situations. Thus, while it is always useful to identify any environmental factors that may be exacerbating the situation, this needs to be supplemented with teaching of appropriate alternative behaviors and skills.

CONCLUSION AND FUTURE DIRECTIONS

In sum, it is important to consider ER deficits when addressing severe behavioral disturbance and situations of apparent psychiatric comorbidity in ASD. Poor regulation of emotions is likely to underlie many of the observed manifestations of both internalizing and externalizing concerns. Further, impaired ER can exacerbate problems with attention, communication, problem-solving, and social interaction.⁵⁸ Above, we outlined many characteristics of ASD that may directly contribute to impaired ER in this population, and how ER impairment then acts as a proximal risk factor for the expression of psychopathology in clients with ASD. Unfortunately, research on ER in ASD is in its infancy. Further research is needed to better understand the specific mechanisms involved.

Measures that are sensitive to ER deficits in ASD need to be developed and validated. Accurate assessment is critical, both in terms of its role in treatment planning,¹³ and in determining outcome in clinical trials. For many reasons, standard measures developed for other populations may not be clinically sensitive or appropriate for use in ASD. Standard questionnaires are often verbally loaded with items such as “complains of” that make them inapplicable to the limited- and non-verbal ASD population. Further, ER problems may manifest differently in ASD.^{15,38,59} Thus, in order to improve both our understanding and treatment of ER problems in ASD, assessment may need to be addressed first.

Given the growing body of research implicating ER deficits in many of the psychiatric and behavioral problems manifested by clients with ASD, we propose that a focus on ER and its possible expression in people with ASD be considered in research on treatment development and evaluation. Although we described some treatment strategies that incorporate ER as a component of the intervention, there is very little clinical trials research in ASD explicitly focused on ER. Further, some treatments that effectively improve ER in other populations, such as acceptance and mindfulness-based interventions, have yet to be tested in ASD. There remains a critical need to develop and study effective treatments for the range of manifestations of ER concerns in ASD. The field of affective neuroscience has much to offer in this respect. It may be possible, for instance, to develop interventions that target ER capacity directly. Doing so may have cascading benefits on multiple expressed problems (e.g., irritability, anxiety). Although this type of basic and applied research is needed across the lifespan in ASD, it is notable that there have been no published ER studies of adolescents with ASD. Given that this is a developmental period characterized by a heightened risk for psychopathology and emotional reactivity⁶⁰, hormonal and neural changes that promote cognitive flexibility and increase the saliency of social incentives (e.g., peer approval),⁶¹ and complex and often highly ambiguous social challenges that require

effective regulation of emotions for successful navigation, understanding the role of ER impairment in adolescence may be a priority.

Acknowledgments

Funding:

Dr. Mazefsky: NICHD K23CHD060601

Dr. White: NIMH K01MH079945

References

1. Thompson, RA. Monographs of the Society for Research in Child Development. Chicago, Illinois: University of Chicago Press; 1994. Emotion regulation: A theme in search of definition; p. 25-52.
2. Gross JJ. The emerging field of emotion regulation: an integrative review. *Rev Gen Psychol.* 1998; 2(5):271–299.
3. Cole PM, Martin SE, Dennis TA. Emotion regulation as a scientific construct: methodological challenges and directions for child development research. *Child Dev.* 2004; 75(2):317–33. [PubMed: 15056186]
4. Eisenberg N, Spinrad TL. Emotion-related regulation: sharpening the definition. *Child Dev.* 2004; 75(2):334–9. [PubMed: 15056187]
5. Campos JJ, Frankel CB, Camras L. On the nature of emotion regulation. *Child Dev.* 2004; 75(2): 377–94. [PubMed: 15056194]
6. Siener S, Kerns KA. Emotion regulation and depressive symptoms in preadolescence. *Child Psychiatry Hum Dev.* 2012; 43(3):414–30. [PubMed: 22160271]
7. Rieffe C, Oosterveld P, Terwogt MM, Mootz S, Van Leeuwen E, Stockmann L. Emotion regulation and internalizing symptoms in children with autism spectrum disorders. *Autism.* 2011; 15(6):655–70. [PubMed: 21733959]
8. Cisler JM, Olatunji BO, Feldner MT, Forsyth JP. Emotion Regulation and the Anxiety Disorders: An Integrative Review. *J Psychopathol Beh Assess.* 2010; 32(1):68–82.
9. Aldao A, Nolen-Hoeksema S, Schweizer S. Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clin Psychol Rev.* 2010; 30(2):217–37. [PubMed: 20015584]
10. Mazefsky CA, Herrington J, Siegel M, et al. The Role of Emotion Regulation in Autism Spectrum Disorders. *J Am Acad Child Psychiatry.* 2013 in press.
11. Mazefsky CA, Pelphrey KA, Dahl RE. The need for a broader approach to emotion regulation research in autism. *Child Dev Perspect.* 2012; 6(1):92–97. [PubMed: 22639681]
12. Konstantareas MM, Stewart K. Affect regulation and temperament in children with Autism Spectrum Disorder. *J Autism Dev Dis.* 2006; 36(2):143–54.
13. Mazefsky, CA.; White, SW. The role of assessment in guiding treatment planning for youth with ASD. In: Scarpa, A.; White, S.; Attwood, T., editors. *Promising CBT Interventions for Children and Adolescents with High Functioning ASDs.* New York, NY: Guildford Publications;
14. Mazefsky, CA.; Filipink, R.; Link, J.; Lubetsky, MJ. Medical evaluation and co-morbid psychiatric disorders. In: Lubetsky, MJ.; Handen, BL.; McGonigle, JJ., editors. *Autism Spectrum Disorder.* Oxford University Press; 2011.
15. Mazefsky CA, Oswald DP, Day TN, Eack SM, Minshew NJ, Lainhart JE. ASD, a psychiatric disorder, or both? Psychiatric diagnoses in adolescents with high-functioning ASD. *J Clin Child Adol Psychol.* 2012; 41(4):516–23.
16. Mazefsky CA. Managing Problem Emotions and Behaviors in Children With ASD: An Assessment-Driven Three-Step Approach. *Perspectives on Language Learning and Education.* 2011; 19:38–47.
17. Berthoz S, Hill EL. The validity of using self-reports to assess emotion regulation abilities in adults with autism spectrum disorder. *Eur Psychiatry.* 2005; 20(3):291–8. [PubMed: 15935431]

18. Mazefsky CA, Kao J, Oswald DP. Preliminary evidence suggesting caution in the use of psychiatric self-report measures with adolescents with high-functioning autism spectrum disorders. *Res Autism Spectr Dis*. 2011; 5(1):164–174.
19. Rieffe C, Meerum Terwogt M, Kotronopoulou K. Awareness of single and multiple emotions in high-functioning children with autism. *J Autism Dev Dis*. 2007; 37(3):455–65.
20. Ciarrochi J, Heaven PCL, Supavadeeprasit S. The link between emotion identification skills and socio-emotional functioning in early adolescence: a 1-year longitudinal study. *J Adol*. 2008; 31(5): 565–82.
21. Barrett LF, Gross J, Christensen TC, Benvenuto M. Knowing what you're feeling and knowing what to do about it: Mapping the relation between emotion differentiation and emotion regulation. *Cogn Emot*. 2001; 15(6):713–724.
22. Saarni, C. *The Development of Emotional Competence*. New York, NY: Guilford Press/Guilford Publications, Inc; 1999.
23. Samson AC, Huber O, Gross JJ. Emotion regulation in Asperger's syndrome and high-functioning autism. *Emotion*. 2012; 12(4):659–65. [PubMed: 22642342]
24. Williams DL, Goldstein G, Minshew NJ. Neuropsychologic functioning in children with autism: further evidence for disordered complex information-processing. *Child Neuropsychol*. 2006; 12(4–5):279–98. [PubMed: 16911973]
25. Baron-Cohen, S. *Mindblindness: An Essay on Autism and Theory of Mind*. Palatino, CA: MIT press; 1997.
26. Murray MJ. Attention-deficit/Hyperactivity Disorder in the context of Autism spectrum disorders. *Curr Psychiatry Rep*. 2010; 12(5):382–8. [PubMed: 20694583]
27. Dunn W, City K. The Impact of Sensory Processing Abilities on the Daily Lives of Young Children and Their Families: A Conceptual Model. *Infants Young Child*. 1997; 9(4):24–35.
28. Rogers SJ, Ozonoff S. Annotation: what do we know about sensory dysfunction in autism? A critical review of the empirical evidence. *J Child Psychol Psychiatry*. 2005; 46(12):1255–68.
29. Monk CS. The development of emotion-related neural circuitry in health and psychopathology. *Dev Psychopath*. 2008; 20(4):1231–50.
30. Herrington, JD.; Schultz, R. *Understanding Neuropsychiatric Disorders: Insights From Neuroimaging*. Cambridge: Cambridge University Press; 2010. *Neuroimaging of developmental disorders*.
31. Mazefsky CA, Minshew NJ. Clinical pearl: The spectrum of autism-from neuronal connections to behavioral expression. *Virtual Mentor*. 2010; 12(11):867–872. [PubMed: 23186793]
32. Mazefsky CA, Folstein SE, Lainhart JE. Overrepresentation of mood and anxiety disorders in adults with autism and their first-degree relatives: what does it mean? *Autism Res*. 2008; 1(3): 193–7. [PubMed: 19360666]
33. Mazefsky CA, Conner CM, Oswald DP. Association between depression and anxiety in high-functioning children with autism spectrum disorders and maternal mood symptoms. *Autism Res*. 2010; 3(3):120–7. [PubMed: 20578069]
34. Gerdts J, Bernier R. The broader autism phenotype and its implications on the etiology and treatment of autism spectrum disorders. *Autism Res Treat*. 2011; 2011:545901. [PubMed: 22937250]
35. Nolen-Hoeksema S, Watkins ER. A Heuristic for Developing Transdiagnostic Models of Psychopathology: Explaining Multifinality and Divergent Trajectories. *Perspect Psychol Science*. 2011; 6(6):589–609.
36. Nummenmaa L, Engell AD, Von dem Hagen E, Henson RNA, Calder AJ. Autism spectrum traits predict the neural response to eye gaze in typical individuals. *Neuro Image*. 2012; 59(4):3356–63. [PubMed: 22062191]
37. Redcay E. The superior temporal sulcus performs a common function for social and speech perception: implications for the emergence of autism. *Neurosci Biobehav Rev*. 2008; 32(1):123–42. [PubMed: 17706781]
38. Leyfer OT, Folstein SE, Bacalman S, et al. Comorbid psychiatric disorders in children with autism: interview development and rates of disorders. *J Autism Dev Dis*. 2006; 36(7):849–61.

39. Lang R, Regester A, Lauderdale S, Ashbaugh K, Haring A. Treatment of anxiety in autism spectrum disorders using cognitive behaviour therapy: A systematic review. *Dev Neurorehab.* 2010; 13(1):53–63.
40. Lickel A, Maclean WE, Hepburn ABS. Assessment of the Prerequisite Skills for Cognitive Behavioral Therapy in Children with and Without Autism Spectrum Disorders. *J Autism Dev Dis.* 2012:992–1000.
41. Scarpa, A.; White, SW.; Attwood, T., editors. *Cognitive-Behavioral Interventions for Children and Adolescents with High-Functioning Autism Spectrum Disorders.* New York, NY: Guilford Publications;
42. Attwood T. Cognitive Behaviour Therapy for Children and Adults with Asperger's Syndrome. *Behav Change.* 2004; 21(3):147–161.
43. Reaven J, Blakeley-Smith A, Culhane-Shelburne K, Hepburn S. Group cognitive behavior therapy for children with high-functioning autism spectrum disorders and anxiety: a randomized trial. *J Child Psychol Psychiatry.* 2012; 53(4):410–419. [PubMed: 22435114]
44. Scarpa, A.; Wells, A.; Attwood, T. *Exploring Feelings for Young Children with High-functioning Autism or Asperger's Disorder.* London: Jessica Kingsley Publishers; 2013.
45. Scarpa A, Reyes NM. Improving Emotion Regulation with CBT in Young Children with High Functioning Autism Spectrum Disorders: A Pilot Study. *Behav Cog Psychother.* 2011; 39(4):495–500.
46. Linehan, MM. *Cognitive-behavioral treatment of borderline personality disorder.* Guilford Press/Guilford Publications, Inc; 1993. p. 558
47. Hartmann, K.; Urbano, M.; Manser, K.; Okwara, L. Modified Dialectical Behavior Therapy to Improve Emotion Regulation in Autism Spectrum Disorders. In: Richardson, C.; Wood, R., editors. *Autism Spectrum Disorders.* Nova Science Publishers; 2012. p. 41-72.
48. Vøllestad J, Nielsen MB, Nielsen GH. Mindfulness- and acceptance-based interventions for anxiety disorders: a systematic review and meta-analysis. *Br J Clin Psychol.* 2012; 51(3):239–60. [PubMed: 22803933]
49. Kumar S, Feldman G, Hayes A. Changes in Mindfulness and Emotion Regulation in an Exposure-Based Cognitive Therapy for Depression. *Cogn Ther Res.* 2008; 32(6):734–744.
50. Arch JJ, Eifert GH, Davies C, Vilardaga JCP, Rose RD, Craske MG. Randomized clinical trial of cognitive behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for mixed anxiety disorders. *J Consult Clin Psychol.* 2012; 80(5):750–65. [PubMed: 22563639]
51. Aman MG, Lam KSL, Van Bourgondien ME. Medication patterns in patients with autism: temporal, regional, and demographic influences. *J Child Adol Psychopharm.* 2005; 15(1):116–26.
52. Oswald DP, Sonenklar NA. Medication use among children with autism spectrum disorders. *J Child Adol Psychopharm.* 2007; 17(3):348–55.
53. Siegel M. Psychopharmacology of autism spectrum disorder: evidence and practice. *Child Adol Psychiatr Clin N Am.* 2012; 21(4):957–73.
54. Williamson ED, Martin A. Psychotropic medications in autism: practical considerations for parents. *J Autism Dev Dis.* 2012; 42(6):1249–55.
55. Buron, KD.; Curtis, M. *The Incredible 5-Point Scale: Assisting Students with Autism Spectrum Disorders in Understanding Social Interactions and Controlling Their Emotional Responses.* Shawnee Mission, KS: Autism Asperger Publishing Company; 2003.
56. Mazefsky, CA.; Handen, BL. Addressing behavioral and emotional challenges in school-aged children with ASD. In: Lubetsky, M.; McGonigle, J.; Handen, B., editors. *Autism Spectrum Disorder.* Oxford University Press; 2011.
57. Iwata BA, Dorsey MF, Slifer KJ, Bauman KE, Richman GS. Toward a functional analysis of self-injury. *J App Beh Anal.* 1994; 27(2):197–209.
58. Laurent AC, Otr L, Rubin E. Challenges in emotional regulation in asperger syndrome and high-functioning autism. *Top Lang Dis.* 2004; 4(24):286–297.
59. Magnuson KM, Constantino JN. Characterization of depression in children with autism spectrum disorders. *J Dev Beh Ped.* 2011; 32:332–340.
60. Dahl RE, Gunnar MR. Heightened stress responsiveness and emotional reactivity during pubertal maturation: implications for psychopathology. *Dev Psychopath.* 2009; 21(1):1–6.

61. Crone EA, Dahl RE. Understanding adolescence as a period of social-affective engagement and goal flexibility. *Nat RevNeurosci*. 2012; 13(9):636–50.

Key Points

- Emotion regulation involves modulating the temporal features, intensity, or valence of one's emotions in the service of adaptive or goal-directed behavior
- Disrupted ER may be inherent in ASD
- Impaired ER may be a more parsimonious explanation than psychiatric comorbidity for severe behavioral disturbances observed in ASD
- Few interventions have been developed to explicitly target ER processes in ASD
- ER may be addressed (even if not labeled as such) in some existing psychosocial treatments used in ASD, including the provision of positive behavioral supports, enhancing emotional language, and modified cognitive-behavioral therapy
- Areas of future need include the development and validation of measures to assess ER in ASD for treatment planning and evaluation purposes, as well as the development of interventions to promote ER that incorporate the unique characteristics of ASD




Fig. 1. Characteristics of ASD that may contribute to emotion dysregulation