



Asbjørn Mohr Drewes, MD, PhD, DMSc, Professor, Series Editor

Animal models of pancreatitis: Can it be translated to human pain study?

Jing-Bo Zhao, Dong-Hua Liao, Thomas Dahl Nissen

Jing-Bo Zhao, Dong-Hua Liao, Mech-Sense, Department of Gastroenterology and Surgery, Aalborg University Hospital, DK 9000 Aalborg, Denmark

Jing-Bo Zhao, Dong-Hua Liao, Institute of Clinical Medicine, Aarhus University, DK 8200 Aarhus N, Denmark

Jing-Bo Zhao, Dong-Hua Liao, The College of Bioengineering, Chongqing University, Chongqing 400016, China

Thomas Dahl Nissen, Biomedical Research Laboratory, Aalborg University Hospital, DK 9000 Aalborg, Denmark

Author contributions: Zhao JB, Liao DH and Nissen TD designed, drafted, revised and finally approved this paper.

Correspondence to: Jing-Bo Zhao, MD, PhD, Mech-Sense, Department of Gastroenterology and Surgery, Aalborg University Hospital, Sdr. Skovvej 15, DK 9000 Aalborg, Denmark. jz@rn.dk

Telephone: +45-9-9326907 Fax: +45-9-9326801

Received: June 13, 2013 Revised: August 12, 2013

Accepted: August 20, 2013

Published online: November 14, 2013

are also addressed.

© 2013 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Animal model; Pancreatitis; Visceral pain; Mechanism

Core tip: Choosing the right model of pancreatitis is difficult and the scientific rationale needs to be carefully considered. Furthermore, no model of pancreatitis parallels all classical symptoms and the question under investigation is of importance when choosing a model. One of the main symptoms of chronic pancreatitis is visceral pain and in order to improve the pain treatment and obtain more knowledge about the physiology behind the pancreatitis associated visceral pain, animal models of pancreatitis associated visceral pain are needed.

Abstract

Chronic pancreatitis affects many individuals around the world, and the study of the underlying mechanisms leading to better treatment possibilities are important tasks. Therefore, animal models are needed to illustrate the basic study of pancreatitis. Recently, animal models of acute and chronic pancreatitis have been thoroughly reviewed, but few reviews address the important aspect on the translation of animal studies to human studies. It is well known that pancreatitis is associated with epigastric pain, but the understanding regarding to mechanisms and appropriate treatment of this pain is still unclear. Using animal models to study pancreatitis associated visceral pain is difficult, however, these types of models are a unique way to reveal the mechanisms behind pancreatitis associated visceral pain. In this review, the animal models of acute, chronic and un-common pancreatitis are briefly outlined and animal models related to pancreatitis associated visceral pain

Zhao JB, Liao DH, Nissen TD. Animal models of pancreatitis: Can it be translated to human pain study? *World J Gastroenterol* 2013; 19(42): 7222-7230 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v19/i42/7222.htm> DOI: <http://dx.doi.org/10.3748/wjg.v19.i42.7222>

INTRODUCTION

Pancreatitis represents a common disorder of the gastrointestinal tract. Acute pancreatitis (AP) has an incidence ranged from 4.9 to 35 per 100000 populations^[1], whereas chronic pancreatitis (CP) has an incidence from 2.4 to 4.4 per 100000 populations^[2]. The etiology of this disease is complex and so far a variety of environmental factors including alcohol abuse, nicotine habits, hereditary factors, efferent duct obstructions, immunological factors and rare metabolic factors have all been described. However,

the pathophysiology of AP and CP remains poorly defined^[3]. As a result appropriate therapies are still limited, and prognosis has not improved to date, which is mainly due to the lack of a satisfactory animal model of pancreatitis^[4,5].

It is well known that pancreatitis is associated with visceral pain, however, the understanding of pain signaling related to pancreatitis is poor^[6]. In order to facilitate the development of new pharmaceutical treatments for AP and CP, characterization of the mediators and receptors or ion channels on the sensory nerve terminals and the pathways of the pain signaling are needed. Therefore, in this aspect, the animal models of pancreatitis are needed in parallel in order to explore the mechanism behind pancreatitis associated visceral pain, as this is difficult to study in humans.

In this review, we briefly outline the animal models of acute, chronic and un-common pancreatitis as well as animal models related to pancreatitis associated visceral pain.

ANIMAL MODELS OF ACUTE PANCREATITIS

AP is an inflammatory condition of the pancreas characterized clinically by abdominal pain and elevated levels of pancreatic enzymes in the blood^[7]. Other characteristics of AP include edema, acinar cell necrosis, hemorrhage, and severe inflammation of the pancreas. Severe AP may lead to systemic inflammatory response syndrome and multi-organ dysfunction syndrome, which account for the high mortality rates of AP^[8,9]. As it is difficult to study AP in the clinic, animal studies are important in order to understand the pathogenesis of AP, however an AP model which is strictly comparable to human AP is still needed. The current animal models of AP have contributed to our knowledge of mechanisms involved in early cellular events, pathogenesis and pathophysiology of AP^[10,11]. We have illustrated the summary of existing AP animal models in Table 1^[12-59]. Details of different AP animal models including advantages, disadvantages and clinical relevance can be found in a recently published review^[4]. From a methodological aspect, selecting the appropriate AP animal model depends on the objectives of each study as different animal models are targeted to different AP features. For developing the effective treatment for AP in the clinic, continued investigation of AP animal models are needed.

ANIMAL MODELS OF CHRONIC PANCREATITIS

A recently published review^[5] has described the most frequently used and best established models for CP in animals. The majority of the animal models are rodent models, since mice and rats are easy to handle and there is a steadily increasing number of genetic models ob-

Table 1 Different animal models of acute pancreatitis

Methods	Models and examples
Non-invasive	Hormone-induced
	Acute caerulein pancreatitis of rats ^[12] , mice ^[13] , dogs ^[14] , and syrian hamsters ^[15]
	Trinidadian scorpion toxin induced acute pancreatitis in dogs ^[16]
	Alcohol-induced: rats ^[17-19] , cats ^[20] and dogs ^[21]
	Immune-mediated
	Ovalbumin in rabbit ^[22]
	Foreign serum in mice ^[23] and rat ^[24]
	Spontaneous model of autoimmune acute pancreatitis mice ^[25]
	Diet-induced: Fed a choline-deficient diet containing ethionine in mice ^[26]
	Gene knockout: Interleukin (IL)-1 and tumour necrosis factor- α ^[27] , IL-6 ^[28] , IL-10 ^[29] , chemoattractant cytokine receptor-1 ^[30] , neurokinin-1 receptor ^[31] , intercellular adhesion molecule 1 (ICAM-1) ^[32] , metallothionein-1 ^[33] , cathepsin B ^[34] , mouse a2-macroglobulin and murinoglobulin ^[35] , complement factor C5a ^[36] , granulocyte-macrophage colony-stimulating factor ^[37] and phospholipase A2 ^[38]
Invasive	<i>L</i> -arginine-induced: Administration of a large dose of <i>L</i> -arginine in rats ^[39,40]
	Closed duodenal loop (CDL): Dog ^[41] and rat ^[42,43]
	Antegrade pancreatic duct perfusion: Cat ^[44] and rat ^[45]
	Various compounds infusion into the pancreatic duct: Rat ^[46] and dog ^[47]
	Combined intraductal glycodeoxycholic acid with intravenous caerulein: Rat ^[48]
	Vascular-induced
	Impairment of pancreatic circulation in dogs ^[49]
	To occlude pancreatic arteries in rats ^[50]
	Occlusion of pancreatic veins in dogs ^[51] and in rats ^[52]
	Complete but reversible ischaemia of the pancreas by occluding different arteries using microvascular clips: Rats ^[53] and canine ^[54]
	Duct ligation
	Ligating the distal bile duct at the level of the duodenum ^[55]
	Combined pancreatic duct ligation with the secretory stimulation, secretin in dogs ^[56]
	Combining duct ligation with both secretory stimulation and minimal arterial blood ^[57]
	Duct-ligated opossums models ^[58]
Transient obstruction of the sphincter of Oddi (SO) in Australian brush tailed possums ^[59]	

tained by gene deletion or transgenic expression of genetic variants. In the same way for animal models of AP, the models of CP can be classified into noninvasive or nonsurgical models and invasive or surgical models. Table 2 summarizes different animal models of CP^[60-100].

In the non-invasive models, repetitive caerulein injections are amongst the most widely used models. Firstly, caerulein injections are relatively easy to perform and show a high reliability and reproducibility. Secondly, other compounds mediating injury such as lipopolysaccharides or cyclosporin A can easily be added to the design. Thirdly, serial caerulein injections can be performed in transgenic or knockout animals. It is likely that there are dose and frequency dependency for caerulein. The most translational models include repetitive injections of *L*-arginine, which appears to produce CP similar to that in humans^[70-72]. In this model, fibrotic tissues are progressively

Table 2 Different animal models of chronic pancreatitis

Methods	Models and examples
Non-invasive	Caerulein-induced Serial caerulein injections in mice ^[60] and rats ^[61] Combination of repetitive caerulein injections with toxins and other agents such as lipopolysaccharides ^[62] , cyclosporin A ^[63] , dibutyltin dichloride ^[64] and Alcohol ^[65-67] Intraperitoneal caerulein injections are administered in genetically transformed mice such as TRX-1 transgenic mice ^[68,69]
	Arginine-induced A single <i>L</i> -arginine injection in rat ^[70] Serial <i>L</i> -arginine injections ^[70-72] Alcohol feeding-induced: Lieber-DeCarli formula ^[73-76] Genetic models: Wistar Bonn/Kobori (WBN/Kob) rats ^[77-79] , R122H transgenic mice ^[80] , SPINK3-deficient (SPINK3-/-) mice ^[81] , CFTR-deficient (cfrmlUNC) mice ^[82] and CFTR(-/-) pigs ^[83] ; Kif3a-deficient mice ^[84] ; PERK-deficient (PERK-/-) mice ^[85] ; Interleukin 1- β transgenic mice ^[86]
Invasive	Sodium taurocholate-induced: Retrograde infusion of sodium taurocholate (NaTc) into the pancreatic duct system of the rat ^[87]
	Oleic acid-induced: Retrograde infusion of oleic acid ^[72,88-91] , viscous solution of zein ^[92] , mixture of zein-oleic acid, or viscous solution consisting of zein-oleic acid-linoleic acid ^[93,94] into rat pancreatic duct
	Congestion of pancreatic fluid flow: Combination of transient stasis of pancreatic juice flow and mild pancreatic duct injury ^[95]
	Duct ligation model Ligation of the common bile duct close to the duodenum pancreatic tissue in dogs ^[96] , mouse ^[97] and pigs ^[98] Incomplete pancreatic duct ligation in canine ^[99] Occlusion with two different tissue glues in the rat ^[100]

replaced with adipose tissue. Due to the high impact of alcohol consumption as a risk factor on the pathogenesis in human pancreatic diseases, alcohol has frequently been used to trigger CP in animal models^[73,74]. However, it is still being considered whether a model for CP induced by alcohol alone is feasible or satisfactory. The combination of alcohol feeding with caerulein injections exacerbates the course of pancreatitis and consequently increases pancreatic fibrosis and the loss of parenchyma.

Genetic animal models of CP are suitable for different studies. It is well known that activation of trypsinogen is one of the key events in the early phase of pancreatitis, and therefore genetic abnormalities found in the trypsinogen gene and in its inhibitors might be of particular importance of which R122H transgenic mice^[80] are a good example. Transgenic expression of the R122H mutation of murine trypsin 4 in the pancreas of mice led to progressive fibrosis and chronic inflammation of the pancreas. Repetitive inductions of experimental pancreatitis with supramaximal doses of cerulein resulted in extensive deposition of collagen in periacinar and perilobular spaces of this transgenic animal. However other genetic models might also help us to understand how CP develops^[77-79,81,83-86,101].

Invasive animal models can also be used to induce CP. As an example, retrograde infusion of sodium taurocholate (NaTc) into the pancreatic duct^[46] or intraductal

infusion of NaTc^[72] can generate pancreatitis, however the structure of the pancreatic tissue will return to an almost normal state after 14 d. Retrograde infusion of oleic acid^[72,88-91], viscous solution of zein^[92], a mixture of zein-oleic acid or a viscous solution consisting of zein-oleic acid-linoleic acid^[93,94] into rat pancreatic duct will cause severe pancreatic atrophy with irregular fibrosis and fat replacement over a period of 6 mo. However, these models of pancreatitis appear quite distinct from CP in humans. As one factor alone is inadequate to cause persistent pancreatic injury, a combination of transient stasis of pancreatic juice flow and mild pancreatic duct injury is a well established and reliable method to generate CP in animal models^[95]. It is well known that pancreatic ductal hypertension contributes to the pathogenesis of CP; therefore animal models can also be generated by complete obstruction of the pancreatic duct^[96-98], incomplete pancreatic duct ligation^[99] and occlusion with different tissue glues^[100]. Yamamoto *et al.*^[102] developed an animal model with pancreatic ductal hypertension and demonstrated that this plays an important role in the onset and development of CP in rats. However, models for CP based on duct obstruction are not common and there is only a minority of studies examining the morphological and biochemical changes of the pancreas after duct ligation^[41,103,104].

ANIMAL MODELS OF UN-COMMON PANCREATITIS

Un-common types of pancreatitis can include autoimmune pancreatitis (AIP), hereditary pancreatitis^[105], groove pancreatitis^[106], tropical pancreatitis, pancreatitis in ectopic or heterotopic pancreatic tissue, ascariis-induced pancreatitis, pancreatitis in cystic fibrosis, pancreas divisum, annular pancreas, pancreatic cancer manifesting as AP, and duodenal villous adenoma with pancreatitis. With exception of AIP and hereditary pancreatitis, no relevant animal models were found for other un-common pancreatitis. Furthermore, hereditary pancreatitis animal models were mentioned in the genetic animal models of CP above. Therefore only animal models of AIP are briefly introduced in this section.

To date, several animal models of AIP have been described. The first model involves the adoptive transfer of amylase-specific (an antigen mainly located in acinar cells) CD4⁺ T cells and results in pancreatitis in naive syngenic recipient animals^[107]. Notably, the histological lesions of this model mimic the lobulocentric inflammatory reaction in type 1 AIP. A model developed by immunization of neonatally thymectomized mice with CA (an antigen mainly located on the pancreatic epithelium) and later transfer of CD4⁺ lymphocytes resulted in a duct-centric pattern of pancreatitis resembling type 2 AIP^[108]. In another model, NTx-NFS/sld mice spontaneously developed sialoadenitis in which a-fodrin was involved as an autoantigen, as reported in some patients with Sjogren syndrome and AIP^[109]. Transforming growth factor- β

(TGF β) appears to be an important regulatory factor in maintaining immune homeostasis. Loss of TGF β signaling contributes to AIP in TGF β dominant negative mutant mice^[110].

Recently two animal models for AIP were proposed. The WBN/Kob rat model, associated with congenital decreased peripheral Tregs spontaneously develops sialoadenitis, thyroiditis, sclerosing cholangitis and tubulointerstitial nephritis^[111]. Although the target antigens remain unclear, CD8⁺ cells may be the effector cell in this rat model^[112]. Another recently described animal model of AIP is the Treg-deficient NOD mouse^[113]. CD28KO mice spontaneously develop AIP that closely resembles the human disease^[113]. More recently, Haruta *et al*^[114] investigated the possible involvement of chronic, persistent exposure to avirulent bacteria in the pathogenesis of AIP using C57BL/6 mice.

Existing animal models for AIP have several limitations. In most models the disease is induced by adoptive transfer of autoreactive cells and/or antibodies rather than spontaneous development of the disease with identical antigen specificity. The distribution of lesions produced in animal models for AIP is also variable. This may be attributed to the diversity of target antigens, different methods of immune staining and different mouse strains. In addition, typical histopathological findings of AIP (*e.g.*, lymphoplasmacytic infiltration with fibrosis, obliterative phlebitis and GELs) are rarely observed in animal models. Thus, there is a need to develop spontaneous animal models with identical autoantigens and typical histopathological findings for AIP.

VISCERAL PAIN IN ANIMAL MODELS OF CHRONIC PANCREATITIS

One of the main clinical symptoms of CP in humans is pain, occurring either in episodes or as a constant disabling pain^[115,116]. Hence, an important goal of treatment for CP is to relieve the pain. The analgesic treatment is often inadequate as the pathophysiology behind CP as well as the mechanisms behind the accompanying pain is not yet fully understood^[117]. As described in the previous sections, no single animal model displays all aspects of CP and each of the different models display histological similarities to the human condition to various degrees. In order to improve the pain treatment and obtain more knowledge about the physiology behind CP associated pain, animal models of CP associated pain are needed.

Rat models of CP where pancreatic nociception was investigated, have been established through invasive, non-invasive and spontaneous models^[118,119]. In these models pancreatic pain has been shown through both mechanical and thermal stimulation of the abdomen (referred pain^[120]) as well as direct electrical stimulation of the pancreas^[118,121]. These models had histopathological similarities to the human disease and had progressive fibrosis and inflammation. Furthermore, the models showed correlation between nociceptive behaviour and increased

expression of nerve growth factor (NGF) in the pancreas and calcitonin gene-related peptide (CGRP), substance P (SP), proteinase-activated receptor 2 (PAR2), and brain-derived neurotrophic factor (BDNF) in thoracic dorsal root ganglion and spinal cord segments^[118,122-124]. Increased expression of NGF, CGRP, SP and BDNF has also been shown in human patients with CP^[125-127].

Several animal models have investigated the mechanisms involved in pain accompanying CP. Takamido *et al*^[119] reported morphological changes of the nervous system being involved in development of CP pain. This study suggested that elongation of dorsal root ganglia axons and enlargement of intrapancreatic nerve bundles as being a possible mechanism of pain generation in CP. On a supraspinal level, findings have suggested that descending facilitation from the rostral ventromedial medulla plays an important role in persistent pain associated with CP^[128]. Furthermore, recent rat experiments have suggested that spinal microglia becomes activated during CP and has an important role in initiating and maintaining chronic pain^[129].

TRANSLATION OF PANCREATITIS-ASSOCIATED VISCERAL PAIN STUDY FROM ANIMAL TO HUMAN

It may be difficult to use animal models to study pancreatitis associated visceral pain as pain is a subjective experience. However animal models are needed to explore the molecular mechanisms behind pancreatitis associated visceral pain as this is difficult to study in humans. The molecular mechanisms behind the chronic pain associated with CP are poorly understood, but within recent years, animal experiments have suggested some mechanisms that might be involved. The transient receptor potential vanilloid 1 (TRPV1) and transient receptor potential ankyrin 1 (TRPA1) have been shown to be contributing factors to pain in CP^[122,130,131]. It has been shown that CP is accompanied by an increased level of NGF which caused an up-regulation of TRPV1 expression and sensitivity, resulting in hyperalgesia and allodynia^[122,130]. TRPA1 is important in both inflammation and pain in CP and can be sensitized through activation of PAR2^[131].

The mechanisms mentioned above could be used as targets for the development of novel therapeutics, aiming at treating the chronic pain accompanying CP. Neutralizing antibodies against neurotransmitters such as BDNF and NGF^[124,130] or receptor specific antagonists^[122] has proven to reverse the characteristic nociceptive behavioral changes induced by CP in several of the experimental models. Furthermore, inhibition of trypsin or inhibition of microglia activation has also abrogated the pain related behavior seen in response to CP^[123,129]. All these different mechanisms of pain treatment in CP models could have a potential as targets for novel pharmacological treatment of the chronic pain associated with CP in human patients. Also established analgesic drugs such as gaba-

pentin, buprenorphine, and morphine have been tested in animal models of CP^[118,121,132], and shown to have analgesic effect. However, many of these therapeutic approaches need to be tested in humans, before their true potential analgesic treatment of CP pain in humans can be established. It is known that some of these analgesic mechanisms are species specific and specific to the different models of induced CP.

CONCLUSION

Choosing the right model of pancreatitis is difficult and the scientific rationale needs to be carefully considered. Furthermore, no model of pancreatitis parallels all classical symptoms and the question under investigation is of importance when choosing a model. One of the main symptoms of CP is visceral pain and in order to improve the pain treatment and obtain more knowledge about the physiology behind the pancreatitis associated visceral pain, animal models of pancreatitis associated visceral pain are needed.

REFERENCES

- 1 Peery AF, Dellon ES, Lund J, Crockett SD, McGowan CE, Bulsiewicz WJ, Gangarosa LM, Thiny MT, Stizenberg K, Morgan DR, Ringel Y, Kim HP, Dibonaventura MD, Carroll CF, Allen JK, Cook SF, Sandler RS, Kappelman MD, Shaheen NJ. Burden of gastrointestinal disease in the United States: 2012 update. *Gastroenterology* 2012; **143**: 1179-87.e1-3 [PMID: 22885331 DOI: 10.1053/j.gastro.2012.08.002]
- 2 DiMagno MJ, DiMagno EP. Chronic pancreatitis. *Curr Opin Gastroenterol* 2012; **28**: 523-531 [PMID: 22782018 DOI: 10.1097/MOG.0b013e3283567dea]
- 3 Khokhar AS, Seidner DL. The pathophysiology of pancreatitis. *Nutr Clin Pract* 2004; **19**: 5-15 [PMID: 16215091 DOI: 10.1177/011542650401900105]
- 4 Su KH, Cuthbertson C, Christophi C. Review of experimental animal models of acute pancreatitis. *HPB (Oxford)* 2006; **8**: 264-286 [PMID: 18333137 DOI: 10.1080/13651820500467358]
- 5 Aghdassi AA, Mayerle J, Christochowitz S, Weiss FU, Sessler M, Lerch MM. Animal models for investigating chronic pancreatitis. *Fibrogenesis Tissue Repair* 2011; **4**: 26 [PMID: 22133269 DOI: 10.1186/1755-1536-4-26]
- 6 Barreto SG, Saccone GT. Pancreatic nociception--revisiting the physiology and pathophysiology. *Pancreatol* 2012; **12**: 104-112 [PMID: 22487519 DOI: 10.1016/j.pan.2012.02.010]
- 7 Vege SS. Pathogenesis of acute pancreatitis. Available from: URL: <http://www.uptodate.com/contents/pathogenesis-of-acute-pancreatitis#H2> (15-04-2013)
- 8 Chan YC, Leung PS. Acute pancreatitis: animal models and recent advances in basic research. *Pancreas* 2007; **34**: 1-14 [PMID: 17198179 DOI: 10.1097/01.mpa.0000246658.38375.04]
- 9 Steer ML, Saluja A. Experimental acute pancreatitis: studies of the early events that lead to cell injury. In: Go VLW, Gardner JD, Brooks FP, Lebenthal E, DiMagno EP, Scheele GA, editors. *The Pancreas, Biology, Pathobiology and Disease*. 2nd ed. New York: Raven Press, 1993: 489-499
- 10 Fétaud V, Frossard JL, Farina A, Pastor CM, Bühler L, Dumonceau JM, Hadengue A, Hochstrasser DF, Lescuyer P. Proteomic profiling in an animal model of acute pancreatitis. *Proteomics* 2008; **8**: 3621-3631 [PMID: 18686302 DOI: 10.1002/pmic.200800066]
- 11 Pini M, Rhodes DH, Castellanos KJ, Hall AR, Cabay RJ, Chennuri R, Grady EF, Fantuzzi G. Role of IL-6 in the resolution of pancreatitis in obese mice. *J Leukoc Biol* 2012; **91**: 957-966 [PMID: 22427681 DOI: 10.1189/jlb.1211627]
- 12 Watanabe O, Baccino FM, Steer ML, Meldolesi J. Supramaximal caerulein stimulation and ultrastructure of rat pancreatic acinar cell: early morphological changes during development of experimental pancreatitis. *Am J Physiol* 1984; **246**: G457-G467 [PMID: 6720895]
- 13 Niederau C, Ferrell LD, Grendell JH. Caerulein-induced acute necrotizing pancreatitis in mice: protective effects of proglumide, benzotript, and secretin. *Gastroenterology* 1985; **88**: 1192-1204 [PMID: 2984080]
- 14 Renner IG, Wisner JR. Exogenous secretin ameliorates ceruletide induced acute pancreatitis in the dog. *Dig Dis Sci* 1983; **28**: 946 (abstract)
- 15 Adler G, Kern HF, Scheele GA. Experimental models and concepts in acute pancreatitis. In: Go VWL, Gardner JD, Brooks FP, Lebenthal E, Di Magno EP, Scheele GA, editors. *The exocrine pancreas: biology, pathobiology, diseases*. New York: Raven, 1986
- 16 Pantoja JL, Renner IG, Abramson SB, Edmondson HA. Production of acute hemorrhagic pancreatitis in the dog using venom of the scorpion, *Buthus quinquestriatus*. *Dig Dis Sci* 1983; **28**: 429-439 [PMID: 6839906 DOI: 10.1007/BF02430532]
- 17 Siech M, Heinrich P, Letko G. Development of acute pancreatitis in rats after single ethanol administration and induction of a pancreatic juice edema. *Int J Pancreatol* 1991; **8**: 169-175 [PMID: 2033327]
- 18 Weber H, Merkord J, Jonas L, Wagner A, Schröder H, Käding U, Werner A, Dummmler W. Oxygen radical generation and acute pancreatitis: effects of dibutyltin dichloride/ethanol and ethanol on rat pancreas. *Pancreas* 1995; **11**: 382-388 [PMID: 8532655]
- 19 Werner J, Laposata M, Fernández-del Castillo C, Saghir M, Iozzo RV, Lewandrowski KB, Warshaw AL. Pancreatic injury in rats induced by fatty acid ethyl ester, a nonoxidative metabolite of alcohol. *Gastroenterology* 1997; **113**: 286-294 [PMID: 9207289 DOI: 10.1016/S0016-5085(97)70106-9]
- 20 Wedgwood KR, Adler G, Kern H, Reber HA. Effects of oral agents on pancreatic duct permeability. A model of acute alcoholic pancreatitis. *Dig Dis Sci* 1986; **31**: 1081-1088 [PMID: 3757722 DOI: 10.1007/BF01300261]
- 21 Friedman HS, Lowery R, Shaughnessy E, Scorza J. The effects of ethanol on pancreatic blood flow in awake and anesthetized dogs. *Proc Soc Exp Biol Med* 1983; **174**: 377-382 [PMID: 6420795]
- 22 THAL A. Studies on pancreatitis. II. Acute pancreatic necrosis produced experimentally by the arthus sensitization reaction. *Surgery* 1955; **37**: 911-917 [PMID: 14373312]
- 23 Janigan DT, Nevalainen TJ, MacAulay MA, Vethamany VG. Foreign serum-induced pancreatitis in mice. I. A new model of acute pancreatitis. *Lab Invest* 1975; **33**: 591-607 [PMID: 1202281]
- 24 Nevalainen TJ. Pancreatic injury caused by intraductal injection of foreign serum in rat. *Virchows Arch B Cell Pathol* 1978; **27**: 89-98 [PMID: 417460]
- 25 Kanno H, Nose M, Itoh J, Taniguchi Y, Kyogoku M. Spontaneous development of pancreatitis in the MRL/Mp strain of mice in autoimmune mechanism. *Clin Exp Immunol* 1992; **89**: 68-73 [PMID: 1352748 DOI: 10.1111/j.1365-2249.1992.tb06879.x]
- 26 Lombardi B, Estes LW, Longnecker DS. Acute hemorrhagic pancreatitis (massive necrosis) with fat necrosis induced in mice by DL-ethionine fed with a choline-deficient diet. *Am J Pathol* 1975; **79**: 465-480 [PMID: 1094837]
- 27 Denham W, Yang J, Fink G, Denham D, Carter G, Ward K, Norman J. Gene targeting demonstrates additive detrimental effects of interleukin 1 and tumor necrosis factor during pancreatitis. *Gastroenterology* 1997; **113**: 1741-1746 [PMID: 9352880 DOI: 10.1053/gast.1997.v113.pm9352880]
- 28 Suzuki S, Miyasaka K, Jimi A, Funakoshi A. Induction of

- acute pancreatitis by cerulein in human IL-6 gene transgenic mice. *Pancreas* 2000; **21**: 86-92 [PMID: 10881937 DOI: 10.1097/00006676-200007000-00056]
- 29 **Gloor B**, Todd KE, Lane JS, Rigberg DA, Reber HA. Mechanism of increased lung injury after acute pancreatitis in IL-10 knockout mice. *J Surg Res* 1998; **80**: 110-114 [PMID: 9790823 DOI: 10.1006/jsre.1997.5289]
- 30 **Gerard C**, Frossard JL, Bhatia M, Saluja A, Gerard NP, Lu B, Steer M. Targeted disruption of the beta-chemokine receptor CCR1 protects against pancreatitis-associated lung injury. *J Clin Invest* 1997; **100**: 2022-2027 [PMID: 9329966]
- 31 **Grady EF**, Yoshimi SK, Maa J, Valeroso D, Vartanian RK, Rahim S, Kim EH, Gerard C, Gerard N, Bunnett NW, Kirkwood KS. Substance P mediates inflammatory oedema in acute pancreatitis via activation of the neurokinin-1 receptor in rats and mice. *Br J Pharmacol* 2000; **130**: 505-512 [PMID: 10821777]
- 32 **Frossard JL**, Saluja A, Bhagat L, Lee HS, Bhatia M, Hofbauer B, Steer ML. The role of intercellular adhesion molecule 1 and neutrophils in acute pancreatitis and pancreatitis-associated lung injury. *Gastroenterology* 1999; **116**: 694-701 [PMID: 10029629]
- 33 **Fu K**, Tomita T, Sarras MP, De Lisle RC, Andrews GK. Metallothionein protects against cerulein-induced acute pancreatitis: analysis using transgenic mice. *Pancreas* 1998; **17**: 238-246 [PMID: 9788536 DOI: 10.1097/00006676-199810000-00003]
- 34 **Halangk W**, Lerch MM, Brandt-Nedelev B, Roth W, Ruthenbueger M, Reinheckel T, Domschke W, Lippert H, Peters C, Deussing J. Role of cathepsin B in intracellular trypsinogen activation and the onset of acute pancreatitis. *J Clin Invest* 2000; **106**: 773-781 [PMID: 10995788 DOI: 10.1172/JCI9411]
- 35 **Umans L**, Serneels L, Overbergh L, Stas L, Van Leuven F. alpha2-macroglobulin- and murinoglobulin-1- deficient mice. A mouse model for acute pancreatitis. *Am J Pathol* 1999; **155**: 983-993 [PMID: 10487856]
- 36 **Bhatia M**, Saluja AK, Singh VP, Frossard JL, Lee HS, Bhagat L, Gerard C, Steer ML. Complement factor C5a exerts an anti-inflammatory effect in acute pancreatitis and associated lung injury. *Am J Physiol Gastrointest Liver Physiol* 2001; **280**: G974-G978 [PMID: 11292607]
- 37 **Frossard JL**, Saluja AK, Mach N, Lee HS, Bhagat L, Hadenque A, Rubbia-Brandt L, Dranoff G, Steer ML. In vivo evidence for the role of GM-CSF as a mediator in acute pancreatitis-associated lung injury. *Am J Physiol Lung Cell Mol Physiol* 2002; **283**: L541-L548 [PMID: 12169573]
- 38 **Mizuma K**, Schröder T, Kaarne M, Korpela H, Lempinen M. Serum phospholipase A2 in diet-induced pancreatitis. *Eur Surg Res* 1984; **16**: 156-161 [PMID: 6202520]
- 39 **Hegyí P**, Rakonczay Z, Sári R, Góg C, Lonovics J, Takács T, Czako L. L-arginine-induced experimental pancreatitis. *World J Gastroenterol* 2004; **10**: 2003-2009 [PMID: 15237423]
- 40 **Varga IS**, Matkovics B, Czako L, Hai DQ, Kotorman M, Takacs T, Sasvari M. Oxidative stress changes in L-arginine-induced pancreatitis in rats. *Pancreas* 1997; **14**: 355-359 [PMID: 9163781 DOI: 10.1097/00006676-199705000-00005]
- 41 **Pfeffer RB**, Stasior O, Hinton JW. The clinical picture of the sequential development of acute hemorrhagic pancreatitis in the dog. *Surg Forum* 1957; **8**: 248-251 [PMID: 13529599]
- 42 **Nevalainen TJ**, Seppä A. Acute pancreatitis caused by closed duodenal loop in the rat. *Scand J Gastroenterol* 1975; **10**: 521-527 [PMID: 1153948]
- 43 **Orda R**, Hadas N, Orda S, Wiznitzer T. Experimental acute pancreatitis. Inducement by taurocholate sodium-trypsin injection into a temporarily closed duodenal loop in the rat. *Arch Surg* 1980; **115**: 327-329 [PMID: 6153522 DOI: 10.1001/archsurg.1980.01380030073018]
- 44 **Widdison AL**, Alvarez C, Reber HA. The low-pressure duct perfusion model of acute pancreatitis. *Eur Surg Res* 1992; **24** Suppl 1: 55-61 [PMID: 1601024 DOI: 10.1159/000129239]
- 45 **Olazábal A**. Effect of prostaglandins E2 and I2 and of indomethacin on deoxycholic acid-induced damage to the rat bile-pancreatic duct. *Gastroenterology* 1983; **84**: 928-934 [PMID: 6339311]
- 46 **Aho HJ**, Nevalainen TJ, Aho AJ. Experimental pancreatitis in the rat. Development of pancreatic necrosis, ischemia and edema after intraductal sodium taurocholate injection. *Eur Surg Res* 1983; **15**: 28-36 [PMID: 6840152]
- 47 **Musa BE**, Nelson AW, Gillette EL, Ferguson HL, Lumb WV. A model to study acute pancreatitis in the dog. *J Surg Res* 1976; **21**: 51-56 [PMID: 933516 DOI: 10.1016/0022-4804(76)90009-3]
- 48 **Schmidt J**, Rattner DW, Lewandrowski K, Compton CC, Mandavilli U, Knoefel WT, Warshaw AL. A better model of acute pancreatitis for evaluating therapy. *Ann Surg* 1992; **215**: 44-56 [PMID: 1731649 DOI: 10.1097/0000658-199201000-00007]
- 49 **Barzilai A**, Ryback BJ, Medina JA, Toth L, Dreiling DA. The morphological changes of the pancreas in hypovolemic shock and the effect of pretreatment with steroids. *Int J Pancreatol* 1987; **2**: 23-32 [PMID: 3681033]
- 50 **Redha F**, Uhlenschmid G, Ammann RW, Freiburghaus AU. Injection of microspheres into pancreatic arteries causes acute hemorrhagic pancreatitis in the rat: a new animal model. *Pancreas* 1990; **5**: 188-193 [PMID: 2315295 DOI: 10.1097/00006676-199003000-00011]
- 51 **Pfeffer RB**, Lazzarini-Robertson A, Safadi D, Mixer G, Secoy CF, Hinton JW. Gradations of pancreatitis, edematous, through hemorrhagic, experimentally produced by controlled injection of microspheres into blood vessels in dogs. *Surgery* 1962; **51**: 764-769 [PMID: 14486114]
- 52 **Sjövall S**, Holmin T, Evander A, Stenram U. Splenic and gastro-duodenal vein occlusion--influence on the pancreatic gland and on the outcome of experimental pancreatitis. *Int J Pancreatol* 1988; **3**: 143-149 [PMID: 2452221]
- 53 **Hoffmann TF**, Leiderer R, Waldner H, Arbogast S, Messmer K. Ischemia reperfusion of the pancreas: a new in vivo model for acute pancreatitis in rats. *Res Exp Med (Berl)* 1995; **195**: 125-144 [PMID: 8570908 DOI: 10.1007/BF02576782]
- 54 **Broe PJ**, Zuidema GD, Cameron JL. The role of ischemia in acute pancreatitis: studies with an isolated perfused canine pancreas. *Surgery* 1982; **91**: 377-382 [PMID: 6175032]
- 55 **Baxter JN**, Jenkins SA, Day DW, Roberts NB, Cowell DC, Mackie CR, Shields R. Effects of somatostatin and a long-acting somatostatin analogue on the prevention and treatment of experimentally induced acute pancreatitis in the rat. *Br J Surg* 1985; **72**: 382-385 [PMID: 2581647 DOI: 10.1002/bjs.1800720516]
- 56 **Popper HL**, Necheles H. Edema of the pancreas. *Surg Gynecol Obstet* 1942; **74**: 123
- 57 **Popper HL**, Necheles H, Russell KC. Transition of pancreatic edema into pancreatic necrosis. *Surg Gynecol Obstet* 1948; **87**: 79-82 [PMID: 18860330]
- 58 **Senninger N**, Moody FG, Coelho JC, Van Buren DH. The role of biliary obstruction in the pathogenesis of acute pancreatitis in the opossum. *Surgery* 1986; **99**: 688-693 [PMID: 2424109]
- 59 **Chen JW**, Thomas A, Woods CM, Schloithe AC, Toouli J, Saccone GT. Sphincter of Oddi dysfunction produces acute pancreatitis in the possum. *Gut* 2000; **47**: 539-545 [PMID: 10986215 DOI: 10.1136/gut.47.4.539]
- 60 **Neuschwander-Tetri BA**, Burton FR, Presti ME, Britton RS, Janney CG, Garvin PR, Brunt EM, Galvin NJ, Poulos JE. Repetitive self-limited acute pancreatitis induces pancreatic fibrogenesis in the mouse. *Dig Dis Sci* 2000; **45**: 665-674 [PMID: 10759232]
- 61 **Elsässer HP**, Haake T, Grimmig M, Adler G, Kern HF. Repetitive cerulein-induced pancreatitis and pancreatic fibrosis in the rat. *Pancreas* 1992; **7**: 385-390 [PMID: 1594561 DOI: 10.1097/00006676-199205000-00017]

- 62 **Segersvärd R**, Sylván M, Lempinen M, Larsson J, Permert J. Impact of chronic and acute high-fat feeding on acute experimental pancreatitis complicated by endotoxaemia. *Scand J Gastroenterol* 2004; **39**: 74-80 [PMID: 14992565 DOI: 10.1080/00365520310007233]
- 63 **Vaquero E**, Molero X, Tian X, Salas A, Malagelada JR. Myofibroblast proliferation, fibrosis, and defective pancreatic repair induced by cyclosporin in rats. *Gut* 1999; **45**: 269-277 [PMID: 10403741 DOI: 10.1136/gut.45.2.269]
- 64 **Sparmann G**, Merkord J, Jäschke A, Nizze H, Jonas L, Löhr M, Liebe S, Emmrich J. Pancreatic fibrosis in experimental pancreatitis induced by dibutyltin dichloride. *Gastroenterology* 1997; **112**: 1664-1672 [PMID: 9136846 DOI: 10.1016/S0016-5085(97)70049-0]
- 65 **Gukovsky I**, Lugea A, Shahsahebi M, Cheng JH, Hong PP, Jung YJ, Deng QG, French BA, Lungo W, French SW, Tsukamoto H, Pandolfi SJ. A rat model reproducing key pathological responses of alcoholic chronic pancreatitis. *Am J Physiol Gastrointest Liver Physiol* 2008; **294**: G68-G79 [PMID: 17884979 DOI: 10.1152/ajpgi.00006.2007]
- 66 **Deng X**, Wang L, Elm MS, Gabazadeh D, Diorio GJ, Eagon PK, Whitcomb DC. Chronic alcohol consumption accelerates fibrosis in response to cerulein-induced pancreatitis in rats. *Am J Pathol* 2005; **166**: 93-106 [PMID: 15632003 DOI: 10.1016/S0002-9440(10)62235-3]
- 67 **Perides G**, Tao X, West N, Sharma A, Steer ML. A mouse model of ethanol dependent pancreatic fibrosis. *Gut* 2005; **54**: 1461-1467 [PMID: 15870229 DOI: 10.1136/gut.2004.062919]
- 68 **Ohashi S**, Nishio A, Nakamura H, Asada M, Tamaki H, Kawasaki K, Fukui T, Yodoi J, Chiba T. Overexpression of redox-active protein thioredoxin-1 prevents development of chronic pancreatitis in mice. *Antioxid Redox Signal* 2006; **8**: 1835-1845 [PMID: 16987036 DOI: 10.1089/ars.2006.8.1835]
- 69 **Leung PS**, Chan YC. Role of oxidative stress in pancreatic inflammation. *Antioxid Redox Signal* 2009; **11**: 135-165 [PMID: 18837654 DOI: 10.1089/ars.2008.2109]
- 70 **Delaney CP**, McGeeney KF, Dervan P, Fitzpatrick JM. Pancreatic atrophy: a new model using serial intra-peritoneal injections of L-arginine. *Scand J Gastroenterol* 1993; **28**: 1086-1090 [PMID: 8303212 DOI: 10.3109/00365529309098314]
- 71 **Weaver C**, Bishop AE, Polak JM. Pancreatic changes elicited by chronic administration of excess L-arginine. *Exp Mol Pathol* 1994; **60**: 71-87 [PMID: 8070543 DOI: 10.1006/exmp.1994.1007]
- 72 **Yamaguchi T**, Kihara Y, Taguchi M, Nagashio Y, Tashiro M, Nakamura H, Otsuki M. Persistent destruction of the basement membrane of the pancreatic duct contributes to progressive acinar atrophy in rats with experimentally induced pancreatitis. *Pancreas* 2005; **31**: 365-372 [PMID: 16258372 DOI: 10.1097/01.mpa.0000179729.61457.e5]
- 73 **Lieber CS**, DeCarli LM. Alcoholic liver injury: experimental models in rats and baboons. *Adv Exp Med Biol* 1975; **59**: 379-393 [PMID: 1237225]
- 74 **Lieber CS**, Decarli LM. Animal models of ethanol dependence and liver injury in rats and baboons. *Fed Proc* 1976; **35**: 1232-1236 [PMID: 944146]
- 75 **Sarles H**, Lebreuil G, Tasso F, Figarella C, Clemente F, Devaux MA, Fagonde B, Payan H. A comparison of alcoholic pancreatitis in rat and man. *Gut* 1971; **12**: 377-388 [PMID: 4329553 DOI: 10.1136/gut.12.5.377]
- 76 **Tsukamoto H**, Towner SJ, Yu GS, French SW. Potentiation of ethanol-induced pancreatic injury by dietary fat. Induction of chronic pancreatitis by alcohol in rats. *Am J Pathol* 1988; **131**: 246-257 [PMID: 3358454]
- 77 **Ohashi K**, Kim JH, Hara H, Aso R, Akimoto T, Nakama K. WBN/Kob rats. A new spontaneously occurring model of chronic pancreatitis. *Int J Pancreatol* 1990; **6**: 231-247 [PMID: 1698893]
- 78 **Hashimoto T**, Yamada T, Yokoi T, Sano H, Ando H, Nakazawa T, Ohara H, Nomura T, Joh T, Itoh M. Apoptosis of acinar cells is involved in chronic pancreatitis in Wbn/Kob rats: role of glucocorticoids. *Pancreas* 2000; **21**: 296-304 [PMID: 11039475 DOI: 10.1097/00006676-200010000-00012]
- 79 **Mori M**, Fu X, Chen L, Zhang G, Higuchi K. Hereditary pancreatitis model WBN/Kob rat strain has a unique haplotype in the Pdwk1 region on chromosome 7. *Exp Anim* 2009; **58**: 409-413 [PMID: 19654439 DOI: 10.1538/expanim.58.409]
- 80 **Archer H**, Jura N, Keller J, Jacobson M, Bar-Sagi D. A mouse model of hereditary pancreatitis generated by transgenic expression of R122H trypsinogen. *Gastroenterology* 2006; **131**: 1844-1855 [PMID: 17087933 DOI: 10.1053/j.gastro.2006.09.049]
- 81 **Ohmuraya M**, Hirota M, Araki K, Baba H, Yamamura K. Enhanced trypsin activity in pancreatic acinar cells deficient for serine protease inhibitor kazal type 3. *Pancreas* 2006; **33**: 104-106 [PMID: 16804421 DOI: 10.1097/01.mpa.0000226889.86322.9b]
- 82 **Dimagno MJ**, Lee SH, Hao Y, Zhou SY, McKenna BJ, Owyang C. A proinflammatory, antiapoptotic phenotype underlies the susceptibility to acute pancreatitis in cystic fibrosis transmembrane regulator (-/-) mice. *Gastroenterology* 2005; **129**: 665-681 [PMID: 16083720 DOI: 10.1016/j.gastro.2005.05.059]
- 83 **Rogers CS**, Stoltz DA, Meyerholz DK, Ostedgaard LS, Rokhlina T, Taft PJ, Rogan MP, Pezzulo AA, Karp PH, Itani OA, Kabel AC, Wohlford-Lenane CL, Davis GJ, Hanfland RA, Smith TL, Samuel M, Wax D, Murphy CN, Rieke A, Whitworth K, Uc A, Starner TD, Brogden KA, Shilyansky J, McCray PB, Zabner J, Prather RS, Welsh MJ. Disruption of the CFTR gene produces a model of cystic fibrosis in newborn pigs. *Science* 2008; **321**: 1837-1841 [PMID: 18818360 DOI: 10.1126/science.1163600]
- 84 **Lin F**, Hiesberger T, Cordes K, Sinclair AM, Goldstein LS, Somlo S, Igarashi P. Kidney-specific inactivation of the KIF3A subunit of kinesin-II inhibits renal ciliogenesis and produces polycystic kidney disease. *Proc Natl Acad Sci USA* 2003; **100**: 5286-5291 [PMID: 12672950 DOI: 10.1073/pnas.0836980100]
- 85 **Harding HP**, Zeng H, Zhang Y, Jungries R, Chung P, Plesken H, Sabatini DD, Ron D. Diabetes mellitus and exocrine pancreatic dysfunction in perk-/- mice reveals a role for translational control in secretory cell survival. *Mol Cell* 2001; **7**: 1153-1163 [PMID: 11430819 DOI: 10.1016/S1097-2765(01)00264-7]
- 86 **Marrache F**, Tu SP, Bhagat G, Pendyala S, Osterreicher CH, Gordon S, Ramanathan V, Penz-Osterreicher M, Betz KS, Song Z, Wang TC. Overexpression of interleukin-1beta in the murine pancreas results in chronic pancreatitis. *Gastroenterology* 2008; **135**: 1277-1287 [PMID: 18789941 DOI: 10.1053/j.gastro.2008.06.078]
- 87 **Aho HJ**, Nevalainen TJ, Lindberg RL, Aho AJ. Experimental pancreatitis in the rat. The role of phospholipase A in sodium taurocholate-induced acute haemorrhagic pancreatitis. *Scand J Gastroenterol* 1980; **15**: 1027-1031 [PMID: 6165066 DOI: 10.3109/00365528009181808]
- 88 **Mundlos S**, Adler G, Schaar M, Koop I, Arnold R. Exocrine pancreatic function in oleic acid-induced pancreatic insufficiency in rats. *Pancreas* 1986; **1**: 29-36 [PMID: 2437560 DOI: 10.1097/00006676-198601000-00007]
- 89 **Henry JP**, Steinberg WM. Pancreatic function tests in the rat model of chronic pancreatic insufficiency. *Pancreas* 1993; **8**: 622-626 [PMID: 8302800 DOI: 10.1097/00006676-199309000-00015]
- 90 **Yamaguchi T**, Nakamura H, Kihara Y, Taguchi M, Yoshikawa H, Otsuki M. Long-term overexpression of membrane type-1 matrix metalloproteinase and matrix metalloproteinase-2 in oleic acid-induced pancreatitis in rats. *Pancreas* 2002; **24**: 348-356 [PMID: 11961487 DOI: 10.1097/00006676-20020500-00005]
- 91 **Tashiro M**, Nakamura H, Taguchi M, Yamaguchi T, Yoshi-

- kawa H, Fukumitsu K, Kihara Y, Otsuki M. Oleic acid-induced pancreatitis alters expression of transforming growth factor-beta1 and extracellular matrix components in rats. *Pancreas* 2003; **26**: 197-204 [PMID: 12604920 DOI: 10.1097/00006676-200303000-00018]
- 92 Tomita T, Rhodes J, Falscroft J, Doull V, Kimmel JR, Pollock HG. Endocrine pancreas in the rat model of exocrine pancreatic insufficiency. *Pancreas* 1988; **3**: 568-575 [PMID: 3054870 DOI: 10.1097/00006676-198810000-00010]
- 93 Steinberg WM, Burns MK, Henry JP, Nochomovitz LE, Anderson KK. Cerulein induces hyperplasia of the pancreas in a rat model of chronic pancreatic insufficiency. *Pancreas* 1987; **2**: 176-180 [PMID: 3628223 DOI: 10.1097/00006676-198703000-00009]
- 94 Kataoka K, Sasaki T, Yorizumi H, Sakagami J, Kashima K. Pathophysiologic studies of experimental chronic pancreatitis in rats induced by injection of zein-oleic acid-linoleic acid solution into the pancreatic duct. *Pancreas* 1998; **16**: 289-299 [PMID: 9548669 DOI: 10.1097/00006676-199804000-00014]
- 95 Yamaguchi T, Kihara Y, Yamamoto M, Otsuki M. Transient stasis of pancreatic fluid flow together with mild injury of the pancreatic duct cause chronic pancreatitis. *Dig Dis Sci* 2011; **56**: 2326-2332 [PMID: 21274624 DOI: 10.1007/s10620-011-1586-1]
- 96 Churg A, Richter WR. Early changes in the exocrine pancreas of the dog and rat after ligation of the pancreatic duct. A light and electron microscopic study. *Am J Pathol* 1971; **63**: 521-546 [PMID: 5581235]
- 97 Watanabe S, Abe K, Anbo Y, Katoh H. Changes in the mouse exocrine pancreas after pancreatic duct ligation: a qualitative and quantitative histological study. *Arch Histol Cytol* 1995; **58**: 365-374 [PMID: 8527243 DOI: 10.1679/aohc.58.365]
- 98 Boerma D, Straatsburg IH, Offerhaus GJ, Gouma DJ, van Gulik TM. Experimental model of obstructive, chronic pancreatitis in pigs. *Dig Surg* 2003; **20**: 520-526 [PMID: 14534374 DOI: 10.1159/000073688]
- 99 Tanaka T, Ichiba Y, Fujii Y, Itoh H, Kodama O, Dohi K. New canine model of chronic pancreatitis due to chronic ischemia with incomplete pancreatic duct obstruction. *Digestion* 1988; **41**: 149-155 [PMID: 3224767 DOI: 10.1159/000199767]
- 100 Isaksson G, Lundquist I, Ihse I. Effects on the exocrine and endocrine pancreas of duct occlusion with two different tissue glues in the rat. *Eur Surg Res* 1983; **15**: 136-144 [PMID: 6345169 DOI: 10.1159/000128345]
- 101 Cano DA, Sekine S, Hebrok M. Primary cilia deletion in pancreatic epithelial cells results in cyst formation and pancreatitis. *Gastroenterology* 2006; **131**: 1856-1869 [PMID: 17123526 DOI: 10.1053/j.gastro.2006.10.050]
- 102 Yamamoto M, Otani M, Otsuki M. A new model of chronic pancreatitis in rats. *Am J Physiol Gastrointest Liver Physiol* 2006; **291**: G700-G708 [PMID: 16959955 DOI: 10.1152/ajpgi.00502.2005]
- 103 Klein ES, Grateron H, Rudick J, Dreiling DA. Pancreatic intraductal pressure: I. A consideration of regulatory factors. *Am J Gastroenterol* 1983; **78**: 507-509 [PMID: 6881116]
- 104 De Rai P, Franciosi C, Confalonieri GM, Biffi R, Andreoni B, Uggeri F, Malesci A. Effects of somatostatin on acute pancreatitis induced in rats by injection of taurocholate and trypsin into a temporarily closed duodenal loop. *Int J Pancreatol* 1988; **3**: 367-373 [PMID: 2902146]
- 105 Rosendahl J, Bödeker H, Mössner J, Teich N. Hereditary chronic pancreatitis. *Orphanet J Rare Dis* 2007; **2**: 1 [PMID: 17204147 DOI: 10.1186/1750-1172-2-1]
- 106 Tyagi P, Thaper S, Bhatia V, Sharma M. Often a missed type of pancreatitis: groove pancreatitis. *Indian J Gastroenterol* 2012; **31**: 215-216 [PMID: 22829067 DOI: 10.1007/s12664-012-0229-6]
- 107 Zen Y, Fujii T, Harada K, Kawano M, Yamada K, Takahira M, Nakanuma Y. Th2 and regulatory immune reactions are increased in immunoglobulin G4-related sclerosing pancreatitis and cholangitis. *Hepatology* 2007; **45**: 1538-1546 [PMID: 17518371 DOI: 10.1002/hep.21697]
- 108 Okazaki K, Uchida K, Ohana M, Nakase H, Uose S, Inai M, Matsushima Y, Katamura K, Ohmori K, Chiba T. Autoimmune-related pancreatitis is associated with autoantibodies and a Th1/Th2-type cellular immune response. *Gastroenterology* 2000; **118**: 573-581 [PMID: 10702209 DOI: 10.1016/S0016-5085(00)70264-2]
- 109 Haneji N, Nakamura T, Takio K, Yanagi K, Higashiyama H, Saito I, Noji S, Sugino H, Hayashi Y. Identification of alpha-fodrin as a candidate autoantigen in primary Sjögren's syndrome. *Science* 1997; **276**: 604-607 [PMID: 9110981]
- 110 Hahm KB, Im YH, Lee C, Parks WT, Bang YJ, Green JE, Kim SJ. Loss of TGF-beta signaling contributes to autoimmune pancreatitis. *J Clin Invest* 2000; **105**: 1057-1065 [PMID: 10772650 DOI: 10.1172/JCI8337]
- 111 Sakaguchi Y, Inaba M, Tsuda M, Quan GK, Omae M, Ando Y, Uchida K, Okazaki K, Ikehara S. The Wistar Bonn Kobori rat, a unique animal model for autoimmune pancreatitis with extrapancreatic exocrinopathy. *Clin Exp Immunol* 2008; **152**: 1-12 [PMID: 18279444 DOI: 10.1111/j.1365-2249.2008.03588.x]
- 112 Kountouras J, Zavos C, Chatzopoulos D. A concept on the role of Helicobacter pylori infection in autoimmune pancreatitis. *J Cell Mol Med* 2005; **9**: 196-207 [PMID: 15784177 DOI: 10.1111/j.1582-4934.2005.tb00349.x]
- 113 Meagher C, Tang Q, Fife BT, Bour-Jordan H, Wu J, Pardoux C, Bi M, Melli K, Bluestone JA. Spontaneous development of a pancreatic exocrine disease in CD28-deficient NOD mice. *J Immunol* 2008; **180**: 7793-7803 [PMID: 18523243]
- 114 Haruta I, Yanagisawa N, Kawamura S, Furukawa T, Shimizu K, Kato H, Kobayashi M, Shiratori K, Yagi J. A mouse model of autoimmune pancreatitis with salivary gland involvement triggered by innate immunity via persistent exposure to avirulent bacteria. *Lab Invest* 2010; **90**: 1757-1769 [PMID: 20733561 DOI: 10.1038/labinvest.2010.153]
- 115 Ammann RW. Diagnosis and management of chronic pancreatitis: current knowledge. *Swiss Med Wkly* 2006; **136**: 166-174 [PMID: 16633964]
- 116 André-Sandberg A, Hoem D, Gislason H. Pain management in chronic pancreatitis. *Eur J Gastroenterol Hepatol* 2002; **14**: 957-970 [PMID: 12352215 DOI: 10.1097/00042737-200209000-00006]
- 117 DiMaggio EP. Toward understanding (and management) of painful chronic pancreatitis. *Gastroenterology* 1999; **116**: 1252-1257 [PMID: 10220520 DOI: 10.1016/S0016-5085(99)70031-4]
- 118 Vera-Portocarrero LP, Lu Y, Westlund KN. Nociception in persistent pancreatitis in rats: effects of morphine and neuropeptide alterations. *Anesthesiology* 2003; **98**: 474-484 [PMID: 12552208 DOI: 10.1097/00000542-200302000-00029]
- 119 Takamido S, Kataoka Y, Tanano A, Cui Y, Ikeura T, Shimatani M, Kubota Y, Okazaki K, Yamada H. Intrapaneatic axonal hyperbranching of dorsal root ganglia neurons in chronic pancreatitis model rats and its relation to pancreatic pain. *Pancreas* 2006; **33**: 268-279 [PMID: 17003649 DOI: 10.1097/01.mpa.0000240600.72946.23]
- 120 Wesselmann U, Lai J. Mechanisms of referred visceral pain: uterine inflammation in the adult virgin rat results in neurogenic plasma extravasation in the skin. *Pain* 1997; **73**: 309-317 [PMID: 9469520 DOI: 10.1016/S0304-3959(97)00112-7]
- 121 Winston JH, He ZJ, Shenoy M, Xiao SY, Pasricha PJ. Molecular and behavioral changes in nociception in a novel rat model of chronic pancreatitis for the study of pain. *Pain* 2005; **117**: 214-222 [PMID: 16098667 DOI: 10.1016/j.pain.2005.06.013]
- 122 Xu GY, Winston JH, Shenoy M, Yin H, Pendyala S, Pasricha PJ. Transient receptor potential vanilloid 1 mediates hyperalgesia and is up-regulated in rats with chronic pancreatitis. *Gastroenterology* 2007; **133**: 1282-1292 [PMID: 17698068 DOI: 10.1053/j.gastro.2007.06.015]
- 123 Zhang W, Gao J, Zhao T, Wei L, Wu W, Bai Y, Zou D, Li Z.

- Proteinase-activated receptor 2 mediates thermal hyperalgesia and is upregulated in a rat model of chronic pancreatitis. *Pancreas* 2011; **40**: 300-307 [PMID: 21311307 DOI: 10.1097/MPA.0b013e318201cbc1]
- 124 **Hughes MS**, Shenoy M, Liu L, Colak T, Mehta K, Pasricha PJ. Brain-derived neurotrophic factor is upregulated in rats with chronic pancreatitis and mediates pain behavior. *Pancreas* 2011; **40**: 551-556 [PMID: 21499209 DOI: 10.1097/MPA.0b013e318214fb77]
- 125 **Friess H**, Shrikhande S, Shrikhande M, Martignoni M, Kulli C, Zimmermann A, Kappeler A, Ramesh H, Büchler M. Neural alterations in surgical stage chronic pancreatitis are independent of the underlying aetiology. *Gut* 2002; **50**: 682-686 [PMID: 11950816 DOI: 10.1136/gut.50.5.682]
- 126 **Büchler M**, Weihe E, Friess H, Malfertheiner P, Bockman E, Müller S, Nohr D, Beger HG. Changes in peptidergic innervation in chronic pancreatitis. *Pancreas* 1992; **7**: 183-192 [PMID: 1372738 DOI: 10.1097/00006676-199203000-00009]
- 127 **Zhu ZW**, Friess H, Wang L, Zimmermann A, Büchler MW. Brain-derived neurotrophic factor (BDNF) is upregulated and associated with pain in chronic pancreatitis. *Dig Dis Sci* 2001; **46**: 1633-1639 [PMID: 11508661]
- 128 **Vera-Portocarrero LP**, Xie JY, Kowal J, Ossipov MH, King T, Porreca F. Descending facilitation from the rostral ventromedial medulla maintains visceral pain in rats with experimental pancreatitis. *Gastroenterology* 2006; **130**: 2155-2164 [PMID: 16762636 DOI: 10.1053/j.gastro.2006.03.025]
- 129 **Liu PY**, Lu CL, Wang CC, Lee IH, Hsieh JC, Chen CC, Lee HF, Lin HC, Chang FY, Lee SD. Spinal microglia initiate and maintain hyperalgesia in a rat model of chronic pancreatitis. *Gastroenterology* 2012; **142**: 165-173.e2 [PMID: 21963786 DOI: 10.1053/j.gastro.2011.09.041]
- 130 **Zhu Y**, Colak T, Shenoy M, Liu L, Pai R, Li C, Mehta K, Pasricha PJ. Nerve growth factor modulates TRPV1 expression and function and mediates pain in chronic pancreatitis. *Gastroenterology* 2011; **141**: 370-377 [PMID: 21473865 DOI: 10.1053/j.gastro.2011.03.046]
- 131 **Cattaruzza F**, Johnson C, Leggit A, Grady E, Schenk AK, Cevikbas F, Cedron W, Bondada S, Kirkwood R, Malone B, Steinhoff M, Bunnett N, Kirkwood KS. Transient receptor potential ankyrin 1 mediates chronic pancreatitis pain in mice. *Am J Physiol Gastrointest Liver Physiol* 2013; **304**: G1002-G1012 [PMID: 23558009 DOI: 10.1152/ajpgi.00005.2013]
- 132 **Liao XZ**, Zhou MT, Mao YF, Xu H, Chen H, Sun JH, Xiong YC. Analgesic effects of gabapentin on mechanical hypersensitivity in a rat model of chronic pancreatitis. *Brain Res* 2010; **1337**: 104-112 [PMID: 20417627 DOI: 10.1016/j.brainres.2010.04.035]

P- Reviewer: Chow WK S- Editor: Gou SX L- Editor: A
E- Editor: Zhang DN





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza,

315-321 Lockhart Road, Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045