

Enduring legacy of a toxic fan via episodic redistribution of California gold mining debris

Michael Bliss Singer^{a,b,1}, Rolf Aalto^c, L. Allan James^d, Nina E. Kilham^e, John L. Higson^a, and Subhajt Ghoshal^f

^aDepartment of Earth and Environmental Sciences, University of St. Andrews, St. Andrews KY16 9AL, United Kingdom; ^bEarth Research Institute, University of California, Santa Barbara, CA 93106; ^cDepartment of Geography, College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4RJ, United Kingdom; ^dDepartment of Geography, University of South Carolina, Columbia, SC 29208; ^eDepartment of Geography and Global Studies, Sonoma State University, Rohnert Park, CA 94928; and ^fComputational Modeling Center, Air Products and Chemicals, Inc., Allentown, PA 18195

Edited by Robert E. Dickinson, The University of Texas at Austin, Austin, TX, and approved September 30, 2013 (received for review February 6, 2013)

The interrelationships between hydrologically driven evolution of legacy landscapes downstream of major mining districts and the contamination of lowland ecosystems are poorly understood over centennial time scales. Here, we demonstrate within piedmont valleys of California's Sierra Nevada, through new and historical data supported by modeling, that anthropogenic fans produced by 19th century gold mining comprise an episodically persistent source of sediment-adsorbed Hg to lowlands. Within the enormous, iconic Yuba Fan, we highlight (i) an apparent shift in the relative processes of fan evolution from gradual vertical channel entrenchment to punctuated lateral erosion of fan terraces, thus enabling entrainment of large volumes of Hg-laden sediment during individual floods, and (ii) systematic intrafan redistribution and downstream progradation of fan sediment into the Central Valley, triggered by terrace erosion during increasingly long, 10-y flood events. Each major flood apparently erodes stored sediment and delivers to sensitive lowlands the equivalent of ~10–30% of the entire postmining Sierran Hg mass so far conveyed to the San Francisco Bay-Delta (SFBD). This process of protracted but episodic erosion of legacy sediment and associated Hg is likely to persist for >10⁴ y. It creates, within an immense swath of river corridor well upstream of the SFBD, new contaminated floodplain surfaces primed for Hg methylation and augments/replenishes potential Hg sources to the SFBD. Anticipation, prediction, and management of toxic sediment delivery, and corresponding risks to lowland ecology and human society globally, depend on the morphodynamic stage of anthropogenic fan evolution, synergistically coupled to changing frequency of and duration extreme floods.

mercury | lowland contamination | flood frequency | climate change | morphodynamics

Hg poses contamination risks to food webs globally (1, 2), particularly where sediment-bound Hg is delivered to topographically low, depositional environments with high methylation potential. Sediment is the vehicle for ~97% of transition metal mass loading to oceans (3), and the geomorphic configuration of stored contaminated sediment on land has important implications for its subsequent remobilization (4) and delivery to the base of food webs. Sediment displaced by industrial mines is typically contaminated by toxic metals such as Hg (5–7), which threaten lowland ecosystems downstream of mining sites around the world (e.g., refs. 6–9). This contaminated sediment often infills confined river valleys (10) as massive anthropogenic fans, similar to natural alluvial fans, which undergo subsequent incision once the supply of mining sediment is stopped. Such fans are likely to increase in ubiquity and scale across the globe as demand increases for mined metals and rare earth elements.

Nineteenth century hydraulic gold mining in California's Sierra Nevada foothills (1853–1884) contemporaneously delivered ~1.1 km³ of Hg-contaminated sediment (10, 11) to downstream valleys at concentrations >10-fold higher than geographical “hot spots” of atmospheric Hg deposition (12). These immense mining sediment slugs choked Sierra rivers, which aggraded tens of meters in confined piedmont valleys, and formed anthropogenic

fans grading to the margins of the Central Valley, including the ~252 × 10⁶-m³ Yuba Fan (10) (Fig. 1 *A* and *B*). Since that time, sediment-bound Hg has contaminated fish and waterfowl of the San Francisco Bay-Delta (SFBD) (13, 14). However, the dominant modern geographical sources of Hg, as well as the processes, patterns, and time scale of its delivery to this downstream ecosystem, are debated (15–17), partly due to the lack of a generic understanding of postmining fan evolution. It is unclear whether there are modern sources of Hg to this system and by what set of physical and biogeochemical processes Hg may penetrate these sensitive ecosystems. The legacy impacts of historical mining are therefore not fully comprehended, in part, because of incomplete knowledge of the magnitude and frequency of processes contributing to fan evolution and sediment exhaustion.

Following cessation of hydraulic mining, Sierra valley aggradation was succeeded by vertical channel incision into piedmont fans, producing historical terraces along channels (10). James (18) demonstrated that ~90% of the original sediment deposit delivered to the Bear River valley (Fig. 1*A*) remained in terrace storage >100 y after mining and hypothesized that modern floods are capable of transporting it downstream. Apart from Gilbert's classic work on bed-level change (10), generalizable theory for flood-based evolution of anthropogenic fans is absent, especially over the critical management time scale of decades to centuries in this and other fluvial systems (5, 6). This precludes predictions of mining sediment exhaustion time scales and the downstream risks of heavy metal contamination to food webs.

To fill this research gap, we analyzed high-resolution historical topographical and bathymetric data collected in 1906 and 1999, and interpreted them here within the context of: historical US Geological Survey (USGS) streamflow (Q) measurements and

Significance

This paper is of fundamental interest to the millions of residents living at the downstream end of this and other global river basins beset by industrial metals mining. Sediment-bound Hg has contaminated food webs of the San Francisco Bay-Delta, but the dominant geographical sources of Hg to downstream ecosystems in this and similar river basins are debated. Likewise, the processes by which Hg is delivered to lowlands and the patterns of its floodplain deposition are poorly understood. This research addresses a gap in generic theory of postmining fan evolution that enables anticipation, prediction, and management of contamination risk to food webs.

Author contributions: M.B.S. designed research; M.B.S., R.A., L.A.J., N.E.K., J.L.H., and S.G. performed research; M.B.S. contributed new reagents/analytic tools; M.B.S., R.A., and L.A.J. analyzed data; and M.B.S., R.A., and L.A.J. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: bliss@eri.ucsb.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1302295110/-DCSupplemental.




Fig. 1. Yuba Fan. (A) HgT sediment sampling locations (red circles). The longitudinal transect along which mining sediment travels from the Sierra to the SFBD is shown in yellow. Sample sites A–F are described in the main text. Base: Shuttle Radar Topography Mission (SRTM) 1 arc-second. Yuba Fan delineation and hypothesized patterns of sediment redistribution and export (B) and longitudinal elevation profiles of the Yuba River channel bed, 1906 and 1999 (C). (Insets) Plots show cross-sectional change at sites A and B. *Fan incision rates for the early (10) and late 20th century are based on trends in water surface elevation. Supporting methodological description and data on slowing incision are provided in *SI Methods* and *Table S1*. ASL, above sea level.

flood frequency; National Aeronautics and Space Administration flood imagery; and our newly developed geochemical dataset that includes total Hg concentration (HgT), Zr/Ni, and ¹³⁷Cs, supported by physically based mathematical modeling of terrace erosion and flux of mining sediment during large floods (*SI Methods*).

Results and Discussion

Terrestrial/bathymetric topographical data, collected in 1906 and 1999, reveal that the Yuba River longitudinal profile has incised systematically over the entire fan in the past century, manifesting as substantial valley-floor lowering (i.e., erosion of the channel and adjacent floodplains) and terrace erosion in the upper fan

and as vertical channel incision with modest (<1 m) floodplain aggradation in the lower fan (Fig. 1C). The riverbed level of the middle fan (“Yuba Gold Fields”) has been affected by the presence of Daguerre Point and Barrier Dams, built to trap mining sediment. Barrier Dam was destroyed by a large flood in 1907 (10), resulting in a marked adjustment of the thalweg long profile (Fig. 1C). The reservoir behind Daguerre Point Dam (built in 1910) remains completely full of Hg-laden sediment (19), so it suppresses vertical incision within the Yuba Gold Fields and has negligible sediment trap efficiency during floods.

Nevertheless, the overall centennial pattern of change appears to be systematic incision and redistribution of sediment from the

Our analyses highlight the importance of anthropogenic fans in the apparent lateral-expansion stage of morphodynamic evolution as primary stores, sources, and downstream exporters of mining sediment and associated contaminants. Assuming that 90% of the Yuba Fan remains in storage [equivalent to estimates for the Bear River (18)], ~0.1% of mining sediment (both fine and coarse) is evacuated per decade by a single large flood event such as those outlined here, implying a source of contaminated sediments to lowlands that will last for $>10^4$ y (*SI Methods*). This is consistent with a right-skewed model of sediment exhaustion (31) punctuated by increasingly heavy-tailed flood frequency (28, 29), which could slightly shorten mining sediment exhaustion times.

Based on modeled fluxes for flood events and assuming (*SI Methods*) conservatively low HgT, equivalent to concentrations in recent lowland flood deposits (Fig. 2*F*), the recent major floods exported large masses of Hg from the Yuba Fan into the Central Valley (Fig. 4*B* and *C*). Such per flood event-based contaminant flux, predominantly derived from historical terraces relative to channel/bar sediments (Fig. 5*A*), represents a substantial fraction of the cumulative Hg mass (~200 kg) that is estimated to have reached the SFBD since mining began (15). Therefore, most Hg exported from anthropogenic fans is apparently still stored in lowland floodplains and floodways upstream of the SFBD. It manifests as an ~70-km long swath of Hg contamination from the Yuba Fan into the Central Valley (Fig. 5*A*).

This interpretation is corroborated by an independent geochemical proxy for mine-derived sediments, Zr/Ni (15). Hydraulic mining sediments in the Sierra largely derive from highly weathered Eocene sediments from which gold was extracted, and thus have high Zr/Ni ratios due to the selective removal of Ni and relative immobility of Zr (15). We analyzed our samples for these elements and found a clear down-valley trend in Zr/Ni (Fig. 5*B*). The HgT data show much higher variability, primarily as a consequence of sampling at various topographical positions (Fig. 2), but roughly follow a similar trend. Each elemental signature of hydraulic mining is apparently diluted downstream through the Yuba Fan, the scatter in HgT notwithstanding, and is relatively concentrated in floodplain deposits near the fan margin (Fig. 5). These data support the redistribution of mining sediment from the upper to lower Yuba Fan and its punctuated export from the entire fan into the Central Valley (Fig. 1*B*). They are also consistent with Hg isotope ratios in shallow SFBD sediments, which suggest a persistent upstream (Sierran) Hg source (16, 32).

Lowland floodplains are potential hotspots for Hg methylation of mining debris. Fine-grained Hg from hydraulic mining is as-

sociated with sulfide minerals (19), which are prone to oxidation (to mercury sulfate) during transport. Once delivered to lowlands, these sediments are redeposited, producing contaminated floodplain surfaces that undergo frequent wetting/drying cycles. These cycles may stimulate resident sulfate-reducing bacteria to methylate the newly arrived Hg near the sediment–water interface at very high rates (33, 34), posing risks to food webs in this and other ecosystems beset by mining-induced displacement of contaminated sediment.

The enduring legacy of mining on a grand landscape scale entails persistent lateral erosion of anthropogenic fans that continues well after the slowing of initial vertical incision and return of the channel to its original bed level. This process stage of fan evolution apparently sustains the evacuation of legacy sediment, triggered by increasingly potent floods that effectively deliver toxic sediment slugs downstream into sensitive lowlands, thus augmenting a major potential source of food web contamination in tidal wetlands of the SFBD.

These links between upland legacy mining sediment and lowland contamination, which are germane to issues of river management such as licensing for mining or dams in this and other basins around the world, are enabled by the topographical position of stored sediment and the frequency/magnitude of flood events capable of transporting contaminated sediment over tens of kilometers. Our findings suggest dangerous synergies between the morphodynamic stage of fan evolution, regional shifts in climate, and contamination risks to lowland ecosystems and human populations.

Methods

This paper presents the culmination of a multipronged approach, including assessment of historical topographical datasets, investigation of historical streamflow measurements at USGS gauging stations to determine rates of incision, flood frequency analysis, interpretation of HgT chemostratigraphy, Zr/Ni geochemistry as an independent proxy for mining sediment, and use of ^{137}Cs to establish age control for younger deposits. These approaches were combined with terrace erosion modeling based on slope stability tied to hyporheic infiltration, partitioning of failed material into suspended sediment load, and numerical modeling suspended flux through the fan (*SI Methods*).

Topographical longitudinal and cross-section profiles were extracted from historical maps and recent LiDAR tied to photogrammetry and sonar data (35). Stratified sediment samples for HgT ($n = 176$) were extracted from channel bars and bank exposures at field-identified sedimentary units ($n = 105$) typically in vertical sections. Samples were sieved through $<63\text{-}\mu\text{m}$ stainless-steel sieves to compare a common sediment population across all sites. The resulting material was digested at a USGS Hg laboratory in Menlo Park, California. Digested samples were harvested and analyzed on a Tekran




Fig. 5. Downstream patterns of geochemistry within and beyond the Yuba Fan. Measured HgT (*A*) and Zr/Ni (*B*) in sediments (*Dataset S1*). Power relationships fit to all HgT (including all stratigraphic positions from each historical terrace) and Zr/Ni data within the Yuba Fan are given in Fig. S1. Colors in *A* refer to geographical domains listed at the top of the figure. Shaded ovals indicate elevated geochemical concentrations downstream of the margins of anthropogenic fans.

Series 2600 Automated Mercury Analysis System according to standard procedures for quality assurance/quality control (QA/QC) (US Environmental Protection Agency method 1631, revision E) using International Atomic Energy Agency (IAEA) 405 as certified reference material. Zr/Ni analyses were conducted by Inductively Coupled Plasma Mass Spectrometry (ICP-MS) and X-Ray Fluorescence (XRF) at Activation (ACT) Laboratories Ltd. and the University of St. Andrews. The ^{137}Cs data were generated in the Exeter Radiometry Laboratory with high-purity germanium (HPGe) spectrometers.

Physically based terrace erosion modeling derives from the infinite slope stability model for riverbanks (36), wherein factor of safety (FoS) was driven by stage-discharge recorded at the upstream gauging station. This boundary flow fed into a 2D numerical model of infiltration via the Richards equation following the Dupuit–Forchheimer approximation. FoS was computed based on field-measured failure geometry and grain sizes and modeled water content. Following failure, the bed grain size distribution (GSD) was updated based on contribution from measured terrace GSD depending on the depth of failure. Sediment flux was computed iteratively for the cross-section to quantify the suspended load (Fig. 4A), based on daily flood stage and updated GSD by the Singer–Dunne equation (37). Flow and SSC through the Yuba River and its floodplain were modeled using TELEMAC 2-D (Laboratoire

National d'Hydraulique, Paris, France) and Sisyphé (a coproperty of the Centre d'Études Maritimes et Fluviales, Université de Technologie de Compiègne, Électricité de France, and the Société Grenobloise d'Études et d'Application Hydrauliques), a hydrodynamic and washload transport modeling approach (38), on a finite element mesh derived from high-resolution topographical data for the channel and overbank areas. Suspended sediment input at the upstream boundary in the model of the Yuba Fan (and corresponding uncertainty) was conservatively approximated as a point source based on an empirical relationship between sediment concentration and discharge (*SI Methods*).

ACKNOWLEDGMENTS. We thank E. Buer, M. Megison, J. Touzel, J. Wooten, and A. Newborn, for field help; M. Marvin-DiPasquale for use of the USGS mercury laboratory; M. Cox, S. Franklin, J. Grapes, E. Buer, E. Kakourous, L. Kieu, A. Calder, and E. Singer for laboratory assistance; and the University of Bristol's School of Geographical Sciences for providing TELEMAC 2-D to N.E.K. This work was supported by National Science Foundation Grants BCS-0521663 and EAR-1226741 (to M.B.S.), Grant BCS-0520933 (to L.A.J.), and Grant BCS-0521774 (to R.A.). M.B.S. also received support from The Carnegie Trust for Scottish Universities and The Royal Society of Edinburgh. J.L.H. was supported by a Natural Environment Research Council Fellowship (United Kingdom).

- Mergler D, et al. (2007) Methylmercury exposure and health effects in humans: A worldwide concern. *AMBIO* 36(1):3–11.
- Cristol DA, et al. (2008) The movement of aquatic mercury through terrestrial food webs. *Science* 320(5874):335.
- Gibbs RJ (1977) Transport phases of transition metals in the Amazon and Yukon Rivers. *Geol Soc Am Bull* 88(6):829–843.
- Miller JR (1997) The role of fluvial geomorphic processes in the dispersal of heavy metals from mine sites. *J Geochem Explor* 58(2-3):101–118.
- Lece S, Pavlovsky R, Schlomer G (2008) Mercury contamination of active channel sediment and floodplain deposits from historic gold mining at Gold Hill, North Carolina, USA. *Environmental Geology* 55(1):113–121.
- Miller J, et al. (1999) Effects of the 1997 flood on the transport and storage of sediment and mercury within the Carson River valley, west-central Nevada. *J Geol* 107(3): 313–327.
- Knighon AD (1989) River adjustment to changes in sediment load: The effects of tin mining on the Ringarooma River, Tasmania, 1875–1984. *Earth Surface Processes and Landforms* 14(4):333–359.
- Swanson KM, et al. (2008) Sediment load and floodplain deposition rates: Comparison of the Fly and Strickland rivers, Papua New Guinea. *Journal of Geophysical Research Earth Surface* 113(1):F01S03, 10.1029/2006JF000623.
- Maurice-Bourgoin L, Aalto R, Guyot JL (2002) Sediment-associated mercury distribution within a major Amazonian tributary: Century-scale contamination history and importance of flood plain accumulation. *The Structure, Function and Management Implications of Fluvial Sedimentary Systems*, eds Dyer FJ, Thoms MC, Olley JM (IAHS Press, Wallingford, UK), pp 161–168.
- Gilbert GK (1917) Hydraulic-Mining Debris in the Sierra Nevada, *US Geological Survey Professional Paper* (USGC, Menlo Park, CA), Vol 105.
- Alpers CN, Hunerlach MP, May JT, Hothem RL (2005) *Mercury Contamination from Historic Gold Mining in California* (US Geological Survey, Sacramento, CA), US Geological Survey Fact Sheet F52005–3014.
- Nater EA, Grigal DF (1992) Regional trends in mercury distribution across the Great-Lakes states, North Central USA. *Nature* 358(6382):139–141.
- Eagles-Smith CA, Ackerman JT, De La Cruz SEW, Takekawa JY (2009) Mercury bioaccumulation and risk to three waterbird foraging guilds is influenced by foraging ecology and breeding stage. *Environ Pollut* 157(7):1993–2002.
- Greenfield BK, Jahn A (2010) Mercury in San Francisco Bay forage fish. *Environ Pollut* 158(8):2716–2724.
- Bouse RM, et al. (2010) Mercury-contaminated hydraulic mining debris in San Francisco Bay. *San Francisco Estuary and Watershed Science Journal* 8(1):28.
- Gehrke GE, Blum JD, Marvin-DiPasquale M (2011) Sources of mercury to San Francisco Bay surface sediment as revealed by mercury stable isotopes. *Geochim Cosmochim Acta* 75(3):691–705.
- Hornberger MI, Luoma SN, van Geen A, Fuller C, Anima R (1999) Historical trends of metals in the sediments of San Francisco Bay, California. *Mar Chem* 64(1-2):39–55.
- James LA (1989) Sustained storage and transport of hydraulic gold mining sediment in the Bear River, California. *Ann Assoc Am Geogr* 79(4):570–592.
- Hunerlach MP, Alpers CN, Marvin-DiPasquale M, Taylor HE, DeWild JF (2004) *Geochemistry of Mercury and Other Trace Elements in Fluvial Tailings Upstream of Daguerre Point Dam, Yuba River, California, August 2001* (US Geological Survey, Reston, VA) US Geological Survey Scientific Investigations Report 2004–5165.
- Adler LL (1980) Adjustment of the Yuba River, California, to the influx of hydraulic mining debris, 1849–1979. PhD dissertation (University of California, Los Angeles).
- James LA, Singer MB, Ghoshal S, Megison M (2009) Historical channel changes in the lower Yuba and Feather Rivers, California: Long-term effects of contrasting river-management strategies. *Management and Restoration of Fluvial Systems with Broad Historical Changes and Human Impacts* (Geological Society of America, Boulder, CO), Geological Society of America Special Publication 451, eds James LA, Rathburn SL, Whittecar GR, pp 57–81.
- Nicholas AP, Quine TA (2007) Modeling alluvial landform change in the absence of external environmental forcing. *Geology* 35(6):527–530.
- James LA (2004) Decreasing sediment yields in northern California: Vestiges of hydraulic gold-mining and reservoir trapping. *International Symposium on Sediment Transfer through the Fluvial System* (IAHS, Wallingford, UK), pp 235–244.
- Kilham NE, Roberts D, Singer MB (2012) Remote sensing of suspended sediment concentration during turbid flood conditions on the Feather River, California: A modeling approach. *Water Resour Res* 48(1):W01521, 10.1029/2011WR010391.
- Singer MB, Aalto R, James LA (2008) Status of the lower Sacramento Valley flood-control system within the context of its natural geomorphic setting. *Natural Hazards Review* 9(3):104–115.
- Walling DE, Owens PN, Leeks GJL (1998) The role of channel and floodplain storage in the suspended sediment budget of the River Ouse, Yorkshire, UK. *Geomorphology* 22(3-4):225–242.
- Rumsby BT, Macklin MG (1994) Channel and floodplain response to recent abrupt climate change: The Tyne basin, Northern England. *Earth Surface Processes and Landforms* 19(6):499–515.
- Dettinger M (2011) Climate change, atmospheric rivers, and floods in California—A multimodel analysis of storm frequency and magnitude changes. *J Am Water Resour Assoc* 47(3):514–523.
- Das T, Dettinger M, Cayan D, Hidalgo H (2011) Potential increase in floods in California's Sierra Nevada under future climate projections. *Clim Change* 109(Suppl 1):71–94.
- Snyder NP, et al. (2004) Estimating accumulation rates and physical properties of sediments behind a dam: Englebright Lake, Yuba River, northern California. *Water Resour Res* 40(11):W11301, 10.1029/2004WR003279.
- James, LA (1993) Sustained reworking of hydraulic mining sediment in California: G.K. Gilbert's sediment wave model reconsidered. *Zeitschrift für Geomorphologie, Supplementbände* 88:49–66.
- Donovan PM, Blum JD, Yee D, Gehrke GE, Singer MB (2013) An isotopic record of mercury in San Francisco Bay sediment. *Chemical Geology* 349–350:87–98.
- Compeau GC, Bartha R (1985) Sulfate-reducing bacteria: Principal methylators of mercury in anoxic estuarine sediment. *Appl Environ Microbiol* 50(2):498–502.
- Gilmour CC, Henry EA, Mitchell R (1992) Sulfate stimulation of mercury methylation in fresh-water sediments. *Environ Sci Technol* 26(11):2281–2287.
- James LA, Hodgson ME, Ghoshal S, Latiolais MM (2012) Geomorphic change detection using historic maps and DEM differencing: The temporal dimension of geospatial analysis. *Geomorphology* 137(1):181–198.
- Simon A, Curini A, Darby SE, Langendoen EJ (2000) Bank and near-bank processes in an incised channel. *Geomorphology* 35(3-4):193–217.
- Singer MB, Dunne T (2004) Modeling decadal bed-material flux based on stochastic hydrology. *Water Resour Res* 40(3):W03302, 10.1029/2003WR002723.
- Hervouet JM (2000) TELEMAC modeling system: An overview. *Hydrol Processes* 14(13): 2209–2210.