

# Genetically Engineered Transvestites Reveal Novel Mating Genes in Budding Yeast

Lori B. Huberman and Andrew W. Murray<sup>1</sup>

Molecular and Cellular Biology and Faculty of Arts and Sciences Center for Systems Biology, Harvard University, Cambridge, Massachusetts 02138

**ABSTRACT** Haploid budding yeast has two mating types, defined by the alleles of the *MAT* locus, *MATa* and *MAT $\alpha$* . Two haploid cells of opposite mating types mate by signaling to each other using reciprocal pheromones and receptors, polarizing and growing toward each other, and eventually fusing to form a single diploid cell. The pheromones and receptors are necessary and sufficient to define a mating type, but other mating-type-specific proteins make mating more efficient. We examined the role of these proteins by genetically engineering “transvestite” cells that swap the pheromone, pheromone receptor, and pheromone processing factors of one mating type for another. These cells mate with each other, but their mating is inefficient. By characterizing their mating defects and examining their transcriptomes, we found *Afb1* (**a-factor barrier**), a novel *MAT $\alpha$* -specific protein that interferes with **a-factor**, the pheromone secreted by *MATa* cells. Strong pheromone secretion is essential for efficient mating, and the weak mating of transvestites can be improved by boosting their pheromone production. Synthetic biology can characterize the factors that control efficiency in biological processes. In yeast, selection for increased mating efficiency is likely to have continually boosted pheromone levels and the ability to discriminate between partners who make more and less pheromone. This discrimination comes at a cost: weak mating in situations where all potential partners make less pheromone.

**B**IOLOGICAL processes are typically defined by the genes that are necessary and sufficient for function. However, in many cases, this minimal gene set does not encompass all the proteins involved in a process, and additional proteins promote biological efficiency. Finding these additional proteins may require the detection of subtle phenotypes, making it hard to know if all the genes involved in a process have been identified. One way to answer this question is to reengineer a pathway and ask whether the synthetic version fully mimics the natural function. Here, we show that this form of synthetic biology illuminates how cells of the budding yeast *Saccharomyces cerevisiae* mate efficiently.

Budding yeast can be stably maintained as haploids or diploids. Haploids mate when two cells of opposite mating types signal to each other using reciprocal pheromones and receptors, polarize and grow toward each other, and eventually fuse to form a single diploid. Yeast has two mating types,

**a** and  $\alpha$  (Figure 1A), determined by two alternative alleles at the *MAT* locus, *MATa* and *MAT $\alpha$* , which encode different transcription factors (Herskowitz 1988). These factors regulate the expression of mating-type-specific genes, many of which are involved with the production and detection of the pheromones that yeast cells use to signal to one another. The pheromones (**a**- and  $\alpha$ -factor) are detected by G-protein-coupled receptors. *MATa* cells express **a-factor** (Betz and Duntze 1979), which is secreted through an ATP binding cassette transporter (*Ste6*) (McGrath and Varshavsky 1989) and the  $\alpha$ -factor receptor (*Ste2*) (Blumer *et al.* 1988; Dohlman and Thorner 2001). *MAT $\alpha$*  cells express  $\alpha$ -factor (Kurjan and Herskowitz 1982; Singh *et al.* 1983) and the **a-factor** receptor (*Ste3*) (Hagen *et al.* 1986; Dohlman and Thorner 2001). Pheromone binding activates a signaling pathway that produces three responses: cell polarization, cell cycle arrest in G1, and increased transcription of pheromone response genes (Bardwell 2005).

Bender and Sprague (1989) used mutations that alter pheromone and receptor expression to show that a cell's mating type is determined by which pheromones and receptors it expresses. Although pheromone secretion and detection are the essential elements for mating, additional, mating-type-specific genes make mating more efficient. One of these is the


Copyright © 2013 by the Genetics Society of America

doi: 10.1534/genetics.113.155846

Manuscript received August 3, 2013; accepted for publication September 30, 2013; published Early Online October 11, 2013.

Supporting information is available online at <http://www.genetic.org/lookup/suppl/doi:10.1534/genetics.113.155846/-/DC1>.

<sup>1</sup>Corresponding author: Northwest Science Building Room 469, 52 Oxford St., Cambridge, MA 02138. E-mail: amurray@mcb.harvard.edu



**Figure 1** Yeast cells expressing the pheromone and receptor of the opposite mating type are capable of forming zygotes with cells of their original mating type. (A) *MATa* cells express the pheromone **a**-factor, the  $\alpha$ -factor receptor Ste2, and the  $\alpha$ -factor protease Bar1. *MAT $\alpha$*  cells express the pheromone  $\alpha$ -factor and the **a**-factor receptor Ste3. (B) *MAT $\alpha$ -playing-a* cells are *MAT $\alpha$*  cells that express **a**-factor instead of  $\alpha$ -factor, Ste2 instead of Ste3, and the  $\alpha$ -factor protease Bar1. *MATa-playing- $\alpha$*  cells are *MATa* cells that express  $\alpha$ -factor instead of **a**-factor, Ste3 instead of Ste2, and lack Bar1 (*bar1 $\Delta$* ). (C) The indicated diploid strains were grown in YPD, and pictures were taken using DIC with  $\times 20$  magnification. (Right) Abnormal morphologies indicative of cells secreting pheromones that they can respond to. (D) Mating efficiency of the indicated crosses. Mating efficiencies are the percentage of diploids that form colonies on double dropout plates relative to the number of colonies formed on single dropout plates. Error bars are standard deviations. Matings were performed as described in *Materials and Methods*.

*MATa*-specific  $\alpha$ -factor protease, Bar1 (Sprague and Herskowitz 1981; MacKay *et al.* 1988), which helps *MATa* cells detect an  $\alpha$ -factor gradient and polarize toward *MAT $\alpha$*  partners (Jackson and Hartwell 1990; Barkai *et al.* 1998). Yeast cells also express mating-type-specific agglutinins, which help cells attach to mating partners (Cappellaro *et al.* 1991) in liquid but individually have little effect on mating efficiency on solid media (Lipke *et al.* 1989; Roy *et al.* 1991; de Nobel *et al.* 1995). Evidence for the final, characterized *MATa*-specific gene was produced by Bender and Sprague (1989) who noted that cells expressing *MATa*-specific proteins and Ste3 were unable to mount a pheromone response. The gene responsible for this was later identified as *ASG7*, which terminates pheromone signaling after mating has occurred and allows diploid cells to escape from the G1 arrest of their parental haploid cell (Kim *et al.* 2000; Roth *et al.* 2000).

Bender and Sprague (1989) used mutations at *MAT* and exogenous promoters to manipulate pheromone and receptor expression. As a result, any quantitative defects in mating could reflect incorrect levels of pheromone and receptor

expression or the accessory role of other genes in mating. To distinguish these possibilities, we constructed “transvestite” strains: genetically engineered strains that have a wild-type allele at *MAT* but express the pheromone, pheromone receptor, and proteins responsible for secreting or processing pheromones that are normally induced by the other *MAT* allele (Figure 1B). These strains should mate well if we have swapped all the genes required for efficient mating and expressed them at the right level. Mating defects in these engineered cells indicate the presence of additional, uncharacterized, mating-type-specific proteins or incorrect expression of the known mating genes.

By studying these genetically engineered cells, we learned more about the requirements for efficient mating. *MATa*-playing- $\alpha$  cells (*MATa* cells that express  $\alpha$ -factor and Ste3) mate threefold less efficiently than genuine *MAT $\alpha$*  cells. Their main defect is low  $\alpha$ -factor secretion: increasing  $\alpha$ -factor production makes them mate almost as well as genuine *MAT $\alpha$*  cells. In contrast, *MAT $\alpha$ -playing-a* cells (*MAT $\alpha$*  cells that express **a**-factor, Ste6, Ste2, and Bar1) mate 60-fold less

efficiently than genuine *MATa* cells. These transvestites have two defects: they express a novel, *MATa*-specific *a*-factor blocker, which we named Afb1 (*a*-factor barrier), and they show a transient as opposed to a prolonged arrest when exposed to *a*-factor. Our manipulations reveal that mating is not robust to reduced levels of pheromone production.

## Materials and Methods

### Yeast strains and culturing

Supporting Information Table S1 lists the strains that we used. All strains were derived from the W303 wild-type background (*ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1*) using standard genetic techniques. All media were prepared as described (Sherman *et al.* 1974) and contained 2% wt/vol of glucose. Cells were grown in synthetic complete (SC) media (2% glucose) or yeast extract peptone dextrose (YPD) (2% glucose) at 30° in culture tubes on roller drums or on agar plates or at room temperature (25°) for time-lapse microscopy. Mating assays used agar plates containing SC without adenine (SC-ade), SC without uracil (SC-ura), or SC without adenine and uracil (SC-ade-ura). Bovine serum albumin (BSA) was used to reduce the nonspecific absorption of *a*-factor to glass and plastic surfaces. A 10% wt/vol stock was prepared in deionized water and then diluted into media to 0.1% wt/vol. Synthetic *a*-factor (Biosynthesis, Lewisville, TX) was suspended in dimethyl sulfoxide and then diluted into either YPD + 0.1% BSA or SC + 0.1% BSA at the appropriate concentration. Yeast extract was obtained from EMD Millipore (Billerica, MA). Peptone and yeast nitrogen base were obtained from BD (Franklin Lakes, NJ). Bacto-agar was obtained from US Biological (Swampscott, MA). Unless otherwise noted, all chemicals were obtained from Sigma-Aldrich (St. Louis).

### Quantitative mating assay

Quantitative mating assays were modified from Reid and Hartwell (1977). Briefly, cells were grown to log phase ( $\sim 5 \times 10^6$  cells/ml). Cells ( $5 \times 10^6$ ) were harvested from each strain, mixed at a 1:1 ratio, sonicated, and filtered onto a 0.22- $\mu$ m nitrocellulose filter (Millipore). Filters were placed on a YPD plate and incubated at 30° for 5 hr. To assay for the initial ratio of the haploid cells, a  $2.5 \times 10^{-5}$  dilution of the initial mating mixture was plated onto SC-ade and SC-ura plates. After 5 hr, cells were washed off the filters into 1 ml of deionized water and then plated onto SC-ade, SC-ura, and SC-ade-ura plates at appropriate dilutions to produce  $\sim 400$  colonies per plate. SC dropout plates were incubated for 2 days before counting the colonies on each plate. Mating efficiencies were determined by dividing the number of colonies on the SC-ade-ura plate by the number of colonies on whichever of the SC-ade or SC-ura plates plated after the mating incubation had fewer colonies. Three technical replicates were done of each mating assay and averaged for a single biological replicate.

Error bars are the standard deviation of at least five biological replicates. Statistical significance was determined using Student's *t*-test.

### Bioassay for *a*-factor production

The bioassay for *a*-factor production was modified from Gonçalves-Sá and Murray (2011). For details see File S1.

### Shmooring index

Cells were grown to log phase ( $\sim 5 \times 10^6$  cells/ml), washed into YPD + 0.1% BSA with various concentrations of synthetic *a*-factor added at  $5 \times 10^5$  cells/ml, and then incubated at 30° on a roller drum for 2 hr. After incubation, the cells were sonicated, fixed using 60% ethanol at -20°, and resuspended into 20% glycerol in phosphate-buffered saline (PBS). Culture tubes were BSA-coated by incubating overnight at 4° with PBS + 2% BSA. The PBS + 2% BSA was poured out immediately prior to the use of the culture tube. At least 200 cells were counted to determine the percentage of cells shmooring. Error bars are standard deviations. Statistical significance was determined using Student's *t*-test.

### Microscopy

Microscopy was done at room temperature using a Nikon Ti-E inverted microscope with either a  $\times 20$  Plan Apo VC 0.75NC air lens or a  $\times 60$  Plan Apo VC 1.4NA oil lens, and images were acquired with a Photometrics CoolSNAP HQ camera (Roper Scientific). Time-lapse photography was done using Metamorph 7.7 (Molecular Devices). For details see File S2.

### Halo assay

Halo assays were modified from Sprague (1991). Cells whose *a*-factor production was to be measured were grown to saturation in YPD at 30°. For halo assays on individual strains,  $4.5 \times 10^8$  cells of each strain were pelleted and resuspended in 20  $\mu$ l of deionized water. For halo assays on cell mixtures, cells were mixed at a 1:8 ratio (*MATa* wild-type:cell type of interest) with a final cell count of  $4.5 \times 10^8$ . Cells were pelleted and resuspended in 20  $\mu$ l of deionized water. Ten microliters of each strain or strain mix was spotted onto YPD plates and incubated overnight ( $\sim 24$  hr) at 30°. Supersensitive *MATa sst2 $\Delta$*  cells grown to stationary phase were then sprayed over the cell spots using a martini atomizer (item 900432, Oenophilia, Hillsborough, NC). Plates were incubated overnight ( $\sim 18$  hr) at 30°, and pictures were taken using a Panasonic (Secaucus, NJ) Lumix DMC-TZ5 camera.

### RNA isolation and sequencing

Cells were grown to log phase ( $5 \times 10^6$  cells/ml) in YPD + 0.1% BSA at 30°. Ten milliliters of the culture was harvested by spinning at 4°, washed in 1 ml RNase-free ice-cold water, pelleted, and flash-frozen in dry ice. *a*-Factor (10 nM) was added to the remaining culture, incubated for 2 hr at 30°, and harvested in the same manner. RNA was isolated as described by Collart and Oliviero (2001) and dissolved in 1 mM sodium

citrate, pH 6.4. RNase-free chemicals were obtained from Invitrogen (Carlsbad, CA) except for chloroform, which was obtained from VWR (Radnor, PA).

RNA libraries were prepared using the Illumina TruSeq kit (<http://www.illumina.com>) and sequenced using an Illumina HiSeq2000 with 50-bp single-end reads with 89× mean coverage across the genome.

### Sequence analysis

To analyze the sequencing data, the RNA sequences were aligned to the S288C reference genome r64 (downloaded from the Saccharomyces Genome Database at <http://www.yeastgenome.org>) using TopHat (Trapnell *et al.* 2009). We then used Cufflinks (Trapnell *et al.* 2010) to look for genes with significantly different levels of gene expression between *MATa* *bar1Δ* cells and *MATα*-playing-*a* *P<sub>BAR1</sub>-BAR1* cells. Significant differences in expression were identified using the default setting in Cufflinks, which tests the observed log-fold change in gene expression against the null hypothesis of no difference between the two samples with a false discovery rate of 0.05 (Trapnell *et al.* 2010). The data discussed in this publication have been deposited in Gene Expression Omnibus at the National Center for Biotechnology Information (NCBI) (Edgar *et al.* 2002) and are accessible through GEO series accession no. GSE49372 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE49372>).

## Results

### Transvestite cells can mate

To find genes required for efficient mating, we constructed two types of transvestite strains. *MATa*-playing- $\alpha$  cells are *MATa* cells that have been engineered to produce  $\alpha$ -factor and the *a*-factor receptor by replacing the open reading frame of *STE2* with *STE3* and the open reading frames of the two *a*-factor genes with those of the two  $\alpha$ -factor genes (replacing *MFA1* with *MF $\alpha$ 1* and *MFA2* with *MF $\alpha$ 2*) (Figure 1B). We also deleted *BAR1*, which encodes the  $\alpha$ -factor protease (Sprague and Herskowitz 1981), and *ASG7*, which inhibits signaling from *Ste3* (Roth *et al.* 2000). However, these cells are still *MATa* at the *MAT* locus and thus will have *MATa*-specific expression patterns for all genes except those we manipulated. *MATα*-playing-*a* cells are *MATα* at the *MAT* locus but have been engineered to produce *a*-factor and the  $\alpha$ -factor receptor by replacing the open reading frame of *STE3* with *STE2*, *MF $\alpha$ 1* with *MFA1*, and *MF $\alpha$ 2* with *MFA2* (Figure 1B). We also drove the expression of *BAR1* with an engineered version of the haploid-specific promoter *P<sub>FUS1</sub>* (Ingolia and Murray 2007), which is expressed in both *MATa* and *MATα* cells (Trueheart *et al.* 1987), and the expression of the *a*-factor transporter *STE6* with the *MF $\alpha$ 1* promoter. Since we tested mating efficiency on solid media, we did not manipulate the expression of the mating agglutinins, which function mainly when cells mate in liquid (Lipke *et al.* 1989; Roy *et al.* 1991).


We asked if the manipulated genes are the only mating-type-specific proteins required for efficient mating. Crossing these cells with wild-type cells of their original mating type (e.g., *MATα*-playing-*a* crossed to *MATα*) carries a potential caveat. When two wild-type cells mate, the combination of the transcription factors expressed from *MATa* and *MATα* turns off the pheromone and receptor genes of both mating types (Haber 1998), and the zygotes escape pheromone-induced G1 arrest (Roth *et al.* 2000). But zygotes produced by crossing a transvestite to a wild-type cell of the same mating type will keep expressing pheromones and receptors from both mating types (since both parents have the same *MAT* locus), raising the concern that these zygotes respond to their own pheromones and remain arrested in G1. To measure the mating efficiency of these crosses, we selected for viable diploids by crossing transvestite cells and wild-type cells with complementary nutritional requirements. We obtained both *MATa*-playing- $\alpha$ /*MATa* and *MATα*-playing-*a*/*MATα* diploids. Most of these diploids progress normally through the cell cycle and have normal cell morphology (Figure 1C). We did find an occasional population of *MATα*-playing-*a*/*MATα* diploids with abnormal morphology, suggesting delayed progression through G1, but even these are capable of budding (Figure 1C). The ability of these diploids to bud indicates that it is possible to measure the mating efficiency of transvestites crossed with wild-type cells.

We used quantitative mating assays to measure the mating efficiency of the transvestite cells. Cells of the two mating types are incubated together and then plated on media that distinguishes diploid cells from either parental haploid. When wild-type *MATa* cells are mated with wild-type *MATα* cells, 66% of haploids form diploids (Figure 1D). However, the mating efficiency of the *MATa*-playing- $\alpha$  cells crossed with *MATa* cells is 3-fold lower than that of a wild-type cross, and the mating efficiency of the *MATα*-playing-*a* cells crossed with *MATα* cells is ~60-fold lower than the efficiency of a wild-type cross (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 1D). These mating defects are synergistic: crossing the transvestite strains to each other decreases mating efficiency ~700-fold (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 1D). Our observation that transvestites can mate with unmanipulated strains with the same *MAT* locus confirms earlier work showing that pheromones and receptors define a cell's mating type (Bender and Sprague 1989). But the low mating efficiency of the transvestite crosses implies that there are additional requirements for efficient mating.

### *MATa*-playing- $\alpha$ cells produce too little $\alpha$ -factor

We studied the mating defects of transvestite cells. *MATa*-playing- $\alpha$  cells mate threefold less efficiently than genuine *MATα* cells (Figure 1D). Because the engineered genes in this strain encode the pheromone and pheromone receptor, the best candidates for this difference were the ability of the *MATa*-playing- $\alpha$  cells to respond to *a*-factor and to produce  $\alpha$ -factor.

We began by testing the response to *a*-factor. We made mating mixtures of *MATa*-playing- $\alpha$  cells expressing YFP



**Figure 2** Low mating efficiency of *MATa*-playing- $\alpha$  cells is due to low  $\alpha$ -factor production. (A) *MATa*-playing- $\alpha$  *P<sub>FUS1</sub>-YFP* cells in a mating mixture with *MATa* *P<sub>ACT1</sub>-mCherry* cells. Yellow indicates YFP expression. Red indicates mCherry expression. The orange cell is a diploid expressing both YFP and mCherry. The picture was taken 2.5 hr after mixing the cells using DIC and fluorescence at  $\times 20$  magnification. (B)  $\alpha$ -Factor production is measured by growing cells in YPD, harvesting the supernatant, and exposing *MATa bar1Δ* cells to the supernatant. The shmooving index of the *MATa bar1Δ* cells is measured and then compared to a standard curve, produced with synthetic  $\alpha$ -factor, to determine the amount of  $\alpha$ -factor present in the media. Error bars are standard deviations. (C) Mating efficiency relative to a wild-type cross between *MATa* and *MATα* cells. Matings were performed as described in *Materials and Methods*. Error bars are standard deviations.

under the pheromone-inducible promoter *P<sub>FUS1</sub>* and *MATa* cells expressing mCherry under the *ACT1* promoter and assayed for the expression of YFP in the *MATa*-playing- $\alpha$  cells after 2.5 hr. The expression of YFP in the *MATa*-playing- $\alpha$  cells indicates that they can successfully detect **a**-factor using the **a**-factor receptor and activate pheromone-induced genes (Figure 2A). The two cell types mated to form zygotes that continue to signal to themselves, thus forming diploid cells, which express both YFP under the *FUS1* promoter and mCherry under the *ACT1* promoter (Figure 2A). Since the G-protein and downstream components of the pheromone-signaling pathway should be the same in *MATa* and *MATα* cells (Bardwell 2005), we asked if reduced pheromone production is the cause of the mating defect.

Pheromone production is important for zygote formation (Kurjan 1985; Michaelis and Herskowitz 1988), and *MATa* cells prefer the *MATα* cell that produces the highest amount of  $\alpha$ -factor (Jackson and Hartwell 1990). Since we manipulated pheromone genes to make the transvestites, we measured the pheromone production of *MATa*-playing- $\alpha$  cells using a bioassay. We grew cells in rich medium, filtered out the cells, incubated the medium with *MATa* cells lacking

the  $\alpha$ -factor protease *Bar1*, measured the fraction of cells that arrest and shmoo (the shmooing index), and compared these data to a standard curve generated with synthetic  $\alpha$ -factor. Unstimulated, *MATa*-playing- $\alpha$  cells produce  $\sim 70$  times less  $\alpha$ -factor than *MATα* cells (Student's *t*-test,  $P = 0.01$ ) (Figure 2B). To measure the  $\alpha$ -factor production of stimulated cells, we mixed the  $\alpha$ -factor-producing cells in a 10:1 mixture with *MATa bar1Δ* cells (which produce **a**-factor and do not destroy  $\alpha$ -factor) and measured the  $\alpha$ -factor present in the supernatant. Stimulated *MATa*-playing- $\alpha$  cells produce 20-fold less  $\alpha$ -factor than stimulated wild-type *MATα* cells (Student's *t*-test,  $P = 9 \times 10^{-6}$ ) (Figure 2B).

To test the effect of reduced pheromone production in *MATα* cells, we knocked out *MFα1*, which is the majority  $\alpha$ -factor producer in *MATα* cells (Kurjan 1985). This reduces  $\alpha$ -factor production 12-fold compared to wild-type *MATα* cells in unstimulated cells (Student's *t*-test,  $P = 0.02$ ) and 9-fold in stimulated cells (Student's *t*-test,  $P = 10^{-5}$ ) (Figure 2B). We compared the mating efficiency of *MATα mfa1Δ* cells, which have decreased mating efficiency (Kurjan 1985), to that of *MATa*-playing- $\alpha$  cells and determined that *MATα mfa1Δ* cells mate only 1.5-fold more efficiently than *MATa*-playing- $\alpha$

cells (Student's *t*-test,  $P = 0.004$ ) (Figure 2C). This suggests that the reduced mating efficiency of the *MATa*-playing- $\alpha$  cells is due, at least in part, to low  $\alpha$ -factor production.

To test this hypothesis, we increased  $\alpha$ -factor production in the *MATa*-playing- $\alpha$  cells by expressing *MF $\alpha$ 1* from the *TDH3* promoter. This promoter is not pheromone-regulated, but it is one of the most highly expressed promoters in the yeast genome (Krebs 1953; McAlister and Holland 1985) and should increase  $\alpha$ -factor production to at least wild-type *MAT $\alpha$*  levels. Unstimulated *MATa*-playing- $\alpha$  *P<sub>TDH3</sub>-MF $\alpha$ 1* cells secrete twice as much  $\alpha$ -factor as unstimulated *MAT $\alpha$*  cells (Student's *t*-test,  $P = 0.04$ ), but when stimulated, *MATa*-playing- $\alpha$  *P<sub>TDH3</sub>-MF $\alpha$ 1* cells secrete fivefold more  $\alpha$ -factor than stimulated *MAT $\alpha$*  cells (Student's *t*-test,  $P = 10^{-4}$ ) (Figure 2B), suggesting that  $\alpha$ -factor production is regulated by post-translational mechanisms, such as pheromone maturation and secretion and transcription of the pheromone genes. If low  $\alpha$ -factor production accounts for the weak mating of *MATa*-playing- $\alpha$  cells, *MATa*-playing- $\alpha$  *P<sub>TDH3</sub>-MF $\alpha$ 1* cells should have a mating efficiency approaching that of wild-type *MAT $\alpha$*  cells, which is indeed what we found (Figure 2C). This confirms previous results, which showed that sufficient  $\alpha$ -factor production is important for efficient mating (Kurjan 1985) and shows that the principal defect of *MATa*-playing- $\alpha$  cells is insufficient  $\alpha$ -factor production. This defect could reflect a difference in the strengths of the *MFA1* vs. the *MF $\alpha$ 1* promoter or differences in the translation or processing of  $\alpha$ -factor between *MATa* and *MAT $\alpha$*  cells. Our analysis also shows that there are no additional *MATa*-specific genes, beyond those that we manipulated (*STE2*, *MFA1*, *MFA2*, *BAR1*, and *ASG7*), that interfere with the ability of *MAT $\alpha$*  cells to mate.

### ***AFB1* encodes a novel a-factor barrier protein**


We examined the decreased mating efficiency of the *MAT $\alpha$* -playing-*a* cells. Because pheromone production is important for efficient mating (Kurjan 1985; Michaelis and Herskowitz 1988), we investigated *a*-factor production of *MAT $\alpha$* -playing-*a* cells. Both  $\alpha$ -factor and *a*-factor go through several processing steps before secretion (Betz and Duntze 1979; Kurjan and Herskowitz 1982). But while  $\alpha$ -factor is secreted as a small, unmodified peptide (Kurjan and Herskowitz 1982), mature *a*-factor is modified with a 15-carbon farnesyl group, causing it to be very hydrophobic (Betz and Duntze 1979; Chen *et al.* 1997) and hard to quantify biochemically. We therefore used a bioassay to measure the relative *a*-factor production of the *MAT $\alpha$* -playing-*a* cells: we plated patches of *a*-factor-producing cells and then sprayed the plates with a suspension of *MAT $\alpha$*  cells that were made supersensitive to pheromone by deleting *SST2*, which encodes a GTPase-activating protein that reduces the duration of signaling from the pheromone-activated G protein (Chan and Otte 1982a,b; Dohlman *et al.* 1996; Apanovitch *et al.* 1998). The *a*-factor secreted by the patch of cells arrests the *MAT $\alpha$*  *sst2 $\Delta$*  tester cells in G1, producing a halo of growth inhibition (Chan and Otte 1982a,b); the halo's diameter increases with the amount of *a*-factor

produced by the cell patch (Figure 3A). The halo produced by *MAT $\alpha$* -playing-*a* cells is smaller than that of wild-type *MATa* cells, implying that *MAT $\alpha$* -playing-*a* cells secrete less *a*-factor than wild-type *MATa* cells (Figure 3B).

We considered two explanations for the low *a*-factor secretion of *MAT $\alpha$* -playing-*a* cells: *MAT $\alpha$* -playing-*a* cells secrete less *a*-factor than *MATa* cells or *MAT $\alpha$*  cells secrete a protein that provides a barrier to *a*-factor that is analogous to the *MATa*-specific  $\alpha$ -factor protease, *Bar1*. We tested for the presence of a *MAT $\alpha$* -specific *a*-factor blocker secreted from *MAT $\alpha$* -playing-*a* cells by comparing the halo sizes of two mixtures of cells: *MATa* cells mixed with *MAT $\alpha$* -playing-*a* cells and *MATa* cells mixed with *MATa* cells that lack the genes encoding *a*-factor and thus produce no *a*-factor (*MATa mfa1 $\Delta$  mfa2 $\Delta$* ). If the *MAT $\alpha$* -playing-*a* cells secrete an *a*-factor blocker, we would expect the halo size of the *MATa* cells mixed with *MAT $\alpha$* -playing-*a* cells to be smaller than that of the *MATa* cells mixed with pheromone-less *MATa* cells because the *a*-factor blocker would interfere with the *a*-factor from both the *MAT $\alpha$* -playing-*a* cells and the *MATa* cells. However, if there is no *a*-factor blocker, we would expect the halo size of the *MATa* cells mixed with *MAT $\alpha$* -playing-*a* cells to be larger than that of the *MATa* cells mixed with pheromone-less *MATa* cells because both the *MATa* cells and the *MAT $\alpha$* -playing-*a* cells are capable of secreting *a*-factor. The halo produced by *MATa* cells mixed with *MAT $\alpha$* -playing-*a* cells is smaller than the halo produced by *MATa* cells mixed with pheromone-less *MATa* cells, indicating that *MAT $\alpha$* -playing-*a* cells secrete an *a*-factor blocker (Figure 3C).

We searched for the gene responsible for this activity by comparing the transcriptomes of *MATa* and *MAT $\alpha$* -playing-*a* cells. Although the gene expression of pheromone-stimulated *MATa* cells has been investigated, the extreme hydrophobicity of *a*-factor has made similar experiments on pheromone-stimulated *MAT $\alpha$*  cells difficult (Roberts *et al.* 2000). The *MAT $\alpha$* -playing-*a* cells make it possible to study the transcriptome of pheromone-stimulated cells that are *MAT $\alpha$*  at the *MAT* locus but are stimulated by  $\alpha$ -factor in a controlled fashion. We chose a concentration of pheromone, 10 nM, in a regime in which *MATa bar1 $\Delta$*  and *MAT $\alpha$* -playing-*a* cells with *BAR1* under its endogenous promoter have a similar shmooing index (Figure 4A) to compare the transcriptomes of stimulated and unstimulated *MATa* and *MAT $\alpha$* -playing-*a* cells using RNA sequencing.

Just as *MAT $\alpha$*  cells do not secrete *Bar1* to cleave their own  $\alpha$ -factor (Sprague and Herskowitz 1981), we would not expect *MATa* cells to secrete an *a*-factor blocker to inhibit their own *a*-factor. Thus, we hypothesized that a *MAT $\alpha$* -specific *a*-factor blocker would be expressed more highly in both pheromone-stimulated and unstimulated *MAT $\alpha$* -playing-*a* than *MATa* cells. Ten genes fit this criterion and of these, only one, *YLR040C*, is annotated as encoding a secreted protein that is not already known to be important in mating (Yeast Genome Database at <http://www.yeastgenome.org>) (see Table S2).



**Figure 3** *AFB1* encodes a novel **a-factor** blocker. (A) Halo assays are done by allowing cell patches to grow on YPD overnight and then spraying supersensitive  $MAT\alpha$  *sst2*Δ cells over the cell patches. Where **a-factor** produced by the cell patches has diffused into the YPD, the  $MAT\alpha$  *sst2*Δ cells cannot grow, forming a halo around the cell patch with a size that corresponds to the amount of **a-factor** secretion. (B) Halo assays done on various cell

*YLR040C* was previously identified as an  $\alpha$ -specific gene by its reduced transcription in a *MAT $\alpha$*  cell that lacked the transcription factor *Mat $\alpha$ 1* (Galgoczy *et al.* 2004), which induces expression of  $\alpha$ -specific genes (Strathern *et al.* 1981). *YLR040C* has also been shown to be translated by ribosome profiling (Brar *et al.* 2012) and localized to the cell wall (Hamada *et al.* 1999; Giaever *et al.* 2002). Deletion of *YLR040C* was reported as having no effect on mating (Galgoczy *et al.* 2004). We found that in unstimulated cells the gene is expressed 11-fold more strongly in *MAT $\alpha$ -playing-a* than in *MATa* cells and that its transcription is not significantly induced when *MAT $\alpha$ -playing-a* cells are exposed to pheromone (see Table S2). The protein is conserved in yeasts that experienced the whole-genome duplication around 100 million years ago and is also found in some yeasts, such as *Hansenula polymorpha*, that substantially predate this event (Wolfe and Shields 1997; Dietrich *et al.* 2004; Dujon *et al.* 2004; Kellis *et al.* 2004) (Figure 3D). The experiments described below demonstrate that *YLR040C* encodes a protein that provides an **a-factor barrier** function, leading us to name this gene *AFB1* for **a-factor barrier**.

To determine whether *AFB1* is indeed the **a-factor** blocker, we knocked it out in *MAT $\alpha$ -playing-a* cells. The halos produced by *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells are larger than those of *MAT $\alpha$ -playing-a AFB1* cells, indicating that deleting *AFB1* increases the amount of pheromone secreted from a patch of *MAT $\alpha$ -playing-a* cells (Figure 3B). The halo around the *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells, however, is still smaller than the halo produced by wild-type *MATa* cells, suggesting that *MAT $\alpha$ -playing-a* cells secrete less **a-factor** than wild-type *MATa* cells (Figure 3B). We also placed *AFB1* under a strong (*ACT1*) promoter in *MATa* cells and observed a decrease in halo size compared to wild-type *MATa* cells, indicating that *Afb1* is able to block **a-factor** secreted by *MATa* cells (Figure 3B).

To test whether *Afb1* is responsible for blocking **a-factor** produced by other cells, we made cell mixtures of *MATa* cells with *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells and compared the halo produced by this mixture to the halo produced by the mixtures of *MATa* cells with *MAT $\alpha$ -playing-a* cells and to the halo produced by *MATa* cells with *MATa mfa1 $\Delta$  mfa2 $\Delta$*  cells. As expected from our other results, the mixture of *MATa* cells with *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells has a slightly

larger halo than the *MATa* cells mixed with *MATa mfa1 $\Delta$  mfa2 $\Delta$*  cells and a significantly larger halo than the *MATa* cells mixed with *MAT $\alpha$ -playing-a* cells (Figure 3C). This result indicates that, when *Afb1* is not present in the cell mixtures, the **a-factor** from the wild-type *MATa* cells as well as that from the *MAT $\alpha$ -playing-a* cells is free to interact with the supersensitive *MAT $\alpha$*  cells. Taken together, our results provide strong evidence that *Afb1* has an **a-factor barrier** function.

We asked whether the expression of *AFB1* affected the mating efficiency of *MAT $\alpha$ -playing-a* cells. We crossed *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells with wild-type *MAT $\alpha$*  cells and observed a fivefold increase in mating efficiency over a similar cross with *MAT $\alpha$ -playing-a* cells (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 3E). However, deleting *AFB1* from wild-type *MAT $\alpha$*  cells does not reduce their mating efficiency (Figure 3E), perhaps because small changes in **a-factor** production do not have a large effect on mating efficiency (Michaelis and Herskowitz 1988). We tested this possibility in two ways. The first was to delete *MFA1* from wild-type *MATa* cells. We saw a small decrease in the halo size of *MATa mfa1 $\Delta$*  compared to that of wild-type *MATa* cells, but as previously reported (Michaelis and Herskowitz 1988), the mating efficiency of *MATa mfa1 $\Delta$*  cells was statistically indistinguishable from that of *MATa MFA1* cells (Figure 3, B and E). We also tested the mating efficiency of *MATa* cells with *AFB1* placed under the *ACT1* promoter. These cells produce a smaller halo than wild-type *MATa* cells (Figure 3B) but mate slightly better than wild-type *MATa* cells, implying that there is a range of **a-factor** production that results in efficient mating, at least in the absence of additional mutations (Student's *t*-test,  $P = 0.03$ ) (Figure 3E).

#### ***MAT $\alpha$ -playing-a* cells shmoo but arrest only transiently in the presence of pheromone**


Although the expression of *AFB1* in *MAT $\alpha$ -playing-a* cells was responsible for a portion of the reduced mating efficiency of *MAT $\alpha$ -playing-a* cells, *MAT $\alpha$ -playing-a afb1 $\Delta$*  cells still mate 12-fold less efficiently than wild-type *MATa* cells (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 3E). We hypothesized that the response of *MAT $\alpha$ -playing-a* cells to pheromone could also reduce their mating efficiency.

There are three cellular responses to pheromone stimulation: altered gene expression, cell polarization, and cell

---

patches containing a single cell type. *MATa mfa1 $\Delta$  mfa2 $\Delta$*  is a negative control, and *MATa* is a positive control. White bars indicate the width of the halo. The halo assays were repeated multiple times, and the rank order of the halo sizes is consistent. (C) Halo assays done on cell patches containing two cell types at a 1:8 ratio of *MATa* cells to the experimental cell of interest. White bars indicate the width of the halo. The halo assays were repeated multiple times, and the rank order of the halo sizes is consistent. (D) Sequences were obtained from the Yeast Genome Order Browser (<http://ygod.ucc.ie>) and Blast searches of fungal genomes at NCBI. Sequences were aligned in Jalview using the MAFFT-L-INS-I option, and the aligned core that follows the signal sequence and precedes the GPI anchorage sequence is shown. Amino acids are colored with a scheme that represents chemically similar amino acids in similar colors, and universally conserved amino acids are highlighted in red in the consensus sequence. Full species names: *Saccharomyces cerevisiae*, *Saccharomyces mikatae*, *Saccharomyces kudriavzevii*, *Saccharomyces bayanus*, *Candida glabrata*, *Kazachstania africanus*, *Kazachstania naganishii*, *Naumovzzyma castelli*, *Naumovzzyma dairenensis*, *Tetraspora blattae*, *Tetraspora phaffi*, *Vanderwaltomyzoa polyspora*, *Zygosaccharomyces rouxii*, *Torulasporea delbrueckii*, *Kluveromyces lactis*, *Saccharomyces kluyveri*, *Eremothecium cymbalariae*, and *Hansenula/Ogataea polymorpha*. Of these, *Z. rouxii*, *T. delbrueckii*, *K. lactis*, *S. kluyveri*, *E. cymbalariae*, and *H. polymorpha* did not undergo a whole-genome duplication. (E) Mating efficiency of the indicated crosses relative to a wild-type cross between *MATa* and *MAT $\alpha$*  cells. Matings were performed as described in *Materials and Methods*. Error bars are standard deviations.





**Figure 4** *MATα*-playing-*a* cells shmoo but arrest transiently in the presence of pheromone. (A) Shmooing indices of *MATa bar1Δ* cells and *MATα*-playing-*a P<sub>BAR1</sub>-BAR1* cells exposed to known concentrations of  $\alpha$ -factor. Error bars are standard deviations. (B) Shmooing indices of *MATa* cells and *MATα*-playing-*a P<sub>FUS1</sub>\*-BAR1* cells exposed to known concentrations of  $\alpha$ -factor. Error bars are standard deviations. (C) Mating efficiency relative to a wild-type cross between *MATa* and *MATα* cells. Matings were performed as described in *Materials and Methods*. Error bars are standard deviations. (D) *MATa bar1Δ* cells shmooing and *MATα*-playing-*a bar1Δ* cells shmooing and budding when incubated with SC plus 10 nM  $\alpha$ -factor in a microfluidic chamber. Pictures were taken using DIC with  $\times 60$  magnification 8 hr after the addition of  $\alpha$ -factor. White arrows point to buds.

cycle arrest (Bardwell 2005). We compared the transcriptomes of *MATα*-playing-*a* and *MATa* cells both with and without exposure to  $\alpha$ -factor (see Table S2), excluding those genes, such as *STE3* and *BAR1*, that had been removed during the construction of the strains. Twenty-one genes showed a more than 2-fold variation in both comparisons. Ten genes showed a more than 2-fold variation when comparing the two stimulated cell types but were not significantly different when comparing the unstimulated cells, and another 8 genes showed the opposite pattern. As expected, known  $\alpha$ -specific genes, such as the *MATα*-specific agglutinin gene, *SAG1*, were expressed more strongly in *MATα*-playing-*a* cells than in *MATa* cells, and known *a*-specific genes, such as the *a*-specific agglutinin gene, *AGA2*, were expressed less strongly in *MATα*-playing-*a* cells than in *MATa* cells. Despite our attempts to engineer their expression to match the levels seen in *MATa* cells, three important *a*-specific genes—*STE2*, *MFA1*, and *MFA2*—are expressed at 5-fold, 7-fold, and 120-fold lower levels, respectively, in *MATα*-playing-*a* cells compared to *MATa* cells (see Table S2).

We assayed cell polarization (Segall 1993; Butty *et al.* 1998) by measuring the shmooing index of *MATα*-playing-*a* cells stimulated with known quantities of synthetic  $\alpha$ -factor. We found that *MATα*-playing-*a P<sub>BAR1</sub>-BAR1* cells have a similar shmooing index to *MATa bar1Δ* cells at low concentra-


tions of  $\alpha$ -factor, indicating that the *MATα*-playing-*a* cells are as sensitive to low concentrations of  $\alpha$ -factor as *MATa* cells (Figure 4A) and suggesting that *BAR1* is not expressed in *MATα* cells. Because efficient mating in *MATa*  $\times$  *MATα* crosses depends on the secretion of *Bar1* by the *MATa* cells, we investigated the pheromone response of *MATα*-playing-*a* cells that express *BAR1* from a mutant version of the pheromone-induced *FUS1* promoter (Trueheart *et al.* 1987): the mutant promoter *P<sub>FUS1</sub>\** was selected to have a low basal and a high pheromone-stimulated level of expression (Ingolia and Murray 2007). These cells make fewer shmooos at 50 nM  $\alpha$ -factor than *MATα*-playing-*a P<sub>BAR1</sub>-BAR1* cells (which lack detectable *Bar1* activity) make at 2 nM  $\alpha$ -factor, indicating that *MATα*-playing-*a P<sub>FUS1</sub>\*-BAR1* cells are able to secrete *Bar1* (Student's *t*-test,  $P = 0.02$ ) (Figure 4, A and B). In contrast, the *MATα*-playing-*a P<sub>FUS1</sub>\*-BAR1* cells make significantly more shmooos than wild-type *MATa* cells at each concentration of  $\alpha$ -factor tested (Student's *t*-test,  $P < 0.005$ ) (Figure 4B), suggesting that *MATα*-playing-*a P<sub>FUS1</sub>\*-BAR1* cells secrete less *Bar1* than wild-type *MATa* cells. This difference is unlikely to be the sole remaining cause of the mating defect of *MATα*-playing-*a P<sub>FUS1</sub>\*-BAR1* cells: reducing the expression of *Bar1* in *MATa* cells, by expressing it from the *FUS1*\* promoter, does not reduce their mating efficiency (Figure 4C).

We observed  $\alpha$ -factor-induced cell cycle arrest (Chang and Herskowitz 1990) in a microfluidic device. Pheromone stimulation arrests *MATa* cells in G1 through phosphorylation of *Far1*, a protein that binds to cyclin-dependent kinase/cyclin complexes (Chang and Herskowitz 1990; Tyers and Futcher 1993; Peter and Herskowitz 1994). When *MATa bar1Δ* cells are exposed to 10 nM  $\alpha$ -factor, their cell cycle remains arrested for many hours while they form multiple successive shmoos (Figure 4D and File S3). However, even at this high  $\alpha$ -factor concentration, *MAT $\alpha$ -playing-a bar1Δ* cells form shmoos but arrest only transiently (Figure 4D and File S4).

The transient cell cycle arrest in *MAT $\alpha$ -playing-a* cells could be due to a difference in the response of *MAT $\alpha$*  and *MATa* cells to pheromone stimulation or the inhibition of *Ste2* by *MAT $\alpha$* -specific proteins in the *MAT $\alpha$ -playing-a* cells. We tried to find the responsible genes by looking for differential expression of genes that might have an effect on cell cycle arrest between pheromone-stimulated *MATa* and *MAT $\alpha$ -playing-a* cells. We manipulated the expression of three candidates—*PCL1*, *GYP8*, and *TOS4*—which had at least a twofold difference in expression between stimulated *MATa* and *MAT $\alpha$ -playing-a* cells and a plausible connection to cell cycle control (see Table S2). None of these manipulations altered the pheromone-induced cell cycle arrest of either *MATa* or *MAT $\alpha$ -playing-a* cells (data not shown).

### How robust is mating?

Mating would be robust to variation in pheromone levels if substantial increases or decreases in pheromone expression had no effect on mating efficiency. The mating of the transvestite strains to each other suggests that mating efficiency is not robust to variation in pheromone production. Mating *MATa*-playing- $\alpha$  cells to wild-type *MATa* cells reduces mating frequency 3-fold, and mating *MAT $\alpha$ -playing-a* cells to wild-type *MAT $\alpha$*  cells reduces mating 60-fold, relative to a wild-type *MATa*  $\times$  *MAT $\alpha$*  cross, but the mating frequency of the cross between the two transvestites is reduced 700-fold, suggesting that mating defects are synergistic (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 1D). If this synergism is largely due to reduced pheromone production by the transvestite strains, increasing pheromone production should increase the efficiency of the intertransvestite cross. We increased  $\alpha$ -factor production from *MATa*-playing- $\alpha$  cells by placing *MF $\alpha$ 1* under the control of the *TDH3* promoter and  $\alpha$ -factor production from *MAT $\alpha$ -playing-a* cells by deleting *AFB1*. When crossed to each other, these strains mate 90 times better than the cross between the original *MATa*-playing- $\alpha$  and *MAT $\alpha$ -playing-a* cells. Thus, after improving pheromone production, the intertransvestite cross is only 8-fold less efficient than a standard *MATa*  $\times$  *MAT $\alpha$*  cross (Student's *t*-test,  $P < 10^{-6}$ ) (Figure 5). If reduced pheromone production is the primary cause of the weak mating of the intertransvestite cross, the cross between a *MAT $\alpha$*  strain making less  $\alpha$ -factor and a *MAT $\alpha$ -playing-a* cell should mimic the intertransvestite cross. The mating efficiency of the cross



**Figure 5** Mating is not robust to changes in pheromone production. Mating efficiency relative to a wild-type cross between *MATa* and *MAT $\alpha$*  cells. Matings were performed as described in *Materials and Methods*. Note the logarithmic scale for mating efficiency. Error bars are standard deviations.

between *MAT $\alpha$  mfa1Δ* cells, which produce less  $\alpha$ -factor than wild-type *MAT $\alpha$*  cells, and *MAT $\alpha$ -playing-a* cells is statistically indistinguishable from that of the double transvestite cross (Figure 2B and Figure 5).

## Discussion

Our experiments show that genetic engineering can be used to investigate the factors that control the efficiency of mating in budding yeast. We tested the idea that previous research had found all the genes that control mating efficiency by engineering transvestite strains that switch the mating genes of one mating type for those that are normally expressed in its partner. The behavior of these strains led to two conclusions: there are still more genes that control mating, such as the  $\alpha$ -factor barrier protein *Afb1*, and mating is not robust to reductions in pheromone production.

### Engineering efficient maters

Investigating the pheromone production of the transvestite strains allowed us to account for a significant portion of their mating defects. Unstimulated *MATa*-playing- $\alpha$  cells secrete 70-fold less  $\alpha$ -factor than wild-type *MAT $\alpha$*  cells, and stimulated *MATa*-playing- $\alpha$  cells secrete 20-fold less  $\alpha$ -factor than stimulated *MAT $\alpha$*  cells. Increasing the  $\alpha$ -factor production of the *MATa*-playing- $\alpha$  cells increased their mating efficiency to nearly that of wild type, showing that the main defect of the *MATa*-playing- $\alpha$  cells is low  $\alpha$ -factor production and that the level of  $\alpha$ -factor secretion is important for efficient mating. The observation that  $\alpha$ -factor secretion is still pheromone-inducible, even when  $\alpha$ -factor expression is driven by a strong, constitutive promoter, demonstrates that pheromone processing and export respond to pheromone stimulation. Indeed, *Ste13*, a protein required for the maturation

of  $\alpha$ -factor (Julius *et al.* 1983), is pheromone-induced (Achstetter 1989).

The mating defects of *MAT $\alpha$ -playing-a* cells are more complex. We determined that these cells do not make as much **a**-factor as wild-type *MATa* cells and that at least part of this is due to the expression of the novel **a**-factor blocker *Afb1*. Increasing the **a**-factor production of *MAT $\alpha$ -playing-a* cells by deleting *AFB1* causes a fivefold increase in their mating efficiency, indicating that sufficient **a**-factor expression is important for efficient mating as a *MATa* cell. We were unable to engineer *MAT $\alpha$*  cells to mate efficiently as *MATa* cells. There are two possible explanations for the remaining defect: even after the removal of *Afb1*, the *MAT $\alpha$ -playing-a* cells make less **a**-factor than *MATa* cells, and *MAT $\alpha$ -playing-a* cells arrest only transiently in response to  $\alpha$ -factor. We suspect that both contribute to the reduced mating of *MAT $\alpha$ -playing-a* cells.

Similar pheromone and receptor swaps have been done on other fungi, including *Cryptococcus neoformans* (Stanton *et al.* 2010) and *Ustilago maydis* (Bölker *et al.* 1992). Like the strains we constructed, the engineered versions of these organisms could mate to cells that bore the same genes at the mating-type locus. In *Candida albicans*, **a-a** or  $\alpha$ - $\alpha$  matings can be induced by enhancing autocrine signaling or the presence of the opposite pheromone (Alby *et al.* 2009). Studying the mating defects of engineered transvestites in other fungi should identify additional genes involved in their mating pathways.

#### ***AFB1* encodes a novel *MAT $\alpha$ -specific a-factor barrier protein***

Studies on pheromone-induced genes in *MAT $\alpha$*  cells were hampered by the difficulties in working with **a**-factor. We avoided these by looking at the pattern of gene expression in *MAT $\alpha$ -playing-a* cells, which would still express  $\alpha$ -specific genes but would increase their expression in response to  $\alpha$ - rather than **a**-factor. We argued that novel  $\alpha$ -specific genes would be identified by higher expression in *MAT $\alpha$ -playing-a* than in *MATa* cells. Mixing experiments suggested that *MAT $\alpha$ -playing-a* cells produced an extracellular factor that interfered with the action of **a**-factor, prompting us to look for the secreted product of a *MAT $\alpha$ -specific* gene. This computational sieve produced a single gene, *YLR040C*, which had previously been identified as a gene regulated by the *MAT $\alpha$ -specific* transcription factor, *Mat $\alpha$ 1* (Galgoczy *et al.* 2004). Removing *YLR040C* increased **a**-factor production from *MAT $\alpha$ -playing-a* cells and the mating efficiency of *MAT $\alpha$ -playing-a* cells, leading us to rename *YLR040C* *AFB1* for **a**-factor barrier. There have been previous searches for a protein with an **a**-factor barrier function. The first reported a supersensitive *MAT $\alpha$*  mutant, which mapped to a location on chromosome XII >600 kilobases away from *AFB1* (Steden *et al.* 1989). The second reported the detection of *MAT $\alpha$ -specific* **a**-factor endopeptidase activity, but the gene responsible for this was not identified, and the protein was not purified (Marcus *et al.* 1991). Without being able to manip-

ulate the genes involved in these studies, it is impossible to assess their effect on **a**-factor activity or stability or their relationship to *AFB1*.

Deleting *AFB1* increased the mating efficiency of *MAT $\alpha$ -playing-a* cells. Sequence analysis shows that *Afb1* is conserved in ascomycetes as evolutionarily distant as *H. polymorpha* and contains an N-terminal signal sequence and C-terminal motif that suggests that it is a GPI-anchored protein (Hamada *et al.* 1999) but lacks other detectable motifs. In particular, *Afb1* shows no sequence homology with any other protease but contains a number of conserved aromatic residues (Figure 3D). Our inability to find *Afb1* throughout the ascomycete fungi has two possible interpretations: either the protein evolves too rapidly to be detected by standard tools that use sequence homology to identify orthologs or the protein evolved within one branch of the ascomycete lineage, rather than in its last common ancestor. Unusually rapid evolution of a single protein or independent evolution of the same function in different lineages may also explain why the  $\alpha$ -factor-degrading protease *Bar1* in *S. cerevisiae* is not the closest homolog of the same protein in *C. albicans* (Schaefer *et al.* 2007).

We speculate that *Afb1* acts to bind and sequester **a**-factor rather than to degrade it. The biological function of *Afb1* may mirror that of *Bar1*, which promotes the efficient mating of *MATa* cells by keeping the  $\alpha$ -factor concentration at the plasma membrane within the narrow range needed for accurate pheromone gradient detection (Barkai *et al.* 1998). Since *Afb1* is predicted to be GPI-anchored, it is possible that the function of *Afb1* closely mimics that of *Bar1* trapped in the cell wall of *MATa* cells: creating a pheromone sink that makes it more likely that two cells of the same mating type will avoid each other (Jin *et al.* 2011) and makes it easier to distinguish between two close potential partners (Rappaport and Barkai 2012). It is also possible that *Afb1* in *S. cerevisiae* acts like *Bar1* in *C. albicans* (Alby *et al.* 2009): decreasing the threat of autocrine signaling caused by leaky repression of **a**-factor in *MAT $\alpha$*  cells.

#### ***MAT $\alpha$ -playing-a* cells arrest only transiently in response to pheromone**

In *MATa* cells, exposure to  $\alpha$ -factor leads to a prolonged cell cycle arrest. In contrast, *MAT $\alpha$ -playing-a* cells show only a transient arrest, even though their ability to shmoo is statistically indistinguishable from wild-type *MATa* cells. This result surprised us because *MATa* and *MAT $\alpha$*  cells arrest the cell cycle in the same fashion: by signaling through *Far1* (Peter and Herskowitz 1994; Bardwell 2005).

There are two possible explanations for the transient cell cycle arrest of *MAT $\alpha$ -playing-a* cells. The first is that *MATa* and *MAT $\alpha$*  cells have evolved to respond to pheromone stimulation in subtly different ways and that *MAT $\alpha$*  cells shmoo but do not experience enduring arrest. Although it is important for cells to be in the same phase of the cell cycle during nuclear fusion, it is possible that transient arrest of *MAT $\alpha$*  cells is sufficient to allow for the formation of zygotes, while

a lasting arrest is required for *MATa* cells. Because  $\alpha$ -factor is more diffusible, we suspect that initial signaling is usually from *MAT $\alpha$*  to *MATa* cells, meaning that it is the *MATa* cells that arrest first and thus need to wait until the *MAT $\alpha$*  cells receive a strong enough signal to arrest, implying that fusion would usually occur shortly after the arrest of the *MAT $\alpha$*  cell but at a longer and more variable time after the arrest of the *MATa* cell.

The second possibility is that interactions between *Ste2* and proteins present in the *MAT $\alpha$* -playing-*a* cells make the cells keep cycling, like cells that express both *Ste3* and *Asg7* (Bender and Sprague 1989; Roth *et al.* 2000). We looked for genes that might be responsible for the lack of enduring arrest in *MAT $\alpha$* -playing-*a* cells by focusing on genes that are differentially expressed in pheromone-stimulated *MATa* and *MAT $\alpha$* -playing-*a* cells and might not have been identified in earlier work. Although we tested the effect of deleting or overexpressing several candidate genes individually, we did not find an individual gene responsible for the transient cell cycle arrest in *MAT $\alpha$* -playing-*a* cells.

### Robustness of mating

Characterizing the mating defects of the transvestite strains allowed us to improve our understanding of the pheromone response of *MAT $\alpha$*  cells and study the robustness of mating efficiency to changes in gene expression. We investigated changing the expression levels of three proteins: *Bar1*,  $\alpha$ -factor, and *a*-factor.

The  $\alpha$ -factor protease *Bar1* helps *MATa* cells to detect an  $\alpha$ -factor gradient and choose a mating partner (Sprague and Herskowitz 1981; Jackson and Hartwell 1990; Barkai *et al.* 1998). Reducing *Bar1* expression by using an engineered *FUS1* promoter (Ingolia and Murray 2007) reduces the concentration of  $\alpha$ -factor required to get 50% of the cells to shmoo fourfold. This change appears to have little effect on mating: expressing *BAR1* under *P<sub>FUS1</sub>\** in *MATa* cells leaves mating unimpaired, suggesting that mating efficiency is robust to substantial changes in *Bar1* expression.

Mating efficiency is not robust to reductions in  $\alpha$ -factor secretion, a result that might have been predicted from work that showed that cells make graded responses to increasing levels of pheromone stimulation (Moore 1983; Takahashi and Pryciak 2008). Previous studies have shown that agglutination, shmoo formation, and pheromone-induced transcription increase with increasing  $\alpha$ -factor concentration (Moore 1983; Takahashi and Pryciak 2008). Decreased  $\alpha$ -factor production leads to a fusion defect and thus to a decrease in mating efficiency (Brizzio *et al.* 1996). We show that an  $\sim$ 10-fold reduction in  $\alpha$ -factor production in otherwise wild-type cells, such as *MAT $\alpha$*  *mf $\alpha$ 1 $\Delta$*  cells, results in a 2-fold reduction in mating efficiency when mated to a wild-type partner. Mating *MAT $\alpha$*  *mf $\alpha$ 1 $\Delta$*  cells to a compromised partner, such as the *MAT $\alpha$* -playing-*a* cells, results in a synergistic reduction in mating efficiency. Although reduced levels of *a*-factor production have also been shown to cause a cell fusion defect and a decrease in mating effi-

ciency (Brizzio *et al.* 1996), the precise regulation of *a*-factor production does not appear to be as important to mating efficiency as precise regulation of  $\alpha$ -factor production. *MATa* *mf $\alpha$ 1 $\Delta$*  cells have a mating efficiency that is indistinguishable from that of wild type (Michaelis and Herskowitz 1988), and reducing the *a*-factor production of *MATa* cells by overexpressing *AFB1* actually causes a slight increase in mating efficiency, indicating that the ideal quantity of *a*-factor production may be less than the amount of *a*-factor produced by wild-type *MATa* cells but greater than the amount of *a*-factor produced by *MAT $\alpha$* -playing-*a* cells.

Taken together, these results argue for a molecular arms race in pheromone production. Cells prefer the partner that makes the most pheromone (Jackson and Hartwell 1990), possibly because this is the only indicator of fitness available to a potential mating partner. We speculate that both *MATa* and *MAT $\alpha$*  cells have evolved to produce higher and higher concentrations of pheromone, resulting in the need for proteins such as *Bar1* and *Afb1* to improve gradient detection in dense mating mixtures. Once such functions have evolved, they imply that mutations that reduce pheromone production back to ancestral levels will decrease mating efficiency because the pheromone antagonists overwhelm the lower pheromone levels.

### Acknowledgments

We thank V. Denic, R. Gaudet, C. Hunter, and members of the Murray laboratory for reading and commenting on the manuscript. We also thank J. Gonçalves-Sá for thoughtful discussion and M. Piel for strains. This work was supported by National Institute of General Medical Sciences grant P50GM068763 of the National Centers for Systems Biology Program and a National Science Foundation Graduate Research Fellowship and Ashford Fellowship to L.B.H.

### Literature Cited

- Achstetter, T., 1989 Regulation of  $\alpha$ -factor production in *Saccharomyces cerevisiae*: *a*-factor pheromone-induced expression of the MF $\alpha$ 1 and STE13 genes. *Mol. Cell. Biol.* 9: 4507–4514.
- Alby, K., D. Schaefer, and R. J. Bennett, 2009 Homothallic and heterothallic mating in the opportunistic pathogen *Candida albicans*. *Nature* 460: 890–893.
- Apanovitch, D. M., K. C. Slep, P. B. Sigler, and H. G. Dohlman, 1998 Sst2 is a GTPase-activating protein for Gpa1: purification and characterization of a cognate RGS-G $\alpha$  protein pair in yeast. *Biochemistry (N. Y.)* 37: 4815–4822.
- Bardwell, L., 2005 A walk-through of the yeast mating pheromone response pathway. *Peptides* 26: 339–350.
- Barkai, N., M. D. Rose, and N. S. Wingreen, 1998 Protease helps yeast find mating partners. *Nature* 396: 422–423.
- Bender, A., and G. Sprague, 1989 Pheromones and pheromone receptors are the primary determinants of mating specificity in the yeast *Saccharomyces cerevisiae*. *Genetics* 121: 463–476.
- Betz, R., and W. Duntze, 1979 Purification and partial characterization of *a* factor, a mating hormone produced by mating-type-*a* cells from *Saccharomyces cerevisiae*. *Eur. J. Biochem.* 95: 469–475.

- Blumer, K. J., J. E. Reneke, and J. Thorner, 1988 The STE2 gene product is the ligand-binding component of the  $\alpha$ -factor receptor of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 263: 10836–10842.
- Bölker, M., M. Urban, and R. Kahmann, 1992 The *a* mating type locus of *U. maydis* specifies cell signaling components. *Cell* 68: 441–450.
- Brar, G. A., M. Yassour, N. Friedman, A. Regev, N. T. Ingolia *et al.*, 2012 High-resolution view of the yeast meiotic program revealed by ribosome profiling. *Science* 335: 552–557.
- Brizzio, V., A. E. Gammie, G. Nijbroek, S. Michaelis, and M. D. Rose, 1996 Cell fusion during yeast mating requires high levels of *a*-factor mating pheromone. *J. Cell Biol.* 135: 1727–1739.
- Butty, A. C., P. M. Pryciak, L. S. Huang, I. Herskowitz, and M. Peter, 1998 The role of Far1p in linking the heterotrimeric G protein to polarity establishment proteins during yeast mating. *Science* 282: 1511–1516.
- Cappellaro, C., K. Hauser, V. Mrša, M. Watzele, G. Watzele *et al.*, 1991 *Saccharomyces cerevisiae* *a*- and  $\alpha$ -agglutinin: characterization of their molecular interaction. *EMBO J.* 10: 4081–4088.
- Chan, R. K., and C. A. Otte, 1982a Physiological characterization of *Saccharomyces cerevisiae* mutants supersensitive to G1 arrest by *a* factor and  $\alpha$  factor pheromones. *Mol. Cell. Biol.* 2: 21–29.
- Chan, R. K., and C. A. Otte, 1982b Isolation and genetic analysis of *Saccharomyces cerevisiae* mutants supersensitive to G1 arrest by *a* factor and  $\alpha$  factor pheromones. *Mol. Cell. Biol.* 2: 11–20.
- Chang, F., and I. Herskowitz, 1990 Identification of a gene necessary for cell cycle arrest by a negative growth factor of yeast: FAR1 is an inhibitor of a G1 cyclin, CLN2. *Cell* 63: 999–1011.
- Chen, P., S. K. Sapperstein, J. D. Choi, and S. Michaelis, 1997 Biogenesis of the *Saccharomyces cerevisiae* mating pheromone *a*-factor. *J. Cell Biol.* 136: 251–269.
- Collart, M. A., and S. Oliviero, 2001 Preparation of yeast RNA. *Curr. Protoc. Mol. Biol.* 23: 13.12.1–13.12.5.
- de Nobel, H., J. Pike, P. N. Lipke, and J. Kurjan, 1995 Genetics of *a*-agglutinin function in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* 247: 409–415.
- Dietrich, F. S., S. Voegeli, S. Brachat, A. Lerch, K. Gates *et al.*, 2004 The *Ashbya gossypii* genome as a tool for mapping the ancient *Saccharomyces cerevisiae* genome. *Science* 304: 304–307.
- Dohlman, H. G., and J. W. Thorner, 2001 Regulation of G protein-initiated signal transduction in yeast: paradigms and principles. *Annu. Rev. Biochem.* 70: 703–754.
- Dohlman, H. G., J. Song, D. Ma, W. E. Courchesne, and J. Thorner, 1996 Sst2, a negative regulator of pheromone signaling in the yeast *Saccharomyces cerevisiae*: expression, localization, and genetic interaction and physical association with Gpa1 (the G-protein  $\alpha$  subunit). *Mol. Cell. Biol.* 16: 5194–5209.
- Dujon, B., D. Sherman, G. Fischer, P. Durrens, S. Casaregola *et al.*, 2004 Genome evolution in yeasts. *Nature* 430: 35–44.
- Edgar, R., M. Domrachev, and A. E. Lash, 2002 Gene expression omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* 30: 207–210.
- Galgoczy, D. J., A. Cassidy-Stone, M. Llinás, S. M. O'Rourke, I. Herskowitz *et al.*, 2004 Genomic dissection of the cell-type-specification circuit in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 101: 18069–18074.
- Giaever, G., A. M. Chu, L. Ni, C. Connelly, L. Riles *et al.*, 2002 Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* 418: 387–391.
- Gonçalves-Sá, J., and A. Murray, 2011 Asymmetry in sexual pheromones is not required for ascomycete mating. *Curr. Biol.* 21: 1337–1346.
- Haber, J. E., 1998 Mating-type gene switching in *Saccharomyces cerevisiae*. *Annu. Rev. Genet.* 32: 561–599.
- Hagen, D. C., G. McCaffrey, and G. F. Sprague, Jr., 1986 Evidence the yeast STE3 gene encodes a receptor for the peptide pheromone *a* factor: gene sequence and implications for the structure of the presumed receptor. *Proc. Natl. Acad. Sci. USA* 83: 1418–1422.
- Hamada, K., H. Terashima, M. Arisawa, N. Yabuki, and K. Kitada, 1999 Amino acid residues in the  $\omega$ -minus region participate in cellular localization of yeast glycosylphosphatidylinositol-attached proteins. *J. Bacteriol.* 181: 3886–3889.
- Herskowitz, I., 1988 Life cycle of the budding yeast *Saccharomyces cerevisiae*. *Microbiol. Rev.* 52: 536–553.
- Ingolia, N. T., and A. W. Murray, 2007 Positive-feedback loops as a flexible biological module. *Curr. Biol.* 17: 668–677.
- Jackson, C. L., and L. H. Hartwell, 1990 Courtship in *S. cerevisiae*: choose mating partners by responding to the strongest pheromone signal. *Cell* 63: 1039–1051.
- Jin, M., B. Errede, M. Behar, W. Mather, S. Nayak *et al.*, 2011 Yeast dynamically modify their environment to achieve better mating efficiency. *Sci. Signal.* 4: ra54.
- Julius, D., L. Blair, A. Brake, G. Sprague, and J. Thorner, 1983 Yeast  $\alpha$  factor is processed from a larger precursor polypeptide: the essential role of a membrane-bound dipeptidyl aminopeptidase. *Cell* 32: 839–852.
- Kellis, M., B. W. Birren, and E. S. Lander, 2004 Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature* 428: 617–624.
- Kim, J., E. Bortz, H. Zhong, T. Leeuw, E. Leberer *et al.*, 2000 Localization and signaling of G $\beta$  subunit Ste4p are controlled by *a*-factor receptor and the *a*-specific protein Apg7p. *Sci. Signal.* 20: 8826–8835.
- Krebs, E. G., 1953 Yeast glyceraldehyde-3-phosphate dehydrogenase I. Electrophoresis of fractions precipitated by nucleic acid. *J. Biol. Chem.* 200: 471–478.
- Kurjan, J., 1985  $\alpha$ -Factor structural gene mutations in *Saccharomyces cerevisiae*: effects on  $\alpha$ -factor production and mating. *Mol. Cell. Biol.* 5: 787–796.
- Kurjan, J., and I. Herskowitz, 1982 Structure of a yeast pheromone gene (MF $\alpha$ ): a putative  $\alpha$ -factor precursor contains four tandem copies of mature  $\alpha$ -factor. *Cell* 30: 933–943.
- Lipke, P., D. Wojciechowicz, and J. Kurjan, 1989 AG  $\alpha$ 1 is the structural gene for the *Saccharomyces cerevisiae*  $\alpha$ -agglutinin, a cell surface glycoprotein involved in cell-cell interactions during mating. *Mol. Cell. Biol.* 9: 3155–3165.
- MacKay, V. L., S. Welch, M. Y. Insley, T. R. Manney, J. Holly *et al.*, 1988 The *Saccharomyces cerevisiae* BAR1 gene encodes an exported protein with homology to pepsin. *Proc. Natl. Acad. Sci. USA* 85: 55–59.
- Marcus, S., C. Xue, F. Naider, and J. Becker, 1991 Degradation of *a*-factor by a *Saccharomyces cerevisiae*  $\alpha$ -mating-type-specific endopeptidase: evidence for a role in recovery of cells from G1 arrest. *Mol. Cell. Biol.* 11: 1030–1039.
- McAlister, L., and M. J. Holland, 1985 Differential expression of the three yeast glyceraldehyde-3-phosphate dehydrogenase genes. *J. Biol. Chem.* 260: 15019–15027.
- McGrath, J. P., and A. Varshavsky, 1989 The yeast STE6 gene encodes a homologue of the mammalian multidrug resistance P-glycoprotein. *Nature* 340: 400–404.
- Michaelis, S., and I. Herskowitz, 1988 The *a*-factor pheromone of *Saccharomyces cerevisiae* is essential for mating. *Mol. Cell. Biol.* 8: 1309–1318.
- Moore, S. A., 1983 Comparison of dose-response curves for  $\alpha$  factor-induced cell division arrest, agglutination, and projection formation of yeast cells. Implication for the mechanism of  $\alpha$  factor action. *J. Biol. Chem.* 258: 13849–13856.
- Peter, M., and I. Herskowitz, 1994 Direct inhibition of the yeast cyclin-dependent kinase Cdc28-Cln by Far1. *Science* 265: 1228–1231.

- Rappaport, N., and N. Barkai, 2012 Disentangling signaling gradients generated by equivalent sources. *J. Biol. Phys.* 38: 267–278.
- Reid, B. J., and L. H. Hartwell, 1977 Regulation of mating in the cell cycle of *Saccharomyces cerevisiae*. *J. Cell Biol.* 75: 355–365.
- Roberts, C. J., B. Nelson, M. J. Marton, R. Stoughton, M. R. Meyer *et al.*, 2000 Signaling and circuitry of multiple MAPK pathways revealed by a matrix of global gene expression profiles. *Sci. Signal.* 287: 873–880.
- Roth, A. F., B. Nelson, C. Boone, and N. G. Davis, 2000 *Asg7p-Ste3p* inhibition of pheromone signaling: regulation of the zygotic transition to vegetative growth. *Sci. Signal.* 20: 8815–8825.
- Roy, A., C. Lu, D. Marykwas, P. Lipke, and J. Kurjan, 1991 The AGA1 product is involved in cell surface attachment of the *Saccharomyces cerevisiae* cell adhesion glycoprotein *a*-agglutinin. *Mol. Cell. Biol.* 11: 4196–4206.
- Schaefer, D., P. Côte, M. Whiteway, and R. J. Bennett, 2007 Barrier activity in *Candida albicans* mediates pheromone degradation and promotes mating. *Eukaryot. Cell* 6: 907–918.
- Segall, J. E., 1993 Polarization of yeast cells in spatial gradients of  $\alpha$  mating factor. *Proc. Natl. Acad. Sci. USA* 90: 8332–8336.
- Sherman, F., G. Fink, and C. Lawrence, 1974 *Methods in Yeast Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Singh, A., E. Y. Chen, J. M. Lugovoy, C. N. Chang, R. A. Hitzeman *et al.*, 1983 *Saccharomyces cerevisiae* contains two discrete genes coding for the  $\alpha$ -factor pheromone. *Nucleic Acids Res.* 11: 4049–4063.
- Sprague, G. F. Jr., 1991 Assay of yeast mating reaction. *Methods Enzymol.* 194: 77–93.
- Sprague, G. F. Jr., and I. Herskowitz, 1981 Control of yeast cell type by the mating type locus. I. Identification and control of expression of the *a*-specific gene *BAR1*. *J. Mol. Biol.* 153: 305–321.
- Stanton, B. C., S. S. Giles, M. W. Staudt, E. K. Kruzel, and C. M. Hull, 2010 Allelic exchange of pheromones and their receptors reprograms sexual identity in *Cryptococcus neoformans*. *PLoS Genet.* 6: e1000860.
- Steden, M., R. Betz, and W. Duntze, 1989 Isolation and characterization of *Saccharomyces cerevisiae* mutants supersensitive to G1 arrest by the mating hormone *a*-factor. *Mol. Gen. Genet.* 219: 439–444.
- Strathern, J., J. Hicks, and I. Herskowitz, 1981 Control of cell type in yeast by the mating type locus: the  $\alpha$ 1- $\alpha$ 2 hypothesis. *J. Mol. Biol.* 147: 357–372.
- Takahashi, S., and P. M. Pryciak, 2008 Membrane localization of scaffold proteins promotes graded signaling in the yeast MAP kinase cascade. *Curr. Biol.* 18: 1184–1191.
- Trapnell, C., L. Pachter, and S. L. Salzberg, 2009 TopHat: discovering splice junctions with RNA-seq. *Bioinformatics* 25: 1105–1111.
- Trapnell, C., B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan *et al.*, 2010 Transcript assembly and quantification by RNA-seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* 28: 511–515.
- Trueheart, J., J. D. Boeke, and G. R. Fink, 1987 Two genes required for cell fusion during yeast conjugation: evidence for a pheromone-induced surface protein. *Mol. Cell. Biol.* 7: 2316–2328.
- Tyers, M., and B. Futcher, 1993 *Far1* and *Fus3* link the mating pheromone signal transduction pathway to three G1-phase *Cdc28* kinase complexes. *Mol. Cell. Biol.* 13: 5659–5669.
- Wolfe, K. H., and D. C. Shields, 1997 Molecular evidence for an ancient duplication of the entire yeast genome. *Nature* 387: 708–712.

Communicating editor: M. D. Rose

# GENETICS

**Supporting Information**

<http://www.genetic.org/lookup/suppl/doi:10.1534/genetics.113.155846/-/DC1>

## **Genetically Engineered Transvestites Reveal Novel Mating Genes in Budding Yeast**

Lori B. Huberman and Andrew W. Murray

### File S1

To measure unstimulated  $\alpha$ -factor production, cells were grown to log phase ( $\sim 5 \times 10^6$  cells/mL), washed into YPD + 0.1% BSA, and resuspended in BSA-coated culture tubes at  $5 \times 10^6$  cells/mL in YPD + 0.1% BSA. The supernatant of the media was then harvested after a 15 minute, 30 minute, or 120 minute incubation at 30°C on a roller drum using BSA-coated 1mL syringes (BD, Franklin Lakes, NJ) and filtered through 0.45 $\mu$ m BSA-coated syringe filters (Pall Corporation, Port Washington, NY).  $5 \times 10^5$  log phase *MATa bar1 $\Delta$*  cells were then incubated in 1x, 0.5x, and 0.2x dilutions of the supernatant for 2 hours, sonicated, fixed using 60% ethanol at -20°C, and resuspended into 20% glycerol in phosphate buffered saline (PBS). The percent of the cells shmooing was counted and compared to a calibration curve produced by treating *MATa bar1 $\Delta$*  cells with different concentrations of synthetic  $\alpha$ -factor.

Stimulated  $\alpha$ -factor production was measured in the same way except that the cells being tested were grown to log phase ( $\sim 5 \times 10^6$  cells/mL), counted, washed into YPD + 0.1% BSA, and  $5 \times 10^6$  cells/mL of the cells being tested were mixed with  $5 \times 10^5$  cells/mL *MATa bar1 $\Delta$*  cells. After 2 hours incubation at 30°C on roller drums, the cells were washed into fresh YPD + 0.1% BSA and put into BSA-coated culture tubes. The supernatant was then harvested and analyzed as described above.

To generate a standard curve, *MATa bar1 $\Delta$*  cells were incubated in integer values of synthetic  $\alpha$ -factor concentrations between 0nM and 7nM, and the best fit line was generated with an  $R^2$  of 0.9 to determine a constant (K) for the relationship between  $\alpha$ -factor concentration and the percentage of shmooos ( $K=0.07$ ).  $\alpha$ -factor production in molecules/cell/minute was then determined by  $\frac{N \times K \times \% \text{ shmooos} \times \text{volume}}{r \times 2^D \times t \times l}$ , where  $N$  is Avagadro's constant,  $r$  is the starting number of cells,  $D$  is the number of doublings expected,  $t$  is the incubation time, and  $l$  is the dilution. The expected number of doublings was calculated by the incubation time divided by an expected 90 minute doubling time for yeast.

Culture tubes and syringes were BSA-coated by incubating overnight at 4°C with PBS + 2% BSA. The PBS + 2% BSA was poured out immediately prior to use of the culture tube or syringe. Filters were BSA-coated by filtering 1mL of PBS + 2% BSA through them prior to use. At least 200 cells were counted to determine the percentage of cells shmooing. Error bars are standard deviations. Statistical significance was determined using Student's  $t$ -test.



## File S2

Pictures of the *MATa/a* and *MATa/α* diploids were acquired by growing cells to log phase ( $\sim 5 \times 10^6$  cells/mL) at 30°C in YPD. Cells were then loaded into a microfluidics chamber (CellAsic, Hayward, CA) (Lee *et al.* 2008), which was pretreated by perfusing YPD through the chamber at 34kPa for 10 minutes. Once cells were loaded, YPD was perfused through the microfluidics chamber at 14kPa. Pictures were taken using a 20x Plan Apo VC 0.75NC air lens after 10 hours growth using differential interference contrast with a 10ms exposure.

Pictures and movies of various strains responding to 10nM  $\alpha$ -factor were produced in a similar way. Cells were grown to log phase ( $\sim 5 \times 10^6$  cells/mL) at 30°C in YPD, washed into SC + 0.1% BSA, and loaded into a microfluidics chamber, which had been pretreated by perfusing PBS + 2% BSA through the chamber at 34kPa for 10 minutes and then SC + 0.1% BSA through the chamber at 34kPa for 10 minutes. Once the cells were loaded, SC + 0.1% BSA + 10nM  $\alpha$ -factor was perfused through the chamber at 14kPa. Pictures were taken every 10 minutes for 8 hours using a 60x Plan Apo VC 1.4NA oil lens with differential interference contrast with a 10ms exposure.

For pictures of the *MATa*-playing- $\alpha$  cells responding to  $\alpha$ -factor, *MATa P<sub>ACT1</sub>-mCherry* cells were mixed with *MATa*-playing- $\alpha$  cells on a filter as described for the quantitative mating assay. Cells were washed off the filters into SC after 2.5 hours and placed directly onto glass slides (Corning, Tewksbury, MA). Pictures were taken immediately using a 20x Plan Apo VC 0.75NC air lens with a 10ms exposure for differential interference contrast and a 300ms exposure for fluorescent images.

## LITERATURE CITED

Lee, P. J., N. C. Helman, W. A. Lim and P. J. Hung, 2008 A microfluidic system for dynamic yeast cell imaging. *BioTechniques* **44**: 91-95.

### File S3

#### ***MATa* cells are capable of enduring arrest in response to pheromone stimulation.**

*MATa bar1Δ* cells were grown in a microfluidics chamber for 8 hours perfused with SC plus 10nM  $\alpha$ -factor and imaged using DIC every 10 minutes with 60x magnification.

File S3 is available for download as an avi file at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.155846/-/DC1>.

### File S4

#### ***MAT $\alpha$ -playing-a* cells arrest only transiently in response to pheromone stimulation.**

*MAT $\alpha$ -playing-a bar1Δ* cells were grown in a microfluidics chamber for 8 hours perfused with SC plus 10nM  $\alpha$ -factor and imaged using DIC every 10 minutes with 60x magnification.

File S4 is available for download as an avi file at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.155846/-/DC1>.

**Table S1 Strains used in this study**

<b>Strain Name</b>	<b>Genotype (all cells are in the W303 background)</b>
LBHY29	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY41	<i>MATα ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY44	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 bar1Δ::HphMX4 MFA1-HIS3::mfa1Δ MFA2-TRP1::mfa2Δ STE3-NatMX4::ste2Δ asg7Δ::URA3(Kluyveromyces lactis) ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY47	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 P<sub>MFA1</sub>-STE6-URA3 MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ P<sub>FUS1</sub>*-BAR1-KanMX6 ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY49	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 P<sub>MFA1</sub>-STE6-ura3<sup>+</sup> MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ P<sub>FUS1</sub>*-BAR1-KanMX6 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY89	<i>MATα mfa1Δ::NatMX4 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
LBHY92	<i>MATα mfa1Δ::NatMX4 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
LBHY93	<i>MATα mfa1Δ::NatMX4 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY98	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 bar1Δ::HphMX4 MFA1-HIS3::mfa1Δ MFA2-TRP1::mfa2Δ STE3-NatMX4::ste2Δ asg7Δ::URA3(K. lactis) P<sub>ACT1</sub>-yCerulean @ ADE2 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY108	<i>MATα P<sub>ACT1</sub>-mCherry-HIS3MX6 @ P<sub>ACT1</sub> ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
LBHY156	<i>MATα mfa1Δ::KanMX6 mfa2Δ::HphMX4 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
LBHY177	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ bar1Δ::KanMX6 ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY286	<i>MATα mfa1Δ::KanMX6 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY290	<i>MATα P<sub>FUS1</sub>*-BAR1-KanMX6 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY316	<i>MATα/α P<sub>FUS1</sub>-YFP @ LEU2/leu2-112 P<sub>MFA1</sub>-STE6-URA3/P<sub>STE6</sub>-STE6 MFA1-TRP1::mfa1Δ/MFA1 MFA2-HIS3::mfa2Δ/MFA2 STE2-NatMX4::ste3Δ/STE3 P<sub>FUS1</sub>*-BAR1-KanMX6/ bar1Δ::ADE2 SPA2/ SPA2-YFP:HIS3 ade2-1/ade2-1 can1-100 /can1-100 his3-11,15/his3-11,15 trp1-1/trp1-1 ura3-1/ura3-1</i>
LBHY318	<i>MATα/a P<sub>FUS1</sub>-YFP @ LEU2/ leu2-112 bar1Δ::HphMX4/BAR1 MFA1-HIS3::mfa1Δ/MFA1 MFA2-TRP1::mfa2Δ/MFA2 STE3-NatMX4::ste2Δ/STE2 asg7Δ::URA3(K. lactis)/ASG7 SPA2-CFP:KanMX6/SPA2 ADE2/ade2-1 can1-100/can1-100 his3-11,15/P<sub>FUS1</sub>-YFP @ HIS3 trp1-1/trp1-1 ura3-1/ura3-1</i>
LBHY346	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 P<sub>MFA1</sub>-STE6-URA3 MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ P<sub>FUS1</sub>*-BAR1-KanMX6 afb1Δ::HphMX4 ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY350	<i>MATα afb1Δ::HphMX4 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY352	<i>MATα afb1Δ::HphMX4 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY395	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 bar1Δ::HphMX4 MFA1-HIS3::mfa1Δ MFA2-TRP1::mfa2Δ STE3-NatMX4::ste2Δ asg7Δ::URA3(K. lactis) P<sub>TDH3</sub>-MFA1:KanMX6 @ P<sub>TDH3</sub> ade2-1 can1-100 his3-11,15 trp1-1 ura3-1</i>
LBHY397	<i>MATα P<sub>ACT1</sub>-AFB1:KanMX6 @ P<sub>ACT1</sub> ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
LBHY409	<i>MATα P<sub>ACT1</sub>-AFB1:KanMX6 @ P<sub>ACT1</sub> ade2-1 can1-100 his3-11,15 leu2-112 trp1-1</i>
LBHY410	<i>MATα P<sub>FUS1</sub>-YFP @ LEU2 P<sub>MFA1</sub>-STE6-ura3<sup>+</sup> MFA1-TRP1::mfa1Δ MFA2-HIS3::mfa2Δ STE2-NatMX4::ste3Δ P<sub>FUS1</sub>*-BAR1-KanMX6 afb1Δ::HphMX4 can1-100 his3-11,15 trp1-1 ura3-1</i>

MP 381	<i>MAT<math>\alpha</math> bar1<math>\Delta</math>::ADE2 SPA2-YFP:HIS3 ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1</i>
MP 384	<i>MAT<math>\alpha</math> bar1<math>\Delta</math>::ADE2 SPA2-YFP:HIS3 ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1</i>
MP 420	<i>MAT<math>\alpha</math> SPA2-CFP:KanMX6 P<sub>FUS1</sub>-YFP @ HIS3 can1-100 leu2-112 trp1-1 ura3-1</i>
W303	<i>MAT<math>\alpha</math> ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1 (W303 wildtype)</i>
W303	<i>MAT<math>\alpha</math> ade2-1 can1-100 his3-11,15 leu2-112 trp1-1 ura3-1 (W303 wildtype)</i>

---

All strains are from this study except for MP 381, MP 384, and MP 420, which are from M. Piel and the W303 wildtype strains, which are from Thomas and Rothstein 1989 (Thomas and Rothstein 1989).

#### LITERATURE CITED

Thomas, B. J., and R. Rothstein, 1989 Elevated recombination rates in transcriptionally active DNA. *Cell* **56**: 619-630.

**Table S2 Genes identified as differentially expressed in MAT $\alpha$  and MAT $\alpha$ -playing-a cells using RNA sequencing**

Gene	Log <sub>2</sub> Fold Change:	
	Unstimulated MAT $\alpha$ -playing-a	Stimulated MAT $\alpha$ -playing-a
	÷	÷
	Unstimulated MAT $\alpha$	Stimulated MAT $\alpha$
<i>MAT<math>\alpha</math>1</i>	<b>7.56*</b>	<b>7.86*</b>
<i>YCR097W-A</i>	<b>3.97*</b>	<b>6.96*</b>
<i>YDR008C</i>	<b>4.30*</b>	<b>5.70*</b>
<i>SAG1</i>	<b>3.63*</b>	<b>5.49*</b>
<i>YCL065W</i>	<b>2.91*</b>	<b>5.33*</b>
<i>TRP1</i>	<b>4.35*</b>	<b>5.31*</b>
<i>MAT<math>\alpha</math>2</i>	<b>1.28*</b>	<b>3.08*</b>
<i>YLR040C</i>	<b>3.41*</b>	<b>2.38*</b>
<i>YLR041W</i>	<b>3.21*</b>	<b>2.03*</b>
<i>YML131W</i>	0.66*	0.69*
<i>TPO2</i>	-0.74*	0.53*
<i>YPR158C-D</i>	-0.61*	-0.48*
<i>ADE17</i>	-0.75*	-0.48*
<i>BAP2</i>	-0.94*	-0.54*
<i>FSF1</i>	-0.64*	-0.58*
<i>ILV3</i>	-0.75*	-0.58*
<i>GDH1</i>	-0.87*	-0.60*
<i>FIT3</i>	-0.84*	-0.63*
<i>ALD5</i>	<b>-1.03*</b>	-0.88*
<i>BAT1</i>	<b>-1.70*</b>	<b>-1.20*</b>
<i>OAC1</i>	<b>-2.27*</b>	<b>-1.57*</b>
<i>LEU1</i>	<b>-2.47*</b>	<b>-1.82*</b>
<i>ADE2</i>	<b>-1.38*</b>	<b>-1.93*</b>
<i>GYP8</i>	<b>-1.02*</b>	<b>-2.05*</b>
<i>STE2</i>	<b>-2.67*</b>	<b>-2.18*</b>
<i>MFA1</i>	<b>-1.23*</b>	<b>-2.80*</b>
<i>YNL146C-A</i>	<b>-2.57*</b>	<b>-4.00*</b>
<i>MFA2</i>	<b>-5.51*</b>	<b>-6.85*</b>
<i>ASG7</i>	<b>-2.68*</b>	<b>-9.11*</b>
<i>AGA2</i>	<b>-7.29*</b>	<b>-9.28*</b>
<b>Genes whose expression changed significantly only in pheromone-stimulated cells</b>		
<i>YHR145C</i>	-0.61	<b>3.63*</b>

<i>YCR041W</i>	<b>2.14</b>	<b>3.53*</b>
<i>PCL1</i>	-0.12	<b>1.85*</b>
<i>SNL1</i>	0.05	<b>1.75*</b>
<i>YLR346C</i>	<b>1.20</b>	<b>1.70*</b>
<i>TOS6</i>	-0.03	<b>1.30*</b>
<i>INO1</i>	0.16	<b>1.30*</b>
<i>PST1</i>	-0.06	<b>1.16*</b>
<i>OYE3</i>	-0.22	<b>1.05*</b>
<i>TOS4</i>	-0.04	<b>1.03*</b>
<i>YOX1</i>	-0.12	0.97*
<i>IMD2</i>	0.02	0.95*
<i>PIR3</i>	0.47	0.95*
<i>LEU2</i>	0.21	0.91*
<i>SPO11</i>	0.05	0.88*
<i>MCD1</i>	0.07	0.87*
<i>HES1</i>	0.29	0.85*
<i>GRE2</i>	0.13	0.80*
<i>AAD16</i>	0.00	0.80*
<i>YJL218W</i>	0.41	0.79*
<i>YGL230C</i>	0.05	0.78*
<i>AAD6</i>	0.03	0.77*
<i>EGT2</i>	0.12	0.76*
<i>YJR154W</i>	0.19	0.74*
<i>MCH2</i>	0.35	0.73*
<i>HTA2</i>	-0.17	0.73*
<i>YDR134C</i>	0.06	0.69*
<i>PIR1</i>	-0.11	0.69*
<i>KDX1</i>	0.52	0.68*
<i>HSP31</i>	0.22	0.68*
<i>YGR035C</i>	0.17	0.66*
<i>APL1</i>	0.07	0.66*
<i>YNR064C</i>	0.02	0.64*
<i>YBR071W</i>	-0.04	0.63*
<i>HAL1</i>	0.48	0.62*
<i>YOL159C</i>	0.04	0.62*
<i>CRG1</i>	0.22	0.62*
<i>YNL134C</i>	-0.04	0.60*
<i>AAD10</i>	-0.15	0.60*
<i>SVS1</i>	-0.09	0.59*

<i>SPI1</i>	0.44	0.58*
<i>YDR034W-B</i>	-0.51	0.57*
<i>RNR1</i>	0.07	0.56*
<i>SRL3</i>	0.08	0.56*
<i>YNL058C</i>	0.21	0.55*
<i>YKE4</i>	-0.02	0.54*
<i>ACA1</i>	0.28	0.52*
<i>SSA4</i>	-0.10	0.50*
<i>HHF1</i>	-0.09	0.50*
<i>DSE4</i>	0.29	0.48*
<i>OAZ1</i>	0.13	0.47*
<i>SEO1</i>	0.24	0.47*
<i>NIS1</i>	-0.11	0.47*
<i>YEH1</i>	0.01	0.46*
<i>HXT1</i>	-0.08	-0.47*
<i>PHO12</i>	-0.24	-0.50*
<i>YOL103W-B</i>	-0.31	-0.50*
<i>SIT1</i>	-0.46	-0.51*
<i>PHO11</i>	-0.24	-0.52*
<i>ENA1</i>	0.00	-0.52*
<i>LEU9</i>	-0.22	-0.52*
<i>HIS4</i>	-0.50	-0.52*
<i>YPL257W-B</i>	-0.31	-0.53*
<i>ALT2</i>	-0.20	-0.54*
<i>PDH1</i>	0.08	-0.56*
<i>PRY2</i>	-0.29	-0.57*
<i>HXK1</i>	-0.03	-0.57*
<i>TAT1</i>	-0.35	-0.57*
<i>YJR003C</i>	-0.15	-0.66*
<i>THI22</i>	-0.26	-0.81*
<i>FET3</i>	-0.32	-0.83*
<i>AXL1</i>	-0.52	-0.93*

---

**Genes whose expression changed significantly only in unstimulated cells**

---

<i>RDN5-2</i>	<b>2.90*</b>	<b>1.33</b>
<i>RDN5-5</i>	<b>2.75*</b>	0.73
<i>RDN5-4</i>	<b>2.66*</b>	<b>1.02</b>
<i>RDN5-3</i>	<b>2.51*</b>	0.93
<i>RDN5-1</i>	<b>2.34*</b>	0.99
<i>RDN5-6</i>	<b>2.34*</b>	0.99

<i>YLR042C</i>	0.66*	0.31
<i>HER1</i>	0.64*	0.15
<i>HAP4</i>	0.55*	0.23
<i>YJL171C</i>	-0.59*	-0.20
<i>DRE2</i>	-0.59*	-0.19
<i>HMX1</i>	-0.64*	-0.32
<i>GAP1</i>	-0.64*	-0.25
<i>ARN2</i>	-0.65*	-0.41
<i>FUS1</i>	-0.65*	-0.06
<i>MAE1</i>	-0.70*	-0.15
<i>AGA1</i>	-0.85*	-0.15
<i>FIT2</i>	-0.89*	-0.40
<i>ISU2</i>	-0.92*	-0.28
<i>YGP1</i>	<b>-1.04*</b>	-0.10
<i>DIC1</i>	<b>-1.26*</b>	-0.28

---

\* Indicates statistical significance determined by the default settings in Cufflinks ( $p < 0.001$ ) (Trapnell *et al.* 2010). Numbers in bold indicate that there is a greater than two-fold change between the expression of the gene in *MAT $\alpha$* -playing-a and *MATa* cells.

#### LITERATURE CITED

Trapnell, C., B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan *et al.*, 2010 Transcript assembly and quantification by RNA-seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**: 511-515.