



Regulation of reactive oxygen species-mediated abscisic acid signaling in guard cells and drought tolerance by glutathione

Shintaro Munemasa¹, Daichi Muroyama¹, Hiroki Nagahashi², Yoshimasa Nakamura¹, Izumi C. Mori³ and Yoshiyuki Murata^{1*}

¹ Division of Agricultural and Life Science, Graduate School of Environmental and Life Science, Okayama University, Okayama, Japan

² Faculty of Agriculture, Okayama University, Okayama, Japan

³ Institute of Plant Science and Resources, Okayama University, Kurashiki, Japan

Edited by:

Nobuyuki Uozumi, Tohoku University, Japan

Reviewed by:

Shan Lu, Nanjing University, China
Jin Chen, Michigan State University, USA

*Correspondence:

Yoshiyuki Murata, Division of Agricultural and Life Science, Graduate School of Environmental and Life Science, Okayama University, 1-1-1 Tsushima-Naka, Okayama 7008530, Japan
e-mail: muta@cc.okayama-u.ac.jp

The phytohormone abscisic acid (ABA) induces stomatal closure in response to drought stress, leading to reduction of transpirational water loss. A thiol tripeptide glutathione (GSH) is an important regulator of cellular redox homeostasis in plants. Although it has been shown that cellular redox state of guard cells controls ABA-mediated stomatal closure, roles of GSH in guard cell ABA signaling were largely unknown. Recently we demonstrated that GSH functions as a negative regulator of ABA signaling in guard cells. In this study we performed more detailed analyses to reveal how GSH regulates guard cell ABA signaling using the GSH-deficient Arabidopsis mutant *cad2-1*. The *cad2-1* mutant exhibited reduced water loss from rosette leaves. Whole-cell current recording using patch clamp technique revealed that the *cad2-1* mutation did not affect ABA regulation of S-type anion channels. We found enhanced activation of Ca²⁺ permeable channels by hydrogen peroxide (H₂O₂) in *cad2-1* guard cells. The *cad2-1* mutant showed enhanced H₂O₂-induced stomatal closure and significant increase of ROS accumulation in whole leaves in response to ABA. Our findings provide a new understanding of guard cell ABA signaling and a new strategy to improve plant drought tolerance.

Keywords: abscisic acid, glutathione, reactive oxygen species, guard cell, stomata

INTRODUCTION

A phytohormone abscisic acid (ABA) induces closing of stomatal pores on leaf epidermis, resulting in reduction of transpirational water loss. The central ABA signaling module is composed of ABA receptors PYR/PYL/RCAR, clade A type 2C protein phosphatases (PP2Cs), and subclass 2 of Snf1-related kinases (SnRK2s) and regulates downstream targets (Fujii et al., 2009; Ma et al., 2009; Park et al., 2009) including ion channels (Geiger et al., 2009; Lee et al., 2009).

Activation of slow type (S-type) anion channels is a key step for ABA signaling in guard cells and drives depolarization of plasma membrane of guard cells, which in turn evokes K⁺ extrusion (Linder and Raschke, 1992; Schroeder and Keller, 1992). A guard cell plasma membrane protein SLAC1 represents the S-type anion channel activity (Negi et al., 2008; Vahisalu et al., 2008). It has been demonstrated that ABA activation of S-type anion channels is mediated via a cytosolic Ca²⁺-dependent pathway (Marten et al., 2007; Siegel et al., 2009). ABA activates hyperpolarization-activated Ca²⁺-permeable cation (I_{Ca}) channels in the plasma membrane of guard cells (Schroeder and Hagiwara, 1990; Hamilton et al., 2000; Pei et al., 2000; Kwak et al., 2003) and induces elevation of cytosolic free Ca²⁺ concentration ([Ca²⁺]_{cyt}) in guard cells (McAinsh et al., 1990; Schroeder and Hagiwara, 1990; Gilroy et al., 1991; Allan et al., 1994; Grabov and Blatt, 1998; Allen et al., 1999; Marten et al., 2007). The [Ca²⁺]_{cyt}

signals are decoded by Ca²⁺ sensor proteins such as calcium dependent protein kinases (CDPKs). Electrophysiology experiments using *Xenopus* oocyte demonstrated that Arabidopsis CDPKs, CPK6, CPK21, and CPK23, directly phosphorylate and activate SLAC1 channel (Geiger et al., 2010; Brandt et al., 2012).

It has been suggested that guard cell ABA signaling involves redox regulation. Reactive oxygen species (ROS) including hydrogen peroxide (H₂O₂) serve as a key mediator of ABA activation of I_{Ca} channels (Pei et al., 2000; Murata et al., 2001; Kwak et al., 2003). Exogenous application of H₂O₂ activates I_{Ca} channels and evokes guard cell [Ca²⁺]_{cyt} increases (Pei et al., 2000). Plasma membrane NAD(P)H oxidases are responsible for ABA-induced ROS production in guard cells (Kwak et al., 2003). Arabidopsis glutathione peroxidase 3 (AtGPX3) was shown to function as both a ROS scavenger and a ROS signal transducer in ABA signaling (Miao et al., 2006). Emerging evidences suggest that ROS production by apoplastic enzymes such as cell-wall bound peroxidases is also involved in induction of stomatal closure (An et al., 2008; Khokon et al., 2011; Hossain et al., 2013).

Glutathione (GSH) is the most abundant non-protein thiol compound in plants and a key regulator of cellular redox homeostasis. GSH is involved in various physiological processes including growth, development, and defense response to biotic and abiotic stresses (May et al., 1998; Noctor and Foyer, 1998). Previously we reported that ABA as well as methyl jasmonate

(MeJA) decreases the GSH contents of guard cells (Akter et al., 2010; Okuma et al., 2011). Arabidopsis GSH-deficient mutants, *chl-1* and *cad2-1* exhibit enhanced ABA-induced and MeJA-induced stomatal closure and a membrane permeable derivative of GSH, GSH monoethyl ester (GSHmee) restored the stomatal phenotype of *chl-1* and *cad2-1* mutants (An et al., 2008; Akter et al., 2010, 2012, 2013; Okuma et al., 2011), demonstrating that GSH functions as a negative regulator of ABA and MeJA signaling in guard cells. However, the detailed mechanism of how GSH modulates the guard cell responses is still unclear.

In this study, we analyzed GSH regulation of ABA signaling in guard cells using the Arabidopsis GSH-deficient mutant *cad2-1*. The *cad2-1* mutant is deficient in the first GSH biosynthesis enzyme, γ -glutamylcysteine synthetase. We found that the *cad2-1* mutation causes enhanced ROS activation of I_{Ca} channel and ABA-induced ROS accumulation in apoplast. A new signal model for regulation of ROS-mediated ABA signaling by GSH in guard cells is also proposed.

MATERIALS AND METHODS

PLANT MATERIAL AND GROWTH

The Arabidopsis ecotype Columbia (Col) and *cad2-1* mutant plants were grown on soil mixture of 70% (v/v) vermiculite (Asahi-kogyo, Okayama, Japan) and 30% (v/v) Sakata Supermix-A (Sakata Seed Corporation, Yokohama, Japan) in growth chambers at 21°C under a 16-h-light/8-h-dark photoperiod with photon flux density of 80 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Four- to six-week-old plants were used in all experiments.

WATER LOSS ASSAY

Three detached rosette leaves were placed on a plastic tray and the loss in fresh weight was monitored at the indicated times.

STOMATAL APERTURE MEASUREMENTS

Stomatal aperture measurements were performed as described previously (Munemasa et al., 2007; Okuma et al., 2011). Detached rosette leaves were floated on stomatal assay buffer containing 5 mM KCl, 50 μM CaCl_2 , and 10 mM MES-Tris (pH 5.6) for 2 h in the light to induce stomatal opening, followed by the addition of H_2O_2 . After 2-h incubation in the light, the leaves were shredded and epidermal tissues were collected. At least 20 stomatal apertures were measured on each individual experiment.

ELECTROPHYSIOLOGY

Guard cell protoplasts (GCPs) were prepared from Arabidopsis rosette leaves by the enzymatic method, as described previously (Pei et al., 1997). Whole-cell currents were recorded using patch clamp technique, as described previously (Munemasa et al., 2007, 2011). For S-type anion current measurements, the pipette solution contained 150 mM CsCl, 2 mM MgCl_2 , 6.7 mM EGTA, 5.58 mM CaCl_2 (free Ca^{2+} concentration: 2 μM), and 10 mM HEPES-Tris (pH 7.1). The bath solution contained 30 mM CsCl, 2 mM MgCl_2 , 1 mM CaCl_2 , and 10 mM MES-Tris (pH 5.6). 5 mM Mg-ATP was freshly added to the pipette solution before experiments. For I_{Ca} current measurements, the pipette solution contained 10 mM BaCl_2 , 4 mM EGTA, and 10 mM HEPES-Tris

(pH 7.1). The bath solution contained 100 mM BaCl_2 , and 10 mM MES-Tris (pH 5.6). 0.1 mM DTT was added to both pipette and bath solutions freshly before experiments. In both cases, osmolarity was adjusted to 500 mmol/kg (pipette solutions) and 485 mmol/kg (bath solutions) with D-sorbitol.

DETECTION OF ROS ACCUMULATION IN WHOLE LEAVES

Accumulation of H_2O_2 in whole leaves was monitored using 3,3'-diaminobenzidine (DAB) according to Maruta et al. (2010) with slight modifications. Detached rosette leaves were vacuum infiltrated with DAB assay buffer containing 1 mg mL^{-1} DAB, 5 mM KCl, 50 μM CaCl_2 , and 10 mM MES-Tris (pH 5.6). Infiltrated leaves are incubated in DAB assay buffer with or without ABA for 4 h in the light. The leaves were then decolorized by boiling in ethanol. Apoplastic ROS were visualized as a reddish-brown color and quantified using Image-J software (National Institutes of Health, USA).

STATISTICAL ANALYSIS

All statistical significance was analyzed by double-tailed Student's *t*-test. We regarded differences at the level of $P < 0.05$ as significant.

RESULTS

THE *cad2-1* MUTANT SHOWED ENHANCED DROUGHT TOLERANCE

Previously we reported that GSH depletion by the *cad2-1* mutation enhances ABA-induced stomatal closure in Arabidopsis (Okuma et al., 2011). To assess effect of the *cad2-1* mutation on drought tolerance, we monitored water loss from detached rosette leaves. As shown in Figure 1, compared to wild type, the *cad2-1* mutant exhibited reduced water loss from detached rosette leaves ($P < 0.018$ for Col vs. *cad2-1* at 120 min after leaf detachment). This result suggests that the *cad2-1* mutation improves drought tolerance.

EFFECT OF THE *cad2-1* MUTATION ON ABA REGULATION OF S-TYPE ANION CHANNELS.

The *cad2-1* mutation does not affect ABA activation of I_{Ca} channels, suggesting that GSH functions downstream of $[\text{Ca}^{2+}]_{\text{cyt}}$







FIGURE 1 | The *cad2-1* mutant shows reduced transpirational water loss. Fresh weight loss of detached rosette leaves. The bars represent the mean \pm SE values of five independent replicates.

elevation in guard cell ABA signaling (Okuma et al., 2011). Activation of S-type anion channel is mediated by $[Ca^{2+}]_{cyt}$ elevation and considered as a crucial step for ABA-, MeJA-, and CO_2 -induced stomatal closure (Schroeder and Hagiwara, 1989; Pei et al., 1997; Munemasa et al., 2007; Hu et al., 2010). To confirm whether GSH depletion in the *cad2-1* mutant affects ABA regulation of S-type anion channel activity, whole-cell patch-clamp analysis was performed. S-type anion currents were observed in wild-type GCPs pretreated with $10 \mu M$ ABA ($P < 0.023$ for Col Control vs. Col ABA at -135 mV; **Figure 2**). We also found that ABA evoked S-type anion currents in *cad2-1* GCPs to the same extent as in wild-type GCPs ($P < 0.045$ for *cad2-1* Control vs. *cad2-1* ABA at -135 mV; **Figure 2**). Note that in our experimental condition, free Ca^{2+} concentration in the pipette solution was buffered to $2 \mu M$ and no obvious S-type anion currents were observed in both wild-type guard cells and *cad2-1* guard cells without ABA pretreatment.

THE *cad2-1* MUTATION ENHANCED H_2O_2 ACTIVATION OF I_{Ca} CHANNELS

ABA activation of I_{Ca} channels involves ROS as a second messenger (Pei et al., 2000; Murata et al., 2001; Kwak et al., 2003). Depletion of the major redox regulator, GSH, in the *cad2-1* mutant might affect ROS-mediated ABA signaling in guard cells. Previously we demonstrated that ABA activation of I_{Ca} channels is not enhanced in *cad2-1* guard cells (Okuma et al., 2011). However, ROS activation of I_{Ca} channels in *cad2-1* guard cells was not yet analyzed. We examined effects of the *cad2-1* mutation

on ROS regulation of I_{Ca} channels in guard cells. H_2O_2 at 1 mM activates hyperpolarization-activated currents in both wild-type GCPs and *cad2-1* GCPs (**Figure 3**). The H_2O_2 activation of I_{Ca} currents was enhanced in *cad2-1* GCPs compared to wild-type GCPs ($P < 0.018$ for Col H_2O_2 vs. *cad2-1* H_2O_2 at -180 mV; **Figure 3**). To examine effects of the enhanced H_2O_2 activation of *cad2-1* I_{Ca} channels on H_2O_2 -induced stomatal closure, we performed stomatal bioassay. Significant reduction of stomatal apertures was observed in $100 \mu M$ H_2O_2 -treated *cad2-1* mutant ($P < 0.038$ for *cad2-1* Control vs. *cad2-1* H_2O_2 ; **Figure 4**) but not wild type ($P = 0.13$ for Col Control vs. Col H_2O_2 ; **Figure 4**). Moreover, we found that H_2O_2 -induced stomatal closure in the *cad2-1* mutant was significantly attenuated by a membrane permeable derivative of GSH, GSHmee ($P < 0.047$ for H_2O_2 without GSHmee vs. H_2O_2 with GSHmee; **Figure S1**). These results suggest that decreased GSH contents in *cad2-1* guard cells confer enhanced stomatal response to H_2O_2 .




THE *cad2-1* MUTANT SHOWED ABA-INDUCED APOPLASTIC ROS ACCUMULATION IN LEAVES

Accumulation of ROS in guard cell cytosol occurs during ABA-induced stomatal closure (Pei et al., 2000; Kwak et al., 2003). Previously we revealed that ABA-induced ROS accumulation in guard cell cytosol was not altered by GSH depletion (Jahan et al., 2008; Okuma et al., 2011; Akter et al., 2012). In addition to ROS produced by plasma membrane NAD(P)H oxidases, ROS produced by apoplastic enzymes such as cell-wall bound peroxidases participate in regulation of stomatal movement (An et al., 2008; Khokon et al., 2011; Hossain et al., 2013). To examine whether apoplastic ROS accumulation contributes to guard cell ABA signaling, we performed DAB staining experiments, which allows us to monitor ROS produced by apoplastic enzymes as well as ROS produced by NAD(P)H oxidases (Bindschedler et al., 2006; Khokon et al., 2011; Hossain et al., 2013). Wild-type leaves did not exhibit apoplastic ROS accumulation even when treated with high concentration of ABA (50 μ M) ($P = 0.72$ for Col Control vs. Col ABA; **Figure 5**). However, apoplastic ROS accumulation was significantly increased by ABA in *cad2-1* leaves ($P < 0.04$ for *cad2-1* Control vs. *cad2-1* ABA; **Figure 5**). This result suggests that the accumulation of apoplastic ROS contributes to enhanced ABA response of *cad2-1* guard cells.

DISCUSSION

Depletion of GSH enhances ABA-induced stomatal closure (Jahan et al., 2008; Okuma et al., 2011; Akter et al., 2012). However, the mechanism of how GSH regulates guard cell ABA signaling and involvement of GSH in controlling transpirational water loss have been unclear. In this manuscript, we tested involvement of GSH in controlling water loss from leaves and performed the detailed analysis of GSH regulation of guard cell ABA signaling using the GSH-deficient *cad2-1* mutant. We confirmed that the *cad2-1* mutation caused not only enhanced ABA-induced stomatal closure (Okuma et al., 2011) but also reduction of water loss from leaves (**Figure 1**). Hence manipulation of GSH level might provide a new strategy to improve plant drought tolerance.




ABA activation of S-type anion channels is not altered in the *cad2-1* guard cells (**Figure 2**). It has been suggested that the elevated $[Ca^{2+}]_{cyt}$ is required for ABA activation of S-type anion channels and ABA “primes” Ca^{2+} sensitivity of S-type anion channels (Siegel et al., 2009). In this study, we used the pipette solution where free $[Ca^{2+}]$ was buffered to 2 μ M (See Materials and Methods). No obvious S-type anion current was observed in *cad2-1* guard cells as well as wild-type guard cells without ABA pretreatment (**Figure 2**), suggesting that the *cad2-1* mutation does not affect the priming state of Ca^{2+} sensitivity of S-type anion channels.

ABA induces guard cell $[Ca^{2+}]_{cyt}$ elevation via activation of plasma membrane I_{Ca} channels (Hamilton et al., 2000; Pei et al., 2000). Previously we found identical ABA activation of I_{Ca} channels in wild-type and *cad2-1* guard cells (Okuma et al., 2011). ROS mediates ABA activation of I_{Ca} channels (Pei et al., 2000; Murata et al., 2001; Kwak et al., 2003). In this study, we found enhanced response of *cad2-1* I_{Ca} channels to exogenous H_2O_2 (**Figure 3**). It was reported that ozone, an elicitor of ROS, induces biphasic $[Ca^{2+}]_{cyt}$ elevation in seedlings and depletion of GSH by the *cad2-1* mutation and buthionine sulphoximine, an inhibitor of γ -glutamylcysteine synthetase, enhances first peak of the ozone-induced biphasic $[Ca^{2+}]_{cyt}$ response (Evans et al., 2005). These results suggest that GSH pools control sensitivity of plasma membrane Ca^{2+} permeable channels to ROS and downstream Ca^{2+} signals in plant cells.

Consistent with the enhancement of I_{Ca} channel response to ROS, the *cad2-1* mutant exhibits enhanced H_2O_2 -induced stomatal closure (**Figure 4**). Note that depletion of GSH in guard cell cytosol is induced by ABA (Okuma et al., 2011), but not by H_2O_2 (Akter et al., 2013). These results imply that ABA sensitizes guard cells to ROS by decreasing GSH content via a pathway distinct from the ABA-mediated ROS production pathway (**Figure 6**).

Previously we reported identical cytosolic ROS accumulation induced by ABA in cytosol of wild-type and *cad2-1* guard cells using 2',7'-dichlorodihydrofluorescein diacetate (Okuma et al., 2011). It has been suggested that apoplastic ROS signals are also



involved in regulation of stomatal movement (An et al., 2008; Khokon et al., 2011; Hossain et al., 2013). In this study we monitored ROS accumulation in whole leaves using DAB. ABA induced ROS accumulation in whole leaves of the *cad2-1* mutant but not in leaves of wild type (Figure 5). These results suggest that GSH depletion by the *cad2-1* mutation affects ROS homeostasis in apoplastic space rather than that in guard cells during ABA-induced stomatal closure.

Based on the findings obtained in this study, we present one simplified signal model shown in Figure 6. ABA decreases guard cell GSH content via ROS-independent pathway (Akter et al., 2013). The decreased GSH content causes significant ROS accumulation in apoplast (Figure 5) and also sensitizes guard cell I_{Ca} channels to apoplastic ROS (Figure 3) by unknown mechanism, resulting in enhanced stomatal response to ABA. In wild-type leaves, ABA decreases GSH content (Okuma et al., 2011) but does not induce significant apoplastic ROS accumulation (Figure 5), suggesting that the apoplastic ROS signal is employed to modulate ABA responsiveness of guard cells by other stimuli rather than by ABA signaling itself. Molecular mechanisms of how GSH regulates ROS sensitivity to plasma membrane Ca^{2+} permeable channels and apoplastic ROS homeostasis during ABA-induced stomatal closure would be investigated in the future.

ACKNOWLEDGMENTS

We thank Dr. Eiji Okuma (Okayama University) for critical reading of the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://www.frontiersin.org/journal/10.3389/fpls.2013.00472/abstract>

Figure S1 | GSH^{mee} attenuates enhanced stomatal response to H₂O₂ of the *cad2-1* mutant. Detached rosette leaves of the *cad2-1* mutant were incubated on stomatal assay buffer (see Materials and Methods) with or without 10 μ M GSH^{mee} for 2 h in the light, followed by the addition of 100 μ M H₂O₂. Stomatal apertures were measured 2 h after H₂O₂ application. Twenty averages from three independent experiments (60 total stomata per bar) are shown. The bars represent the mean \pm SE values.

REFERENCES

- Akter, N., Okuma, E., Sobahan, M. A., Uraji, M., Munemasa, S., Nakamura, Y., et al. (2013). Negative regulation of methyl jasmonate-induced stomatal closure by glutathione in *Arabidopsis*. *J. Plant Growth Regul.* 32, 208–215. doi: 10.1007/s00344-012-9291-7
- Akter, N., Sobahan, M. A., Hossain, M. A., Uraji, M., Nakamura, Y., Mori, I. C., et al. (2010). The involvement of intracellular glutathione in methyl jasmonate signaling in *Arabidopsis* guard cells. *Biosci. Biotechnol. Biochem.* 74, 2504–2506. doi: 10.1271/bbb.100513
- Akter, N., Sobahan, M. A., Uraji, M., Ye, W., Hossain, M. A., Mori, I. C., et al. (2012). Effects of depletion of glutathione on abscisic acid- and methyl jasmonate-induced stomatal closure in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 76, 2032–2037. doi: 10.1271/bbb.120384
- Allan, A. C., Fricker, M. D., Ward, J. L., Beale, M. H., and Trewavas, A. J. (1994). Two transduction pathways mediate rapid effects of abscisic acid in *Commelina* guard cells. *Plant Cell* 6, 1319–1328. doi: 10.2307/3869829
- Allen, G. J., Kuchitsu, K., Chu, S. P., Murata, Y., and Schroeder, J. I. (1999). *Arabidopsis abi1-1* and *abi2-1* phosphatase mutations reduce abscisic acid-induced cytosolic calcium rises in guard cells. *Plant Cell* 11, 1785–1798.
- An, Z., Jing, W., Liu, Y. L., and Zhang, W. H. (2008). Hydrogen peroxide generated by copper amine oxidase is involved in abscisic acid-induced stomatal closure in *Vicia faba*. *J. Exp. Bot.* 59, 815–825. doi: 10.1093/jxb/erm370
- Bindschedler, L. V., Dewdney, J., Blee, K. A., Stone, J. M., Asai, T., Plotnikov, J., et al. (2006). Peroxidase-dependent apoplastic oxidative burst in *Arabidopsis* required for pathogen resistance. *Plant, J.* 47, 851–863. doi: 10.1111/j.1365-313X.2006.02837.x
- Brandt, B., Brodsky, D. E., Xue, S., Negi, J., Iba, K., Kangasjärvi, J., et al. (2012). Reconstitution of abscisic acid activation of SLAC1 anion channel by CPK6 and OST1 kinases and branched ABI1 PP2C phosphatase action. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10593–10598. doi: 10.1073/pnas.1116590109
- Evans, N. H., McAinsh, M. R., Hetherington, A. M., and Knight, M. R. (2005). ROS perception in *Arabidopsis thaliana*: the ozone-induced calcium response. *Plant J.* 41, 615–626. doi: 10.1111/j.1365-313X.2004.02325.x
- Fujii, H., Chinnusamy, V., Rodrigues, A., Rubio, S., Antoni, R., Park, S. Y., et al. (2009). *In vitro* reconstitution of an abscisic acid signalling pathway. *Nature* 462, 660–664. doi: 10.1038/nature08599
- Geiger, D., Scherzer, S., Mumm, P., Marten, I., Ache, P., Matschi, S., et al. (2010). Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca^{2+} affinities. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8023–8028. doi: 10.1073/pnas.0912030107
- Geiger, D., Scherzer, S., Mumm, P., Stange, A., Marten, I., Bauer, H., et al. (2009). Activity of guard cell anion channel SLAC1 is controlled by drought-stress signaling kinase-phosphatase pair. *Proc. Natl. Acad. Sci. U.S.A.* 106, 21425–21430. doi: 10.1073/pnas.0912021106
- Gilroy, S., Fricker, M. D., Read, N. D., and Trewavas, A. J. (1991). Role of calcium in signal transduction of *Commelina* guard cells. *Plant Cell* 3, 333–344. doi: 10.2307/3869209
- Grabov, A., and Blatt, M. R. (1998). Membrane voltage initiates Ca^{2+} waves and potentiates Ca^{2+} increases with abscisic acid in stomatal guard cells. *Proc. Natl. Acad. Sci. U.S.A.* 95, 4778–4783. doi: 10.1073/pnas.95.8.4778
- Hamilton, D. W. A., Hills, A., Köhler, B., and Blatt, M. R. (2000). Ca^{2+} channels at the plasma membrane of stomatal guard cells are activated by hyperpolarization and abscisic acid. *Proc. Natl. Acad. Sci. U.S.A.* 97, 4967–4972. doi: 10.1073/pnas.080068897
- Hossain, M. S., Ye, W., Hossain, M. A., Okuma, E., Uraji, M., Nakamura, Y., et al. (2013). Glucosinolate degradation products, isothiocyanates, nitriles, and thiocyanates, induce stomatal closure accompanied by peroxidase-mediated reactive oxygen species production in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 77, 977–983. doi: 10.1271/bbb.120928
- Hu, H., Boisson-Dernier, A., Israelsson-Nordström, M., Böhmer, M., Xue, S., Ries, A., et al. (2010). Carbonic anhydrases are upstream regulators of CO₂-controlled stomatal movements in guard cells. *Nat. Cell Biol.* 12, 87–93. doi: 10.1038/ncb2009
- Jahan, M. S., Ogawa, K., Nakamura, Y., Shimoishi, Y., Mori, I. C., and Murata, Y. (2008). Deficient glutathione in guard cells facilitates abscisic acid-induced stomatal closure but does not affect light-induced stomatal opening. *Biosci. Biotechnol. Biochem.* 72, 2795–2798. doi: 10.1271/bbb.80407
- Khokon, A. R., Okuma, E., Hossain, M. A., Munemasa, S., Uraji, M., Nakamura, Y., et al. (2011). Involvement of extracellular oxidative burst in salicylic acid-induced stomatal closure in *Arabidopsis*. *Plant Cell Environ.* 34, 434–443. doi: 10.1111/j.1365-3040.2010.02253.x
- Kwak, J. M., Mori, I. C., Pei, Z. M., Leonhardt, N., Torres, M. A., Dangl, J. L., et al. (2003). NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J.* 22, 2623–2633. doi: 10.1093/emboj/cdg277
- Lee, S. C., Lan, W., Buchanan, B. B., and Luan, S. (2009). A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. *Proc. Natl. Acad. Sci. U.S.A.* 106, 21419–21424. doi: 10.1073/pnas.0910601106
- Linder, B., and Raschke, K. (1992). A slow anion channel in guard cells, activating at large hyperpolarization, may be principal for stomatal closing. *FEBS Lett.* 313, 27–30. doi: 10.1016/0014-5793(92)81176-M
- Ma, Y., Szostkiewicz, I., Korte, A., Moes, D., Yang, Y., Christmann, A., et al. (2009). Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* 324, 1064–1068. doi: 10.1126/science.1172408
- Marten, H., Konrad, K. R., Dietrich, P., Roelfsema, M. R. G., and Hedrich, R. (2007). Ca^{2+} -dependent and -independent abscisic acid activation of plasma

- membrane anion channels in guard cells of *Nicotiana tabacum*. *Plant Physiol.* 143, 28–37. doi: 10.1104/pp.106.092643
- Maruta, T., Tanouchi, A., Tamoi, M., Yabuta, Y., Yoshimura, K., Ishikawa, T., et al. (2010). Arabidopsis chloroplastic ascorbate peroxidase isoenzymes play a dual role in photoprotection and gene regulation under photooxidative stress. *Plant Cell Physiol.* 51, 190–200. doi: 10.1093/pcp/pcp177
- May, M. J., Vernoux, T., Leaver, C., van Montagu, M., and Inze, D. (1998). Glutathione homeostasis in plants: implications for environmental sensing and plant development. *J. Exp. Bot.* 49, 649–667. doi: 10.1093/jexbot/49.321.649
- McAinsh, M. R., Brownlee, C., and Hetherington, A. M. (1990). Abscisic acid-induced elevation of guard-cell cytosolic Ca^{2+} precedes stomatal closure. *Nature* 343, 186–188. doi: 10.1038/343186a0
- Miao, Y., Lv, D., Wang, P., Wang, X. C., Chen, J., Miao, C., et al. (2006). Arabidopsis glutathione peroxidase functions as both a redox transducer and a scavenger in abscisic acid and drought stress responses. *Plant Cell* 18, 2749–2766. doi: 10.1105/tpc.106.044230
- Munemasa, S., Hossain, M. A., Nakamura, Y., Mori, I. C., and Murata, Y. (2011). The Arabidopsis calcium-dependent protein kinase, CPK6, functions as a positive regulator of methyl jasmonate signaling in guard cells. *Plant Physiol.* 155, 553–561. doi: 10.1104/pp.110.162750
- Munemasa, S., Oda, K., Watanabe-Sugimoto, M., Nakamura, Y., Shimoishi, Y., and Murata, Y. (2007). The *coronatine-insensitive 1* mutation reveals the hormonal signaling interaction between abscisic acid and methyl jasmonate in *Arabidopsis* guard cells. Specific impairment of ion channel activation and second messenger production. *Plant Physiol.* 143, 1398–1407. doi: 10.1104/pp.106.091298
- Murata, Y., Pei, Z. M., Mori, I. C., and Schroeder, J. I. (2001). Abscisic acid activation of plasma membrane Ca^{2+} channels in guard cells requires cytosolic NAD(P)H and is differentially disrupted upstream and downstream of reactive oxygen species production in *abi1-1* and *abi2-1* protein phosphatase 2C mutants. *Plant Cell* 13, 2513–2523. doi: 10.1105/tpc.13.11.2513
- Negi, J., Matsuda, O., Nagasawa, T., Oba, Y., Takahashi, H., Kawai-Yamada, M., et al. (2008). CO_2 regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature* 452, 483–486. doi: 10.1038/nature06720
- Noctor, G., and Foyer, C. H. (1998). Ascorbate and glutathione: keeping active oxygen under control. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49, 249–279. doi: 10.1146/annurev.arplant.49.1.249
- Okuma, E., Jahan, M. S., Munemasa, S., Hossain, M. A., Muroyama, D., Islam, M. M., et al. (2011). Negative regulation of abscisic acid-induced stomatal closure by glutathione in *Arabidopsis*. *J. Plant Physiol.* 168, 2048–2055. doi: 10.1016/j.jplph.2011.06.002
- Park, S. Y., Fung, P., Nishimura, N., Jensen, D. R., Fujii, H., Zhao, Y., et al. (2009). Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* 324, 1068–1071. doi: 10.1126/science.1173041
- Pei, Z. M., Kuchitsu, K., Ward, J. M., Schwarz, M., and Schroeder, J. I. (1997). Differential abscisic acid regulation of guard cell slow anion channels in *Arabidopsis* wild-type and *abi1* and *abi2* mutants. *Plant Cell* 9, 409–423. doi: 10.2307/3870491
- Pei, Z. M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G. J., et al. (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* 406, 731–734. doi: 10.1038/35021067
- Schroeder, J. I., and Hagiwara, S. (1989). Cytosolic calcium regulates ion channels in the plasma membrane of *Vicia faba* guard cells. *Nature* 338, 427–430. doi: 10.1038/338427a0
- Schroeder, J. I., and Hagiwara, S. (1990). Repetitive increases in cytosolic Ca^{2+} of guard cells by abscisic acid activation of nonselective Ca^{2+} permeable channels. *Proc. Natl. Acad. Sci. U.S.A.* 87, 9305–9309. doi: 10.1073/pnas.87.23.9305
- Schroeder, J. I., and Keller, B. U. (1992). Two types of anion channel currents in guard cells with distinct voltage regulation. *Proc. Natl. Acad. Sci. U.S.A.* 89, 5025–5029. doi: 10.1073/pnas.89.11.5025
- Siegel, R. S., Xue, S., Murata, Y., Yang, Y., Nishimura, N., Wang, A., et al. (2009). Calcium elevation-dependent and attenuated resting calcium-dependent abscisic acid induction of stomatal closure and abscisic acid-induced enhancement of calcium sensitivities of S-type anion and inward-rectifying K channels in *Arabidopsis* guard cells. *Plant J.* 59, 207–220. doi: 10.1111/j.1365-313X.2009.03872.x
- Vahisalu, T., Kollist, H., Wang, Y. F., Nishimura, N., Chan, W. Y., Valerio, G., et al. (2008). SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature* 452, 487–491. doi: 10.1038/nature06608

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 12 August 2013; paper pending published: 03 October 2013; accepted: 31 October 2013; published online: 20 November 2013.

Citation: Munemasa S, Muroyama D, Nagahashi H, Nakamura Y, Mori IC and Murata Y (2013) Regulation of reactive oxygen species-mediated abscisic acid signaling in guard cells and drought tolerance by glutathione. *Front. Plant Sci.* 4:472. doi: 10.3389/fpls.2013.00472

This article was submitted to *Plant Physiology*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2013 Munemasa, Muroyama, Nagahashi, Nakamura, Mori and Murata. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.