

Optimized gene editing technology for *Drosophila melanogaster* using germ line-specific Cas9

Xingjie Ren^{a,1}, Jin Sun^{a,1}, Benjamin E. Housden^b, Yanhui Hu^b, Charles Roesel^c, Shuailiang Lin^b, Lu-Ping Liu^{a,d}, Zhihao Yang^a, Decai Mao^a, Lingzhu Sun^{a,d}, Qujie Wu^{a,d}, Jun-Yuan Ji^e, Jianzhong Xi^{f,g}, Stephanie E. Mohr^b, Jiang Xu^{a,h,i,2}, Norbert Perrimon^{b,j,2}, and Jian-Quan Ni^{a,2}

^aGene Regulatory Laboratory, School of Medicine, Tsinghua University, Beijing 100084, China; ^bDepartment of Genetics, Harvard Medical School, Boston, MA 02115; ^cDepartment of Biology, Northeastern University, Boston, MA 02115; ^dTsinghua Fly Center, Tsinghua University, Beijing 100084, China; ^eDepartment of Molecular and Cellular Medicine, College of Medicine, Texas A&M Health Science Center, College Station, TX 77843; ^fInstitute of Molecular Medicine and ^gCollege of Engineering, Peking University, Beijing 100871, China; ^hSchool of Basic Medical Sciences, Wuhan University, Wuhan 430071, China; ⁱCollege of Bioengineering, Hubei University of Technology, Wuhan 430068, China; and ^jHoward Hughes Medical Institute, Harvard Medical School, Boston, MA 02115

Contributed by Norbert Perrimon, October 8, 2013 (sent for review September 5, 2013)

The ability to engineer genomes in a specific, systematic, and cost-effective way is critical for functional genomic studies. Recent advances using the CRISPR-associated single-guide RNA system (Cas9/sgRNA) illustrate the potential of this simple system for genome engineering in a number of organisms. Here we report an effective and inexpensive method for genome DNA editing in *Drosophila melanogaster* whereby plasmid DNAs encoding short sgRNAs under the control of the *U6b* promoter are injected into transgenic flies in which Cas9 is specifically expressed in the germ line via the *nanos* promoter. We evaluate the off-targets associated with the method and establish a Web-based resource, along with a searchable, genome-wide database of predicted sgRNAs appropriate for genome engineering in flies. Finally, we discuss the advantages of our method in comparison with other recently published approaches.

nanos-Cas9 | HRMA

Much of our knowledge of the mechanisms underlying biological processes relies on genetic approaches, whereby gene activity is perturbed and the phenotypic consequences of perturbation are analyzed in detail. In recent years, several major advances have been made in the design of methods for specifically and efficiently perturbing genomes. Arguably, the most exciting advances rely on the ability to induce double-strand breaks (DSBs) by targeting a nuclease to a specific genomic sequence. Repair of DSBs by the error-prone nonhomologous end-joining (NHEJ) mechanism allows for the recovery of small deletions; moreover, repair of DSBs by homologous recombination (HR) in the presence of a donor template opens the door to a wide range of specifically engineered changes at the targeted site (1).

Two nuclease-based systems, the zinc-finger nuclease (ZFN) and transcription activator-like effector nuclease (TALEN) systems, work effectively in a number of organisms (2–7). But because these approaches require the production of a construct encoding a unique DNA-binding protein fused to the nuclease domain, they can be both cumbersome and costly. In contrast, the recent approach based on the bacterial CRISPR-associated single-guide RNA (Cas9/sgRNA) system does not require production of specific fusion proteins for each targeted sequence (8–10).

Cas9 was first identified in type II *Streptococcus pyogenes* as an RNA-guided defense system against invading viruses and plasmids (11–13). This adaptive immune-like system contains three components: CRISPR RNA (crRNA), trans-activating CRISPR RNA (tracrRNA), and Cas9. The tracrRNA triggers Cas9 nuclease activity and the crRNA guides Cas9 to cleave the specific foreign dsDNA sequence via base-pairing between the crRNA and the target DNA. Importantly, a single-guide RNA (sgRNA, also known as chiRNA), comprising the minimal crRNA and tracrRNA, can function similarly to the crRNA and tracrRNA, thereby providing a simplified method for genome editing (8–10, 14–20).

Given the great promise of the Cas9/sgRNA method for genome engineering, we set out to test the system in *Drosophila*

melanogaster. Although many genomic engineering tools are already available for *Drosophila* (21, 22), including ZFNs and TALENs (2–7), we reasoned that the Cas9/sgRNA system might provide a more cost-effective approach, allowing its use at a genome scale. Similar to other recent reports (14, 16, 20), we found that the Cas9/sgRNA system works in *Drosophila*. Moreover, we found that injection of sgRNAs into a transgenic line expressing Cas9 in the germ line (*nanos*-Cas9) provides a straightforward, effective, and inexpensive method that can facilitate the production of large mutant collections. We also evaluated the frequency of off-target events associated with Cas9/sgRNA. In addition, to facilitate prediction and testing of off-target events, we established a searchable database of in silico predicted sgRNAs that includes information about predicted off-target events. Finally, we compared our approach with methodologies recently reported by others (14, 16, 20).

Results

Optimization of the Cas9/sgRNA System. To test the efficacy of the Cas9/sgRNA system in inducing germ-line mutations in *D. melanogaster*, we generated a vector that expresses the Cas9 gene under the *nanos* (*nos*) promoter (Fig. 1A). We reasoned that the *nos* promoter would induce strong and restricted expression of the transgene in germ cells, which might help avoid

Significance

Using the recently introduced Cas9/sgRNA technique, we have developed a method for specifically targeting *Drosophila* germ-line cells to generate heritable mutant alleles. We have established transgenic lines that stably express Cas9 in the germ line and compared different promoters and scaffolds of sgRNA in terms of their efficiency of mutagenesis. An overall mutagenesis rate of 74.2% was achieved with this optimized system, as determined by the number of mutant progeny out of all progeny screened. We also evaluated the off-targets associated with the method and established a Web-based resource, as well as a searchable, genome-wide database of predicted sgRNAs appropriate for genome engineering in flies. Our results demonstrate that this optimized Cas9/sgRNA system in *Drosophila* is efficient, specific, and cost-effective and can be readily applied in a semi-high-throughput manner.

Author contributions: J. Xu, N.P., and J.-Q.N. designed research; X.R., J.S., B.E.H., L.-P.L., Z.Y., D.M., L.S., Q.W., J. Xu, N.P., and J.-Q.N. performed research; Y.H., C.R., S.L., J.-Y.J., and J. Xi contributed new reagents/analytic tools; X.R., J.S., B.E.H., L.-P.L., Z.Y., D.M., L.S., Q.W., J. Xu, and J.-Q.N. analyzed data; and S.E.M., J. Xu, N.P., and J.-Q.N. wrote the paper.

The authors declare no conflict of interest.

¹X.R. and J.S. contributed equally to this work.

²To whom correspondence may be addressed. E-mail: jiangxu@mail.tsinghua.edu.cn, perrimon@receptor.med.harvard.edu, or nijq@mail.tsinghua.edu.cn.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1318481110/-DCSupplemental.

Table 1. Heritable mutation rates using nos-Cas9 and sgRNA plasmids

sgRNA	G0 adults					Overall heritable mutation rate (individual), %, median (range) [†]
	Embryos, <i>n</i>	Total <i>n</i>	Survival rate, %	Fertile, <i>n</i>	Germ-line mutants, <i>n</i> (%) [*]	
<i>w1</i>	65	14	21.5	13	4 (30.8)	2.7 (3.1–15.1)
<i>w2</i>	60	15	25.0	13	3 (23.1)	0.6 (1.2–3.2)

^{*}The percentage of germ-line mutants was calculated as the proportion of fertile G0 flies that gave rise to white-eyed progeny.

[†]The overall heritable mutation rate was calculated as the number of white-eye F1s divided by the number of all F1s observed. Mutation rates for each G0 fly that generated white-eyed progeny are shown in parentheses and are calculated as the number of white-eyed F1s divided by the total F1s observed from a single G0 fly.

13-nt neighboring protospacer adjacent motif (PAM), the so-called “seed” region, can be tolerated, given that sgRNA with a single mismatch in nucleotides 14–20 can still guide Cas9 to introduce DSB in the genome (8–10, 23). Furthermore, Cas9 can induce a DSB at genomic sequences with as many as five nucleotide mismatches to the sgRNA (18, 24). To evaluate the potential off-target events associated with *w1* and *w2* sgRNAs, we identified all possible locations that share high sequence similarity with the target site and include a PAM sequence (see *Online Resource of sgRNA Designs for Drosophila Genome Editing* and Fig. 3). The tool did not identify any off-target sequences using default conditions, but when search criteria were relaxed, 10 potential off-target sites were found (five for each sgRNA), with the closest match containing six mismatches to the on-target sequence (Fig. 3).

We next PCR-amplified the potential off-target regions from genomic DNA of white-eyed F1 flies with defined deletions generated by injection of both *w1* and *w2* sgRNAs and assayed for the presence of mutations using HRMA or sequencing. Using HRMA, we analyzed eight independent F1 adult flies for each sgRNA and failed to detect any mutations at any of these sites (Table S1). Finally, we sequenced 20 cloned PCR products from each of eight F1 adults per off-target site and did not detect any mutations, further confirming the HRMA results (Fig. S1).

Online Resource of sgRNA Designs for *Drosophila* Genome Editing. To facilitate the use of sgRNAs in *Drosophila*, we established a freely available online resource of precomputed sgRNAs (www.flymai.org/crispr). We developed the resource in two steps. We first extracted all possible 23mers from both the forward and reverse strands of the genomic sequence of *D. melanogaster* based on FlyBase release 5.52 (July 15, 2013). Some 7.6 million of these 23mers end with a PAM sequence (NGG) and have a unique 15-bp sequence, which includes the seed region sequence and PAM sequence, in the genome. We selected this subset of 23mers as potential CRISPR sgRNA designs and aligned them to the genome sequence. We found that 28% of these sgRNA sequences aligned to intergenic regions, 37% aligned to introns, and 35% targeted the exon region of annotated genes [coding sequence (CDS) or noncoding regions; Table 4].

We next systematically analyzed potential off-target sites, which share a certain level of sequence similarity (up to five

mismatches) and end with a PAM sequence (NGG), for each CRISPR sgRNA design. Although the presence of the NGG PAM sequence on the genomic DNA is required for Cas9 recognition, variations in the PAM (e.g., NAG) can be tolerated in bacteria (18, 24); thus, to be cautious, in the off-target analysis we also included the sites ending with NAG. We found that for CRISPR sgRNA designs targeting gene CDS regions within exons, 104,892 (5% of all designs targeting CDS) have no predicted off-targets, whereas 506,998 (24%) have no predicted off-targets in CDS regions of other genes (Table 4). The designs that target a CDS and have no predicted off-targets correspond to 11,950 *Drosophila* genes (i.e., 86% coverage of protein-coding genes). Together, the 611,890 designs that target a CDS and have no predicted off-targets in another CDS correspond to 12,996 protein-coding genes (93% coverage of protein-coding genes).

To make the designs available, we developed a user interface that allows the community to query all relevant CRISPR designs by gene identifier (gene symbol, CG number, or FBgn) or genome coordinates and to view all corresponding sgRNA designs on a genome browser (Fig. S2). To identify CRISPR sgRNAs targeting intergenic regions, such as promoters, users can enter an identifier corresponding to a nearby gene, then use the navigator to find the intergenic region of interest, or can enter genome coordinates directly. To help users select relevant designs, CRISPR sgRNAs with different predicted properties regarding the target region and potential off-targets are separated into different tracks, for example “CRISPRs target CDS region/without any off-target” and “CRISPRs target intron/without any off-target.” By clicking on the sgRNA of interest at the genome browser, the user can access detailed information, such as sequence and potential off-target genes (Fig. S2).

Discussion

We report a simple and effective method for efficiently generating heritable loss-of-function alleles in *D. melanogaster* using the Cas9/sgRNA system. Specifically, we show that injection of sgRNAs into a transgenic line that expresses Cas9 in the germ line (*nos-Cas9*) is an effective and relatively inexpensive technique for genome engineering.

Several recent studies also have reported success in using the Cas9/sgRNA system to edit genomic DNA in *Drosophila* (14,

Table 2. Heritable mutation rates using nos-Cas9 and different sgRNA plasmid constructs

SgRNA promoter	sgRNA scaffold	G0 adults					Overall heritable mutation rate (individual), % median (range) [†]
		Embryos, <i>n</i>	Total <i>n</i>	Survival, %	Fertile, <i>n</i>	Germ-line mutants, <i>n</i> (%) [*]	
<i>U6b</i>	Short	78	36	46.2	28	6 (21.4)	3.2 (1.1–64.6)
<i>U6a</i>	Short	68	17	25.0	5	4 (80.0)	2.1 (1.2–6.8)
<i>nos-mini</i>	Short	56	9	16.1	7	0	0
<i>U6b</i>	Long	64	18	28.1	13	4 (30.8)	1.7 (1.5–10.0)

^{*}The percentage of germ-line mutants was calculated as the proportion of fertile G0 flies that gave rise to white-eyed progeny.

[†]The overall heritable mutation rate was calculated as the number of white-eyed F1s divided by the number of all F1s observed. Mutation rates for each G0 fly that generated white-eyed progeny are shown in parentheses and are calculated as the number of white-eyed F1s divided by the total F1s observed from a single G0 fly.




Fig. 2. Testing the optimized *Drosophila nos-Cas9/sgRNA* system on the *white* locus. (A) Schematic of two sgRNAs targeting the *white* locus, with the 20-nt target sequence underscored and PAM in bold type. Gray boxes represent exons. *w1* sgRNA targets the first exon, and *w2* targets the second. (B) RT-PCR results confirming the expression of Cas9 in {*nos-Cas9*}attP40 and {*nos-Cas9*}attP2 embryos, with actin 5C as an internal control. (C and D) Representative sequencing results showing the indel mutations generated in this study, with *w1* sgRNA (C) and *w2* sgRNA (D). (E) Schematic showing the defined deletion generated by using *w1* and *w2* sgRNAs, along with representative sequencing results showing the break points of the deletions generated. The genomic sequence between the sgRNA target sites is not shown. In C–E, the targeted genomic DNA sequence is underscored, and the NGG PAM sequence is in bold type. Dashed lines represent the locations of genomic deletions detected by sequencing.

16, 20). Gratz et al. (16) coinjected two DNA plasmids, one encoding Cas9 and the other encoding a single sgRNA targeting the X-linked *yellow* (*y*) gene. In that study, Cas9 was expressed under the control of the *hsp70* promoter and 3' UTR, and the sgRNA was under the control of the snRNA *U6b* promoter. The system was effective in generating somatic patches of yellow tissues in 62% of injected G0 males. Furthermore, germ-line transmission, as detected by the presence of at least one yellow offspring, was observed in 6.4% of the G0 males. The overall germ-line transmission rate (i.e., the number of yellow offspring as a percentage of all progeny) was 0.25%. Gratz et al. (16) also showed that when Cas9 was coinjected with a pair of sgRNAs designed to work together to precisely delete *y*, 25% of the injected males produced yellow progeny, with an overall germ-line transmission rate of 1.4%.

A different approach was taken by Bassett et al. (14) and Yu et al. (20), who injected fly embryos with in vitro transcripts encoding Cas9 and sgRNA. Bassett et al. (14) injected a mix of Cas9 mRNA and sgRNA transcripts targeting either *y* or *w*. In their system, both Cas9 and sgRNA were under the control of a bacterial T7 promoter, and a poly-A tail was added post-transcriptionally to the in vitro-transcribed Cas9 mRNA. This system was more effective than the system reported by Gratz et al. (16) at generating somatic mosaics, with 86% of the injected G0 males exhibiting patches of yellow tissues; however,

the adult survival rate was only 3%. For the *w* gene, 25% of injected G0 males had visible phenotypes, and the survival rate was <3%. Although the authors reported that reducing the concentration of mRNA increased the survival rate from 3% to 11%, the proportion of mosaic adults decreased dramatically, from 86% to 10%. Furthermore, germ-line transmission, as detected by the presence of at least one yellow offspring, was observed in 58% of the G0 males, and 34.5% of all offspring contained a mutation in the *y* gene.

Similarly, Yu et al. (20) injected the in vitro synthesized Cas9 mRNA and sgRNA, and reported a similar efficiency of mutagenesis. In that study, the Cas9 mRNA was transcribed under the control of the Sp6 promoter, and transcription of the sgRNA was under the control of the T7 promoter.

Compared with the foregoing methods, the approach that we describe herein has three key advantages. First, we did not observe any visible phenotypes in G0-injected flies, most likely because control by the *nos* promoter limits Cas9 expression to the germ cells. Importantly, the absence of somatic events will circumvent the potentially deleterious effects of inducing mutations into the somatic tissues of G0 animals when genes required for cell viability or developmental processes are targeted. Second, the frequency of germ-line transmission that we observed is much higher than that reported for other methods. Injection of a single sgRNA plasmid into Cas9 transgenic flies

Table 3. Heritable mutation rates using transgenic {*nos-Cas9*}attP2 flies and sgRNA plasmid

sgRNA	Embryos, <i>n</i>	G0 adults			Overall heritable mutation rate (individual), %, median (range) [†]
		Total <i>n</i>	Survival, %	Fertile, <i>n</i>	
<i>w1</i>	61	16	26.2	13	42.9 (3.3–100)
<i>w2</i>	65	17	26.2	17	12.4 (1.1–58.3)
<i>w1 + w2</i>	71	21	29.6	12	74.2 (13.6–100)

*The percentage of germ-line mutants was calculated as the proportion of fertile G0 flies that gave rise to white-eyed progeny.

[†]The overall heritable mutation rate was calculated as the number of white-eyed F1s divided by the number of all F1s observed. Mutation rates for each G0 fly that generated white-eyed progeny are shown in parentheses and were calculated as the number of white-eyed F1s divided by the total number of F1s observed from a single G0 fly.

A

NGG:PAM N:mismatch
 CAGGAGCTATTAATTCGCGGAGG w1 sgRNA target site
 GAAGGGGATTACTTCGCGGAGG w1 potential off-target site: Mo25
 AGTGAATAATAATTCGCGGGG w1 potential off-target site: CG6428
 GTTATCCGTTCAATTCGCGGTGG w1 potential off-target site: pyd (intron)
 CAATATGTTTAATTCGCGCGG w1 potential off-target site: His2Av (intron)

B

TAGTTGGCCGCTCCCTGAACCGG w2 sgRNA target site
 TAGGCATCCAGTCCCTGAACCGG w2 potential off-target site: Nrg
 TGTGGTATGTTCCCTGAACCGG w2 potential off-target site: Vm26Ac
 GGCTTTGAAATCCCTGAACCGG w2 potential off-target site: CTPsyn
 TCTGGAAGTCTCCCTGAACAGG w2 potential off-target site: CG1907
 CCGGCTGTAGCTCCCTGAACAGG w2 potential off-target site: fru (intron)

Fig. 3. No off-targets were detected using the optimized *Drosophila* Cas9/sgRNA system to generate heritable mutations. (A and B) Potential off-target sites for w1 sgRNA (A) and w2 sgRNA (B). Genomic regions that have at least 11-nt homology to the sgRNA seed region and a neighboring PAM sequence were sequenced from F1 white-eyed flies generated by the optimized system. Mismatches between the potential off-targets and the targeted region are underscored. The PAM sequences are in bold type.

produced a G0 rate of 93.3% (28 of 30) and an overall heritable mutation rate as high as 42.9%. In addition, we found that injection of a pair of sgRNAs can lead to the generation of defined genome deletion of the region between the targets, with a further improved overall heritable mutation rate (74.2%; Table 3). Indeed, after injection of two sgRNAs, 91.7% (11 of 12) of fertile G0 flies produced F1 mutants. Among all mutant-producing fertile G0 flies, 18.2% (2 of 11) produced mutant progeny at a rate of 50–80%, 18.2% (2 of 11) did so at a rate of 80–100%, and 54.5% (6 of 11) generated only mutant progeny (100%). We found a much higher survival rate of G0 flies (26.2%) after injection of only w1 sgRNA plasmid. This rate compares favorably with a previously reported 3% survival obtained after direct injection of Cas9 mRNA and sgRNA (14). At 81%, the G0 fertility rate of the optimized system was also higher than the 72% rate found with the Cas9 mRNA method. Third, the direct material costs associated with our method are roughly 10–20% of those associated with direct injection of Cas9 mRNA, because our approach avoids the need for in vitro transcription. In addition, because of its high efficiency and low lethality, our approach requires microinjection of only approximately 20 embryos (roughly 1/15th of that in previously reported methods) to achieve the same level of mutagenesis, further contributing to the cost-effectiveness of our approach.

To be useful, a genome engineering technique must be specific. Experience with other mutagenesis methods and previous reports with CRISPRs have taught us that off-targets are a major concern. Previous in vitro studies have indicated that sgRNA/Cas9 specificity extends past a 7- to 12-bp seed sequence and that up to five mismatches are tolerated depending on their positions along the sgRNA (23, 24). Using various methods, we detected no off-targets in either CDS or intergenic regions (Fig. S1), suggesting that the Cas9 protein is expressed at appropriate levels in the transgenic animals, and that specificity can be

improved by designing sgRNAs to avoid highly similar off-targets.

Several CRISPR Web tools are available online; however, these do not provide much information about potential off-targets. We have developed a user interface that allows the community to query all relevant sgRNA designs by gene identifier (gene symbol, CG number, or FBgn) or genome coordinates, and also to view all corresponding sgRNA designs on a genome browser (Fig. S2). To facilitate the use of sgRNAs in *Drosophila*, we have established a freely available online resource of precomputed sgRNAs (www.flyrnai.org/crispr) (Table 4). The designs are organized into subset “tracks” based on off-target potential. Moreover, a detail page that includes a summary list of potential off-targets can be viewed for each design.

Our simplified Cas9/sgRNA system will facilitate the generation of large mutant collections, including both loss-of-function indel mutations and defined deletions. Fig. S3 shows a flowchart designed to expedite the production of such collections in a semi-high-throughput manner. In addition to generation of deletions via NHEJ after production of DSB, based on a previous report (16), we expect that using our system with a donor DNA will facilitate HR-mediated knock-in approaches, such as the introduction of visible markers. This would expand the applications of our approach, including easier mutation identification (e.g., after knock-in of a marker), introduction of site-directed mutagenesis in endogenous loci, and in-frame insertion of ectopic tags into genomic sequence to detect gene expression patterns.

Materials and Methods

Cas9 and sgRNA Plasmid Design. To express Cas9 in *Drosophila* germ cells, we constructed a *nos* cassette plasmid containing approximately 700 base pairs of the *nos* promoter, the *nos* 5' UTR, and the *nos* 3' UTR. The attB donor sequence and a *vermillion*⁺ marker were also included in the *nos* cassette plasmid. The Cas9 coding sequence was then excised from pDNA3.1-Cas9 (25) and cloned into the XbaI and AvrII restriction sites of the *nos* cassette plasmid.

Two sgRNA scaffolds were designed: a short scaffold, as described by Mali et al. (10), and a long scaffold with 10 extra base pairs between the crRNA and the tracrRNA modules (Fig. 1C). To express sgRNA in vivo, we selected three different regulatory sequences: *U6a* (CR31379) and *U6b* (CR32867) (26, 27), both of which direct transcription of *Drosophila* U6 snRNAs, and *nos-mini*, which contains the *nos* promoter and a minimal 5' UTR (28, 29). We used the 404-bp upstream and 93-bp downstream regulatory sequences of *U6a* to construct *U6a-sgRNA-short* and *U6a-sgRNA-long*. Similarly, we used the 400-bp upstream and 95-bp downstream regulatory sequences of *U6b* to construct *U6b-sgRNA-short* and *U6b-sgRNA-long*. The *nos* 700-bp promoter, 80-bp minimal 5' UTR, and 900-bp 3' UTR were used to construct *nos-minisgRNA-short* and *nos-minisgRNA-long*. In all six versions of the sgRNA plasmid, the 20-bp target sequence is inserted into two BbsI sites at the beginning of the sgRNA scaffold. A 680-bp fragment was placed between the BbsI sites as a marker for detecting incomplete enzyme digestion. The sequence of the *nos-Cas9* plasmid is shown in Fig. S4, and partial sequences of the six different versions of the sgRNA constructs used are shown in Fig. S5.

Fly Stocks. All flies were cultured on standard cornmeal food at 25 °C. The *{nos-Cas9}attP40* and *{nos-Cas9}attP2* fly stocks were established according to a previously described protocol (30). In brief, *nos-Cas9* was inserted into *attP40* on the second chromosome or *attP2* on the third chromosome by microinjection into *y[1] sc[1] v[1] P{y[+t.7]}=nos-phiC31\int.NLS}X; P{y[+t.7]}=CaryP}attP40* or *y[1] sc[1] v[1] P{y[+t.7]}=nos-phiC31\int.NLS}X; P{y[+t.7]}=CaryP}attP2* fly embryos, respectively, followed by screening for the

Table 4. Statistics for the sgRNA resource

CRISPR category	All CRISPRs	No off-target	No off-target in CDS regions	Off-target in CDS regions
All CRISPRs	7,566,724	300,034	2,894,160	4,372,530
CRISPRs target intergenic region	2,110,255	76,182	907,314	1,126,759
CRISPRs target gene region	5,456,469	223,852	1,985,158	3,247,459
CRISPRs target intron region	2,786,650	95,393	1,231,172	1,460,085
CRISPRs target exon region	2,669,819	128,459	753,986	1,787,374
CRISPRs target exon-CDS region	2,107,960	104,892	506,998	1,496,070
CRISPRs target exon-noncoding region	561,859	23,567	246,988	291,304

vermillion⁺ marker present in the *nos-Cas9* plasmid. *Canton-S* and *y[1]w[67c23]* flies were used.

Genomic DNA Extraction. Fly genomic DNA was purified via phenol-chloroform extraction. Single flies were homogenized in 400 μ L of lysis buffer (1X PBS, 0.2% SDS, and 200 μ g/mL proteinase K; Roche) and incubated at 50 °C for 1 h, followed by extraction in 400 μ L of phenol-chloroform. The mixture was then centrifuged at 21,000 X g for 20 min at 4 °C, after which the supernatant was transferred to a new tube. An equal volume of isopropanol was added, and the tube was vortexed thoroughly. The mixture was then kept at –20 °C for at least 1 h, followed by centrifugation at 21,000 X g for 20 min at 4 °C. The supernatant was removed, and the pellet was washed with 500 μ L of 75% ethanol, followed by centrifugation at 21,000 X g for 5 min at 4 °C. Finally, the pellet was dried for 10 min and resuspended in 30 μ L of DNase-free water.

Embryo Injection. Plasmid DNA was microinjected into *Drosophila* embryos following standard protocols. To mutagenize flies with Cas9 and sgRNA, the *nos-Cas9* plasmid (250 ng/ μ L) and sgRNA plasmid (250 ng/ μ L) were coinjected as a mixture. For coinjection of the plasmids, the injection solution consisted of 250 ng/ μ L of the *nos-Cas9* plasmid and 250 ng/ μ L of each sgRNA plasmid. When using transgenic Cas9 flies, a single sgRNA plasmid at 250 ng/ μ L was injected or two sgRNA plasmids at 250 ng/ μ L each were coinjected.

Screening of Mutations. To score for germ-line mutations, all G0 adult flies that developed from injected embryos were crossed to *y[1]w[67c23]* flies. The F1 progeny were then screened for white eye color over a 6-d period. If the G0 fly was a female, then all progeny were screened. If the G0 fly was a male, then only female progeny were screened. Individual heritable mutation rates were calculated for each single G0 as the number of mutant F1s out of the number of screened progeny. The overall heritable mutation rate was calculated as the number of all mutant F1s out of the number of all screened progeny for a given sgRNA target. Mutagenesis events were confirmed by sequence analysis of F1 adults; the detection primers are listed in Table S2.

Off-Target Analysis. To investigate the possibility of off-target cleavage by Cas9/sgRNA, we searched the fly genome for potential off-targets containing a match to the sgRNA sequence of at least 11 nt followed by the PAM sequence, as well as a match of 13 or 14 nt but without a neighboring PAM sequence. Primers flanking the potential off-targets were used to PCR-amplify these regions for analysis by HRMA and sequencing. For sequencing analysis, genomic DNA from a single fly was used as template, and the defined DNA fragment was amplified by specific primers carrying EcoRI and XbaI (Table S2). The PCR products can be directly sequenced by one of the

primers or cloned into the linearized VALIUM1 vector (30) by the same set of restriction enzymes, and then sequenced by specific primers (Table S2).

HRMA. For both on-target and off-target HRMA assays, PCR was performed using specific primers (Table S2, off-target sequencing primers) to amplify regions surrounding the target site from genomic DNA isolated from either injected embryos (for on-target analysis) or F1 adults (for off-target analysis). PCR products were diluted 1:500,000 before a second round of PCR using nested primers (Table S2, off-target HRM primers) to generate short amplicons (70–150 bp) including the target sites. Melting curves were generated by heating from 55 °C to 95 °C and measuring fluorescence in the presence of Evagreen DNA dye at increments of 0.1 °C. Both amplification and melt curve steps were performed using a BioRad CFX Real-Time System qPCR machine and BioRad Precision Melt Supermix. Genomic DNA isolated from the uninjected parent strain (*y[1] sc[1] v[1] P[y[+7.7]=nos-phiC31int.NLS]X; P[y[+7.7]=CaryP]attP2*) was used to generate control samples. HRMA data were analyzed to identify melt curves that differed significantly from controls using custom scripts that will be reported in detail separately. The HRMA analysis data are provided in Table S1.

RT-PCR. Total RNA was isolated from 0- to 1-h *Drosophila* embryos using the AxyPrep Multisource Total RNA MiniPrep Kit (Axygen). A total of 3 μ g RNA was used to create cDNA, using the GoldScript cDNA Kit (Invitrogen) according to the manufacturer's protocol, followed by PCR to amplify the target sequence. The *Actin 5C* gene served as an internal control. Primer sequences are listed in Table S2.

Assembly of Online Resource. A Perl module was developed in-house to identify all possible CRISPR sequences from both strands of genome sequences downloaded from FlyBase release 5.52 (July 16, 2013). The BLAST program was used to identify all possible off-target sites, and a second Perl module was developed in-house to annotate on-target and off-target sites, as well as to format GFF3 files to upload genome browse. JBrowse was used and configured to display all CRISPR designs.

ACKNOWLEDGMENTS. We thank Son Nguyen and Ting Wu for helpful discussion. This work was supported by the Tsinghua-Peking Center for Life Sciences, the Ministry of Science and Technology of China 973 Program (Grant 2013CB835100), the PhD Programs Foundation of the Chinese Ministry of Education (Grant 20121018577 and 20120002110056), and the National Institutes of Health, National Institute of General Medical Sciences (Grants R01 GM084947 and R01 GM067761, to N.P.). N.P. is an investigator of the Howard Hughes Medical Institute.

- Gaj T, Gersbach CA, Barbas CF, 3rd (2013) ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol* 31(7):397–405.
- Bibikova M, Golic M, Golic KG, Carroll D (2002) Targeted chromosomal cleavage and mutagenesis in *Drosophila* using zinc-finger nucleases. *Genetics* 161(3):1169–1175.
- Gao G, McMahon C, Chen J, Rong YS (2008) A powerful method combining homologous recombination and site-specific recombination for targeted mutagenesis in *Drosophila*. *Proc Natl Acad Sci USA* 105(37):13999–14004.
- Gong WJ, Golic KG (2003) Ends-out, or replacement, gene targeting in *Drosophila*. *Proc Natl Acad Sci USA* 100(5):2556–2561.
- Huang J, Zhou W, Dong W, Watson AM, Hong Y (2009) Directed, efficient, and versatile modifications of the *Drosophila* genome by genomic engineering. *Proc Natl Acad Sci USA* 106(20):8284–8289.
- Liu J, et al. (2012) Efficient and specific modifications of the *Drosophila* genome by means of an easy TALEN strategy. *J Genet Genomics* 39(5):209–215.
- Rong YS, Golic KG (2000) Gene targeting by homologous recombination in *Drosophila*. *Science* 288(5473):2013–2018.
- Cong L, et al. (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339(6121):819–823.
- Jinek M, et al. (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337(6096):816–821.
- Mali P, et al. (2013) RNA-guided human genome engineering via Cas9. *Science* 339(6121):823–826.
- Bhaya D, Davison M, Barrangou R (2011) CRISPR-Cas systems in bacteria and archaea: Versatile small RNAs for adaptive defense and regulation. *Annu Rev Genet* 45:273–297.
- Terns MP, Terns RM (2011) CRISPR-based adaptive immune systems. *Curr Opin Microbiol* 14(3):321–327.
- Wiedenheft B, Sternberg SH, Doudna JA (2012) RNA-guided genetic silencing systems in bacteria and archaea. *Nature* 482(7385):331–338.
- Bassett AR, Tibbit C, Ponting CP, Liu JL (2013) Highly efficient targeted mutagenesis of *Drosophila* with the CRISPR/Cas9 system. *Cell Rep* 4(1):220–228.
- Friedland AE, et al. (2013) Heritable genome editing in *C. elegans* via a CRISPR-Cas9 system. *Nat Methods* 10(8):741–743.
- Gratz SJ, et al. (2013) Genome engineering of *Drosophila* with the CRISPR RNA-guided Cas9 nuclease. *Genetics* 194(4):1029–1035.
- Hwang WY, et al. (2013) Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol* 31(3):227–229.
- Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA (2013) RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat Biotechnol* 31(3):233–239.
- Wang H, et al. (2013) One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* 153(4):910–918.
- Yu Z, et al. (2013) Highly efficient genome modifications mediated by CRISPR/Cas9 in *Drosophila*. *Genetics* 195(1):289–291.
- Venken KJ, Bellen HJ (2005) Emerging technologies for gene manipulation in *Drosophila melanogaster*. *Nat Rev Genet* 6(3):167–178.
- Venken KJ, Bellen HJ (2007) Transgenesis upgrades for *Drosophila melanogaster*. *Development* 134(20):3571–3584.
- Hsu PD, et al. (2013) DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol* 31(9):827–832.
- Fu Y, et al. (2013) High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. *Nat Biotechnol* 31(9):822–826.
- Chang N, et al. (2013) Genome editing with RNA-guided Cas9 nuclease in zebrafish embryos. *Cell Res* 23(4):465–472.
- Wakiyama M, Matsumoto T, Yokoyama S (2005) *Drosophila* U6 promoter-driven short hairpin RNAs effectively induce RNA interference in Schneider 2 cells. *Biochem Biophys Res Commun* 331(4):1163–1170.
- Das G, Henning D, Reddy R (1987) Structure, organization, and transcription of *Drosophila* U6 small nuclear RNA genes. *J Biol Chem* 262(3):1187–1193.
- Van Doren M, Williamson AL, Lehmann R (1998) Regulation of zygotic gene expression in *Drosophila* primordial germ cells. *Curr Biol* 8(4):243–246.
- Li Y, Minor NT, Park JK, McKearin DM, Maines JZ (2009) Bam and Bgcn antagonize Nanos-dependent germ-line stem cell maintenance. *Proc Natl Acad Sci USA* 106(23):9304–9309.
- Ni JQ, et al. (2008) Vector and parameters for targeted transgenic RNA interference in *Drosophila melanogaster*. *Nat Methods* 5(1):49–51.