

# Identifying multiple causative genes at a single GWAS locus

Michael J. Flister,<sup>1,2</sup> Shirng-Wern Tsaih,<sup>1</sup> Caitlin C. O'Meara,<sup>3</sup> Bradley Endres,<sup>1,2</sup> Matthew J. Hoffman,<sup>1,2</sup> Aron M. Geurts,<sup>1,2</sup> Melinda R. Dwinell,<sup>1,2</sup> Jozef Lazar,<sup>1,4</sup> Howard J. Jacob,<sup>1,2,5,6</sup> and Carol Moreno<sup>1,2</sup>

<sup>1</sup>Human and Molecular Genetics Center, Medical College of Wisconsin, Milwaukee, Wisconsin 53226, USA; <sup>2</sup>Department of Physiology, Medical College of Wisconsin, Milwaukee, Wisconsin 53226, USA; <sup>3</sup>Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA; <sup>4</sup>Department of Dermatology, <sup>5</sup>Department of Pediatrics, Medical College of Wisconsin, Milwaukee, Wisconsin 53226, USA

Genome-wide association studies (GWAS) are useful for nominating candidate genes, but typically are unable to establish disease causality or differentiate between the effects of variants in linkage disequilibrium (LD). Additionally, some GWAS loci might contain multiple causative variants or genes that contribute to the overall disease susceptibility at a single locus. However, the majority of current GWAS lack the statistical power to test whether multiple causative genes underlie the same locus, prompting us to adopt an alternative approach to testing multiple GWAS genes empirically. We used gene targeting in a disease-susceptible rat model of genetic hypertension to test all six genes at the *Agtrap-Plod1* locus (*Agtrap*, *Mthfr*, *Cln6*, *Nppa*, *Nppb*, and *Plod1*) for blood pressure (BP) and renal phenotypes. This revealed that the majority of genes at this locus (five out of six) can impact hypertension by modifying BP and renal phenotypes. Mutations of *Nppa*, *Plod1*, and *Mthfr* increased disease susceptibility, whereas *Agtrap* and *Cln6* mutations decreased hypertension risk. Reanalysis of the human *AGTRAP-PLOD1* locus also implied that disease-associated haplotype blocks with polygenic effects were not only possible, but rather were highly plausible. Combined, these data demonstrate for the first time that multiple modifiers of hypertension can cosegregate at a single GWAS locus.

[Supplemental material is available for this article.]

Genome-wide association studies (GWAS) are able to nominate loci for complex diseases, but are largely unable to identify the causative variant(s) for multiple reasons (Zuk et al. 2012; Chatterjee et al. 2013). First, GWAS loci typically contain several candidate genes and are unable to distinguish between the effects of SNPs in linkage disequilibrium (LD). Second, the possibility exists that multiple causative SNPs influence the overall phenotypic variance at a single GWAS locus (Yang et al. 2012); however, most GWAS lack the statistical power to test this possibility due to sample size (Zuk et al. 2012). For example, Zuk et al. (2012) estimated that association of a single SNP at a relatively strong GWAS locus requiring 4900 test subjects (50% power and genome-wide significance level of  $\alpha = 5 \times 10^{-8}$ ) would require ~450,000 individuals to achieve the same power when analyzing two SNPs simultaneously. Other estimates suggest that the full predictive power of many GWAS might not be met until sample sizes increase to  $>10^6$  individuals (Chatterjee et al. 2013). Of the handful of loci that do show multi-SNP effects (Galameau et al. 2010; Lango Allen et al. 2010; Ripke et al. 2011; Sanna et al. 2011; Sklar et al. 2011; Yang et al. 2012), all are of readily quantifiable traits with high heritability and low experimental variability (e.g., height), whereas none have been identified in more complex disease phenotypes (e.g., hypertension and renal disease). This raises the question whether genetic interactions within GWAS loci could be far more pervasive, but are simply missed by underpowered GWAS or lost in the noise of more complex disease phenotypes.

Here, we aimed to push past the limitations of current GWAS to identify causative gene(s) and test whether multiple genes within a GWAS locus contribute to the overall disease variance. We phenotyped rats with mutations in the *Agtrap-Plod1* locus, which has been associated with blood pressure (BP) or renal disease in 11 human studies (Supplemental Table 1; Kato et al. 2000; Jiang et al. 2004; Zhang et al. 2005; Levy et al. 2009; Newton-Cheh et al. 2009a,b; Chen et al. 2010; Tomaszewski et al. 2010; Johnson et al. 2011; Liu et al. 2011; Fung et al. 2012), yet the genetic mechanism(s) underlying this locus were completely unknown. We used a recent technological advancement—zinc-finger nuclease (ZFN) mutagenesis (Geurts et al. 2009)—to introduce damaged alleles into each of the six *Agtrap-Plod1* genes (*Agtrap*, *Mthfr*, *Cln6*, *Nppa*, *Nppb*, and *Plod1*) in a rodent model of genetic hypertension, the SS (SS/JrHsdMcwi) rat. The uniform genetic background allowed us to assign (+), (−), or (=) phenotypic effects to each of the genes at the *Agtrap-Plod1* locus, which is not possible using mouse knockout models with mixed genetic backgrounds that confound inter-strain comparisons due to genetic heterogeneity (Hunter 2012). Thus, by leveraging the genetic homogeneity of our SS strains, we were able to attribute the phenotypic contributions of multiple genes at the *Agtrap-Plod1* locus that were previously unattainable in human population studies. We then retrospectively analyzed

## Corresponding author E-mail jacob@mcw.edu

Article published online before print. Article, supplemental material, and publication date are at <http://www.genome.org/cgi/doi/10.1101/gr.160283.113>.

© 2013 Flister et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 3.0 Unported), as described at <http://creativecommons.org/licenses/by-nc/3.0/>.

human data sets (e.g., HapMap, 1000 Genomes, and ENCODE) to demonstrate that interplay between multiple causative genes within the human *AGTRAP-PLOD1* locus were not only possible, but rather were highly plausible.

## Results

### Strain generation and phenotyping strategy

We generated six mutant SS strains (*Agtrap*, *Mthfr*, *Clcn6*, *Nppa*, *Nppb*, and *Plod1*) by ZFN mutagenesis for phenotypic comparison with wild-type (WT) littermates. The target sequences for ZFNs, the sequence of mutant alleles, and the predicted functional consequences are provided in Supplemental Table 2. All ZFN-targeted alleles were predicted to cause deleterious mutations by frameshift and/or by disrupting a functional domain of the protein (Supplemental Table 2). Strains were tested for eight cardiovascular disease (CVD)-associated phenotypes following a 10-d high-salt challenge (see Methods and Table 1). Because BP and renal phenotypes are most commonly reported in human studies (Kato et al. 2000; Jiang et al. 2004; Zhang et al. 2005; Levy et al. 2009; Newton-Cheh et al. 2009a,b; Chen et al. 2010; Tomaszewski et al. 2010; Johnson et al. 2011; Liu et al. 2011; Fung et al. 2012), we focused specifically below on the (+), (−), and (=) effects of mutant alleles on BP and urinary protein excretion (a commonly used index for renal damage).

### Blood pressure phenotypes

To test BP, 8- to 9-wk-old males from each gene-targeted strain and WT littermates were placed on a 4% NaCl diet for 10 d, and BP was recorded by radiotelemetry. Of note, all rat strains developed hypertension (>140/90 mmHg) in response to a 4% NaCl diet (Fig. 1). Only *Plod1*, *Nppa*, and *Clcn6* mutations significantly modified BP (i.e., the severity of hypertension) in the SS rat on a 4% NaCl diet compared with WT (Fig. 1A–C). The *Nppa* mutation increased mean arterial pressure (MAP) ( $\Delta + 27$  mmHg,  $P < 0.001$ ), systolic BP (SBP) ( $\Delta + 46$  mmHg,  $P < 0.001$ ), and diastolic BP (DBP) ( $\Delta + 20$  mmHg,  $P < 0.05$ ) compared with WT. The BP changes coincided with increased heart weight ( $1.62 \pm 0.03$  g vs.  $1.13 \pm 0.03$  g,  $P < 0.001$ ), but did not significantly change renal phenotypes (Table 1). This is consistent with increased MAP and heart weight in *Nppa* knockout mice (John et al. 1995) and with decreased circulating NPPA levels associated with human hypertension (Marques et al. 2010). Compared with WT, *Plod1* mutation significantly increased only SBP ( $\Delta + 17$  mmHg,  $P < 0.01$ ), whereas *Clcn6* mutation decreased DBP ( $\Delta - 22$  mmHg,  $P < 0.001$ ) and MAP ( $\Delta - 14$  mmHg,  $P < 0.05$ ) but did not significantly affect SBP (Fig. 1A–C). These data indicate that *NPPA*, *PLOD1*, and *CLCN6* could have additive or subtractive effects on BP in human, depending on the mode of inheritance and functional consequence of the causal variant(s) (i.e., gain or loss of function).

### Renal phenotypes

The *AGTRAP-PLOD1* locus has also been linked with renal function (Supplemental Table 1) but the causative gene(s) are unknown, prompting us to determine if renal phenotypes were secondary to hypertension or independent but residing within the same LD block. On a 4% NaCl diet, urinary protein excretion was increased by mutation of *Plod1* ( $147 \pm 12$  mg/day,  $P < 0.001$ ) and *Mthfr* ( $132 \pm 23$  mg/day,  $P < 0.05$ ) compared with WT ( $91 \pm 5$  mg/day), whereas mutation of *Clcn6* ( $56 \pm 5$  mg/day,  $P < 0.05$ ) and *Agtrap* ( $54 \pm 9$  mg/day,  $P < 0.05$ ) attenuated proteinuria (Fig. 1D). Compared with WT, the changes in proteinuria in *Plod1* and *Clcn6*

mutant rats are likely secondary to BP (Fig. 1A–C). In contrast, the BP in *Mthfr* and *Agtrap* mutant strains did not differ significantly from WT on a 4% NaCl diet (Fig. 1), indicating that the significant changes in proteinuria (relative to WT) were independent of BP in these strains. Of these genes, only *Mthfr* has been suggested as a mediator of renal damage, largely in response to homocysteine-mediated oxidative stress in the kidney (Yi and Li 2008). *Plod1* knockout mice are prone to aortic aneurism (Takaluoma et al. 2007); however, *Plod1* has not yet been associated with renal phenotypes. This is also the first report of a protective effect on proteinuria by *Agtrap*, a mediator of the renin-angiotensin system (Oppermann et al. 2010).

### Functional annotation of the human *AGTRAP-PLOD1* locus

Our data show that multiple genes in the *Agtrap-Plod1* locus modify BP and renal phenotypes in the SS rat (Fig. 1A–D), prompting us to test whether similar mechanisms could be inherited at the homologous human *AGTRAP-PLOD1* locus. First, we determined whether SNPs, genes, and reported phenotypes could be connected by LD in human using data from HapMap (The International HapMap Consortium 2003) and 1000 Genomes (The 1000 Genomes Project Consortium 2012). Genes were considered linked to the reported phenotype if an associated lead SNP or a SNP in LD ( $r^2 > 0.6$ , HapMap CEU population) caused a nonsynonymous mutation or was associated with a change in gene expression (Supplemental Table 3). When taking into account all 226 SNPs in LD, we found that four out of six genes (*MTHFR*, *CLCN6*, *NPPA*, and *NPPB*) can be linked to BP or renal phenotypes at the *AGTRAP-PLOD1* locus (Fig. 1E). Collectively, this strengthens the possibility that multiple modifiers of BP (*NPPA*, *PLOD1*, and *CLCN6*) and renal phenotypes (*MTHFR*, *PLOD1*, *CLCN6*, and *AGTRAP*) can cosegregate at the same GWAS locus.

Using ENCODE data (The ENCODE Project Consortium 2011), we tested more specifically whether CVD-associated haplotypes were likely to influence multiple genes at the *AGTRAP-PLOD1* locus, rather than the “best-fit” candidate gene that is typically reported. Across nine different haplotype blocks, three (1.3%) SNPs caused nonsynonymous mutations (nSNPs), 13 (5.8%) SNPs were linked to gene expression (eSNP), and 16 (7.1%) SNPs were predicted to interrupt a functional motif (fSNP), whereas 197 SNPs (85.8%) had limited functional evidence or no annotated function (Fig. 2A; Supplemental Table 3). Out of the nine total haplotype blocks, six were linked with differential expression or nSNPs in multiple genes (Fig. 2B; Supplemental Table 3). One BP-associated haplotype block (led by rs5063) contained two nSNPs, two eSNPs, and five fSNPs that are linked to *NPPA*, *NPPB*, *CLCN6*, and *MTHFR* (Fig. 2C). Another haplotype block contained two lead SNPs (rs1801131 and rs4846049) that were independently associated with BP and renal phenotypes (Fig. 2D). This haplotype block included one nSNP in *MTHFR* and five fSNPs that were associated with *MTHFR* and *CLCN6* expression (Supplemental Table 3). One fSNP in particular (rs12121543) is predicted to disrupt STAT1 binding in proximity to *MTHFR* and *CLCN6* regulatory regions (Fig. 2D), fitting with previously reported enrichment of STAT1 binding at the *AGTRAP-PLOD1* locus (Johnson et al. 2011).


## Discussion

The goal of this study was to identify mechanism(s) that underlie changes in BP and proteinuria at a GWAS locus (*AGTRAP-PLOD1*)

**Table 1.** Cardiovascular phenotypes of *Plod1*-*Agtrap* mutant rats

Trait	Strain												
	<i>Plod1</i>	<i>P-value</i>	<i>Nppb</i>	<i>P-value</i>	<i>Nppa</i>	<i>P-value</i>	<i>Cln6</i>	<i>P-value</i>	<i>Mthfr</i>	<i>P-value</i>	<i>Agtrap</i>	<i>P-value</i>	WT
MAP	143 ± 3	ns	133 ± 2	ns	163 ± 6	<0.001	123 ± 1	<0.05	130 ± 4	ns	131 ± 3	ns	136 ± 2
DBP	126 ± 5	ns	111 ± 3	ns	143 ± 5	<0.05	101 ± 1	<0.001	110 ± 4	ns	120 ± 5	ns	123 ± 3
SBP	165 ± 4	<0.01	155 ± 2	ns	195 ± 5	<0.001	149 ± 2	ns	152 ± 6	ns	139 ± 2	ns	149 ± 2
Heart rate	351 ± 5	ns	352 ± 10	ns	343 ± 5	ns	337 ± 6	ns	330 ± 12	ns	350 ± 6	ns	339 ± 4
Protein excretion	147 ± 12	<0.001	97 ± 16	ns	73 ± 8	ns	56 ± 5	<0.05	132 ± 23	<0.05	54 ± 9	<0.05	91 ± 5
Creatinine	9.0 ± 0.4	ns	11.5 ± 0.5	ns	10.1 ± 0.3	ns	8.3 ± 0.3	<0.01	12.1 ± 0.4	ns	10.3 ± 0.9	ns	11.5 ± 0.4
Na excretion	8.1 ± 0.7	ns	7.4 ± 1.1	ns	8.6 ± 1.0	ns	5.6 ± 0.9	<0.05	9.7 ± 1.2	ns	9.3 ± 1.0	ns	10.0 ± 0.6
Kidney weight	1.49 ± 0.06	ns	1.45 ± 0.02	ns	1.40 ± 0.04	ns	1.53 ± 0.04	ns	1.61 ± 0.08	ns	1.37 ± 0.05	ns	1.46 ± 0.02
Heart weight	1.14 ± 0.03	ns	1.20 ± 0.03	ns	1.62 ± 0.03	<0.001	1.27 ± 0.04	<0.01	1.24 ± 0.06	<0.05	1.08 ± 0.03	ns	1.13 ± 0.02
Body weight	298 ± 6	<0.05	330 ± 4	ns	299 ± 5	<0.05	282 ± 5	<0.001	333 ± 10	ns	327 ± 4	ns	321 ± 5

Data are presented as mean values ± SE (*n* = 8–25 animals per strain). *P*-values were determined by a one-way ANOVA followed by Holm-Sidak post-hoc test.



**Figure 1.** Mean arterial pressure (MAP) (A), systolic blood pressure (SBP) (B), and diastolic blood pressure (DBP) (C) measured by radiotelemetry in conscious WT and mutant SS rat strains on 4% NaCl diet ( $n = 8-25$  per strain). Data are presented as mean BP  $\pm$  SEM. (\*)  $P < 0.05$ , (\*\*)  $P < 0.01$ , and (\*\*\*)  $P < 0.001$  vs. WT as determined by one-way ANOVA followed by a Holm-Sidak post-hoc test. (D) Rats were placed in metabolic cages overnight to acclimate, followed by a 24-h urine collection. Data are presented as mean protein excretion  $\pm$  SEM ( $n = 8-25$  rats per strain). (\*)  $P < 0.05$  and (\*\*\*)  $P < 0.001$  vs. WT as determined by one-way ANOVA followed by a Holm-Sidak post-hoc test. (E) BP and renal damage are linked with multiple *AGTRAP-PLOD1* locus genes in human association studies and mutant SS rat strains. For human, genes were considered linked to a phenotype if an associated SNP or SNPs in LD ( $r^2 > 0.6$ ) with an associated SNP caused nonsynonymous mutations or were significantly associated with expression of a gene. Blue lines indicate decreased BP and/or proteinuria associated with a gene in the rat. Red lines indicate increased BP and/or proteinuria associated with a gene in the rat.

with high genetic complexity, as indicated by multiple independent associations with BP and renal phenotypes in the human population (Kato et al. 2000; Jiang et al. 2004; Zhang et al. 2005; Levy et al. 2009; Newton-Cheh et al. 2009a,b; Chen et al. 2010; Tomaszewski et al. 2010; Johnson et al. 2011; Liu et al. 2011; Fung et al. 2012). By combining rat phenotype data (Fig. 1; Table 1) with multiple human data sets (Supplemental Tables 1, 3), we were able to draw previously unforeseen conclusions, foremost that multiple genes at a single GWAS locus can influence clinically relevant hypertension phenotypes. Although this has been implied statistically by GWAS for other traits previously (Yang et al. 2012), it had not been tested experimentally in a physiological setting for hypertension. Specific to this locus, we present the first evidence that *Nppa*, *Clcn6*, *Mthfr*, *Plod1*, and *Agrap* mutations cause divergent CVD phenotypes and have the ability to modify renal phenotypes independently of BP (Fig. 1). Combined, these data suggest that the “best-fit” candidate gene interpretations of GWAS are in some cases only a simplified view of the complex genetic architecture underlying individually associated loci. Moreover, the complex combinations of alleles and haplotypes, due to genetic heterogeneity in human, would have the potential to impact the overall disease susceptibility and association of the *AGTRAP-PLOD1* locus with BP and increase the challenge of finding all causative variants.


#### Challenges to identifying hypertension loci with multiple causative variants

GWAS loci that contain multiple interacting alleles have not yet been reported for human hypertension. This is likely attributed to

several confounding factors that limit the sensitivity of GWAS for multi-SNP associations at a single locus. Firstly, current hypertension GWAS are likely too small (i.e., underpowered) to detect interplay between multiple causative genes at the same locus. Power limitations are due in part to a multi-SNP association requiring  $>10$  times the sample size to achieve the same statistical power of a single SNP association (Zuk et al. 2012). This can be further complicated by frequency and effect size of the individual SNPs (e.g., rare SNPs with large effect sizes will have more combined power). Secondly, BP measurements are notoriously variable due to patient stress (e.g., “white coat effect”) and other environmental factors (Mancia et al. 2009), which might artificially inflate or deflate BP readings and obscure true SNP associations. We propose that this “phenotypic noise” likely masks the detection of multi-SNP hypertension GWAS loci, and therefore multiple causative SNPs cannot be ruled out in the absence of systematic experimental testing, as we have demonstrated here.

#### Gene editing in disease sensitive rat strains—alternate approach to characterizing GWAS loci

We were able to test a “multi-SNP hypothesis” by limiting variability using inbred disease-susceptible ZFN rat strains in a controlled experimental setting. By introducing target gene mutations on an inbred SS background by gene editing with ZFNs, our rat strains are isogenic except for the target-gene mutation. Because these strains are otherwise identical, we are able to compare (+), (–), or (=) effects of target-genes on BP and renal phenotypes, without the potential influence of other genetic heterogeneity. In



**Figure 2.** (A) Proportions of functional consequences of all SNPs in LD ( $r^2 > 0.6$ ) with the SNPs associated with BP and renal phenotypes at the *AGTRAP-PLOD1* locus (see Supplemental Table 1). (B) Genes linked with lead SNPs of haplotypes associated with BP and renal phenotypes at the *AGTRAP-PLOD1* locus. A lead SNP was considered linked to a gene if the lead SNP or SNPs in LD ( $r^2 > 0.6$ ) with a lead SNP caused nonsynonymous mutations or were significantly associated with expression of a gene. (C) An example of a BP-associated nSNP in *NPPA* (rs5063) that is in LD ( $r^2 > 0.6$ ) with an nSNP in *MTHFR* (rs2274976). Both are also in LD with fSNPs correlated with the expression of *MTHFR*, *CLCN6*, and *NPPB*. Note that a total of two nSNPs (boxed), two eSNPs (blue), and five fSNPs (green) are in LD with rs5063. (D) An example of an fSNP (rs12121543) that is in LD with two SNPs (rs1801131 and rs4846049) associated with BP and renal phenotypes. An overview of the region between genes *MTHFR* and *CLCN6* shows that rs12121543 falls in a transcriptional active region (vertical black bar overlapping histone modification, DNase hypersensitivity, and ChIP-seq peaks). rs12121543 is predicted to change a conserved nucleotide in a consensus STAT1-binding site and is associated with significant differences in *MTHFR* and *CLCN6* expression compared with the major allele. (A, C, D) nSNP, nonsynonymous SNP; eSNP, expression SNP; and fSNP, functional SNP.

contrast, existing mouse KO models are on a variety of strain backgrounds and therefore carry substantial heterogeneity in addition to the target gene, which can influence comparison of phenotypes between models (Hunter 2012). Using gene editing on a disease-susceptible rat background also provides multiple endogenous variants that are required for pathogenesis of complex diseases, such as hypertension. In other words, all strains developed salt-sensitive hypertension due to the SS background, but to varying degrees and with differences in end-organ damage depending on the modifying effects of the *Agtrap-Plod1* locus (Fig. 1; Table 1).

Having first identified the causative genes underlying a GWAS locus by traditional ZFN-mediated gene editing (Fig. 1), we postulate that gene-editing technology (e.g., ZFN or TALEN) could then be used to ask more specific questions in future studies. Testing-combined haplotype effects could be achieved by editing multiple genes in the same pedigree. This would require sequential gene editing over multiple generations, as it is prohibitive to generate “multi-mutants” by cross-breeding together single mutants in close proximity. For example, recombination between two genes separated by only 150 kb (the approximate size of the *Agtrap-Plod1*

locus) would require >1000 offspring (assuming 1 Mb = 1 cM in the rat). It is also possible to humanize alleles using gene editing in the rat, which would likely be more representative of the hypomorphic alleles and smaller effect sizes observed in human GWAS. Thus, gene-editing technology could be used to test humanized alleles as single variants or whole haplotype blocks, which would likely be the gold standard to predicting human haplotype function in an experimental model of a complex disease.

Gene editing in the rat could also be used to explore additional factors that are generally treated as covariates in large GWAS, including genetic background, gender, and dietary or environmental stressors. To date, we have demonstrated the ability to edit genes on multiple backgrounds (e.g., FHH, SHR, and SS-BN<sup>13</sup> consomic) (<http://rgd.mcw.edu/wg/physgenknockouts>), enabling the possibility of testing GWAS-nominated haplotypes in experimental models with inherently different disease etiologies (e.g., high renin hypertension in SHR vs. low renin hypertension in SS). Testing other experimental parameters, such as diet (e.g., high fat, high glucose, or caloric restriction), environmental stressors (e.g., sleep deprivation), response to pharmacological inhibitors (e.g., ACE inhibitor or diuretics), and biomarkers (e.g., circulating *Nppa*

and homocysteine) will offer significant insight into the molecular underpinnings of GWAS loci. Finally, gender differences in genetic hypertension are becoming increasingly apparent, but not yet fully understood. Thus, future comparisons between males and females of gene-edited rat strains will also likely offer significant insight into gender discrepancies in hypertension risk that is associated with GWAS loci.

## Conclusions

Our data demonstrated for the first time that multiple causative alleles for BP and renal phenotypes can cosegregate at the same hypertension GWAS locus. Of the six *Agtrap-Plod1* genes, only three (*Agtrap*, *Nppa*, and *Mthfr*) were previously reported to have BP or renal phenotypes (John et al. 1995; Yi and Li 2008; Wakui et al. 2013), but none had been systematically examined in a uniform genetic background that is susceptible to hypertension. Using a ZFN rat strategy, we found that *Nppa*, *Clcn6*, *Mthfr*, *Plod1*, and *Agtrap* mutations caused divergent CVD phenotypes and have the ability to modify renal phenotypes independently of BP (Fig. 1). Additionally, *Clcn6* was identified here as a key mediator of BP and proteinuria for the first time. Differentiating this role from *MTHFR* was not previously possible using only human association data, because both genes share a common promoter that separates them by only 47 bp (Supplemental Fig. 1). Analysis of disease-associated human haplotypes suggested that several human haplotypes likely carry multiple causative SNPs. Combined, these data offer new insight into regulation of BP and renal phenotypes at the *Agtrap-Plod1* locus and provide rationale for empirically testing the individual and combined haplotype effects of human SNPs on *AGTRAP-PLOD1* gene function and overall association with CVD risk.

## Methods

### Generation of ZFN-mutated rat strains

Mutant SS rat strains were generated by microinjection of custom CompoZr ZFNs (Sigma) into one-cell SS/JrHsdMcwi (SS) rat embryos and then implanted into pseudopregnant Sprague Dawley females, as described previously (Geurts et al. 2010). Supplemental Table 2 provides the strain designations, target sites, DNA recognition helices of ZFNs, and description of gene mutation. F<sub>0</sub> generation animals were screened for ZFN-induced mutations by PCR amplification with the vendor-supplied primers (Supplemental Table 2) and assayed with the SURVEYOR Mutation Detection kit (Transgenomic, Inc.). Mutant alleles were then confirmed by Sanger sequencing. F<sub>0</sub> animals were back-crossed to the parental strain, and multiple pairs of N2 offspring were intercrossed to establish a breeding colony, thereby reducing any chance of off-target effects to <1%. Expression of mutant and WT alleles were assessed by RT-PCR using primers provided in Supplemental Figure 2.

### Blood pressure phenotyping

All strains were maintained on a 14/10-h light-dark cycle with ad libitum access to normal chow (Teklad Low Salt diet #7034, Harlan Laboratories, Inc.) and water. MAP of 8- to 9-wk-old male rats that had been placed on a 4% NaCl diet for 10 d were recorded by radiotelemetry at 10-sec intervals, every 2 min, and averaged over 4 h. Telemeters (TA11PA-C40, Data Sciences) were subcutaneously implanted with a catheter inserted into the abdominal aorta via the femoral artery.

### Renal function and serum biochemistry

Immediately after MAP recordings, rats were placed in metabolic cages and allowed to acclimate for 24 h, followed by a 24-h urine collection. Urine samples were then cleared of insoluble particulate by centrifugation at 3500 rpm and measured for total protein by Bradford assay (Bio-Rad). Sodium and potassium were measured by flame photometry (IL943, Instrumentation Laboratory); creatinine was measured by autoanalyzer (VET ACE Alera, Alfa Wassermann) using the Jaffe method.

### Analysis of human SNPs and haplotype blocks

LD (defined here as  $r^2 > 0.6$ ) for human SNPs from the CEU cohort from the HapMap (<http://hapmap.ncbi.nlm.nih.gov/>; The International HapMap Consortium 2003) and 1000 genome (<http://browser.1000genomes.org/index.html>; The 1000 Genomes Project Consortium 2012) projects were calculated using SNAP (<http://www.broadinstitute.org/mpg/snap/ldsearch.php>; Johnson et al. 2008). SNPs associated with gene regulatory mechanisms were curated from the ENCODE study (The ENCODE Project Consortium 2011) using the RegulomeDB database (<http://www.regulomedb.org/>; Boyle et al. 2012).

### Statistical analysis

Statistical analyses were performed using Sigma Plot 11.0 software. Data are presented as mean  $\pm$  SE. BP and renal phenotypes of mutant strains were compared with WT (control) by one-way ANOVA followed by a Holm-Sidak post-hoc test. Because MAP, SBP, DBP, and proteinuria failed the equal variance test, these data were log-transformed before one-way ANOVA followed by a Holm-Sidak post-hoc test. For all statistical tests,  $P < 0.05$  was considered significant.

### Competing interest statement

The Medical College of Wisconsin could one day receive royalties on sales of genetically modified rat strains through a license agreement with Sigma Advanced Genetic Engineering (SAGE).

### Acknowledgments

We thank R. Schilling, J. Klotz, M. Grzybowski, S. Kalloway, J. Foeckler, C. Hansen, and M. Casati for excellent technical support. We also acknowledge all members of the PhysGen Knockout team. M.J.F. is supported by an NHLBI training grant (5T32HL007792). A.M.G. is supported by a NIH Director's New Innovator Award (DP2OD008396). This study was supported by NHLBI grant 5RC2HL101681 to H.J.J.

### References

- The 1000 Genomes Project Consortium. 2012. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**: 56–65.
- Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, Karczewski KJ, Park J, Hitz BC, Weng S, et al. 2012. Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* **22**: 1790–1797.
- Chatterjee N, Wheeler B, Sampson J, Hartge P, Chanock SJ, Park JH. 2013. Projecting the performance of risk prediction based on polygenic analyses of genome-wide association studies. *Nat Genet* **45**: 400–405.
- Chen M, Chen X, Guo Y, Shi R, Zhang G. 2010. Brain natriuretic peptide rs198388 polymorphism and essential hypertension in Hunan Han people. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* **35**: 1207–1213.

- The ENCODE Project Consortium. 2011. A user's guide to the encyclopedia of DNA elements (ENCODE). *PLoS Biol* **9**: e1001046.
- Fung MM, Salem RM, Lipkowitz MS, Bhatnagar V, Pandey B, Schork NJ, O'Connor DT. 2012. Methylenetetrahydrofolate reductase (MTHFR) polymorphism A1298C (Glu429Ala) predicts decline in renal function over time in the African-American Study of Kidney Disease and Hypertension (AASK) Trial and Veterans Affairs Hypertension Cohort (VAHC). *Nephrol Dial Transplant* **27**: 197–205.
- Galarneau G, Palmer CD, Sankaran VG, Orkin SH, Hirschhorn JN, Lettre G. 2010. Fine-mapping at three loci known to affect fetal hemoglobin levels explains additional genetic variation. *Nat Genet* **42**: 1049–1051.
- Geurts AM, Cost GJ, Freyvert Y, Zeitler B, Miller JC, Choi VM, Jenkins SS, Wood A, Cui X, Meng X, et al. 2009. Knockout rats via embryo microinjection of zinc-finger nucleases. *Science* **325**: 433.
- Geurts AM, Cost GJ, Remy S, Cui X, Tesson L, Usal C, Menoret S, Jacob HJ, Anegón I, Buelow R. 2010. Generation of gene-specific mutated rats using zinc-finger nucleases. *Methods Mol Biol* **597**: 211–225.
- Hunter KW. 2012. Mouse models of cancer: Does the strain matter? *Nat Rev Cancer* **12**: 144–149.
- The International HapMap Consortium. 2003. The International HapMap Project. *Nature* **426**: 789–796.
- Jiang S, Hsu YH, Xu X, Xing H, Chen C, Niu T, Zhang Y, Peng S. 2004. The C677T polymorphism of the methylenetetrahydrofolate reductase gene is associated with the level of decrease on diastolic blood pressure in essential hypertension patients treated by angiotensin-converting enzyme inhibitor. *Thromb Res* **113**: 361–369.
- John SW, Kregge JH, Oliver PM, Hagaman JR, Hodgins JB, Pang SC, Flynn TG, Smithies O. 1995. Genetic decreases in atrial natriuretic peptide and salt-sensitive hypertension. *Science* **267**: 679–681.
- Johnson AD, Handsaker RE, Pulit S, Nizzari MM, O'Donnell CJ, de Bakker PIW. 2008. SNAP: A web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* **24**: 2938–2939.
- Johnson T, Gaunt TR, Newhouse SJ, Padmanabhan S, Tomaszewski M, Kumari M, Morris RW, Tzoulaki I, O'Brien ET, Poulter NR, et al. 2011. Blood pressure loci identified with a gene-centric array. *Am J Hum Genet* **89**: 688–700.
- Kato N, Sugiyama T, Morita H, Nabika T, Kurihara H, Yamori Y, Yazaki Y. 2000. Genetic analysis of the atrial natriuretic peptide gene in essential hypertension. *Clin Sci (Lond)* **98**: 251–258.
- Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, Willer CJ, Jackson AU, Vedantam S, Raychaudhuri S, et al. 2010. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* **467**: 832–838.
- Levy D, Ehret GB, Rice K, Verwoert GC, Launer LJ, Dehghan A, Glazer NL, Morrison AC, Johnson AD, Aspelund T, et al. 2009. Genome-wide association study of blood pressure and hypertension. *Nat Genet* **41**: 677–687.
- Liu C, Li H, Qi Q, Lu L, Gan W, Loos RJ, Lin X. 2011. Common variants in or near *FGF5*, *CYP17A1* and *MTHFR* genes are associated with blood pressure and hypertension in Chinese Hans. *J Hypertens* **29**: 70–75.
- Mancia G, Bombelli M, Facchetti R, Madotto F, Quarti-Trevano F, Polo Friz H, Grassi G, Sega R. 2009. Long-term risk of sustained hypertension in white-coat or masked hypertension. *Hypertension* **54**: 226–232.
- Marques FZ, Campain AE, Yang YH, Morris BJ. 2010. Meta-analysis of genome-wide gene expression differences in onset and maintenance phases of genetic hypertension. *Hypertension* **56**: 319–324.
- Newton-Cheh C, Johnson T, Gateva V, Tobin MD, Bochud M, Coin L, Najjar SS, Zhao JH, Heath SC, Eyheramendy S, et al. 2009a. Genome-wide association study identifies eight loci associated with blood pressure. *Nat Genet* **41**: 666–676.
- Newton-Cheh C, Larson MG, Vasán RS, Levy D, Bloch KD, Surti A, Guiducci C, Kathiresan S, Benjamin EJ, Struck J, et al. 2009b. Association of common variants in NPPA and NPPB with circulating natriuretic peptides and blood pressure. *Nat Genet* **41**: 348–353.
- Oppermann M, Gess B, Schweda F, Castrop H. 2010. Atrap deficiency increases arterial blood pressure and plasma volume. *J Am Soc Nephrol* **21**: 468–477.
- Ripke S, Sanders AR, Kendler KS, Levinson DF, Sklar P, Holmans PA, Lin DY, Duan J, Ophoff RA, Andreassen OA, et al. 2011. Genome-wide association study identifies five new schizophrenia loci. *Nat Genet* **43**: 969–976.
- Sanna S, Li B, Mulas A, Sidore C, Kang HM, Jackson AU, Piras MG, Usala G, Maninchedda G, Sassu A, et al. 2011. Fine mapping of five loci associated with low-density lipoprotein cholesterol detects variants that double the explained heritability. *PLoS Genet* **7**: e1002198.
- Sklar P, Ripke S, Scott LJ, Andreassen OA, Cichon S, Craddock N, Edenberg HJJ, Nurnberger JI, Rietschel M, Blackwood D, et al. 2011. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near *ODZ4*. *Nat Genet* **43**: 977–983.
- Takaluoma K, Hyry M, Lantto J, Sormunen R, Bank RA, Kivirikko KI, Myllyharju J, Soininen R. 2007. Tissue-specific changes in the hydroxylysine content and cross-links of collagens and alterations in fibril morphology in lysyl hydroxylase 1 knock-out mice. *J Biol Chem* **282**: 6588–6596.
- Tomaszewski M, Debiec R, Braund PS, Nelson CP, Hardwick R, Christofidou P, Denniff M, Codd V, Rafelt S, van der Harst P, et al. 2010. Genetic architecture of ambulatory blood pressure in the general population: Insights from cardiovascular gene-centric array. *Hypertension* **56**: 1069–1076.
- Wakui H, Tamura K, Masuda SI, Tsurumi-Ikeya Y, Fujita M, Maeda A, Ohsawa M, Azushima K, Uneda K, Matsuda M, et al. 2013. Enhanced angiotensin receptor-associated protein in renal tubule suppresses angiotensin-dependent hypertension. *Hypertension* **61**: 1203–1210.
- Yang J, Ferreira T, Morris AP, Medland SE, Madden PA, Heath AC, Martin NG, Montgomery GW, Weedon MN, Loos RJ, et al. 2012. Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat Genet* **44**: 369–375, S361–S363.
- Yi F, Li PL. 2008. Mechanisms of homocysteine-induced glomerular injury and sclerosis. *Am J Nephrol* **28**: 254–264.
- Zhang S, Mao G, Zhang Y, Tang G, Wen Y, Hong X, Jiang S, Yu Y, Xu X. 2005. Association between human atrial natriuretic peptide Val7Met polymorphism and baseline blood pressure, plasma trough irbesartan concentrations, and the antihypertensive efficacy of irbesartan in rural Chinese patients with essential hypertension. *Clin Ther* **27**: 1774–1784.
- Zuk O, Hechter E, Sunyaev SR, Lander ES. 2012. The mystery of missing heritability: Genetic interactions create phantom heritability. *Proc Natl Acad Sci* **109**: 1193–1198.

Received May 11, 2013; accepted in revised form September 3, 2013.