



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Human bocavirus 1 infects commercially available primary human airway epithelium cultures productively



Xuefeng Deng^a, Yi Li^b, Jianming Qiu^{a,*}

^a Department of Microbiology, Molecular Genetics and Immunology, University of Kansas Medical Center, Kansas City, KS, USA

^b Bioengineering Department, Wuhan Bioengineering Institute, Luoyang Economy Development Zone, Wuhan, China

ABSTRACT

Article history:

Received 31 July 2013

Received in revised form

29 September 2013

Accepted 1 October 2013

Available online 14 October 2013

Keywords:

Parvovirus

Human bocavirus

Human bocavirus 1 (HBoV1), a human parvovirus, belongs to the genus *Bocavirus* of the *Parvoviridae* family. It causes wheezing in young children with acute respiratory tract infections. HBoV1 has been shown to infect polarized human airway epithelium (HAE) made in house, and induces airway epithelial damage. In this study, two commercially available HAE cultures, EpiAirway and MucilAir HAE, were examined for HBoV1 infection. Both HAE cultures support fully productive HBoV1 infection. Infected EpiAirway and MucilAir HAE cultures showed loss of cilia, disruption of the tight junction barrier, and a significant decrease in transepithelial electrical resistance. Notably, HBoV1 persistent infection was demonstrated by maintaining HBoV1-infected EpiAirway HAE for as long as 50 days. After 2 days post-infection, progeny virus was produced consistently daily at a level of over 2×10^8 viral genome copies per culture (0.6 cm^2). This study is the first to use commercial sources of HAE cultures for HBoV1 infection. The availability of these cultures will enable a wide range of laboratories to study HBoV1 infection.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Human bocavirus 1 (HBoV1) was discovered in 2005 by large scale sequencing in nasopharyngeal aspirates from young children (Allander et al., 2005). It is a respiratory virus associated with acute respiratory tract infections in young children (Allander et al., 2007; Brodzinski and Ruddy, 2009; Don et al., 2010; Garcia-Garcia et al., 2010; Gendrel et al., 2007; Jartti et al., 2011; Kahn, 2008; Proenca-Modena et al., 2011; Schildgen et al., 2008). Acute respiratory tract infection is one of the leading causes of hospitalization of young children in developed countries (Brodzinski and Ruddy, 2009; Lopez et al., 2006; Shay et al., 1999). HBoV1 infection shows frequently persistence and co-infections with other respiratory viruses (Jartti et al., 2011; Kesebir et al., 2006; Manning et al., 2006; Wang et al., 2010). However, acute HBoV1 infection, diagnosed by a high virus load in respiratory samples, viremia, detection of HBoV1-specific IgM, or an increase in the levels of HBoV1-specific IgG antibodies, and detection of HBoV1 mRNA in nasopharyngeal aspirates, results in respiratory illness (Allander et al., 2007; Christensen et al., 2010, 2013; Deng et al., 2012; Don et al., 2010; Kantola et al., 2008; Nascimento-Carvalho et al.,

2012; Soderlund-Venermo et al., 2009; Wang et al., 2010). Life-threatening HBoV1 infections in pediatric patients, which were associated with high virus loads or diagnostic HBoV1-specific antibodies (Edner et al., 2011; Korner et al., 2011; Ursic et al., 2011), have also been described. Moreover, a recent longitudinal study in children (from infants to puberty) documented a clear association between acute primary HBoV1 infection and respiratory symptoms (Meriluoto et al., 2012).

The virus was tentatively classified as a member of genus *Bocavirus* in the subfamily *Parvovirinae* of the *Parvoviridae* family (Tijssen et al., 2012). Parvoviruses are small, non-enveloped icosahedral viruses with linear single-stranded DNA genomes. The genus *Bocavirus* includes bovine parvovirus type 1 (BPV1) (Chen et al., 1986), minute virus of canines (MVC) (Sun et al., 2009), and other tentative members that are closely related to HBoV1, including porcine bocavirus (Zhai et al., 2010), gorilla bocavirus (Kapoor et al., 2010a) and HBoV genotypes 2–4 (Arthur et al., 2009; Kapoor et al., 2009, 2010b).

A pseudostratified and well-differentiated primary human airway epithelium (HAE) culture model has been used widely to infect respiratory RNA viruses from the apical surface, e.g., influenza viruses (Ilyushina et al., 2012; Matrosovich et al., 2004; Zeng et al., 2011), parainfluenza virus (Zhang et al., 2005, 2011), respiratory syncytial virus (Villeneuve et al., 2012; Zhang et al., 2002), human coronaviruses (Pyrce et al., 2010; Sims et al., 2005; Wang et al., 2000), and human rhinovirus type C (Hao et al., 2012). Although respiratory DNA viruses have been reported to infect HAE, they infect HAE

* Corresponding author at: MS3029, University of Kansas Medical Center, 3901 Rainbow Boulevard, Kansas City, KS 66160, USA. Tel.: +1 913 588 4329; fax: +1 913 588 7295.

E-mail address: jqiu@kumc.edu (J. Qiu).

only from the basolateral surface, e.g., adenovirus (Zabner et al., 1997). In 2008, Dijkman et al. demonstrated that HBoV1 infects apically and replicates in HAE made in house (Dijkman et al., 2009). In 2012, an infectious clone of HBoV1 was established, which generates HBoV1 progeny virions from HEK293 cells transfected with this clone. Moreover, these HBoV1 virions infected HAE made in house productively, from the apical surface (Huang et al., 2012) as well as from the basolateral surface (Deng et al., 2013), which leads to airway epithelial damage.

In this report, two commercially available HAE cultures, i.e., EpiAirway and MucilAir HAE purchased from MatTek Co. (MA, USA) and Epithelix SàRL (Geneva, Switzerland), respectively, were tested for HBoV1 infection. Both HAE cultures can be infected by HBoV1 and caused hallmarks of airway epithelial damage.

2. Materials and methods

2.1. Polarized primary HAE cultures

EpiAirway HAE, which was purchased from MatTek (Ashland, MA, USA), was cultured in a Millicell insert of 0.6 cm² (Millipore, Billerica, MA, USA). MucilAir HAE was obtained from Epithelix SàRL (Geneva, Switzerland), and was maintained in a Costar Transwell insert of 0.33 cm² (Corning, NY, USA). Both HAE cultures were derived from healthy human primary tracheobronchial epithelial cells (derived from individual donors) cultured in an air-liquid interphase (ALI) with their respective property media.

2.2. Virus and infection

HBoV1 virions were collected from apical washes of HBoV1-infected primary B-HAE (Huang et al., 2012), and were used for infection as described previously (Deng et al., 2013).

For apical infection of EpiAirway HAE, HBoV1 virions were diluted in 200 μ l of EpiAirway medium to achieve an MOI of 100 viral genomic copy numbers (gc)/cell, and were applied to the apical chamber. For apical infection of MucilAir HAE, HBoV1 was diluted in 100 μ l of MucilAir medium to reach an MOI of 100 gc/cell. After incubation for 2 h, the apical chamber of the infected HAE ALI was washed three times with 400 μ l and 200 μ l of PBS for EpiAirway and MucilAir HAE, respectively. The cultures were maintained continuously at an ALI. To determine virus release kinetics, 200 μ l and 150 μ l aliquots of PBS, for EpiAirway and MucilAir HAE, respectively, were added to the apical chamber of the HAE culture at various time points, incubated for 15 min at 37 °C, and were harvested as apical washes. Meanwhile, an aliquot of 100 μ l of medium from the basolateral chamber of the HAE culture was collected, followed by replacement with 100 μ l of fresh medium. All the harvested aliquots were stored at 4 °C for quantification of viral DNA as gc.

For basolateral infection, HBoV1 virions were diluted in the ALI medium (1 ml and 0.5 ml for EpiAirway and MucilAir, respectively) in the basolateral chamber of the HAE cultures. The cultures were incubated at 37 °C and 5% CO₂ for 2 h, and then the basolateral inoculums were removed and washed three times with PBS (1 ml and 0.5 ml for EpiAirway and MucilAir, respectively), followed by supplementation of fresh media. Progeny virion release was monitored daily by quantification of viral gc in samples collected from the basolateral chamber at a volume of 100 μ l.

2.3. Real time quantitative PCR (qPCR) analysis of HBoV1 genome copy numbers

Aliquots of 100 μ l of apical sample and 50 μ l of basolateral sample, respectively, were incubated with 25 units of benzonase (Sigma, St. Louis, MO, USA) for 2 h at 37 °C. The treated samples were

digested with 20 μ l of proteinase K (15 mg/ml; Amresco, Solon, OH, USA) at 56 °C for 10 min. Viral DNA was extracted using the QIAamp blood mini kit (Qiagen, Valencia, CA, USA), and eluted in 100 μ l and 50 μ l of deionized H₂O for apical and basolateral samples, respectively. The extracted DNA samples were quantified with respect to the numbers of HBoV1 gc by the qPCR method described previously (Deng et al., 2013; Huang et al., 2012; Lin et al., 2007).

2.4. Immunofluorescence analysis

Immunofluorescence analysis of HBoV1-infected HAE was performed as described previously (Deng et al., 2013; Huang et al., 2012). A rat anti-HBoV1 NS1 polyclonal antibody (Chen et al., 2010) was developed previously in-house (Chen et al., 2010). Anti-ZO-1 (Invitrogen, Grand Island, NY, USA) and anti- β -tubulin IV (Sigma, St. Louis, MO, USA) antibodies were used for detecting the tight junction and cilia, respectively.

2.5. Measurement of the transepithelial electrical resistance (TEER)

The TEER of both mock- and HBoV1-infected HAE cultures was measured using an epithelial volt-ohm meter (Millipore, Billerica, MA, USA) at the indicated days p.i. as described previously (Deng et al., 2013; Huang et al., 2012).

3. Results

3.1. HBoV1 infects EpiAirway HAE persistently

EpiAirway HAE was first infected with the apically washed HBoV1 virions at a multiplicity of infection (MOI) of 100 gc/cell. Progeny virions released from the apical surface at 1×10^6 gc/ μ l at 3 days post-infection (p.i.), and remained as a plateau along with the infection for 50 days (Fig. 1A). Virus was not detected in mock-infected EpiAirway HAE. Since 200 μ l of phosphate buffered saline (pH 7.4; PBS) was used to wash the progeny virions from the apical surface every day, the total virus release from one HAE culture was approximately 2×10^8 gc per day. Virus was also released from the basolateral surface, but at a level of ~ 1 –2 log lower than that from the apical surface (Fig. 1A).

Consistent with the virus release kinetics, the transepithelial electrical resistance (TEER) of HBoV1-infected HAE started to drop off at 4 days p.i., gradually lowering to a level of 390 Ω cm² at 12 days p.i., 306 Ω cm² at 28 days p.i., and to a level close to 200 Ω cm² at 48 days p.i. (Fig. 1B). Compared with the mock-infected counterpart, the TEER decreased ~ 3.5 -fold by the end of infection (at 48 days p.i., Fig. 1B).

Of note, a few of the epithelial cells were washed off in the apical chamber of HBoV1-infected HAE, and many of which were NS1-positive (Fig. 1C), suggesting that HBoV1 infection induces epithelial cell death. By the end of the infection (at 50 days p.i.), β -tubulin IV (a marker of cilia (Matrosovich et al., 2004; Villenave et al., 2012)), the tight junction protein zona occludens-1 (ZO-1) (Gonzalez-Mariscal et al., 2003), as well as HBoV1 NS1 expression, was detected. Although only one third of the cells in the HBoV1-infected EpiAirway HAE expressed NS1, the infected HAE showed no expression of β -tubulin IV and a disassociation of the ZO-1 (Fig. 1D).

In addition, EpiAirway HAE was infected at the basolateral surface. A gradual virus release from both the apical and basolateral surface was observed from 1 to 11 days p.i. Virus release from apical infection reached a plateau ($\sim 2 \times 10^6$ gc/ μ l) at 12 days p.i.; however, there was ~ 1 –1.5 log less virus released from the basolateral surface during infection between 10 and 26 days p.i. (Fig. 2A).




Fig. 1. HBoV1 infects EpiAirway HAE productively and persistently. EpiAirway HAE was infected with the apically washed HBoV1 at an MOI of 100 $gc/\mu l$. (A) At the indicated days p.i., HBoV1 virions were collected from both the apical and basolateral chambers and quantified by qPCR. Averages and standard deviations of the viral $gc/\mu l$ are shown. (B) The TEER of HBoV1-infected EpiAirway HAE was monitored at the indicated days p.i. Averages and standard deviations of the TEER are shown. (C) At 11 days p.i., the apical chambers of three infected EpiAirway HAE cultures were washed with 200 μl of PBS to collect the apically washed cells. A total of 600 μl of the sample was concentrated to 200 μl , which was then cytospun onto a slide followed by immunofluorescence analysis with an anti-HBoV1 NS1 antibody. Confocal images were taken with an Eclipse C1 Plus confocal microscope (Nikon, Melville, NY, USA) controlled by Nikon EZ-C1 software, at a magnification of 40 \times as indicated. (D) At 50 days p.i., insert membranes of the infected EpiAirway HAE were fixed and analyzed by an immunofluorescence assay with an anti-HBoV1 NS1 antibody with co-staining of anti- β -tubulin IV or anti-ZO1 antibody. Confocal images were taken at a magnification of 40 \times . Nuclei were stained with DAPI (blue).




Fig. 2. Analyses of EpiAirway HAE infected with HBoV1 basolaterally. EpiAirway HAE was infected with HBoV1 at an MOI of 100 gc/μl from the basolateral surface. (A) At the indicated days p.i., both apically washed and basolaterally collected samples were quantified by qPCR as gc/μl, which are plotted to the days p.i. and shown as averages and standard deviations. (B) The TEER of HBoV1-infected HAE was monitored at the indicated days p.i. Averages and standard deviations are shown. (C) At the end of the infection, insert membranes of the HAE cultures were fixed and stained with anti-HBoV1 NS1 and anti-β-tubulin IV or anti-ZO1 antibodies. Nuclei were stained with DAPI (blue). The cells were visualized by confocal microscopy at a magnification of 40×.




Fig. 3. Kinetics of HBoV1-infected MucilAir HAE. MucilAir HAE was infected with HBoV1 from either the apical (A) or basolateral (B) surface at an MOI of 100 gc/μl. (A and B) At the indicated days p.i., progeny virions were collected from both the apical and basolateral chambers. HBoV1 virions were quantified by qPCR as viral gc/μl. Averages and standard deviations of the viral gc/μl are shown. (C) The TEER of HBoV1-infected HAE was monitored at the indicated days p.i. Averages and standard deviations of the detected TEER are shown.

Basolateral infection also caused significant epithelial damage represented by the gradual decrease in TEER (Fig. 2B), cilia loss and destruction of the tight junction (Fig. 2C).

Taken together, these results suggest that HBoV1 infects EpiAirway HAE both apically and basolaterally, and that the infection is persistent and causes airway epithelial damage.

3.2. HBoV1 infects MucilAir HAE cultures

MucilAir HAE was infected with HBoV1 either apically or basolaterally at an MOI of 100 gc/cell. Virus production after infection was monitored by collecting samples from both the apical and basolateral chambers of the infected HAE.

During apical infection, progeny virions were released apically at a level of 2×10^4 gc/μl at 3 days p.i., reached a peak of 7×10^6 gc/μl at 9 days p.i., and gradually decreased to 2×10^5 gc/μl at 26 days p.i. (Fig. 3A). Virus was also released from the basolateral surface, but at a level of 1–2 log less than that from the apical surface

(Fig. 3A). Mock-infected HAE did not release any virions from either the apical or basolateral surfaces. During basolateral infection, virions were released much more slowly in early infection (at 1–10 days p.i.), reached a peak (1×10^6 and 1×10^5 gc/μl, respectively, from the apical and basolateral surfaces) at 12–13 days p.i., and gradually decreased to levels of 8×10^4 and 5×10^3 gc/μl by the end of infection (Fig. 3B).

The TEER was detected to monitor the epithelial barrier function of the HAE affected by HBoV1 infection. Although the TEER of the MucilAir HAE was initially low (300–400 Ω cm²), after HBoV1 infection, it decreased further by ~2- to 3-fold, gradually to levels of 130 and 180 Ω cm² during apical and basolateral infection, respectively, at 26 days p.i. (Fig. 3C), which is consistent with the virus release kinetics (Fig. 3A and B).

At 26 days p.i., at least one third of the cells in the infected HAE expressed HBoV1 NS1 (Fig. 4A). Either apically- or basolaterally infected MucilAir HAE lost cilia and the tight junction, as shown by staining with anti-β-tubulin IV (Fig. 4B) and anti-ZO1 (Fig. 4C)




Fig. 4. HBoV1 infection of MucilAir HAE causes airway epithelial damage. At the end of the infection (26 days p.i.), insert membranes of the infected MucilAir HAE cultures were fixed and stained with an anti-HBoV1 NS1 antibody (A), or co-stained with anti-HBoV1 NS1 and anti- β -tubulin IV antibody (B) and with anti-HBoV1 NS1 and anti-ZO-1 antibodies (C). Nuclei were stained with DAPI (blue). The stained cells were visualized by confocal microscopy at magnifications of 40 \times and 100 \times , as indicated.

antibodies, respectively, and showed an obvious nucleus enlargement (Fig. 4B and C, DAPI), strongly suggesting airway epithelial damage.

4. Discussion

Pseudostratified and polarized HAE cultures have been reported for productive HBoV1 infection (Deng et al., 2013; Dijkman et al., 2009; Huang et al., 2012), but they were made in house, and thus were not available for most of the researchers in the field. In this report, HBoV1 infection was studied in polarized primary HAE cultures, MucilAir and EpiAirway HAE, purchased from MatTek Co. (Ashland, MA, USA) and Epithelix SàRL (Geneva, Switzerland), respectively. HBoV1 progeny virions were released from both infected MucilAir and EpiAirway HAE cultures, but are at an average of $\sim 1-2 \times 10^8$ gc per insert. These results suggest that the commercially available HAE cultures support productive HBoV1 infection. Thus, our study provides valuable information to culture HBoV1 and study HBoV1 infection in primary HAE.

Notably, HBoV1 infection can be persistent in EpiAirway HAE for at least 50 days p.i. Previous studies have identified the presence of episomal covalently closed circular (ccc) viral DNA in patients infected with HBoV1-3 (Kang et al., 2009; Kapoor et al., 2011; Windisch et al., 2013; Zhao et al., 2012). However, whether the cccDNA is the viral genomes persisting in HBoV1-infected EpiAirway HAE awaits further investigation, which will reveal mechanisms underlying HBoV1 persistent infection. The TEER of airway epithelium often indicates the extent of polarization and was used as a quality control of polarized HAE cultures from the manufacturers. Previously, primary B-HAE cultures made in house was used, which have a TEER of $>1000 \Omega \text{ cm}^2$ (Huang et al., 2012). However, both MucilAir and EpiAirway HAE had a low TEER (400 and $800 \Omega \text{ cm}^2$, respectively) when we received them. Nevertheless, HBoV1 infection disrupts the polarity of both the MucilAir and EpiAirway HAE cultures infected with HBoV1. The TEER of the infected MucilAir and EpiAirway HAE decreased by 2- to 3-fold, similar to the decrease in TEER in HBoV1-infected HAE made in house (Deng et al., 2013). Notably, in the HAE cultures we tested, there was a higher expression level of cilia in MucilAir HAE than in EpiAirway HAE (Figs. 2C and 4B, compare the β -tubulin). Apparently, the high expression level of cilia did not contribute a high level of virus production from infected MucilAir HAE. Interestingly, by the end of infection, all the cilia in the infected HAE of both types were lost, although only one third of the cells showed NS1 expression, suggesting that HBoV1 infection may release cytotoxic cytokines into the cultures as do RNA respiratory viruses (Vareille et al., 2011), which warrants further investigation.

Because of the nature of the MucilAir and EpiAirway HAE insert membrane materials, further histology analyses of the airway epithelia were not successful in either frozen section or paraffin-embedded section (data not shown). However, for the MucilAir HAE, the polyethylene terephthalate (PET) membrane of the Transwell insert provides a better cell visualization that allowed us to take confocal images of directly fixed infected cells on the insert membrane at a high magnification (Fig. 4B and C), which show much clearer nucleus enlargement of the cells and destruction of the epithelial tight junction in the infected HAE. MucilAir HAE has been used to infect influenza virus (Brookes et al., 2011), respiratory syncytial virus and human rhinovirus (Huang et al., 2011), and EpiAirway HAE has been reported to infect influenza virus (Triana-Baltzer et al., 2010), parainfluenza virus (Moscona et al., 2010; Palermo et al., 2009; Palmer et al., 2012), human rhinovirus (Sharma et al., 2010), and respiratory syncytial virus (Donalisio et al., 2012). MucilAir HAE was cultured in a Transwell insert of 0.33 cm^2 (Costar), while EpiAirway HAE was cultured in a Millicell

insert of 0.6 cm^2 (Millipore). Both HAE cultures are suitable model systems to study HBoV1 for many laboratories, which may lack experience to make primary HAE in house.

In conclusion, this study supports strongly that HBoV1 is a highly infectious respiratory virus. Two sources of HAE cultures, MucilAir and EpiAirway HAE, are commercially available for productive HBoV1 infection, which will facilitate the study of HBoV1 worldwide. More importantly, the airway barrier function was destroyed in both HBoV1-infected MucilAir and EpiAirway HAE. For the first time, HBoV1 infection was demonstrated to be persistent, for as long as 50 days p.i., in an in vitro model of human airway epithelia.

Acknowledgements

We thank Dr. Sarah Tague for help in taking confocal microscopy images and members of the Qiu lab for valuable discussion.

References

- Allander, T., Jartti, T., Gupta, S., Niesters, H.G., Lehtinen, P., Osterback, R., Vuorinen, T., Waris, M., Bjerkner, A., Tiveljung-Lindell, A., van den Hoogen, B.G., Hyyppia, T., Ruuskanen, O., 2007. Human bocavirus and acute wheezing in children. *Clin. Infect. Dis.* 44, 904–910.
- Allander, T., Tammi, M.T., Eriksson, M., Bjerkner, A., Tiveljung-Lindell, A., Andersson, B., 2005. Cloning of a human parvovirus by molecular screening of respiratory tract samples. *Proc. Natl. Acad. Sci. U.S.A.* 102, 12891–12896.
- Arthur, J.L., Higgins, G.D., Davidson, G.P., Givney, R.C., Ratcliff, R.M., 2009. A novel bocavirus associated with acute gastroenteritis in Australian children. *PLoS Pathog.* 5, e1000391.
- Brodzinski, H., Ruddy, R.M., 2009. Review of new and newly discovered respiratory tract viruses in children. *Pediatr. Emerg. Care* 25, 352–360.
- Brookes, D.W., Miah, S., Lackenby, A., Hartgroves, L., Barclay, W.S., 2011. Pandemic H1N1 2009 influenza virus with the H275Y oseltamivir resistance neuraminidase mutation shows a small compromise in enzyme activity and viral fitness. *J. Antimicrob. Chemother.* 66, 466–470.
- Chen, A.Y., Cheng, F., Lou, S., Luo, Y., Liu, Z., Delwart, E., Pintel, D., Qiu, J., 2010. Characterization of the gene expression profile of human bocavirus. *Virology* 403, 145–154.
- Chen, K.C., Shull, B.C., Moses, E.A., Lederman, M., Stout, E.R., Bates, R.C., 1986. Complete nucleotide sequence and genome organization of bovine parvovirus. *J. Virol.* 60, 1085–1097.
- Christensen, A., Døllner, H., Shanke, L.H., Krokstad, S., Moe, N., Nordbø, S.A., 2013. Detection of spliced mRNA from human bocavirus 1 in clinical samples from children with respiratory tract infections. *Emerg. Infect. Dis.* 19, 574–580.
- Christensen, A., Nordbø, S.A., Krokstad, S., Rognlien, A.G., Dollner, H., 2010. Human bocavirus in children: mono-detection, high viral load and viraemia are associated with respiratory tract infection. *J. Clin. Virol.* 49, 158–162.
- Deng, X., Yan, Z., Luo, Y., Xu, J., Cheng, Y., Li, Y., Engelhardt, J., Qiu, J., 2013. In vitro modeling of human bocavirus 1 infection of polarized primary human airway epithelia. *J. Virol.* 87, 4097–4102.
- Deng, Y., Gu, X., Zhao, X., Luo, J., Luo, Z., Wang, L., Fu, Z., Yang, X., Liu, E., 2012. High viral load of human bocavirus correlates with duration of wheezing in children with severe lower respiratory tract infection. *PLoS One* 7, e34353.
- Dijkman, R., Koekkoek, S.M., Molenkamp, R., Schildgen, O., van der Hoek, L., 2009. Human bocavirus can be cultured in differentiated human airway epithelial cells. *J. Virol.* 83, 7739–7748.
- Don, M., Soderlund-Venermo, M., Valent, F., Lahtinen, A., Hedman, L., Canciani, M., Hedman, K., Korppi, M., 2010. Serologically verified human bocavirus pneumonia in children. *Pediatr. Pulmonol.* 45, 120–126.
- Donalisio, M., Rusnati, M., Cagno, V., Civra, A., Bugatti, A., Giuliani, A., Pirri, G., Volante, M., Papotti, M., Landolfo, S., Lembo, D., 2012. Inhibition of human respiratory syncytial virus infectivity by a dendrimeric heparan sulfate-binding peptide. *Antimicrob. Agents Chemother.* 56, 5278–5288.
- Edner, N., Castillo-Rodas, P., Falk, L., Hedman, K., Soderlund-Venermo, M., Allander, T., 2011. Life-threatening respiratory tract disease with human bocavirus-1 infection in a four-year-old child. *J. Clin. Microbiol.* 50, 531–532.
- García-García, M.L., Calvo, C., Falcon, A., Pozo, F., Perez-Brena, P., De Cea, J.M., Casas, I., 2010. Role of emerging respiratory viruses in children with severe acute wheezing. *Pediatr. Pulmonol.* 45, 585–591.
- Gendrel, D., Guedj, R., Pons-Catalano, C., Emirian, A., Raymond, J., Rozenberg, F., Lebon, P., 2007. Human bocavirus in children with acute asthma. *Clin. Infect. Dis.* 45, 404–405.
- Gonzalez-Mariscal, L., Betanzos, A., Nava, P., Jaramillo, B.E., 2003. Tight junction proteins. *Prog. Biophys. Mol. Biol.* 81, 1–44.
- Hao, W., Bernard, K., Patel, N., Ulbrandt, N., Feng, H., Svabek, C., Wilson, S., Stracener, C., Wang, K., Suzich, J., Blair, W., Zhu, Q., 2012. Infection and propagation of human rhinovirus C in human airway epithelial cells. *J. Virol.* 86, 13524–13532.
- Huang, Q., Deng, X., Yan, Z., Cheng, F., Luo, Y., Shen, W., Lei-Butters, D.C., Chen, A.Y., Li, Y., Tang, L., Soderlund-Venermo, M., Engelhardt, J.F., Qiu, J., 2012. Establishment

- of a reverse genetics system for studying human bocavirus in human airway epithelia. *PLoS Pathog.* 8, e1002899.
- Huang, S., Wiszniewski, L., Constant, S., 2011. The use of *In Vitro* 3D cell models in drug development for respiratory diseases. In: Kapetanović, I. (Ed.), *Drug Discovery and Development – Present and Future*. InTech, Rijeka, Croatia, pp. 169–190.
- Ilyushina, N.A., Bovin, N.V., Webster, R.G., 2012. Decreased neuraminidase activity is important for the adaptation of H5N1 influenza virus to human airway epithelium. *J. Virol.* 86, 4724–4733.
- Jartti, T., Hedman, K., Jartti, L., Ruuskanen, O., Allander, T., Soderlund-Venermo, M., 2011. Human bocavirus—the first 5 years. *Rev. Med. Virol.* 22, 46–64.
- Kahn, J., 2008. Human bocavirus: clinical significance and implications. *Curr. Opin. Pediatr.* 20, 62–66.
- Kang, W., Wang, L., Harrell, H., Liu, J., Thomas, D.L., Mayfield, T.L., Scotti, M.M., Ye, G.J., Veres, G., Knop, D.R., 2009. An efficient rHSV-based complementation system for the production of multiple rAAV vector serotypes. *Gene Ther.* 16, 229–239.
- Kantola, K., Hedman, L., Allander, T., Jartti, T., Lehtinen, P., Ruuskanen, O., Hedman, K., Soderlund-Venermo, M., 2008. Serodiagnosis of human bocavirus infection. *Clin. Infect. Dis.* 46, 540–546.
- Kapoor, A., Hornig, M., Asokan, A., Williams, B., Henriquez, J.A., Lipkin, W.I., 2011. Bocavirus episome in infected human tissue contains non-identical termini. *PLoS One* 6, e21362.
- Kapoor, A., Mehta, N., Esper, F., Poljsak-Prijatelj, M., Quan, P.L., Qaisar, N., Delwart, E., Lipkin, W.I., 2010a. Identification and characterization of a new bocavirus species in gorillas. *PLoS One* 5, e11948.
- Kapoor, A., Simmonds, P., Slikas, E., Li, L., Bodhidatta, L., Sethabutr, O., Triki, H., Bahri, O., Oderinde, B.S., Baba, M.M., Bukbuk, D.N., Besser, J., Bartkus, J., Delwart, E., 2010b. Human bocaviruses are highly diverse, dispersed, recombination prone, and prevalent in enteric infections. *J. Infect. Dis.* 201, 1633–1643.
- Kapoor, A., Slikas, E., Simmonds, P., Chiochansin, T., Naeem, A., Shaikat, S., Alam, M.M., Sharif, S., Angez, M., Zaidi, S., Delwart, E., 2009. A newly identified bocavirus species in human stool. *J. Infect. Dis.* 199, 196–200.
- Kesebir, D., Vazquez, M., Weibel, C., Shapiro, E.D., Ferguson, D., Landry, M.L., Kahn, J.S., 2006. Human bocavirus infection in young children in the United States: molecular epidemiological profile and clinical characteristics of a newly emerging respiratory virus. *J. Infect. Dis.* 194, 1276–1282.
- Korner, R.W., Soderlund-Venermo, M., van Koningsbruggen-Rietschel, S., Kaiser, R., Malecki, M., Schildgen, O., 2011. Severe human bocavirus infection, Germany. *Emerg. Infect. Dis.* 17, 2303–2305.
- Lin, F., Zeng, A., Yang, N., Lin, H., Yang, E., Wang, S., Pintel, D., Qiu, J., 2007. Quantification of human bocavirus in lower respiratory tract infections in China. *Infect. Agent Cancer* 2, 3.
- Lopez, A.D., Mathers, C.D., Ezzati, M., Jamison, D.T., Murray, C.J., 2006. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 367, 1747–1757.
- Manning, A., Russell, V., Eastick, K., Leadbetter, G.H., Hallam, N., Templeton, K., Simmonds, P., 2006. Epidemiological profile and clinical associations of human bocavirus and other human parvoviruses. *J. Infect. Dis.* 194, 1283–1290.
- Matrosovich, M.N., Matrosovich, T.Y., Gray, T., Roberts, N.A., Klenk, H.D., 2004. Human and avian influenza viruses target different cell types in cultures of human airway epithelium. *Proc. Natl. Acad. Sci. U.S.A.* 101, 4620–4624.
- Meriluoto, M., Hedman, L., Tanner, L., Simell, V., Makinen, M., Simell, O., Mykkanen, J., Korpelainen, J., Ruuskanen, O., Ilonen, J., Knip, M., Simell, O., Hedman, K., Soderlund-Venermo, M., 2012. Association of human bocavirus 1 infection with respiratory disease in childhood follow-up study, Finland. *Emerg. Infect. Dis.* 18, 264–271.
- Moscona, A., Porotto, M., Palmer, S., Tai, C., Aschenbrenner, L., Triana-Baltzer, G., Li, Q.X., Wurtman, D., Niewiesk, S., Fang, F., 2010. A recombinant sialidase fusion protein effectively inhibits human parainfluenza viral infection in vitro and in vivo. *J. Infect. Dis.* 202, 234–241.
- Nascimento-Carvalho, C.M., Cardoso, M.R., Meriluoto, M., Kemppainen, K., Kantola, K., Ruuskanen, O., Hedman, K., Soderlund-Venermo, M., 2012. Human bocavirus infection diagnosed serologically among children admitted to hospital with community-acquired pneumonia in a tropical region. *J. Med. Virol.* 84, 253–258.
- Palermo, L.M., Porotto, M., Yokoyama, C.C., Palmer, S.G., Mungall, B.A., Greengard, O., Niewiesk, S., Moscona, A., 2009. Human parainfluenza virus infection of the airway epithelium: viral hemagglutinin-neuraminidase regulates fusion protein activation and modulates infectivity. *J. Virol.* 83, 6900–6908.
- Palmer, S.G., Porotto, M., Palermo, L.M., Cunha, L.F., Greengard, O., Moscona, A., 2012. Adaptation of human parainfluenza virus to airway epithelium reveals fusion properties required for growth in host tissue. *MBio* 3, e00137–e212.
- Proenca-Modena, J.L., Gagliardi, T.B., Escremim de, P.F., Iwamoto, M.A., Criado, M.F., Camara, A.A., Acrani, G.O., Cintra, O.A., Cervi, M.C., de Paula Arruda, L.K., Arruda, E., 2011. Detection of human bocavirus mRNA in respiratory secretions correlates with high viral load and concurrent diarrhea. *PLoS One* 6, e21083.
- Pyrk, K., Sims, A.C., Dijkman, R., Jebbink, M., Long, C., Deming, D., Donaldson, E., Vabret, A., Baric, R., van der Hoek, L., Pickles, R., 2010. Culturing the unculturable: human coronavirus HKU1 infects, replicates, and produces progeny virions in human ciliated airway epithelial cell cultures. *J. Virol.* 84, 11255–11263.
- Schildgen, O., Muller, A., Allander, T., Mackay, I.M., Volz, S., Kupfer, B., Simon, A., 2008. Human bocavirus: passenger or pathogen in acute respiratory tract infections? *Clin. Microbiol. Rev.* 21, 291–304.
- Sharma, M., Schoop, R., Hudson, J.B., 2010. The efficacy of Echinacea in a 3-D tissue model of human airway epithelium. *Phytother. Res.* 24, 900–904.
- Shay, D.K., Holman, R.C., Newman, R.D., Liu, L.L., Stout, J.W., Anderson, L.J., 1999. Bronchiolitis-associated hospitalizations among US children, 1980–1996. *J. Am. Med. Assoc.* 282, 1440–1446.
- Sims, A.C., Baric, R.S., Yount, B., Burkett, S.E., Collins, P.L., Pickles, R.J., 2005. Severe acute respiratory syndrome coronavirus infection of human ciliated airway epithelia: role of ciliated cells in viral spread in the conducting airways of the lungs. *J. Virol.* 79, 15511–15524.
- Soderlund-Venermo, M., Lahtinen, A., Jartti, T., Hedman, L., Kemppainen, K., Lehtinen, P., Allander, T., Ruuskanen, O., Hedman, K., 2009. Clinical assessment and improved diagnosis of bocavirus-induced wheezing in children, Finland. *Emerg. Infect. Dis.* 15, 1423–1430.
- Sun, Y., Chen, A.Y., Cheng, F., Guan, W., Johnson, F.B., Qiu, J., 2009. Molecular characterization of infectious clones of the minute virus of canines reveals unique features of bocaviruses. *J. Virol.* 83, 3956–3967.
- Tijssen, P., Agbandje-McKenna, M., Almendral, J.M., Bergoin, M., Flegel, T.W., Hedman, K., Kleinschmidt, J., Li, Y., Pintel, D.J., Tattersall, P., 2012. Family Parvoviridae. In: King, A.M., Lefkowitz, E., Adams, M.J., Carstens, E.B. (Eds.), *Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of Viruses*. Elsevier, pp. 405–425.
- Triana-Baltzer, G.B., Babizki, M., Chan, M.C., Wong, A.C., Aschenbrenner, L.M., Campbell, E.R., Li, Q.X., Chan, R.W., Peiris, J.S., Nicholls, J.M., Fang, F., 2010. DAS181, a sialidase fusion protein, protects human airway epithelium against influenza virus infection: an in vitro pharmacodynamic analysis. *J. Antimicrob. Chemother.* 65, 275–284.
- Ursic, T., Steyer, A., Kopriva, S., Kalan, G., Krivec, U., Petrovec, M., 2011. Human bocavirus as the cause of a life-threatening infection. *J. Clin. Microbiol.* 49, 1179–1181.
- Vareille, M., Kieninger, E., Edwards, M.R., Regamey, N., 2011. The airway epithelium: soldier in the fight against respiratory viruses. *Clin. Microbiol. Rev.* 24, 210–229.
- Villanave, R., Thavagnanam, S., Sarlang, S., Parker, J., Douglas, I., Skibinski, G., Heaney, L.G., McKaigue, J.P., Coyle, P.V., Shields, M.D., Power, U.F., 2012. In vitro modeling of respiratory syncytial virus infection of pediatric bronchial epithelium, the primary target of infection in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 109, 5040–5045.
- Wang, G., Deering, C., Macke, M., Shao, J., Burns, R., Blau, D.M., Holmes, K.V., Davidson, B.L., Perlman, S., McCray Jr., P.B., 2000. Human coronavirus 229E infects polarized airway epithelia from the apical surface. *J. Virol.* 74, 9234–9239.
- Wang, K., Wang, W., Yan, H., Ren, P., Zhang, J., Shen, J., Deubel, V., 2010. Correlation between bocavirus infection and humoral response, and co-infection with other respiratory viruses in children with acute respiratory infection. *J. Clin. Virol.* 47, 148–155.
- Windisch, W., Schildgen, V., Malecki, M., Lenz, J., Brockmann, M., Karagiannidis, C., Schildgen, O., 2013. Detection of H20V DNA in idiopathic lung fibrosis, Cologne, Germany. *J. Clin. Virol.* 58, 325–327.
- Zabner, J., Freimuth, P., Puga, A., Fabrega, A., Welsh, M.J., 1997. Lack of high affinity fiber receptor activity explains the resistance of ciliated airway epithelia to adenovirus infection. *J. Clin. Invest.* 100, 1144–1149.
- Zeng, H., Pappas, C., Katz, J.M., Tumpey, T.M., 2011. The 2009 pandemic H1N1 and triple-reassortant swine H1N1 influenza viruses replicate efficiently but elicit an attenuated inflammatory response in polarized human bronchial epithelial cells. *J. Virol.* 85, 686–696.
- Zhai, S., Yue, C., Wei, Z., Long, J., Ran, D., Lin, T., Deng, Y., Huang, L., Sun, L., Zheng, H., Gao, F., Zheng, H., Chen, S., Yuan, S., 2010. High prevalence of a novel porcine bocavirus in weanling piglets with respiratory tract symptoms in China. *Arch. Virol.* 155, 1313–1317.
- Zhang, L., Bukreyev, A., Thompson, C.I., Watson, B., Peebles, M.E., Collins, P.L., Pickles, R.J., 2005. Infection of ciliated cells by human parainfluenza virus type 3 in an in vitro model of human airway epithelium. *J. Virol.* 79, 1113–1124.
- Zhang, L., Collins, P.L., Lamb, R.A., Pickles, R.J., 2011. Comparison of differing cytopathic effects in human airway epithelium of parainfluenza virus 5 (W3A), parainfluenza virus type 3, and respiratory syncytial virus. *Virology* 421, 67–77.
- Zhang, L., Peebles, M.E., Boucher, R.C., Collins, P.L., Pickles, R.J., 2002. Respiratory syncytial virus infection of human airway epithelial cells is polarized, specific to ciliated cells, and without obvious cytopathology. *J. Virol.* 76, 5654–5666.
- Zhao, H., Zhao, L., Sun, Y., Qian, Y., Liu, L., Jia, L., Zhang, Y., Dong, H., 2012. Detection of a bocavirus circular genome in fecal specimens from children with acute diarrhea in Beijing, China. *PLoS One* 7, e48980.