

# Nifedipine Treatment Reduces Resting Calcium Concentration, Oxidative and Apoptotic Gene Expression, and Improves Muscle Function in Dystrophic *mdx* Mice

Francisco Altamirano<sup>1,2,9</sup>, Denisse Valladares<sup>1,9</sup>, Carlos Henríquez-Olguín<sup>1</sup>, Mariana Casas<sup>1,4</sup>, Jose R. López<sup>2,3</sup>, Paul D. Allen<sup>2,3</sup>, Enrique Jaimovich<sup>1\*</sup>

**1** Centro de Estudios Moleculares de la Célula, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile, **2** Department of Molecular Biosciences, School of Veterinary Medicine, University of California Davis, Davis, California, United States of America, **3** Department of Anesthesiology, Perioperative and Pain Medicine, Brigham & Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States of America, **4** Programa de Fisiología y Biofísica, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile

## Abstract

Duchenne Muscular Dystrophy (DMD) is a recessive X-linked genetic disease, caused by mutations in the gene encoding dystrophin. DMD is characterized in humans and in *mdx* mice by a severe and progressive destruction of muscle fibers, inflammation, oxidative/nitrosative stress, and cell death. In *mdx* muscle fibers, we have shown that basal ATP release is increased and that extracellular ATP stimulation is pro-apoptotic. In normal fibers, depolarization-induced ATP release is blocked by nifedipine, leading us to study the potential therapeutic effect of nifedipine in *mdx* muscles and its relation with extracellular ATP signaling. Acute exposure to nifedipine (10  $\mu$ M) decreased  $[Ca^{2+}]_r$ , NF- $\kappa$ B activity and iNOS expression in *mdx* myotubes. In addition, 6-week-old *mdx* mice were treated with daily intraperitoneal injections of nifedipine, 1 mg/Kg for 1 week. This treatment lowered the  $[Ca^{2+}]_r$  measured *in vivo* in the *mdx vastus lateralis*. We demonstrated that extracellular ATP levels were higher in adult *mdx flexor digitorum brevis* (FDB) fibers and can be significantly reduced after 1 week of treatment with nifedipine. Interestingly, acute treatment of *mdx* FDB fibers with apyrase, an enzyme that completely degrades extracellular ATP to AMP, reduced  $[Ca^{2+}]_r$  to a similar extent as was seen in FDB fibers after 1-week of nifedipine treatment. Moreover, we demonstrated that nifedipine treatment reduced mRNA levels of pro-oxidative/nitrosative (iNOS and gp91<sup>phox</sup>/p47<sup>phox</sup> NOX2 subunits) and pro-apoptotic (Bax) genes in *mdx* diaphragm muscles and lowered serum creatine kinase (CK) levels. In addition, nifedipine treatment increased muscle strength assessed by the inverted grip-hanging test and exercise tolerance measured with forced swimming test in *mdx* mice. We hypothesize that nifedipine reduces basal ATP release, thereby decreasing purinergic receptor activation, which in turn reduces  $[Ca^{2+}]_r$  in *mdx* skeletal muscle cells. The results in this work open new perspectives towards possible targets for pharmacological approaches to treat DMD.

**Citation:** Altamirano F, Valladares D, Henríquez-Olguín C, Casas M, López JR, et al. (2013) Nifedipine Treatment Reduces Resting Calcium Concentration, Oxidative and Apoptotic Gene Expression, and Improves Muscle Function in Dystrophic *mdx* Mice. PLoS ONE 8(12): e81222. doi:10.1371/journal.pone.0081222

**Editor:** James M. Ervasti, University of Minnesota, United States of America

**Received:** June 26, 2013; **Accepted:** October 9, 2013; **Published:** December 9, 2013

**Copyright:** © 2013 Altamirano et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Fondo de Investigación Avanzado en Areas Prioritarias 15010006, Fondo Nacional de Desarrollo Científico y Tecnológico 1110467, ACT1111 and AFM14562 (EJ, MC), National Institutes of Health AR43140, AR052534 (PDA and JRL), Program U-INICIA VID 2011, grant U-INICIA 02/12M; University of Chile (MC) supported this work. FA and DV were recipients of a doctoral fellowship and a doctoral thesis support grants AT-24100066 (FA), AT-24110211 (DV) from Comisión Nacional de Ciencia y Tecnología (CONICYT). FA thanks Vicerrectoría Asuntos Académicos and MECESUP UCH0714 (Universidad de Chile) and REDES 120003 (CONICYT) for travel support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: ejaimovi@med.uchile.cl

<sup>9</sup> These authors contributed equally to this work.

## Introduction

Duchenne muscular dystrophy (DMD) is a severe neuromuscular disorder characterized by the absence of the dystrophin protein and is the most common form of muscular dystrophy with a frequency of 1 in 3500 male births [1]. DMD is caused by a recessive X-linked mutation in dystrophin gene located to Xp21 [2]. Usually patients lose the ability to walk between 6–12 years old due to severe muscle damage and contractures that results in wheelchair dependence [1]. Since DMD is a progressive disease,

most patients die in their twenties due either to respiratory failure or cardiac dysfunction [3].

Resting intracellular calcium concentration ( $[Ca^{2+}]_i$ ) is elevated in myotubes from *mdx* mice, a murine model for DMD [4]. This increased  $[Ca^{2+}]_i$  depends on sarcolemmal  $Ca^{2+}$  entry, as well as  $Ca^{2+}$  leak from sarcoplasmic reticulum (SR), through type 1 ryanodine receptors (RyR1) and inositol tri-phosphate receptors (IP<sub>3</sub>R). In turn, elevated  $[Ca^{2+}]_i$  modulates NF- $\kappa$ B activation, leading to an up-regulation of inducible nitric oxide synthase (iNOS) in dystrophic myotubes [4].

The L-type voltage gated  $\text{Ca}^{2+}$  channels (Dihydropyridine receptors, DHPRs) are also implicated in DMD pathology [5–7]. Their roles as voltage sensors for both excitation-contraction and excitation-transcription coupling, and as their role as L-type  $\text{Ca}^{2+}$  channels in skeletal muscle cells, make them key regulators of intracellular  $[\text{Ca}^{2+}]_i$  [8–10]. The DHPR is also an important modulator of ATP release through Pannexin1 channels in skeletal muscle fibers [9,10]. ATP release elicited by electrical stimuli is blocked with the DHPR antagonist nifedipine as well as with the agonist (–)S-BayK 8644 in FDB fibers [10], suggesting that DHPR could directly control ATP release and that this event is independent of L-type current. An interaction between dystrophin and DHPR has been proposed in transverse tubular system of skeletal muscle fibers [6,11], suggesting that there is DHPR dysregulation in dystrophic skeletal muscle cells. More recently, we showed that basal levels of ATP release are importantly increased in *mdx* muscle fibers [12]. In dystrophic fibers, extracellular ATP stimulation was pro-apoptotic, inducing the transcription of Bax, BIM and PUMA and increasing the levels of activated Bax and cytosolic cytochrome C [12]. These data suggest the potential for involvement of the ATP pathway in the activation of mechanisms related with cell death in muscular dystrophy, opening new perspectives towards possible targets for pharmacological therapies.

A double-blinded controlled clinical trial with nifedipine was carried out in 1987, and showed that there was no significant beneficial effect of nifedipine treatment [13]. In this study, patient selection was restricted to semiology and classical human genetics, since dystrophin was not cloned until 1987 [2]. DMD has been considered stereotyped in its clinical presentation, evolution and severity [14]. However, an inverse correlation between severity of disease and residual amount of dystrophin (related to different mutations) has been found [15]. More recently, a retrospective, single institution, long-term (>10 years) study was done in 75 DMD patients with a complete lack of dystrophin (determined by western blot) and genotyping, showed that DMD can be divided into 4 sub-phenotypes with different cognitive and motor outcomes [16], showing the complexity and heterogeneity of DMD.

Based on our previous results showing that extracellular ATP levels are elevated in *mdx* fibers and that ATP stimulation is pro-apoptotic, here we revisited the potential therapeutic effect of this drug in the *mdx* mouse model. In the present study, we found that acute treatment of *mdx* myotubes with nifedipine reduced  $[\text{Ca}^{2+}]_i$ , NF- $\kappa\text{B}$  activity and iNOS expression. Likewise, intraperitoneal injections of nifedipine for 1 week, reduced *in vivo*  $[\text{Ca}^{2+}]_i$  in *vastus lateralis* muscles, decreased serum creatine kinase levels and increased *in vivo* muscle strength in *mdx* mice. In FDB muscle fibers isolated from the same nifedipine treated *mdx* mice, basal ATP release was reduced and the gene expression of pro-oxidative and pro-apoptotic genes in diaphragm (the most seriously affected muscle) were down-regulated. The present findings demonstrate that nifedipine treatment can effectively modify the observed changes associated with muscle pathology in *mdx* mice and opens a new pharmacologic approach for treatment of patients with DMD.

## Materials and Methods

### Myotube Cultures

Primary myoblasts were isolated from 5–6 week old male wild type C57BL/6 and *mdx* mice and myotubes were differentiated as described previously [4,17].

### Adult FDB Fiber Isolation

*Flexor digitorum brevis* (FDB) muscles were dissected from 5–6 week old male mice and intact muscle fibers were obtained by enzymatic digestion of the whole muscle with collagenase type 4 (Worthington, Lakewood, NJ) for 90 min at 37°C followed by mechanical dissociation with fire polished Pasteur pipettes. Isolated fibers were seeded in matrigel-coated dishes in DMEM supplemented with 10% horse serum and used for experimentation 20–24 hours after isolation.

### Drug Treatment Protocol

Male 5–6 week-old *wt* and *mdx* mice were injected daily for 1 week with either nifedipine solution (1 mg/Kg body weight) or saline intraperitoneally. Nifedipine (Sigma-Aldrich) solution was prepared in dark at 17 mg/mL in absolute ethanol and then was diluted in sterile saline solution (0.9% NaCl) at 0.2 mg/mL for injections.

### Determination of $[\text{Ca}^{2+}]_i$

Double-barreled  $\text{Ca}^{2+}$ -selective microelectrodes were prepared and calibrated as previously described [4]. Pulled microelectrodes were backfilled with the neutral carrier ETH 129 (Fluka-Sigma-Aldrich) and then with pCa7 solution. Only those electrodes with a linear relationship between pCa3 and pCa7 (Nernstian response, 29.5 mV and 30 mV per pCa unit at 23°C and 37°C, respectively) were used experimentally.

Myotubes or muscle fibers were impaled with the double-barreled  $\text{Ca}^{2+}$  selective microelectrodes and potentials were recorded via high impedance amplifier (WPI Duo-773) [18].  $[\text{Ca}^{2+}]_i$  measurements in myotubes were made with or without nifedipine (10  $\mu\text{M}$ ) in Krebs-Ringer solution (in mM: 140 NaCl, 5 KCl, 2.5  $\text{CaCl}_2$ , 1  $\text{MgSO}_4$ , 5 glucose, and 10 Hepes/Tris, pH 7.4) at 23°C, as previously described [4].

The measurements of  $[\text{Ca}^{2+}]_i$  in muscle fibers were done *in vivo* in the *vastus lateralis*. Mice were anesthetized with ketamine 100 mg/Kg and xylazine 5 mg/Kg and kept euthermic with a feedback-heating pad (ATC-1000 Temperature controller, WPI, Sarasota, FL). A small incision was made in the skin in the anterior part of the left leg, the *vastus lateralis* muscle was identified and fascia was partially removed. The superficial fibers were exposed and locally perfused with Krebs Ringer solution.

Measurements of  $[\text{Ca}^{2+}]_i$  in isolated adult FDB fibers were done after fibers were incubated in Krebs Ringer solution with or without apyrase 2 U/mL (grade VII from potato, Sigma-Aldrich) for 10 min at room temperature.

### NF- $\kappa\text{B}$ Luciferase Reporter Activity

Both *wt* and *mdx* myoblasts were transduced with a lentivirus containing 5 tandem NF- $\kappa\text{B}$  binding site repeats cloned upstream of a luciferase reporter gene and populations that stably expressed the transgene were selected using G418, as described previously [4]. Myoblasts stably expressing this reporter were completely normal and differentiated into myotubes after 3–4 days similar to untransduced cells. Myotubes were treated with or without nifedipine for 6 h in differentiation media. Luciferase activity was determined using a dual-luciferase reporter assay system (Promega) according to the manufacturer's instructions, and light detection was carried out in a Berthold F12 luminometer. Results were normalized with total protein and the relations "luciferase activity/mg protein" were shown.

## mRNA Quantitation

Total RNA was isolated from myotubes or diaphragm muscles from both the nifedipine- or saline-treated groups with TRIzol<sup>®</sup> reagent (Invitrogen) according to the manufacturer's protocol. cDNA was prepared by reverse transcription (RT) reaction of 1 µg of total RNA using random primers. Real time PCR was performed as previously described [4] using the following primers: *bax* 5'-GCTGACATGTTTGCTGATGG-3' and 5'-GATCAGCTCGGGCACCTTTAG-3' *bim* 5'-CGACAGTCTCAGGAGGAACC-3' and 5'-CATTTGCAAACACCCTCCTT-3' *gp91<sup>phox</sup>* 5'-TCACATCCTCTACCAAAACC-3' and 5'-CCTTTATTTTTGCCCATCT-3' *p47<sup>phox</sup>* 5'-AGAACA-GAGTCATCCCACAC-3' and 5'-GCTACGTTATTC'TTGC-CATC-3' *iNOS* 5'-CAGCTCAAGAGCCAGAAACG-3' and 5'-TTACTCAGTGCCAGAAGCTG-3' *gapdh* 5'-CTCATGACCA-CAGTCCATGC-3' and 5'-TTCAGCTCTGGGATGACCTT-3' *18S rRNA* 5'-GGGCCCGAAGCGTTACTTT-3' and 5'-TTGCGCCGGTCCAAGAATTT-3'.

## ATP Detection Using a Luciferin/Luciferase Assay

FDB fibers from nifedipine- and saline-treated mice were prepared as described above. Because media replacement causes a mechanical stimulus that itself induces ATP release [19,20], we measured ATP release for up to 9 min beginning 30 min after media change. ATP concentrations were measured with the CellTiter-Glo<sup>®</sup> Luminescent Cell Viability Assay (Promega, Madison, WI, USA) as reported [9]. Data were calculated as pmol extracellular ATP/µg total RNA. Normalization by total RNA instead of total protein was chosen because fibers were seeded on a Matrigel-coated dish (containing a large amount of protein), which may affect the protein determination associated only to fibers.

## Serum Creatine Kinase Determinations

Blood samples were obtained by cardiac puncture in anesthetized mice. Blood was collected in a sterile test tube, allowed to clot on ice for 30 min and then centrifuged at 3000 rpm for 10 minutes. Creatine kinase (CK) levels were determined using the UV-kinetic method (Teco Diagnostics) according to the manufacturer instructions. ΔAbs/min were used to calculate CK enzymatic activity and the results were expressed as International Kilo Units per liter (KUI/L).

## Inverted Grid-hanging Test

Muscle strength of mouse limbs was tested with an inverted grid-hanging test [21]. Mice were placed individually on the center of a 21×21 cm wire grid (wire width ≈0.1 cm and spacing 0.5 cm), mounted 35 cm above a table. After gently inverting the grid, the mouse hanging time was recorded (grip latency). This procedure was repeated three times and the average hanging time values were calculated for each mouse.

## Forced Swimming Test

A 1-liter beaker (11 cm diameter and 15 cm height) filled with water (23°C) was used as swimming pool to assess the exercise tolerance of saline- and nifedipine-treated mice [22]. First, a weight (10% of their body weight) was attached to the tail of a mouse, which was then gently placed in the water, and the time at which the mouse was unable to maintain complete buoyancy was recorded. At this time the mice were immediately removed from the swimming pool, dried gently with paper towels and returned to their cages.

## Statistical Analysis

Data of *n* experiments were expressed as mean ± S.E.M. The significance of difference among treatments was evaluated using a two-tailed *t* test for unpaired data or ANOVA- followed by Tukey's *t*-test. A *P* value <0.05 was considered statistically significant.

## Ethics Approval

All procedures for animal experimentation were done in accordance with guidelines approved by the Bioethical Committee at the Facultad de Medicina, Universidad de Chile and the IACUCs at Harvard Medical School and University of California at Davis.


## Results

### Nifedipine Reduces [Ca<sup>2+</sup>]<sub>r</sub> in Dystrophic *mdx* Myotubes


Myotubes were incubated in Krebs Ringer solution with or without nifedipine (10 µM) for 10 min and [Ca<sup>2+</sup>]<sub>r</sub> was measured with Ca<sup>2+</sup> selective microelectrodes, in both *wt* and *mdx* myotubes. [Ca<sup>2+</sup>]<sub>r</sub> observed in *mdx* myotubes was significantly higher compared to *wt* myotubes (315±8 vs 112±2 nM *P*<0.001) (Figure 1). There was a significant reduction in the [Ca<sup>2+</sup>]<sub>r</sub> in nifedipine-treated *mdx* myotubes compared with untreated *mdx* myotubes (254±12, *P*<0.001). Nifedipine treatment did not modify [Ca<sup>2+</sup>]<sub>r</sub> in *wt* myotubes (119±1 nM, *P*>0.05).

### NF-κB Activity and iNOS Expression were Reduced After Nifedipine Incubation

We previously reported that the increases in NF-κB activity and iNOS expression are modulated by [Ca<sup>2+</sup>]<sub>r</sub> in *mdx* myotubes [4]. To study the effect of nifedipine treatment in NF-κB activity and iNOS expression, we used a NF-κB luciferase reporter and real



**Figure 1. Nifedipine incubation reduces [Ca<sup>2+</sup>]<sub>r</sub> in *mdx* myotubes.** Myotubes were incubated with 10 µM Nifedipine for 10 min at room temperature in Krebs Ringer solution and [Ca<sup>2+</sup>]<sub>r</sub> was measured using double-barreled Ca<sup>2+</sup> selective microelectrodes. Data are expressed as mean ± S.E.M. *Wt* (n = 10), *mdx* (n = 27), *Wt*+Nife (n = 10) and *mdx*+Nife (n = 10). \*\*\**P*<0.001, ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g001



**Figure 2. NF- $\kappa$ B activity and iNOS expression in both *wt* and *mdx* myotubes.** **A.** NF- $\kappa$ B activity was studied with a luciferase reporter. Nifedipine treatment (10  $\mu$ M for 6 h) reduced NF- $\kappa$ B activity in *mdx*, without any significant effect in *wt* myotubes (n=6–12). **B.** mRNA levels of iNOS were determined by real time PCR after 6 h of nifedipine treatment (10  $\mu$ M) (n=6–7). Data are expressed as mean  $\pm$  S.E.M., \* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.001, ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g002

time PCR, respectively (see *Materials and Methods*). After incubation with nifedipine (10  $\mu$ M, 6 h) NF- $\kappa$ B activity was reduced by 33% ( $P$ <0.05) and iNOS mRNA levels were diminished by 61% ( $P$ <0.01) in *mdx* myotubes, without any significant effect in *wt* myotubes ( $P$ >0.05) (Figure 2).

#### Nifedipine Decreases the $[Ca^{2+}]_r$ in *mdx* Muscles

To establish if nifedipine can reduce  $[Ca^{2+}]_r$  in *mdx*, both *wt* and *mdx* mice were injected intraperitoneally daily for 1 week with either 1 mg/Kg nifedipine solution or saline.  $[Ca^{2+}]_r$  measured in *in vivo* in the superficial fibers of *vastus lateralis* muscles was significantly higher in *mdx* muscles compared with *wt* muscles in the saline treated group (Figure 3,  $320 \pm 13$  vs  $111 \pm 3$  nM,  $P$ <0.001). Nifedipine treatment was able to significantly reduce  $[Ca^{2+}]_r$  in *mdx* mice ( $236 \pm 8$  nM,  $P$ <0.001), but had no effect in *wt* muscles ( $108 \pm 1$  nM,  $P$ >0.05).

#### Nifedipine Treatment Reduces ATP Release in Adult Fibers

Recently, we reported that depolarization-induced ATP release is modulated by DHPR activity and can be blocked in skeletal muscle cells by nifedipine [10]. Moreover, extracellular ATP levels are higher at resting conditions in *mdx* skeletal muscle fibers [12]. It is very well established that extracellular ATP can signal through purinergic receptors and modulate the intracellular  $Ca^{2+}$  concentration [23]. In order to establish a correlation between extracellular ATP and the alterations in the  $[Ca^{2+}]_r$ , we measured ATP release in isolated FDB fibers from both *wt* and *mdx* mice after 1-week of nifedipine or saline treatment. Because media replacement causes a mechanical stimulus that itself induces ATP release [19,20], we measured ATP release after 30 min of media change and then up to 9 min. ATP release from *mdx* fibers was higher compared to *wt* fibers at every studied time point (Figure 4A). Nifedipine treatment reduced ATP release in both *wt* and *mdx* fibers, with a more pronounced effect in *mdx* fibers. At


t = 30 min after media change average extracellular ATP in fibers from saline-treated *mdx* mice was higher than in saline-treated *wt* fibers ( $1031 \pm 22$  vs  $540 \pm 93$  pmol ATP/ $\mu$ g RNA),  $P$ <0.01) (Figure 4B). ATP release was significantly decreased in fibers isolated from nifedipine-treated *mdx* ( $442 \pm 18$  pmol ATP/ $\mu$ g RNA,  $P$ <0.001) compared to untreated *mdx* fibers. Similar results were observed after 9 min (Figure 4C), showing that extracellular ATP levels were higher in *mdx* fibers but can be diminished near to *wt* levels after 1-week of nifedipine treatment.

#### Degradation of Extracellular ATP Reduces $[Ca^{2+}]_r$ in *mdx* Adult Fibers

In order to determine the participation of extracellular ATP on  $[Ca^{2+}]_r$  in adult *mdx* fibers, we treated isolated FDB fibers with apyrase (2 U/mL) in Krebs Ringer solution for 10 min. Apyrase is an enzyme that rapidly metabolizes extracellular ATP to AMP [9]. Apyrase treatment significantly reduced  $[Ca^{2+}]_r$  from  $296 \pm 11$  to  $195 \pm 8$  nM ( $P$ <0.001) in *mdx* fibers, without any significant effect in *wt* fibers ( $112 \pm 6$  to  $100 \pm 2$  nM,  $P$ >0.05) (Figure 5).

#### NADPH Oxidase Subunits and iNOS Expression in *mdx* Mice

Dystrophic muscles are characterized by an increase in pro-oxidative gene expression, such as iNOS and NADPH oxidase subunits (gp91<sup>phox</sup>, p67<sup>phox</sup> and rac1) [4,24,25]. Because the diaphragm has been described as the most severely affected muscle in the *mdx* mouse [26] it was used to determine the antioxidant effects of nifedipine treatment. For these studies we determined mRNA levels of iNOS and the NADPH oxidase subunits gp91<sup>phox</sup> and p47<sup>phox</sup>, from whole diaphragm lysates using real time PCR. We found that iNOS mRNA levels were 2.4-fold and gp91<sup>phox</sup> were 3.9-fold higher in *mdx* diaphragms compared to *wt*, with no significant difference in the expression of p47<sup>phox</sup> between the two groups (Figure 6). Nifedipine treatment reduced the mRNA levels of iNOS, gp91<sup>phox</sup> and p47<sup>phox</sup> in *mdx* diaphragms by 86%



**Figure 3. Nifedipine treatment reduces muscle  $[\text{Ca}^{2+}]_r$  in vivo in *mdx* mice.** Mice were treated with daily intraperitoneal injections of nifedipine 1 mg/Kg for 1-week or saline. **A.** Experimental setup to measure  $[\text{Ca}^{2+}]_r$  in vivo with  $\text{Ca}^{2+}$  selective microelectrodes in *vastus lateralis* muscle. **B.** Averaged data from  $[\text{Ca}^{2+}]_r$  determinations with or without nifedipine treatment. Data are expressed as mean  $\pm$  S.E.M. from *n* fibers in *N* mice, \*\*\* $P < 0.001$ , ANOVA-Tukey's.

doi:10.1371/journal.pone.0081222.g003

( $P < 0.001$ ), 91% ( $P < 0.01$ ) and 80% ( $P < 0.05$ ), respectively, with no significant effect in *wt* muscles.

#### Nifedipine Treatment Normalizes Pro-apoptotic Genes Expression in *mdx* Mice


Necrosis is probably one of the major contributors to DMD pathology [27]. However, there is some evidence that suggests that apoptotic pathways could also be important [28,29]. To determine if nifedipine can modify pro-apoptotic gene expression, we determined mRNA levels of both Bax and BIM in diaphragm muscles using real-time PCR. Bax mRNA levels were higher in *mdx* compared with *wt* mice (2.0-fold,  $P < 0.05$ ) (Figure 7) and nifedipine administration significantly diminished mRNA levels of Bax in *mdx* diaphragm by 73% ( $P < 0.001$ ) with no significant effect

in *wt* diaphragm. We did not find any differences in BIM mRNA levels between any of the studied groups.

#### Nifedipine Treatment Diminishes Serum CK Levels and Improves Muscle Strength in *mdx* Mice

To elucidate whether nifedipine treatment could reduce muscle damage, we measured serum CK levels in both *wt* and *mdx* mice after 1 week of either saline or nifedipine treatment. CK levels were significantly higher in saline treated *mdx* mice relative to saline treated *wt* mice ( $15.9 \pm 2.6$  vs  $0.7 \pm 0.3$  KUI/L,  $P < 0.001$ ) (Figure 8A). Nifedipine reduced CK levels in *mdx* mice ( $6.8 \pm 2.5$  KUI/L,  $P < 0.05$ ), without any significant effect in *wt* mice ( $1.5 \pm 0.2$  KUI/L,  $P > 0.05$ ).


In order to determine whether nifedipine treatment might increase muscle strength we used the inverted grid-hanging test



**Figure 4. Extracellular ATP concentration in FDB fibers isolated from either nifedipine- or saline-treated *wt* and *mdx* mice.** **A.** Time course of extracellular ATP levels after media change. ATP concentration was measured with CellTiter-Glo<sup>®</sup> Luminescent Cell Viability Assay. Average extracellular ATP at 30 min (**B**) and at 39 min (**C**) after media change are shown in the figure. Data are expressed as mean  $\pm$  S.E.M. FDB fibers were cultured from *n* = mice are indicated in the figure. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , ANOVA-Tukey's.

doi:10.1371/journal.pone.0081222.g004





**Figure 5. Apyrase treatment reduces [Ca<sup>2+</sup>]<sub>i</sub> in FDB adult fibers.** Adult fibers were isolated from *wt* and *mdx* mice and incubated in Krebs Ringer solution with or without apyrase (2 U/mL) for 10 min at room temperature. [Ca<sup>2+</sup>]<sub>i</sub> was determined by double-barreled Ca<sup>2+</sup> selective microelectrodes. Data are expressed as mean ± S.E.M. from *n* fibers (indicated in the figure) from three different cultures. n.s., no significant difference, \*\*\**P*<0.001, ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g005

and the forced swimming test. Mice were treated daily with nifedipine (1 mg/Kg) or saline for 1 week, and the functional test was performed one day after cessation of treatment. Saline treated *mdx* mice had a significantly shorter hanging time compared to *wt* mice (23±3 s vs 44±1 s, *P*<0.01 Figure 8B). Nifedipine treatment increased the *mdx* hanging time by 61% (37±3 s, *P*<0.05 compared to saline-treated mice (Figure 8B).

Similar results were observed using the forced swimming test. Under this experimental condition, *wt* swimming time was significantly higher compared to saline-treated *mdx* mice (550±25 s vs 398±15 s, *P*<0.05, Figure 8C). In this case nifedipine treatment for 1 week normalized the swimming time


in *mdx* mice increasing it to 594±64 s (*P*<0.01 compared to saline-treated mice, Figure 8C).

## Discussion


Our data show that [Ca<sup>2+</sup>]<sub>i</sub> was elevated in *mdx* myotubes and to a similar extent in adult skeletal muscle fibers from *mdx* mice. Acute exposure of *mdx* myotubes to nifedipine decreased [Ca<sup>2+</sup>]<sub>i</sub>, NF-κB activity and iNOS expression. Likewise, in *mdx* mice, nifedipine treatment for 1-week lowered *in vivo* [Ca<sup>2+</sup>]<sub>i</sub> in the *vastus lateralis*, reduced ATP release in FDB fibers, diminished mRNA levels of pro-oxidative/apoptotic genes in the diaphragm, lowered serum CK levels and improved muscle function as assessed by both inverted grid-hanging test and forced swimming test.

Although still controversial, it is generally accepted that dystrophic skeletal muscle cells have elevated [Ca<sup>2+</sup>]<sub>i</sub> [4,30–32]. Our data support these findings showing that [Ca<sup>2+</sup>]<sub>i</sub> was elevated in *mdx* myotubes and to a similar extent in skeletal muscles from *mdx* mice, measured either in isolated FDB fibers or determined *in vivo*, in anesthetized mice. Moreover, we showed that either acute nifedipine treatment of myotubes or chronic nifedipine treatment of mice for 1-week reduced [Ca<sup>2+</sup>]<sub>i</sub>. Nifedipine is known as specific inhibitor of the DHPR [33]. In addition to its role as an L-type Ca<sup>2+</sup> channel the DHPR operates as voltage sensor for both excitation-contraction and excitation-transcription coupling in adult muscle fibers [8–10]. Dystrophic *mdx* skeletal muscle cells have a dysregulated excitation-contraction coupling as evidenced by reduced calcium transients evoked by single action potentials compared with *wt* fibers [34,35]. This has been used to explain the muscle weakness observed in DMD patients. On the other hand, there is a controversy related to whether there are alterations of L-type Ca<sup>2+</sup> currents in *mdx* muscles, with some studies showing normal maximum currents [5] while others show decreased maximum current [6,7]. No differences have been found in charge movement between normal and dystrophic skeletal muscle fibers [5,36]. In addition, a leftward-shift of the voltage dependence of L-type Ca<sup>2+</sup> current has been described [5].

Our data showed that extracellular ATP levels were higher in *mdx* fibers compared to *wt* fibers. Chronic nifedipine treatment reduces ATP release in adult *mdx* FDB fibers to a level similar to *wt* fibers. Moreover, acute enzymatic ablation of extracellular ATP-



**Figure 6. iNOS and NADPH oxidase subunits gene expression in diaphragm muscles.** Diaphragms were dissected from nifedipine- and saline-treated mice and mRNA levels were assessed by real time PCR. **A.** iNOS, **B.** gp91<sup>phox</sup> and **C.** p47<sup>phox</sup> expression. Data are expressed as mean ± S.E.M. Diaphragms were obtained from *n*=mice as indicated in the figure. \**P*<0.05, \*\**P*<0.01, \*\*\**P*<0.001, ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g006




**Figure 7. Bax and BIM gene expression in diaphragm muscles.** Diaphragms were dissected from nifedipine- and saline-treated mice and mRNA levels were assessed by real time PCR. **A.** Bax, **B.** BIM mRNA levels. Data are expressed as mean  $\pm$  S.E.M. Diaphragms were obtained from *n* = mice as indicated in the figure, \* $P$  < 0.05, \*\*\* $P$  < 0.001, ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g007

ADP with apyrase treatment reduces the  $[Ca^{2+}]_i$  in *mdx* fibers from 296 nM to 195 nM, showing that elevated extracellular ATP is an important modulator of  $[Ca^{2+}]_i$  in *mdx* skeletal muscle fibers. Extracellular ATP can signal through  $P_2X$  (inotropic) and  $P_2Y$  (metabotropic) purinergic receptors [23,37], the latter pathway leading to  $Ca^{2+}$  release through inositol tri-phosphate receptors ( $IP_3R$ ). We previously reported that  $IP_3$  levels were higher in both human DMD and *mdx* cell lines [38] and that either phospholipase C (U-73122) or  $IP_3R$  (Xestospongine C) inhibition could significantly reduce  $[Ca^{2+}]_i$  in *mdx* myotubes [4].


Pannexin-1 channels are involved in ATP release after depolarization in myotubes and adult muscle fibers [9,10]. Based on co-immunoprecipitation and a proximity ligation assay [10], an

interaction between DHPR and Pannexin-1 channels has been proposed. Recently, we have shown that the DHPR is an important modulator of ATP release, and that it is necessary for fast-to-slow phenotype transition in FDB adult muscle fibers stimulated at 20 Hz [10]. ATP release through pannexin-1 channels observed after 20 Hz electrical stimulation of adult muscle fibers is inhibited by nifedipine [10], suggesting that the DHPR could directly control ATP release via Pannexin-1 channels.

Due to a chronic state of fiber injury, dystrophic muscle would be expected to contain high levels of extracellular ATP. In addition, ATP release to the extracellular medium was shown to be elevated in muscle fibers isolated from *mdx* mice [12]. Several



**Figure 8. Nifedipine treatment reduced serum CK and increases muscle function in *mdx* mice.** **A.** Blood samples were collected by cardiac puncture under anesthesia from both nifedipine- or saline-treated mice. CK activities were determined by the UV kinetic method. **B.** Averaged hanging time obtained in the inverted grid-hanging test in saline- and nifedipine treated mice. **C.** Averaged swimming time obtained in the forced swimming test in nifedipine- or saline- treated *mdx* mice. Data are expressed as mean  $\pm$  S.E.M. *n* = mice is indicated in the figure, \* $P$  < 0.05, \*\*\* $P$  < 0.001 ANOVA-Tukey's. doi:10.1371/journal.pone.0081222.g008



**Figure 9. Proposed model for ATP-mediated effects in dystrophic skeletal muscle.** In dystrophic skeletal muscle fibers there is an increase in basal ATP release, through Pannexin1 channels that is modulated by the DHPR [12]. Extracellular ATP increases the  $[Ca^{2+}]_i$  in dystrophic muscle fibers through the activation of purinergic receptors ( $P_2X$ , inotropic and  $P_2Y$ , metabotropic). This leads to the expression of pro-apoptotic and pro-inflammatory genes, increasing the muscle damage observed in dystrophic skeletal muscle cells. Nifedipine treatment reduces the basal ATP release and reduces  $[Ca^{2+}]_i$ , resulting in less pro-inflammatory and pro-apoptotic gene expression and subsequently reduces muscle damage as indicated by a decrease in blood CK and an increase in muscle function assessed by the inverted grid-hanging test and the force swimming test. doi:10.1371/journal.pone.0081222.g009

alterations have been found in ATP signaling in dystrophic skeletal muscle cells. In an immortalized myoblast cell line derived from *mdx* mouse, addition of exogenous ATP to the media induced a large increase in cytosolic  $Ca^{2+}$  concentration compared with its *wt* counterpart [39]. This increased susceptibility to ATP was associated with changes in expression and function of  $P_2X$  channels and to pathogenic  $Ca^{2+}$  entry in dystrophic muscles. Furthermore, enhanced expression of the  $P_2X_4$  and  $P_2X_7$  receptors have been related to macrophage invasion in *mdx* mice [40].

Here we determined the expression of iNOS and NADPH oxidase subunits in diaphragm, because this muscle has been described as the most severely affected muscle in the *mdx* mouse [26]. Nifedipine reduced iNOS, gp91<sup>phox</sup> and p47<sup>phox</sup> mRNA levels in *mdx* diaphragm after 1-week of treatment. We previously demonstrated that iNOS expression was higher in *mdx* myotubes which we attributed to an up-regulated NF- $\kappa$ B activity mediated by the elevated  $[Ca^{2+}]_i$  [4]. Nifedipine treatment reduced iNOS expression in diaphragm, likely due to the same mechanism.

Recently, it has been shown that dihydropyridines have anti-inflammatory and anti-oxidant effects. Nifedipine caused concentration-dependent inhibitory effects on sarcolemmal lipid peroxidation *in vitro* [41]. Toma *et al.*, 2011 suggest that amlodipine, a third generation dihydropyridine L-type calcium channel blocker, may improve endothelial dysfunction in diabetics through anti-oxidant and anti-inflammatory mechanisms. The authors stimu-

lated human endothelial cells with irreversible glycosylated low-density lipoproteins (AGE-LDL), as an *in vitro* model mimicking the diabetic condition. Their results show that amlodipine reduced the expression of NADPH oxidase subunits (p22<sup>phox</sup> and NOX4) and iNOS, and diminished oxidative/nitrosative stress. Moreover, amlodipine had anti-inflammatory effects reducing the activation of MCP-1, VCAM-1, p38 MAPK and NF- $\kappa$ B [42]. In another study, it has been shown that amlodipine (3 mg/kg/day) decrease the expression of NADPH subunits (p47<sup>phox</sup> and rac1), NADPH activity and the expression of inflammatory factor in atherosclerotic lesions [43].

Necrosis is probably the major contributor to DMD pathology [27]. However, there is some evidence that suggests that apoptotic pathways could also be important [28,29]. We observed that nifedipine reduced significantly Bax expression in *mdx* diaphragms, suggestive of a protective anti-apoptotic effect. Bax was abundantly expressed in the *mdx* masseter muscles compared to *wt* muscles at 3 weeks after birth [29]. We have demonstrated that exogenous ATP increases mRNA levels of several pro-apoptotic genes in skeletal muscle fibers, including Bax, BIM and PUMA. Moreover, ATP induced Bax activation and cytochrome C release in *mdx* fibers, which is related with apoptotic cell death [12].

A previous double-blind controlled clinical trial with nifedipine did not demonstrate a beneficial therapeutic response on mean muscle strength, joint contractures, time-functional tests, pulmonary function, or creatine kinase levels in 105 patients between 3–



27 years of age at the end of the study who were presumed to have DMD [13]. Patients were given nifedipine 0.75 mg/kg/day in three divided oral doses for 6 months and then 1.5–2 mg/Kg/day for the final 12 months of the study [13]. A total of 97 patients completed the trial. The power of the study to detect a difference between groups was calculated using the decrease of average muscle strength over time as the primary outcome measure. Average muscle strength score were calculated with a modified manual testing scale (MMT) [14] and the study had a power greater than 0.99 to detect a slowing of the illness to 25% of its original progression. The diagnostic criteria were based on semiology and classical human genetics, however three patients classified as Becker dystrophy were included in the study. Moreover the one patient aged 27 years old was considered to be unusual for DMD [44]. The data presented showed that the average muscle strength score as slightly improved without any significant difference. The trial used an unpaired analysis and it may have been more appropriate to use a paired analysis so that before and after changes in individuals could be compared [44].

As we mentioned before, there is an inverse correlation between severity of disease and dystrophin expression [15] and even patients with a complete lack of dystrophin DMD can be divided into 4 sub-phenotypes with different cognitive and motor outcomes [16]. This shows the high variability in the natural history of DMD. Phenotypic variations have shown to compromise results of clinical trials [45]. Desguerre *et al.*, suggest that trials, which are in danger of being inconclusive due to lack of precise knowledge of DMD's natural history, would strongly benefit from accurate selection of clinically homogeneous patient subsets [16].

High CK levels observed in DMD patients are attributed to an enhanced membrane permeability and muscle damage [46]. In our study, nifedipine treatment significantly reduced CK levels in *mdx* mice after 1-week, but did not normalize it. These data are supported by a study using verapamil, another L-type calcium channel blocker, which showed a significant reduction in the release of CK and LDH from diaphragm and gastrocnemius muscles [47]. In the previous nifedipine clinical trial, authors did not observe any significant difference in averaged CK levels after treatment, reporting that patients from nifedipine group had an

average CK level of 2681 U/liter at the start and a value of 2045 U/liter at the end [13]. High levels of CK in DMD are well-known, and after a rise in infancy, they remain at high levels with an abrupt decrease at around 10 years old due to loss of muscle mass [48]. Thus averaged data from a broad age group could easily miss a significant difference when and if one existed.

Reduction of  $[Ca^{2+}]_r$  and in the expression of pro-oxidative/apoptotic proteins, could be beneficial to dystrophic muscles and may explain a reduction in muscle damage that a decrease in serum CK would indicate. Moreover, an increase in the hanging time and normalization of the forced swimming test were observed in nifedipine-treated *mdx* mice, demonstrating an increase in motor function, probably due to reduced muscle damage or increased regeneration. In summary, these results provide further evidence that  $[Ca^{2+}]_r$  is elevated in *mdx* muscles and this can be modulated *in vivo* by nifedipine administration through a reduction in basal ATP release from dystrophic fibers. Moreover, nifedipine treatment reduced pro-oxidative/apoptotic gene expression with the end result being less muscle damage as evidenced by a significant reduction of serum CK and an increased muscle strength in *mdx* mice (Figure 9). Our study used daily intraperitoneally doses of nifedipine (1 mg/Kg) for 1 week as compared to the clinical study, which used it for a longer time period via the oral route of administration. Moreover, our study was carried out in 5–6 week old *mdx* mice (necrosis/regeneration stage) compared to the clinical study that included patients at different stages of the disease (3–27 years old). Longer studies need to be carried out to assess the positive effects and pitfalls of nifedipine treatment studies in *mdx* mice. However, our results strongly suggest that blockers of the ATP signaling pathway should be tested in DMD patients, as they might be promising in palliating the disease and prolonging muscle function.

## Author Contributions

Conceived and designed the experiments: FA DV JRL PDA EJ. Performed the experiments: FA DV CHO JRL. Analyzed the data: FA DV CHO MC EJ. Contributed reagents/materials/analysis tools: PDA EJ. Wrote the paper: FA MC PDA EJ.

## References

- Blake DJ, Weir A, Newey SE, Davies KE (2002) Function and genetics of dystrophin and dystrophin-related proteins in muscle. *Physiol Rev* 82: 291–329.
- Koenig M, Hoffman EP, Bertelson CJ, Monaco AP, Feener C, et al. (1987) Complete cloning of the Duchenne muscular dystrophy (DMD) cDNA and preliminary genomic organization of the DMD gene in normal and affected individuals. *Cell* 50: 509–517.
- Emery AE (2002) The muscular dystrophies. *Lancet* 359: 687–695.
- Altamirano F, Lopez JR, Henriquez C, Molinski T, Allen PD, et al. (2012) Increased resting intracellular calcium modulates NF-kappaB-dependent inducible nitric-oxide synthase gene expression in dystrophic *mdx* skeletal myotubes. *J Biol Chem* 287: 20876–20887.
- Collet C, Csernoch L, Jacquemond V (2003) Intramembrane charge movement and L-type calcium current in skeletal muscle fibers isolated from control and *mdx* mice. *Biophys J* 84: 251–265.
- Friedrich O, von Wegner F, Chamberlain JS, Fink RH, Rohrbach P (2008) L-type  $Ca^{2+}$  channel function is linked to dystrophin expression in mammalian muscle. *PLoS One* 3: e1762.
- Imbert N, Vandebrouck C, Dupont G, Raymond G, Hassoni AA, et al. (2001) Calcium currents and transients in co-cultured contracting normal and Duchenne muscular dystrophy human myotubes. *J Physiol* 534: 343–355.
- Tanabe T, Beam KG, Powell JA, Numa S (1988) Restoration of excitation-contraction coupling and slow calcium current in dysgenic muscle by dihydropyridine receptor complementary DNA. *Nature* 336: 134–139.
- Buvinic S, Almarza G, Bustamante M, Casas M, Lopez J, et al. (2009) ATP released by electrical stimuli elicits calcium transients and gene expression in skeletal muscle. *J Biol Chem* 284: 34490–34505.
- Jorquera G, Altamirano F, Contreras-Ferrat A, Almarza G, Buvinic S, et al. (2013) Cav1.1 controls frequency-dependent events regulating adult skeletal muscle plasticity. *J Cell Sci*.
- Knudson CM, Hoffman EP, Kahl SD, Kunkel LM, Campbell KP (1988) Evidence for the association of dystrophin with the transverse tubular system in skeletal muscle. *J Biol Chem* 263: 8480–8484.
- Valladares D, Almarza G, Contreras A, Pavez M, Buvinic S, et al. (2013) Electrical stimuli are anti-apoptotic in skeletal muscle via extracellular ATP. Alterations of this signal in *mdx* mice is a likely cause of dystrophy. *PLoS One*. In Press.
- Moxley RT 3rd, Brooke MH, Fenichel GM, Mendell JR, Griggs RC, et al. (1987) Clinical investigation in Duchenne dystrophy. VI. Double-blind controlled trial of nifedipine. *Muscle Nerve* 10: 22–33.
- Brooke MH, Fenichel GM, Griggs RC, Mendell JR, Moxley R, et al. (1983) Clinical investigation in Duchenne dystrophy: 2. Determination of the “power” of therapeutic trials based on the natural history. *Muscle Nerve* 6: 91–103.
- Nicholson LV, Johnson MA, Bushby KM, Gardner-Medwin D, Curtis A, et al. (1993) Integrated study of 100 patients with Xp21 linked muscular dystrophy using clinical, genetic, immunochemical, and histopathological data. Part 3. Differential diagnosis and prognosis. *J Med Genet* 30: 745–751.
- Desguerre I, Christov C, Mayer M, Zeller R, Becane HM, et al. (2009) Clinical heterogeneity of duchenne muscular dystrophy (DMD): definition of sub-phenotypes and predictive criteria by long-term follow-up. *PLoS One* 4: e4347.
- Casas M, Altamirano F, Jaimovich E (2012) Measurement of calcium release due to inositol trisphosphate receptors in skeletal muscle. *Methods Mol Biol* 798: 383–393.

18. Eltit JM, Ding X, Pessah IN, Allen PD, Lopez JR (2012) Nonspecific sarcolemmal cation channels are critical for the pathogenesis of malignant hyperthermia. *FASEB J*.
19. Ho CL, Yang CY, Lin WJ, Lin CH (2013) Ecto-Nucleoside Triphosphate Diphosphohydrolase 2 Modulates Local ATP-Induced Calcium Signaling in Human HaCaT Keratinocytes. *PLoS One* 8: e57666.
20. Yoshida H, Kobayashi D, Ohkubo S, Nakahata N (2006) ATP stimulates interleukin-6 production via P2Y receptors in human HaCaT keratinocytes. *Eur J Pharmacol* 540: 1–9.
21. Kaja S, van de Ven RC, van Dijk JG, Verschuuren JJ, Arahata K, et al. (2007) Severely impaired neuromuscular synaptic transmission causes muscle weakness in the Cacna1a-mutant mouse rolling Nagoya. *Eur J Neurosci* 25: 2009–2020.
22. Razani B, Wang XB, Engelman JA, Battista M, Lagaud G, et al. (2002) Caveolin-2-deficient mice show evidence of severe pulmonary dysfunction without disruption of caveolae. *Mol Cell Biol* 22: 2329–2344.
23. Burnstock G (2006) Purinergic P2 receptors as targets for novel analgesics. *Pharmacol Ther* 110: 433–454.
24. Bellinger AM, Reiken S, Carlson C, Mongillo M, Liu X, et al. (2009) Hypernitrosylated ryanodine receptor calcium release channels are leaky in dystrophic muscle. *Nat Med* 15: 325–330.
25. Whitehead NP, Yeung EW, Froehner SC, Allen DG (2010) Skeletal muscle NADPH oxidase is increased and triggers stretch-induced damage in the *mdx* mouse. *PLoS One* 5: e15354.
26. Stedman HH, Sweeney HL, Shrager JB, Maguire HC, Panettieri RA, et al. (1991) The *mdx* mouse diaphragm reproduces the degenerative changes of Duchenne muscular dystrophy. *Nature* 352: 536–539.
27. Miller JB, Girgenrath M (2006) The role of apoptosis in neuromuscular diseases and prospects for anti-apoptosis therapy. *Trends Mol Med* 12: 279–286.
28. Tews DS, Goebel HH (1997) DNA-fragmentation and expression of apoptosis-related proteins in muscular dystrophies. *Neuropathol Appl Neurobiol* 23: 331–338.
29. Honda A, Abe S, Hiroki E, Honda H, Iwanuma O, et al. (2007) Activation of caspase 3, 9, 12, and Bax in masseter muscle of *mdx* mice during necrosis. *J Muscle Res Cell Motil* 28: 243–247.
30. Allen DG, Gervasio OL, Yeung EW, Whitehead NP (2010) Calcium and the damage pathways in muscular dystrophy. *Can J Physiol Pharmacol* 88: 83–91.
31. Lopez JR, Briceno LE, Sanchez V, Horvart D (1987) Myoplasmic (Ca<sup>2+</sup>) in Duchenne muscular dystrophy patients. *Acta Cient Venez* 38: 503–504.
32. Turner PR, Westwood T, Regen CM, Steinhardt RA (1988) Increased protein degradation results from elevated free calcium levels found in muscle from *mdx* mice. *Nature* 335: 735–738.
33. Neuhaus R, Rosenthal R, Luttgau HC (1990) The effects of dihydropyridine derivatives on force and Ca<sup>2+</sup> current in frog skeletal muscle fibres. *J Physiol* 427: 187–209.
34. Capote J, DiFranco M, Vergara JL (2010) Excitation-contraction coupling alterations in *mdx* and utrophin/dystrophin double knockout mice: a comparative study. *Am J Physiol Cell Physiol* 298: C1077–1086.
35. Woods CE, Novo D, DiFranco M, Vergara JL (2004) The action potential-evoked sarcoplasmic reticulum calcium release is impaired in *mdx* mouse muscle fibres. *J Physiol* 557: 59–75.
36. Hollingworth S, Marshall MW, Robson E (1990) Excitation contraction coupling in normal and *mdx* mice. *Muscle Nerve* 13: 16–20.
37. Abbracchio MP, Burnstock G, Boeynaems JM, Barnard EA, Boyer JL, et al. (2006) International Union of Pharmacology LVIII: update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol Rev* 58: 281–341.
38. Liberona JL, Powell JA, Shenoi S, Petherbridge L, Caviedes R, et al. (1998) Differences in both inositol 1,4,5-trisphosphate mass and inositol 1,4,5-trisphosphate receptors between normal and dystrophic skeletal muscle cell lines. *Muscle Nerve* 21: 902–909.
39. Yeung D, Zablocki K, Lien CF, Jiang T, Arkle S, et al. (2006) Increased susceptibility to ATP via alteration of P2X receptor function in dystrophic *mdx* mouse muscle cells. *FASEB J* 20: 610–620.
40. Yeung D, Kharidia R, Brown SC, Gorecki DC (2004) Enhanced expression of the P2X4 receptor in Duchenne muscular dystrophy correlates with macrophage invasion. *Neurobiol Dis* 15: 212–220.
41. Mak IT, Weglicki WB (1990) Comparative antioxidant activities of propranolol, nifedipine, verapamil, and diltiazem against sarcolemmal membrane lipid peroxidation. *Circ Res* 66: 1449–1452.
42. Toma L, Stancu CS, Sanda GM, Sima AV (2011) Anti-oxidant and anti-inflammatory mechanisms of amlodipine action to improve endothelial cell dysfunction induced by irreversibly glycated LDL. *Biochem Biophys Res Commun* 411: 202–207.
43. Yoshii T, Iwai M, Li Z, Chen R, Ide A, et al. (2006) Regression of atherosclerosis by amlodipine via anti-inflammatory and anti-oxidative stress actions. *Hypertens Res* 29: 457–466.
44. Phillips MF, Quinlivan R (2008) Calcium antagonists for Duchenne muscular dystrophy. *Cochrane Database Syst Rev*: CD004571.
45. Escolar DM, Buysse G, Henricson E, Leshner R, Florence J, et al. (2005) CINTRG randomized controlled trial of creatine and glutamine in Duchenne muscular dystrophy. *Ann Neurol* 58: 151–155.
46. Ozawa E, Hagiwara Y, Yoshida M (1999) Creatine kinase, cell membrane and Duchenne muscular dystrophy. *Mol Cell Biochem* 190: 143–151.
47. Niebrój-Dobosz I, Lukasiuk M (1996) Release of Intracellular Enzymes from Skeletal Muscles and Diaphragm in *Mdx* Mice. *European Journal of Translational Myology* 6: 377–383.
48. Konagaya M, Takayanagi T (1986) Regularity in the change of serum creatine kinase level in Duchenne muscular dystrophy. A study with long-term follow-up cases. *Jpn J Med* 25: 2–8.