

Published in final edited form as:

. 2008 May 1; 40(2): 439–445. doi:10.1657/1523-0430(07-039)[WIESER]2.0.CO;2.

## The water balance of grassland ecosystems in the Austrian Alps

Gerhard Wieser<sup>1</sup>, Albin Hammerle<sup>2</sup>, and Georg Wohlfahrt<sup>2</sup>

<sup>1</sup>Dept. Alpine Timberline Ecophysiology, Federal Research and Training Centre for Forests, Natural Hazards and Landscape, Rennweg 1, 6020 Innsbruck, Austria

<sup>2</sup>Institute of Ecology, University of Innsbruck, Sternwartestraße 15, 6020 Innsbruck, Austria

### Abstract

The altitudinal variation of precipitation, evapotranspiration and runoff was quantified at 16 different grassland sites between 580 and 2550 m a.s.l. in the Austrian Alps. Along this altitudinal transect annual evapotranspiration decreased from roughly 690 mm at low elevation sites to 210 – 220 mm at the upper limit of the alpine grassland belt. A detailed analysis of the data showed that the observed reduction in the annual sum of evapotranspiration could be mainly explained by the altitudinal decline of the length of the snow free period (i.e. the vegetation period). Daily mean sums of evapotranspiration showed no altitudinal trend and averaged 2.2 mm d<sup>-1</sup> independent of elevation, although the leaf area index, growing season mean air temperature, and vapour pressure deficit declined with increasing altitude. As precipitation increased with elevation, evapotranspiration seems to be of secondary importance when compared to runoff. Inter-annual variability of evapotranspiration was fairly low across contrasting dry and wet years (coefficient of variation = 7 %), indicating that even during dry years water availability was not limiting evapotranspiration.

### Introduction

Grasslands are one of the most widespread vegetation types and cover approximately one-fifth of the earth's land surface (Hall and Scurlock, 1991). In the Austrian Alps a variety of grassland ecosystems such as meadows, pastures, alpine grass and sedge mats cover approximately 40 to 45 % of the total land area and extend from the valley floor to the upper limit of the alpine grassland belt at approximately 2600 m a.s.l. (Schiechl and Stern, 1974; Laboratoire des Ecosystèmes Alpines, 1995). Altitudinal alterations in climate (Schröter, 1926; Steinhäuser et al., 1960; Ellenberg, 1996; Franz, 1979; Baumgartner 1980, Fliri, 1975, Veit 2002), edaphic parameters (Franz, 1979) and land use (Cernusca et al., 1999) cause significant changes in species composition, biomass production and canopy structure of grassland ecosystems in the Austrian Alps (Cernusca and Seeber, 1981).

The water balance of montane, subalpine and alpine (sensu Ellenberg, 1996) grassland ecosystems is an important feature, especially in the Austrian Alps where precipitation exceeds the water use of vegetation and the storage capacity of soils. Vegetation is an important factor influencing the annual ecosystem water balance, which is determined by the water input through precipitation (P) which falls as rain, snow, or fog. Some of the P adheres on the above-ground vegetation and evaporates before reaching the ground - the so-called interception (I). The remaining P reaching the ground runs off (R<sub>S</sub>, surface runoff), infiltrates into the soil and percolates through to the water table (R<sub>D</sub>, deep seepage), finally

Corresponding author and proofs to: Gerhard Wieser, Dept. Alpine Timberline Ecophysiology, Federal Research and Training Centre for Forests, Natural Hazards and Landscape, Rennweg 1, 6020 Innsbruck, Austria, Phone ++43-512 573933 5120, Fax ++43-512 57399 5150, Gerhard.Wieser@uibk.ac.at.

leaving the system as base flow. The amount of water which is held against gravitational forces within the soil matrix can be taken up by the plant roots and is finally transpired through the stomata of the leaves into the atmosphere (T, transpiration) and evaporates from the soil surface (E). So, over short periods, there are also changes in the soil water content ( $\Delta W$ ). All the components involved are given in mm ( $= \text{kg m}^{-2}$  of slope parallel ground) and the complete water balance equation of an ecosystem reads:

$$P = E + T + I + R_d + R_s + \Delta W \quad (\text{Eq.1})$$

In view of the fact that it is frequently very difficult to measure all the terms of Eq. 1 and taking into account that over long periods (an entire growing season or year)  $\Delta W$  may be assumed to approach zero, the water balance equation (Eq. 1) can be simplified to:

$$P = ET + R \quad (\text{Eq.2})$$

where ET is the sum of all water vapour fluxes from the system to the atmosphere (E+T+I) and R is the sum of deep seepage and surface runoff.

Earlier estimations of the annual water balance at different altitudes for the European Alps (Baumgartner, 1980) included estimations of ET based on meteorological data (Müller, 1965; Kern, 1975) or estimations of total R from catchment stream flow data (Steinhäusser, 1952, 1970). Since the mid eighties of the last century, lysimeter studies (Körner et al., 1978; Guggenberger, 1980; Wieser, 1983; Scott et al., 2002), the Bowen-ratio method (Cernusca and Seeber, 1981; Staudinger and Rott, 1981; Rosset et al., 2001), and the eddy covariance technique (Wohlfahrt, 2004; Hammerle et al., 2007) have also been used to measure ET for calculating the water balance of grassland ecosystems at various sites within the Austrian Alps.

The objective of this paper is to compile what is presently known about the water balance of grassland ecosystems in the Austrian Alps. We first examine the inter-annual and seasonal variability and the driving forces of ET using data from a long-term mountain grassland study in the Stubai valley. The following section deals with the altitudinal variation of ET and finally we focus on how the overall water balance of grassland ecosystems changes with elevation.

## Study sites and Methods

To characterize the effects of altitude on the various components of the water balance of grassland ecosystems we compare experimental data from 16 sites in the Austrian Alps between 580 and 2550 m a.s.l.. The selected sites differ with respect to geographical location, altitude, exposition, and inclination and (Table 1) and include montane and sub-alpine meadows and pastures below the timberline and alpine grassland ecosystems above the alpine timberline, respectively. Eight sites (stations 1 to 8) were located in the Hohe Tauern mountain range in the southern part of the Austrian Alps. Five stations were in the inner Alpine sector namely in the Ötztaler Alpen (station 9 and 10), the Stubai Alpen (station 11 and 12) and in the Tuxer Voralpen (station 13). In addition, there were also three stations situated along an altitudinal transect in the northern limestone Alps (stations 14 to 16). Figure 1 shows the location of the 16 sites.

A common set of precipitation, ET and runoff measurements was acquired at each site. Precipitation was estimated with Hellmann rain gauges installed 2 m above ground and throughout the growing season also with rain gauges installed slope parallel close to the soil surface (Henning, 1974). The latter are suggested to be more representative for P when focusing on the water availability of canopies during the snow free period (Stigter, 1974).

On average, growing season sums of P estimated slope parallel close to the soil surface were 6-11% higher than P values obtained by Hellmann rain gauges at 2 m above ground.

At all stations (see Table 1) except station 12 (Neustift), ET and total runoff was measured by means of weighting and non-weighting “monolithic” lysimeters (Courtin and Bliss, 1971; Körner et al., 1978; Wieser, 1983; Scott et al., 2002) sunk into the ground. The soil within the lysimeters was undisturbed, pores at the bottom of the lysimeter allowed drainage and forced all runoff into seepage. Evaporation from the snow surface was calculated using Sverdrup’s formula of the sublimation of snow (Müller, 1965). At the two sites in the Ötztaler Alpen and at Neustift ET was quantified using micrometeorological methods (Table 1): At the two sites in the Ötztaler Alpen (station 9 and 10) the Bowen-ratio method (Staudinger and Rott, 1981) was applied in addition to lysimeters; at the study site Neustift, (station 12) the eddy covariance technique was used for determining ET as described in Wohlfahrt (2004) and Hammerle et al. (2007). At Neustift, which is representative for the lower elevation grasslands of the investigated transects, ET has been measured since 2001 and this six year data set will be used to analyse the inter-annual and seasonal variability, as well as the controls on daily ET.

## Results and Discussion

### Inter-annual variability, seasonal trends and driving forces of evapotranspiration

As shown in Fig. 2, daily ET at the low-elevation site Neustift was generally very small from January through March (0.05-0.15 mm d<sup>-1</sup>), when snow usually covers the ground. After snow melt, evaporation rates quickly increased, reaching peak values of 4-5 mm d<sup>-1</sup> during the period May-October, after which ET gradually declined back to wintertime values (Fig. 2). These peak evaporation rates are in good agreement with other studies on well-watered grassland ecosystems (e.g. Meyers, 2001; Rosset et al., 2001; Tasser et al., 2001; Wever et al., 2002).

During 2001-2006 annual ET ranged from 446 mm (2005) to 527 mm (2006) (Fig. 2) or 53 % (2002) to 91 % (2006) of annual P, with coefficients of variation of 7 and 20 %, respectively. The potential driving forces considered (global and net radiation, air and soil temperature, vapour pressure deficit, soil water content and precipitation) explained little of this variability, the best predictor being air temperature, which in a linear regression explained 35 % of the variability in annual ET ( $p = 0.11$ ). Restricting the analysis to the vegetation period, where the majority of ET occurs (Fig. 2), improved regression statistics only slightly (data not shown). Once ET was normalised with precipitation, a statistically significant non-linear relationship was found with precipitation, shown in Fig. 3. There it can be seen that the fraction of precipitation evaporated to the atmosphere increased with decreasing precipitation and that this increase was asymmetric, ie. ET increased disproportional during years with little precipitation. We interpret this behaviour as to indicate that even during dry years, when up to 90% of the precipitation is evaporated, no water stress occurs in these systems, but that in contrast environmental conditions (higher evaporative demand, higher radiation input) lead to increased evaporation rates.

The largest fraction of variability in daily ET (excluding periods of snow cover) was explained by the amount of incident photosynthetically active radiation (Fig. 4,  $r^2 = 0.74$ ,  $p = 0.00$ ), followed by net radiation (data not shown,  $r^2 = 0.70$ ,  $p = 0.00$ ), which were though strongly correlated ( $r^2 = 0.92$ ,  $p = 0.00$ ). Similar correlations have also been observed for the two sites in the Ötztaler Alpen by Staudinger and Rott (1981). This reflects the energy demand for the evaporation of water (Campbell and Norman, 1998), as well as photosynthesis, which is linked to transpiration by stomatal conductance (Larcher, 2001). The good predictive power of these two radiation forcings again indicates that limitations of

transpiration by stomatal closure due to low soil water availability and/or dry air played a minor role at this site. This is corroborated by the dependency of daily ET on the vapour pressure deficit (Fig. 4,  $r^2 = 0.61$ ,  $p = 0.00$ ), whose slope tended to decrease only at very high vapour pressure deficits, as also observed for the sites Seppenbauer, Guttal, and Wallackhaus at the south rim of the Hohen Tauern by Wieser (1983). A step-wise linear regression analysis included incident photosynthetically active radiation and the vapour pressure deficit, the combined linear model explaining 77% of the observed variability in ET ( $p = 0.00$ ).

### Evapotranspiration with respect to altitude

For assessing the altitudinal variation in annual ET we compiled data from all 16 investigated grassland ecosystems (Table 1). These data clearly show, that the annual ET declined from 690 mm in an inner alpine valley floor (580 m a.s.l.; station 14) to 210 – 220 mm at the upper limit of the alpine grassland belt in the inner Alpine sector (2550 m a.s.l., station 16) and in the southern part of the Austrian Alps (2528 m a.s.l.; station 7), respectively (Fig. 5). There were no differences in the vertical gradient of ET between stations located in inner alpine sectors (stations 9 to 13) and in the southern (stations 1 to 8) and northern outer ranges (stations 14 to 16) of the central Austrian Alps (Fig. 5). The observed reduction in ET of 210 mm per 1000 m increase in altitude (Fig. 5;  $r^2 = 0.86$ ,  $p=0,00$  is within the range of earlier estimations for other parts of the central Alps based on catchment stream flow analysis (170 mm km<sup>-1</sup>, Steinhäüßer, 1970; 185 mm km<sup>-1</sup>, Koehl, 1971; 356 mm km<sup>-1</sup> Steinhäüßer 1952) or meteorological calculations (160 mm km<sup>-1</sup>, Müller, 1965; 180 mm km<sup>-1</sup>, Kern, 1975). Snow sublimation during the winter months was only a minor fraction of annual ET and varied between 13 mm at the valley floor and 45 mm at the upper limit of the alpine grassland belt (Fig. 5).

The observed linear altitudinal decline in the annual ET (Fig. 5) can be completely explained by the altitudinal reduction of the vegetation period (Wieser, 1983; Wieser et al., 1984; Menzel et al., 1998; Körner, 2003). When the growing season water loss was divided by the number of snow-free days (see Table 1), the altitudinal gradient disappeared and the daily mean of grassland ET during the growing season ranged from 1.9 to 2.6 mm d<sup>-1</sup> and averaged 2.2 mm d<sup>-1</sup> (Fig. 5; Wieser, 1983; Wieser et al., 1984). The growing season mean ET of the grasslands under study is similar to the value of 2.0 mm d<sup>-1</sup> obtained for mixed forest stands in the Alps between 490 and 1950 m a.s.l. (Matyssek et al., pers. comm.).

The leaf area index of grassland ecosystems declines from peak values of 6-8 in the valley floor (Körner and Mayr, 1981; Wohlfahrt et al., 2001) to less than 2 in the alpine grassland belt (Cernusca and Seeber, 1981). As a consequence of the declining leaf area index, wind velocity (Wohlfahrt and Cernusca, 2002) and aerodynamic conductance were found to be significantly higher in the alpine grassland belt as compared to pastures and meadows at lower elevations (Cernusca and Seeber, 1981). Due to the altitudinal decline in leaf area, evaporation from the soil surface below the vegetation cover increases from 10 % of total ET in the valley floor (Rutter, 1975) up to approximately 25% in the alpine sedge belt (Körner, 1977; Körner et al., 1980). Moreover, due to heat accumulation in the canopy (Körner 2003), low stature alpine grasslands often heat up to a larger extent causing the driving forces for ET to increase (Cernusca and Seeber, 1981). Consequently, high elevation plants display higher leaf conductances (Körner and Mayr, 1981) and higher stomatal densities (Körner et al., 1989) than comparable plants at lower elevations. Periodic soil water shortage at low elevation sites with soil water potentials close to the wilting point cause stomatal restrictions much more often as compared to plants growing above the timberline (Körner and Mayr, 1981), where soil water potentials seldom drop below critical values (Neuwinger-Raschendorfer, 1961; Neuwinger, 1970; 1980; Gunsch, 1972; Wieser et al., 1989), even during the Central Europe-wide extremely dry summer of 2003 (Rebetez et

al., 2006; Wieser and Tausz, 2007). This may be attributed to the higher amount of precipitation (Fig. 6) and to the fact that rain is observed every third or fourth day on average above the timberline (Frei and Schär, 1998).

### The annual water balance with respect to altitude

Annual evapotranspiration normalised with precipitation (E/P ratio) decreased with increasing altitude (Fig. 6 left; all data pooled  $r^2 = 0.83$ ,  $p = 0.00$ ), conversely runoff normalised with precipitation (R/P value) increased with increasing altitude and equalled E/P at approximately 1300 m above sea level. These trends were much more pronounced on the southern outer rim as compared to the inner Alpine region and the Northern limestone Alps. This results from the fact that there is no unique relation between altitude and precipitation in the Alps (Schröter, 1926; Steinhauser et al., 1960; Ellenberg, 1996; Franz, 1979; Baumgartner, 1980; Fliri, 1975; Veit, 2002). On average, annual precipitation increased by approximately 75 mm per 100 m of altitude and by less than 15 mm per 100 m of altitude in the southern outer rims and the northern limestone Alps, respectively (Fig. 6, right). In the inner Alpine region, by contrast, precipitation did not change significantly with altitude and averaged 980 mm irrespective of height above sea level (Fig. 6, right).

### Conclusions

Our compilation of studies on the water balance of grassland ecosystems in the Austrian Alps lead to the following conclusions: (1) even during dry years grassland ecosystems in the Austrian Alps seem not to suffer from water stress, (2) the observed linear altitudinal decline in annual ET can be explained by the altitudinal shortening of the snow free period, and (3) with increasing altitude runoff, at the expense of ET, dominates the water balance.

The major part of studies contained in this compilation was conducted in the late 70ies of the last century. Given recent evidence of climate change impacts on mountain ecosystems, in particular the accelerated warming rates at high altitudes (Kromp-Kolb and Formayer, 2005), it may thus be timely to repeat those measurements in order to detect and quantify possible effects of climate change on the water balance of Austrian Alpine grassland ecosystems.

### References

- Baumgartner, A. Mountain climates from a perspective of forest growth. In: Benecke, U.; Davis, MR., editors. Mountain environments and subalpine forest growth; Proceedings IUFRO Workshop November 1979 Christchurch; New Zealand. New Zealand Forest Service, Wellington. 1980. p. 27-39. Technical Paper No. 70
- Campbell, GS.; Norman, J.M. An introduction to environmental biophysics. 2nd ed.. Springer Verlag; New York: 1998.
- Cernusca, A.; Seeber, MC. Canopy structure, microclimate and energy budget in different alpine plant communities. In: Grace, J.; Ford, ED.; Jarvis, PG., editors. Plants and their Atmospheric Environment; The 21st Symposium of the British Ecological Society Edinburgh; 1979; Oxford: Blackwell; 1981. p. 75-81.
- Cernusca, A.; Tappeiner, U.; Bayfield, N. Land-use changes in European mountain ecosystems. ECOMONT – concepts and results. Blackwell Wissenschaftsverlag; Berlin: 1999. p. 368
- Courtin GM, Bliss LC. A hydric lysimeter to measure evapotranspiration under remote field conditions. Arctic and Alpine Research. 1971; 3:81–89.
- Ellenberg, H. Vegetation Mitteleuropas mit den Alpen in ökologischer, dynamischer und historischer Sicht. Verlag Eugen Ulmer; Stuttgart: 1996.
- Fliri, F. Das Klima der Alpen im Raume von Tirol. Monographien zur Landeskunde Tirols I. Universitätsverlag Wagner; Innsbruck, München: 1975.




- Franz, H. *Ökologie der Hochgebirge*. Ulmer; Stuttgart: 1979.
- Frei, Ch.; Schär, Ch. A precipitation climatology of the Alps from high-resolution rain-gauge observations. *International Journal of Climatology*. 1998; 18:873–900.
- Guggenberger, H. *Untersuchungen zum Wasserhaushalt der alpinen Zwergstrauchheide Patschertkofel* [Ph.D. thesis]. Botany Innsbruck University; 1980.
- Gunsch, J. *Vergleichende ökologische Untersuchungen von Kleinstandorten im Bereich der subalpinen Zirben-Waldgrenze* [Ph.D. thesis]. Botany Innsbruck University; 1972.
- Hall DO, Scurlock JMO. Climate change and productivity of natural grasslands. *Annales of Botany*. 1991; 67(suppl.):49–55.
- Hammerle A, Haslwanger A, Schmitt M, Bahn M, Tappeiner U, Cernusca A, Wohlfahrt G. Eddy covariance measurements of carbon dioxide, latent and sensible energy fluxes above a meadow on a mountain slope. *Boundary-Layer Meteorology*. 2007; 122:397–416. [PubMed: 24465032]
- Henning D. Zur Regenmessung an Hängen. *Wetter und Leben*. 1974; 26:94–97.
- Kern, H. *Schriftenreihe des Bayerischen Landesamtes für Wasserwirtschaft*. Vol. 2. München: 1975.
- Koehl A. Die Landesverdunstung im schweizerischen Rheingebiet. *Jahresbericht der naturforschenden Gesellschaft Graubündens*, N.F. 1971; 94:193–198.
- Körner, Ch. Evapotranspiration und Transpiration verschiedener Pflanzenbestände im alpinen Grasheidegürtel der Hohen Tauern. In: Cernusca, A., editor. *Alpine Grasheide Hohe Tauern. Ergebnisse der Ökosystemstudie 1976*. Vol. 1. Universitätsverlag Wagner; Innsbruck: 1977. p. 47-68. Veröffentlichungen des österreichischen MaB-Hochgebirgsprogramms Hohe Tauern
- Körner, Ch. *Alpine plant life: functional plant ecology of high mountain ecosystems*. 2nd ed.. Springer; Berlin: 2003. p. 349
- Körner, Ch.; Mayr, R. Stomatal behaviour in alpine plant communities between 600 and 2600 metres above sea level. In: Grace, J.; Ford, ED.; Jarvis, PG., editors. *Plants and their Atmospheric Environment; The 21st Symposium of the British Ecological Society Edinburgh*; 1979; Oxford: Blackwell; 1981. p. 205-218.
- Körner, Ch.; Hoflacher, H.; Wieser, G. Untersuchungen zum Wasserhaushalt von Almflächen in Gasteiner-Tal. In: Cernusca, A., editor. *Ökologische Analysen von Almflächen im Gasteiner Tal*. Universitätsverlag Wagner; Innsbruck: 1978. p. 67-79. Veröffentlichungen des Österreichischen MaB-Hochgebirgsprogramms Hohe Tauern, Band 2
- Körner, Ch.; Wieser, G.; Guggenberger, H. Der Wasserhaushalt eines alpinen Rasens in den Zentralalpen. In: Franz, H., editor. *Untersuchungen an alpinen Böden in den Hohen tauern 1974-1978. Stoffdynamik und Wasserhaushalt*. Universitätsverlag Wagner; Innsbruck: 1980. p. 243-264. Veröffentlichungen des Österreichischen MaB-Hochgebirgsprogramms Hohe Tauern, Band 3
- Körner, Ch.; Neumayer, M.; Pelaez Menendez-Riedl, S.; Smeets-Scheel, A. Functional morphology of mountain plants. *Flora*. 1989; 182:353–383.
- Kromp-Kolb, H.; Formayer, H. *Schwarzbuch Klimawandel*. Verlag ecowin; Salzburg: 2005.
- Kronfuss, H. FBVA Bericht. Vol. 100. Wien: 1997. Das Klima einer Hochlagenaufforstung in der subalpinen Höhenstufe. Haggen im Sellraintal bei St. Sigmund, Tirol (Periode 1975-1994).
- Laboratoire des Ecosystèmes Alpines. *Bibliographie de documents pour la carte de la végétation des Alpes (Volumes I à X) et documents de la cartographie écologique (Volumes XI à XXXI)*. Université de Grenoble; 1995. p. 1-83.
- Larcher, W. *Ökophysiologie der Pflanzen: Leben, Leistung und Stressbewältigung der Pflanzen in ihrer Umwelt*. Ulmer; Stuttgart: 2001.
- Menzel, L.; Lang, H.; Rohrmann, M. *Hydrological Atlas of Switzerland, Plate 4.1*. Federal Office of Topography; Bern: 1998. Mean annual evapotranspiration 1973-1992.
- Meyers T. A comparison of summertime water and CO<sub>2</sub> fluxes over rangeland for well watered and drought conditions. *Agricultural and Forest Meteorology*. 2001; 106:205–214.
- Müller W. Zur Schätzung der Gebietsverdunstung im Gebirge. *Arciv für. Meerologie, Geophysik und Bioklimatologie Serie B*. 1965; 13:193–206.
- Neuwinger I. Böden der subalpinen und alpinen Stufe in den Tiroler Alpen. *Mitteilungen der Ostalpin-Dinarischen Gesellschaft für Vegetationskunde*. 1970; 11:135–150.


- Neuwinger I. Erwärmung, Wasserrückhalt und Erosionsbereitschaft subalpiner Böden. *Mitteilungen der Forstlichen Bundesversuchsanstalt Wien*. 1980; 139:113–144.
- Neuwinger-Raschendorfer I. Bodenfeuchtemessungen. *Mitteilungen der Forstlichen Bundesversuchsanstalt Mariabrunn*. 1961; 59:257–264.
- Rebetez M, Mayer H, Dupont O, Schindler D, Gartner K, Kropf JP, Menzel A. Heat and drought 2003 in Europe: a climate synthesis. *Annales of Forest Science*. 2006; 63:569–577.
- Rosset M, Montani M, Tanner M, Fuhrer J. Effects of abandonment on the energy balance and evapotranspiration of wet subalpine grassland. *Agriculture, Ecosystems and Environment*. 2001; 86:277–286.
- Rutter, AJ. The hydrological cycle in vegetation. In: Monteith, JL., editor. *Vegetation and Atmosphere*. Academic Press London; New York, San Francisco: 1975. p. 179–185.
- Schiechl HM, Stern R. Vegetationskartierung – Durchführung und Anwendung in Forschung und Praxis. *Festschrift 100 Jahre Forstliche Bundesversuchsanstalt Wien*. 1974:273–308.
- Schröter, C. *Das Pflanzenleben der Alpen. Eine Schilderung der Hochgebirgsflora*. Verlag Albert Raustein; Zürich: 1926.
- Scott D, Bayfield NG, Cernusca A, Elston DA. Use of a weighing lysimeter system to assess the effects of trampling on evapotranspiration of montane plant communities. *Canadian Journal of Botany*. 2002; 80:675–683.
- Staudinger M, Rott H. Evapotranspiration at two mountain sites during the vegetation period. *Nordic Hydrology*. 1981; 12:207–216.
- Steinhäusser H. Über die Gebietsverdunstung in den österreichischen Südalpen. *Berichte des deutschen Wetterdienstes*. 1952; 35:174–178.
- Steinhäusser H. Gebietsverdunstung und Wasservorrat in verschiedenen Seehöhen Österreichs. *Österreichische Wasserwirtschaft*. 1970; 22:163–170.
- Steinhauser, F.; Eckel, O.; Lauscher, F. *Klimatographie von Österreich*. Springer; Wien: 1960. p. 137–380.
- Stigter CJ. Precipitation measurements above a maize crop. *Archiv für Meteorologie, Geophysik und Bioklimatologie, Ser B*. 1974; 24:95–108.
- Tasser, E.; Tappeiner, U.; Cernusca, A. *Südtirols Almen im Wandel. Ökologische Folgen von Landnutzungsänderungen*. Europäische Akademie; Bozen: 2001. p. 269
- Veit, H. *Die Alpen – Geoökologie und Landschaftsentwicklung*. Ulmer; Stuttgart: 2002.
- Wever LA, Flanagan LB, Carlson PJ. Seasonal and interannual variation in evapotranspiration, energy balance and surface conductance in a northern temperate grassland. *Agricultural and Forest Meteorology*. 2002; 112:31–49.
- Wieser, G. *Bodenwasserhaushalt alpiner Ökosysteme* [Ph.D. thesis]. Botany, Innsbruck University; 1983.
- Wieser, G.; Tausz, M. *Plant Ecophysiology*. Vol. 5. Springer; Dordrecht, The Netherlands: 2007. *Trees at their upper limit. Treelife limitation at the alpine timberline*; p. 232
- Wieser, G.; Körner, Ch.; Cernusca, A. Der Wasserhaushalt von Graslandökosystemen in den österreichischen Alpen. In: Hegg, O., editor. *Verhandlungen Gesellschaft für Ökologie*. Bern: 1984. p. 89–99. Bd. XII/1982
- Wieser, G.; Neuwinger, I.; Cernusca, A. Der Bodenwasserhaushalt alpiner Graslandgesellschaften zwischen 1600 und 2600 m Höhe. In: Cernusca, A., editor. *Struktur und Funktion von Graslandökosystemen im Nationalpark Hohe Tauern*. Universitätsverlag Wagner; Innsbruck: 1989. p. 95–118. Veröffentlichungen des Österreichischen MaB-Hochgebirgsprogramms Hohe Tauern, Band 13
- Wohlfahrt G. Modelling fluxes and scalar concentrations of CO<sub>2</sub>, H<sub>2</sub>O and sensible heat within and above a mountain meadow canopy: a comparison of three Lagrangian models and three parameterisation options for the Lagrangian time scale. *Boundary-Layer Meteorology*. 2004; 113:43–80.
- Wohlfahrt G, Cernusca A. Momentum transfer by a mountain meadow canopy: a simulation analysis based on Massman's (1997) model. *Boundary-Layer Meteorology*. 2002; 103:391–407.

Wohlfahrt G, Sapinsky S, Tappeiner U, Cernusca A. Estimation of plant area index of grasslands from measurements of canopy radiation profiles. *Agricultural and Forest Meteorology*. 2001; 109:1–12.







**FIGURE 1.** Map showing the location of the study sites.  
Refer to Table 1 for station numbers.



**FIGURE 2.** Inter-annual variability of cumulative evapotranspiration (left) and daily evapotranspiration (right) of a low-elevation mountain grassland (study site Neustift, 970 m a.s.l.).




**FIGURE 3.** Annual evapotranspiration normalised with precipitation against precipitation for a low-elevation mountain grassland (study site Neustift, 970 m a.s.l.).




**FIGURE 4.** Daily evapotranspiration of a low-elevation mountain grassland (study site Neustift, 970 m a.s.l.) as a function of incident daily photosynthetically active radiation (left) and the air vapour pressure deficit (right).

Solid lines show best fits to data using a linear ( $r^2 = 0.74$ ,  $p = 0.00$ , left) and a logarithmic ( $r^2 = 0.61$ ,  $p = 0.00$ , right) model, respectively.



**FIGURE 5.** Total annual evapotranspiration ( $r^2 = 0.86$ ,  $p = 0.00$ , solid line, left) and snow sublimation ( $r^2 = 0.62$ ,  $p = 0.00$ , dotted line, left) and daily mean evapotranspiration (right) of grassland ecosystems located on the southern outer rim of the Austrian Alps (●), the inner Alpine region (○), and the Northern limestone Alps (□) between 580 and 2550 m a.s.l. (after Wieser 1983 and Wieser et al. 1984).



**FIGURE 6.** Annual evapotranspiration normalised with precipitation (left) and precipitation (right) with respect to elevation of grassland ecosystems located on the southern outer rim of the Austrian Alps (●), the inner Alpine region (○), and the Northern limestone Alps (□) between 580 and 2550 m a.s.l.

Solid lines show best fits to data using linear models (left: all data pooled;  $r^2 = 0.83$ ,  $p = 0.00$ ; right: southern outer rim of the Austrian Alps:  $r^2 = 0.97$ ,  $p = 0.00$ , Northern limestone Alps:  $r^2 = 0.98$ ,  $p = 0.04$ ).



TABLE 1

## General overview of the study sites

Location and Station Number	Station Name	Latitude-Longitude	Elevation [m]	Exposition	Inclination [°]	Vegetation period [weeks]	Observation period	Experiment	References.
<b>Southern outer rim</b>									
1	Sagritz	46°58'N, 12°55'E	1085	SSW	10	28-30	1978-1979	L	1
2	Heiligenblut	47°03'N, 12°51'E	1300	N	0	27	1978 -1979	L	1
3	Seppenbauer	47°03'N, 12°52'E	1612	S	25	25-26	1978-1980	L	1
4	Guttal	47°04'N, 12°49'E	1912	S	15-20	20-22	1978-1980	L	1
5	Wallackhaus	47°04'N, 12°51'E	2300	S	15-20	14-20	1978-1980	L	1
6	Wehr	47°04'N, 12°51'E	2430	S	5	13-14	1978-1980	L	1
7	Hochtor	47°05'N, 12°51'E	2528	S	10-15	13-14	1978-1980	L	1
8	Stubner Alm	47°07'N, 13°07'E	1815	NE	19	18	1977	L	2
<b>Inner Alpine region</b>									
9	Obergurgl	46°52'N, 11°02'E	1980	WSW	10	13-19	1976-1979	L B	1, 3
10	Hohe Mut	46°51'N, 11°02'E	2550	S	5	9-13	1976-1978	L B	1, 3
11	Haggen	47°13'N, 11°06'E	1805	S	35	22-25	1979-1981	L	4
12	Neustift	47°06'N, 11°19'E	970	-	0	33-40	2001-2006	EC	5
13	Patscherkofel	47°12'N, 11°27'E	2175	SW	5	18	1976	L	6
<b>Northern limestone Alps</b>									
14	Innsbruck	47°16'N, 11°21'E	580	-	0	44	1979-1980	L	1
15	Seegrube	47°18'N, 11°23'E	1960	S	5-10	18	1980	L	1
16	Hafelekar	47°19'N, 11°23'E	2330	S	15-20	18	1980	L	1

L = Lysimeter; B = Bowen-ratio; EC = eddy covariance

<sup>1</sup> Ref: Wieser (1983);<sup>2</sup> Körner et al. (1978);<sup>3</sup> Staudinger and Rott (1981);<sup>4</sup> Kronfuss (1997);<sup>5</sup> Wohlfahrt (2004);<sup>6</sup> Guggenberger (1980)