

Published in final edited form as:


J Org Chem. 2013 November 15; 78(22): . doi:10.1021/jo402010b.

Batch and Flow Photochemical Benzannulations Based on the Reaction of Ynamides and Diazo Ketones. Application to the Synthesis of Polycyclic Aromatic and Heteroaromatic Compounds

Thomas P. Willumstad, Olesya Haze, Xiao Yin Mak, Tin Yiu Lam, Yu-Pu Wang, and Rick L. Danheiser*

Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139

Abstract



Highly substituted polycyclic aromatic and heteroaromatic compounds are produced via a two-stage tandem benzannulation/cyclization strategy. The initial benzannulation step proceeds via a pericyclic cascade mechanism triggered by thermal or photochemical Wolff rearrangement of a diazo ketone. The photochemical process can be performed using a continuous flow reactor which facilitates carrying out reactions on a large scale and minimizes the time required for photolysis. Carbomethoxy ynamides as well as more ketenophilic bisilyl ynamines and *N*-sulfonyl and *N*-phosphoryl ynamides serve as the reaction partner in the benzannulation step. In the second stage of the strategy, RCM generates benzofused nitrogen heterocycles, and various heterocyclization processes furnish highly substituted and polycyclic indoles of types that were not available by using the previous cyclobutenone-based version of the tandem strategy.

INTRODUCTION

The design and invention of improved methods for the synthesis of benzofused nitrogen heterocycles continues to command a great deal of attention from laboratories in synthetic organic chemistry. Driving this interest are the numerous natural products and pharmaceutical compounds whose structures incorporate these heterocyclic systems. The great majority of synthetic approaches to benzofused nitrogen heterocycles involve cyclization of ortho-substituted aniline derivatives, but the application of these strategies to the synthesis of systems bearing a high level of substitution on the benzenoid ring is often problematic due to the challenges associated with the regioselective preparation of the requisite polysubstituted anilines.

Convergent *benzannulation* strategies¹ provide the most attractive approach for the regiocontrolled construction of multiply substituted benzenoid aromatic systems. Our

laboratory has developed two complementary benzannulation strategies based on the reaction of alkynes and vinylketenes^{2,3,4} and we have applied these methods to the synthesis of a number of bioactive natural products.^{5,6} Recently we extended the cyclobutenone-based version of this benzannulation strategy to include ynamine derivatives, demonstrating that the use of certain ynamides as alkyne reaction partners suppresses side reactions that are observed in reactions with simple ynamines. When employed in conjunction with ring closing metathesis, this version of our benzannulation method provides an efficient “tandem” strategy for the synthesis of highly substituted dihydroquinolines, benzazepines, and benzazocines, including a key intermediate in a formal total synthesis of (+)-FR900482.⁷ In the first stage of the tandem strategy, benzannulations using ynamine derivatives **2** (Z = electron withdrawing group) and cyclobutenones **1** furnish anilines having general structure **3** (Scheme 1). Application of ring closing metathesis (or other cyclization methods) to **3** then leads to benzofused heterocycles of type **4** or type **5**, depending on the substitution of the aniline derivative and the cyclization method employed. Further research in our laboratory has focused on the application of various heterocyclization processes in an extension of this “tandem strategy” to the synthesis of indoles bearing a high level of substitution on the six-membered ring.⁸

While the cyclobutenone-based benzannulation depicted in Scheme 1 is effective for the synthesis of a wide range of substituted anilines, polycyclic systems of type **6** present a challenge. As outlined in Scheme 2, the synthesis of polycyclic aniline derivatives with this general structure would require benzannulation with a cyclobutenone of type **7** as the vinylketene precursor. The synthesis of fused-ring cyclobutenones of this type generally requires lengthy and inefficient routes,⁹ and some potential substrates of interest are expected to be prohibitively strained compounds.


In order to extend the benzannulation to include the synthesis of polycyclic systems, we previously developed a “second generation” variant of our strategy in which α -diazo ketones are employed as vinylketene precursors in place of cyclobutenones.^{2b} Diazo ketones of type **8** are readily available (vide infra) by a number of straightforward routes and provide efficient access to the requisite ketenes via Wolff rearrangement.¹⁰ In this paper we report the application of this “second generation” version of our strategy for the first time to reactions in which ynamine derivatives function as the alkyne benzannulation partners. Based on our previous studies, the proposed benzannulation was expected to proceed via the “pericyclic cascade” mechanism outlined in Scheme 3. Irradiation (or thermolysis) of diazo ketones of type **8** would generate the vinylketene **9**, which would be intercepted by the ynamide reaction partner **2** in a regioselective [2 + 2] cycloaddition to furnish vinylcyclobutenone **10**. Reversible electrocyclic cleavage of **10** under the influence of heat or light would then lead to the dienylketene **11**, and rapid 6- π electrocyclic ring closure and tautomerization would finally afford the polysubstituted aniline derivative **12**. In principle, this version of the benzannulation should proceed smoothly not only with cyclic α,β -unsaturated diazo ketones of type **8**, but also with a variety of aryl and hetaryl diazo ketones. For the synthesis of benzofused nitrogen heterocycles, the polycyclic aniline derivatives generated in this fashion would be subjected to various heterocyclization processes providing access to a wide variety of heterocyclic systems that were not available via our previous cyclobutenone-based strategy.

RESULTS AND DISCUSSION

Synthesis of Benzannulation Reaction Partners

The benzannulation reaction partners employed in this investigation were easily prepared by using standard methods. For the synthesis of ynamides,^{11,12} we employed the closely related

copper-promoted *N*-alkynylation methods previously developed in our laboratory¹³ and that of Hsung.¹⁴ Both methods involve the reaction of amine derivatives with alkynyl bromides, which are themselves readily obtained from terminal alkynes or silyl alkynes by reaction with NBS in the presence of catalytic silver nitrate.¹⁵ The Hsung *N*-alkynylation protocol utilizes catalytic CuCN or CuSO₄ in conjunction with diamine ligands and requires elevated temperatures, while our complementary procedure employs 1 equiv of CuI but has the advantage of proceeding at room temperature and thus being applicable to thermally sensitive systems.¹⁶ All of the ynamides required for this study were known compounds with the exception of **15** which was prepared by our *N*-alkynylation method as outlined in eq 1.



(1)

The most popular synthetic routes to α -diazo ketones¹⁷ involve either the reaction of carboxylic acid derivatives with diazo compounds or the reaction of ketones with sulfonyl azides (“diazo transfer”). For the synthesis of the α,β -unsaturated diazo ketone benzannulation partners used in this study, we employed the detrifluoroacetylative diazo transfer protocol previously developed in our laboratory.¹⁸

Initial Photochemical Benzannulation Studies


We initially examined the reaction of the diazo ketone **16** derived from acetylcyclohexene to determine the feasibility of employing ynamides in the photo-Wolff variant of the vinylketene-based benzannulation strategy. Table 1 summarizes our results. Irradiation was carried out in quartz tubes positioned ca. 20 cm from a Hanovia 450 W medium-pressure mercury lamp in a water-cooled quartz immersion well and was continued until TLC analysis indicated complete consumption of the diazo ketone. As we have noted in some prior photochemical benzannulations, at that point the reaction mixture consists of varying amounts of the desired phenolic annulation product and the intermediate vinylcyclobutenone (i.e., **10** in Scheme 3). The build up of colored polymers on the walls of the reaction tube apparently impedes the complete conversion of vinylcyclobutenone to product even on prolonged irradiation. Consequently, the crude reaction products were generally heated to complete the conversion of vinylcyclobutenone to phenol.

The desired benzannulation product **18** was obtained in 48% yield when 1.0 equiv of diazo ketone was used (Table 1, entry 1). We hypothesized that competing processes involving the intermediate vinylketene such as dimerization were limiting the yield, so we explored employing an excess of the diazo ketone for the reaction. In fact, when 2.5 equiv of diazo ketone was used, the yield improved to 67% (entry 2), and the yield also increased dramatically when the reaction was performed using slow addition of the diazo ketone (entry 3). Finally, the desired benzannulation product was generated in 85% yield when both slow addition was employed and the amount of diazo ketone was increased from 1.5 to 2.5 equiv (entry 4).

Dichloromethane was found to be the optimal solvent for the photochemical Wolff rearrangement. In our standard protocol, after irradiation for the indicated time the solution was concentrated and the solvent replaced with toluene. Heating at reflux then resulted in rapid conversion of any remaining intermediate vinylcyclobutenone to the desired phenolic product. For convenience, the benzannulation can be carried out without the need for solvent exchange by employing 1,2-dichloroethane (bp 83 °C), but in this case heating for 24 h is required to complete the reaction (entry 5).

For most benzannulations we found that optimal yields were obtained employing a Hanovia 450 W medium-pressure mercury lamp for irradiation. However, in some cases comparable results were obtained by employing a standard Rayonet photochemical reactor equipped with a circular array of 16 254-nm low-pressure mercury lamps. In the case of the prototypical benzannulation described in Table 1, irradiation for 1 h in a Rayonet reactor using 1 equiv of diazo ketone **16** produced **18** in somewhat lower yield (40%) as compared to reaction with the standard Hanovia protocol (entry 1).

Not unexpectedly, the optimal conditions for effecting benzannulations with aryl diazo ketones differed from the protocol described above for α,β -alkenyl derivatives. In the case of α -diazoacetophenone (**19**), benzannulation proceeded in good yield using only 1.1 equiv of the diazo ketone and without the need for slow addition (eq 2). Wolff rearrangement of **19** generates phenylketene, which has less tendency to undergo dimerization and other unproductive side reactions as compared to the more reactive vinylketene derived from **16**. Consequently, the desired [2 + 2] cycloaddition with ynamide competes more efficiently and the desired naphthylamine **20** is obtained in good yield without the need for slow addition to minimize ketene concentration.




(2)

Reactivity of Ynamine Derivatives in Ketene [2 + 2] Cycloadditions and Benzannulation

The competitive dimerization of vinylketenes presents a particular problem in the “second-generation” photo-Wolff version of the benzannulation due to the fact that the ketene intermediates are generated more rapidly and in higher concentration as compared to generation via thermolysis of cyclobutenones. Although this problem can be mitigated by the use of slow addition and by employing an excess of ketene precursor, these measures are less than ideal and we therefore became interested in replacing carbomethoxy ynamides such as **17** with ynamides that would exhibit greater ketenophilicity in the [2 + 2] cycloaddition step of the pericyclic cascade.


Previously it has been shown that *N*-carbonyl substituents reduce the electron-donating ability of nitrogen atoms to a greater extent than *N*-sulfonyl groups, a consequence of the more significant π -delocalization of the nitrogen lone pair present in the carbonyl derivatives.^{19,20} To confirm our prediction that *N*-sulfonyl ynamides would therefore exhibit increased reactivity in ketene [2 + 2] cycloadditions, we carried out the competition experiment outlined in eq 3. Reaction of an equimolar mixture of ynamides **21**^{13a} and **22**^{14b}

with excess ketene in the presence of an internal standard indicated that the *N*-sulfonyl derivative indeed reacts at a rate twice that of the carbomethoxy ynamide.²¹




(3)

As expected, ynamide **25**^{14e} with the even less electron-withdrawing *phosphoryl* group displays an even higher level of ketenophilicity. In the competition experiment shown in eq 4, we observed that the *N*-phosphoryl ynamide undergoes [2 + 2] cycloaddition at a rate five times faster than the sulfonamide.



(4)


Finally, a third competition experiment comparing the reactivity of the carbomethoxy ynamide and the phosphoryl derivative indicated that the rate of reaction of the phosphoryl ynamide is ca. 8 times that of the carbamate (eq 5).



(5)

N,N-Bis(trialkylsilyl) ynamines such as **27** were another class of ynamine derivatives that we expected to exhibit increased ketenophilicity compared to carbomethoxy derivatives. We anticipated that these electron-rich alkynes would prove highly reactive as benzannulation partners, furnishing aniline derivatives that would also undergo deprotection under exceptionally mild conditions. Of concern to us, however, was the known sensitivity of this class of ynamines to acid and the possibility that they might be unstable in the presence of the phenolic products of the benzannulation.

The feasibility of employing silyl ynamines in the benzannulation was investigated using the known ynamine **27** which we prepared by electrophilic amination of phenylacetylene as described by Würthwein.²² A stratagem for avoiding the potential destruction of the ynamine by phenolic product was devised based on our prior observation that the electrocyclic ring opening of vinylcyclobutenones does not take place when irradiation is conducted through a uranium filter. Uranium filters block wavelengths below 340 nm, allowing photo-Wolff rearrangement to take place but halting the pericyclic cascade at the stage of intermediate **10** (Scheme 3). Heating can then be employed to complete the benzannulation. In the event, application of this protocol to the reaction of the bis(silyl) ynamine **27** with 1.0 equiv of diazo ketone **19** afforded the desired aminonaphthol **28** in 61% overall yield (eq 6).



(6)

Unfortunately, attempts to extend the silyl ynamine benzannulation to include heteroaryl or α,β-alkenyl diazo ketones have not been fruitful, with reactions proceeding in low (<30%) and variable yield. In view of the limited scope of this variant of the benzannulation, the use of silyl ynamines have not been investigated further.²³


Scope of Photochemical Benzannulations Performed in Batch and Continuous Flow

In planning our systematic investigation of the photo-Wolff based ynamide benzannulation, we decided to examine reactions in which irradiation was performed both in batch and under conditions of continuous flow. The advantages of continuous-flow photochemical reactors are now well established.^{24,25} Due to their higher surface-to-volume ratio, flow reactors allow for more uniform and efficient irradiation as compared to conventional batch reactors where light does not penetrate a significant distance into the solution. Conducting photochemical reactions in flow also minimizes any secondary photo processes leading to the decomposition of the desired products. Finally, photochemical reactions conducted in continuous flow are conveniently scaled up, a significant problem and longstanding constraint when conducting photochemistry using standard batch reactors.

In the case of the “second generation” photo-Wolff based benzannulation, we anticipated that the use of a continuous-flow photochemical reactor would increase the efficiency of the reaction, allow for shorter photolysis times, facilitate carrying out large scale reactions, and would lead to an increase in the yield of benzannulations by minimizing side reactions involving photoreactive aniline products. We based the design of our single pass continuous flow reactor on the very practical and easily assembled system described by Booker-Milburn and coworkers.²⁶ UV-transparent fluorinated ethylene propylene (FEP) tubing (0.76 mm i.d., 1.59 mm o.d.) was wound around a standard water-jacketed quartz immersion well into which was placed a 450 W medium pressure mercury lamp. In most cases irradiation was conducted through a Pyrex filter as this served to reduce polymer buildup on the walls of the tubing. A syringe pump was employed to drive the reaction solution (ca. 0.25 M in ynamide and diazo ketone) through the tubing at a flow rate of 0.057-0.32 mL/min (vide infra). The residence time in the irradiated section of tubing ranged from 21 to 33 min and the throughput with this reactor was typically 0.9 mmol/h.²⁷

Table 2 summarizes the results of our investigation of the photochemical benzannulation reaction performed both as a batch process and using the continuous-flow reactor. The photochemical benzannulation of α,β -alkenyl diazo ketone **16** was examined first. As discussed previously, as a batch process the reaction of **16** with *N*-carbomethoxy ynamide **17** is best performed using slow addition of excess diazo ketone to a solution of the ynamide in order to maintain the vinylketene intermediate in low concentration and to thus minimize its dimerization. This slow addition protocol is not easily applied in a continuous flow photolysis, and so for benzannulations using α,β -alkenyl diazo ketones we have found that the more ketenophilic *N*-phosphoryl ynamides such as **25** are the preferred reaction partners. Using **25**, benzannulation with only 1.1 equiv of diazo ketone **16** provides the desired product (**29**) in 66% yield when carried out as a batch process without the need for slow addition. Irradiation of the same benzannulation partners under conditions of continuous flow afforded **29** in comparable yield (entry 2). Importantly, this result established for the first time the feasibility of conducting the photochemical benzannulation under conditions of continuous flow. As discussed earlier, conducting photochemical benzannulations under flow enables the reaction to be performed on multigram scale much more easily than would be the case by employing conventional batch equipment.

It is worth noting with regard to the synthetic utility of *N*-phosphoryl benzannulation products such as **29** that cleavage of the protective group can be achieved either under acidic conditions or by reaction with hydride reagents.²⁸ Deprotection of **29**, for example, is readily accomplished upon heating with Red-Al in toluene and furnishes **37** in 82% yield (eq 7).



(7)


As shown in Table 2, benzannulations involving aryl and hetaryl diazo ketones proceed smoothly with all three classes of ynamides when conducted using either a batch process or in continuous flow. Particularly notable are the syntheses of indoles **31** and **33** (entries 4 and 5), demonstrating the ability of this version of the benzannulation to provide access to indoles bearing multiple substituents on the six-membered ring. In the case of benzannulation with diazo ketone **32**, purification was carried out after converting the air-sensitive phenolic product to the corresponding triflate derivative. This measure facilitated purification and in principle the triflate should serve as a useful handle for the further synthetic elaboration of the product via metal-catalyzed coupling reactions (*vide infra*).

Benzannulation using the *N*-sulfonyl ynamide **22** worked well, but this benzannulation requires irradiation through a uranium filter (>340 nm) in order to prevent photochemical decomposition of the sulfonyl ynamide which most likely takes place via a photo-Fries type process.²⁹ In the case of the batch process, irradiation at this longer wavelength necessitated an increase in the photolysis time up to 50 h from the usual 3 to 8 h employed in reactions of other ynamides. The alternative continuous flow protocol proved to be highly advantageous in this case, as complete conversion could be achieved simply by increasing the residence time from the usual 21 min to 33 min.

Feasibility of Triggering Benzannulations via Thermal Wolff Rearrangement


The next goal in our investigation of the scope of the diazo ketone version of the benzannulation was to determine the feasibility of effecting the initial Wolff rearrangement step under thermal rather than photochemical conditions.¹⁰ If successful, this might allow us to extend this version of the benzannulation to include photochemically unstable ynamides and could also lead to improved yields in the case of benzannulations that generate photochemically unstable products.

Initial experiments using cyclohexenyl diazo ketone **16** were not encouraging. Once again, side reactions involving the intermediate vinylketene competed with the desired [2 + 2] cycloaddition with ynamide, and useful yields were only obtained when the reaction was performed by slow addition of a 2-fold excess of diazo ketone (eq 8).



(8)

Fortunately, both slow addition and the use of excess diazo ketone were not necessary when the more ketenophilic *N*-phosphoryl ynamide **25** was employed in the reaction. As shown in eq 9, heating a solution of diazo ketone **16** and **25** at reflux in toluene affords benzannulation product **29** in essentially the same yield as obtained in the photochemical process (Table 2, entry 2). The application of this thermal Wolff triggered benzannulation to reactions involving photochemically unstable ynamides and aniline products is described in the next section.



(9)

Tandem Ynamide Benzannulation/RCM Strategies

The successful extension of our “second generation” strategy to include ynamides as the alkyne benzannulation partners provides access to polycyclic anilines that were not available via the cyclobutenone-based version of the method. In combination with ring closing metathesis, this new variant of the benzannulation provides an attractive route to a variety of polycyclic benzofused nitrogen heterocyclic systems. For example, as illustrated in Scheme 4, benzannulation using ynamide **15** and α -diazoacetophenone (**19**) leads to the formation of the highly substituted naphthol derivative **38** in good yield. Exposure of **38** to the Grubbs II catalyst **39**³⁰ then affords naphthazocine derivative **40**.


The advantages associated with using the thermal Wolff rearrangement to initiate the benzannulation cascade are illustrated in the tandem benzannulation/RCM sequence shown in Scheme 5. Prior experiments had taught us that *C*-allyl ynamides such as **41** do not participate in photochemical benzannulations in good yield due to the facile cyclization of the *o*-allylphenol products that takes place on irradiation.³¹ This destructive side reaction is avoided when the benzannulation is effected thermally, and ring closing metathesis then provides the benzazepine **44** in excellent yield.³²

Synthesis of Indoles via Tandem Ynamide Benzannulation/Cyclization Strategies

Recently we demonstrated that the combination of the cyclobutenone-based benzannulation with various heterocyclization and annulation processes provides an attractive route to indoles bearing multiple substituents on the benzenoid ring.⁸ A limitation of this tandem strategy, however, is that it cannot provide access to indoles with certain substitution patterns and to indoles fused to additional ring systems. As demonstrated below, the “second generation” variant of the benzannulation strategy addresses these limitations and extends the scope of this chemistry to include the construction of a number of highly substituted polycyclic indoles.

Table 3 illustrates one version of this tandem strategy. Previously we have shown that benzannulations with ynamide **45**⁸ afford products that undergo cyclization to indoles upon exposure in one flask to TBAF and then acid. As shown in Table 3, irradiation of **45** with a variety of α,β -unsaturated diazo ketones proceeds smoothly to afford benzannulation products that in most cases would not have been available via the cyclobutenone-based

method. Treatment in one pot with TBAF and then HCl provides the desired indoles in good to excellent yield. Due to its limited stability, the benzannulation product **54** (entry 4) was best transformed to the corresponding triflate **55** prior to indole formation. In this case, conversion to the indole was accomplished in one step by brief exposure to excess TfOH at room temperature.³³ Particularly notable is the efficient formation of the indole **52**, incorporating an interesting cyclobutarene system.³⁴ Due to their sensitivity to air oxidation, the indole products in entries 2 and 3 were best isolated as their acetate or triflate derivatives. The utility of the triflates as synthetic intermediates for the further elaboration of the indole products is illustrated by the Suzuki-Miyaura coupling shown in eq 10 which we found is best accomplished by using Buchwald's SPhos ligand.³⁵




(10)

N-Allyl ynamides serve as another useful building block in our tandem benzannulation/heterocyclization strategy for the synthesis of highly substituted indoles.⁸ Oxidative cleavage of the allyl double bond following benzannulation affords an aldehyde that undergoes cyclization to generate the desired indole upon treatment with K₂CO₃ or DBU followed by brief exposure to HCl.

Table 4 illustrates the application of this cyclization protocol in conjunction with the "second generation" benzannulation to provide access to very highly substituted indoles whose synthesis would be difficult to achieve by alternative methods. In these experiments to establish the scope of the method the benzannulations were carried out as batch reactions, but application of the continuous flow reactor with *N*-phosphoryl ynamides should allow the benzannulation to be accomplished on multigram scale since the outcome of flow processes is expected to be independent of scale.

The availability of cyclobutarenes such as **66** via this methodology is of particular interest. Thermolysis of cyclobutarenes provides access to *o*-quinodimethanes, whose utility as highly reactive dienes in Diels-Alder cycloadditions is well documented.³⁶ Eq 11 and Scheme 6 illustrate the application of the *o*-quinodimethanes derived from **66** in [4 + 2] cycloadditions with *N*-phenylmaleimide and ethyl propiolate, respectively. Not unexpectedly, little regioselectivity was observed in the Diels-Alder reaction of the unsymmetrical dienophile.



(11)

CONCLUSION

A wide range of benzofused nitrogen heterocyclic compounds are available via the application of our “second generation” benzannulation strategy employed in conjunction with various heterocyclization reactions. The triggering step for the pericyclic cascade involved in the benzannulation is the Wolff rearrangement of an α,β -unsaturated diazo ketone that can be effected photochemically or by heating in refluxing toluene. The photochemical process can be performed by irradiation as a batch process or by using a simple photochemical continuous flow reactor. The benzannulation has also been extended to include several new classes of ynamides that exhibit enhanced reactivity in the key [2 + 2] cycloaddition step of the pericyclic cascade. Ring closing metathesis and several types of nucleophilic cyclization processes are employed to convert the benzannulation products to benzazepines, benzacines, and in particular, to highly substituted indoles and polycyclic indoles that were not available by using previous versions of our tandem benzannulation/cyclization strategy.

EXPERIMENTAL SECTION

General Procedures

All reactions were performed in flame-dried or oven-dried glassware under a positive pressure of argon. Reaction mixtures were stirred magnetically unless otherwise indicated. Air- and moisture-sensitive liquids and solutions were transferred by syringe or cannula and introduced into reaction vessels through rubber septa. Reaction product solutions and chromatography fractions were concentrated by rotary evaporation at ca. 20 mmHg and then at ca. 0.1 mmHg (vacuum pump) unless otherwise indicated. Thin layer chromatography was performed on precoated glass-backed silica gel 60 F-254 0.25 mm plates. Column chromatography was performed on silica gel 60 (230-400 mesh) or basic alumina (80-325 mesh). Slow additions were performed with a syringe pump.

Materials

Commercial grade reagents and solvents were used without further purification except as indicated below. Dichloromethane, diethyl ether, and tetrahydrofuran were purified by pressure filtration through activated alumina. Toluene was purified by pressure filtration through activated alumina and Cu(II) oxide. Acetonitrile, 1,2-dichloroethane, 1,1,1,3,3,3-hexamethyldisilazane, pyridine, triethylamine, 2,2,2-trifluoroethyl trifluoroacetate (TFETFA), and *p*-xylene were distilled under argon from calcium hydride. Triflic anhydride was distilled under argon from P₂O₅. *N*-Bromosuccinimide was recrystallized from boiling


water. 4-DMAP was recrystallized from boiling toluene. Copper (I) iodide was extracted with THF for 24 h in a Soxhlet extractor and then dried under vacuum (0.1 mmHg). Sodium periodate supported on silica gel was prepared according to the procedure of Zhong and Shing.³⁷ Methanesulfonyl azide was prepared by reaction of methanesulfonyl chloride with sodium azide.^{18a} *n*-Butyllithium was titrated using menthol in THF with 1,10-phenanthroline as an indicator.³⁸ Deactivated alumina was prepared by first generating Brockman I grade by heating 150 g of alumina at 250 °C under vacuum (0.050 mmHg) for 12-18 h and then cooling under Ar. The alumina was then transferred to a 500-mL Erlenmeyer with a screw cap and the requisite amount of water was added to obtain the desired grade (Brockman grade II: 3%; grade III: 6%; grade IV: 10%; grade V: 15% w/w water). The alumina was then vigorously shaken for 10-15 min until a free flowing powder was obtained. The powder was allowed to stand at rt for 24 h with occasional mixing prior to use.

Instrumentation

Melting points are uncorrected. ¹H NMR chemical shifts are expressed in parts per million (δ) downfield from tetramethylsilane (with the CHCl₃ peak at 7.27 ppm used as a standard). ¹³C NMR chemical shifts are expressed in parts per million (δ) downfield from tetramethylsilane (with the central peak of CHCl₃ at 77.23 ppm used as a standard). ³¹P NMR chemical shifts are expressed in parts per million (δ) downfield from an external 85% H₃PO₄ standard (0.0 ppm). High-resolution mass spectrometry (HRMS) was performed with a Fourier transform ion cyclotron resonance mass spectrometer.

Construction of the Continuous Flow Reactor

The continuous flow reactor employed in our experiments was constructed based on the design of Booker-Milburn et al.²⁶ Fluorinated ethylene propylene (FEP) tubing, 1.59 mm o.d., 0.76 mm i.d., length = 440 cm (Reactor A) or 1520 cm (Reactor B) was wrapped around a quartz immersion well in tightly packed coils leaving 90 cm of tubing free at each end. The length of tubing wrapped around the well was 260 cm in the case of Reactor A (internal volume ca. 1.2 mL) and 1340 cm in the case of Reactor B (internal volume ca. 6.1 mL). The ends of the tubing were secured to the well with Teflon tape. The top end of the tubing was fitted with a nut, ferrule, and a thread to a female Luer adapter for attachment of a syringe. The bottom end of the tubing was connected through a rubber septum to a 25-mL or 50-mL pear flask equipped with an argon inlet needle and a needle vent. The flask was wrapped in aluminium foil. The immersion well was connected to recirculating tap water via PVC tubing. A Pyrex or uranium filter and a 450 W Hanovia medium-pressure mercury lamp were placed inside the immersion well as shown in the diagram.



Procedures for Preparation of Benzannulation Reaction Partners

1-Bromo-4-(*tert*-butyldimethylsilyloxy)-hex-5-en-1-yne (14)—A 250-mL round-bottomed flask equipped with a rubber septum and argon inlet needle was charged with 4-(*tert*-butyldimethylsilyloxy)-hex-5-en-1-yne³⁹ (2.40 g, 11.4 mmol, 1.0 equiv) and 50 mL of acetone. *N*-bromosuccinimide (2.23 g, 12.6 mmol, 1.1 equiv) was added and the mixture was stirred until all of the NBS dissolved. AgNO₃ (0.194 g, 1.14 mmol, 0.1 equiv) was added and then the reaction mixture was stirred at rt for 16 h. The resulting grey slurry was diluted with 50 mL of H₂O and 80 mL of Et₂O. The aqueous layer was extracted with two 50-mL portions of Et₂O, and the combined organic layers were washed with two 50-mL

portions of satd Na₂S₂O₃ solution and 50 mL of brine, dried over MgSO₄, filtered, and concentrated to afford ca. 3.9 g of green oil. Column chromatography on 20 g of silica (elution with hexanes) afforded 2.60 g (81%) of alkynyl bromide **14** as a colorless oil: IR (thin film) 2957, 2858, 2228, 1473, 1362, 1255, and 1090 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 5.88 (ddd, *J* = 5.6, 10.4, 17.1 Hz, 1H), 5.25 (dt, *J* = 1.5, 17.1 Hz, 1H), 5.11 (dt, *J* = 1.5, 10.4 Hz, 1H), 4.24-4.29 (m, 1H), 2.38 (app qd, *J* = 6.8, 16.5 Hz, 2H), 0.91 (s, 9H), 0.10 (s, 3H), and 0.07 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 140.1, 115.0, 77.6, 72.3, 39.8, 29.7, 26.0, 18.4, -4.4, and -4.7; HRMS (ESI) (*m/z*) [M + Na]⁺ calcd for C₁₂H₂₁BrNaOSi, 311.0443, found: 311.0438.

***N*-(Methoxycarbonyl)-*N*-(2-propenyl)-hex-5-en-4-(*tert*-butyl-dimethylsilanyloxy)-1-ynylamine (**15**)**—

A 50-mL, three-necked, round-bottomed flask fitted with a rubber septum, argon inlet adapter and glass stopper was charged with allyl carbamate **13** (0.160 g, 1.39 mmol, 1.0 equiv) and 6 mL of THF. The solution was cooled to 0 °C, and a solution of KHMDS (1.5 mL, 0.91 M in THF, 1.0 equiv) was then added rapidly dropwise over ca. 4 min. The resulting pale yellow slurry was stirred at 0 °C for 15 min and 3.8 mL of pyridine was then added, followed by CuI (0.265 g, 1.39 mmol, 1.0 equiv) in one portion. The reaction mixture was allowed to warm to rt over 2 h, and a solution of the alkynyl bromide **14** (0.528 g, 1.83 mmol, 1.3 equiv) in 5 mL of THF was then added dropwise via cannula over 25 min. The reaction mixture was stirred at rt for 18 h, diluted with 25 mL of Et₂O, and washed with three 20-mL portions of a 2:1 mixture of brine and concd NH₄OH solution. The combined aqueous layers were extracted with two 15-mL portions of Et₂O and the combined organic phases were washed with 30 mL of brine, dried over MgSO₄, filtered, and concentrated to afford 0.871 g of red-brown oil. Purification by column chromatography on 35 g of silica gel (elution with 3% EtOAc-hexanes) provided 0.222 g (49%) of **15** as a pale yellow oil: IR (thin film) 3084, 2956, 2858, 2265, 1732, 1646, and 1388 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 5.81-5.94 (m, 2H), 5.2105.28 (m, 3H), 5.09 (app dt, *J* = 1.5, 10.4 Hz, 1H), 4.20-4.25 (m, 1H), 3.98-4.08 (m, 2H), 3.79 (s, 3H), 2.40-2.55 (app qd, *J* = 6.0, 16.3 Hz, 2H), 0.90 (s, 9H), 0.08 (s, 3H), and 0.06 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 156.2, 140.4, 132.0, 118.4, 114.6, 75.2, 72.8, 66.9, 54.1, 52.8, 28.5, 26.0, 18.4, -4.4, and -4.7; HRMS (ESI) (*m/z*) [M + Na]⁺ calcd for C₁₇H₂₉NNaO₃Si: 346.1809, found: 346.1807.

1-(*tert*-Butoxycarbonyl)-3-acetylpyrrole—A 50-mL, pear flask equipped with a rubber septum and argon inlet needle was charged with 3-acetylpyrrole⁴⁰ (0.510 g, 4.67 mmol, 1.0 equiv) and 9 mL of CH₂Cl₂. The resulting solution was cooled to 0 °C and a solution of Boc₂O (1.05 g, 4.79 mmol, 1.1 equiv) in 7.5 mL of CH₂Cl₂ was added via cannula over 15 min. After 10 min, a solution of 4-DMAP (0.085 g, 0.70 mmol, 0.15 equiv) and Et₃N (0.67 mL, 4.79 equiv, 1.1 equiv) in 6.5 mL of CH₂Cl₂ was added dropwise via cannula over 20 min. The reaction mixture was stirred at 0 °C for 2.5 h and then washed with two 25-mL portions of water, dried over MgSO₄, filtered, and concentrated to yield 1.07 g of a yellow oil. Column chromatography on 50 g of silica gel (elution with 7% EtOAc-hexanes) furnished 0.952 g (97%) of 1-(*tert*-butoxycarbonyl)-3-acetylpyrrole as a white solid: mp 69-71 °C; IR (neat) 2977, 1752, 1673, 1494, 1396, 1280, 1150 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.80-7.83 (m, 1H), 7.21-7.24 (m, 1H), 6.62-6.65 (m, 1H), 2.45 (s, 3H), 1.63 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ 193.7, 148.4, 128.6, 124.7, 121.5, 111.1, 85.4, 28.1, 27.4; HRMS (ESI) (*m/z*) [M + Na]⁺ calculated for C₁₁H₁₅NNaO₃: 232.0944, found: 232.0945.

1-(*tert*-Butoxycarbonyl)-3-diazoacetylpyrrole (30**)**—A 100-mL, three-necked, round-bottomed flask equipped with two rubber septa, an argon inlet adapter, and a thermocouple was charged with a solution of 1,1,1,3,3,3-hexamethyldisilazane (1.15 mL, 5.45 mmol, 1.2

equiv) in 12 mL of THF and then cooled at 0 °C while *n*-BuLi (2.64 M in hexane, 1.9 mL, 5.02 mmol, 1.1 equiv) was added dropwise over 5 min. After 10 min, the resulting solution was cooled at -78 °C while a solution of 1-(*tert*-butoxycarbonyl)-3-acetylpyrrole (0.951 g, 4.55 mmol, 1.0 equiv) in 12 mL of THF was added dropwise via cannula over 35 min (0.8 mL THF wash). After 45 min, 2,2,2-trifluoroethyl trifluoroacetate (0.73 mL, 5.45 mmol, 1.2 equiv) was added rapidly by syringe. After 10 min, the reaction mixture was diluted with 25 mL of 1 M aq HCl solution and 30 mL of Et₂O. The aqueous phase was extracted with two 15-mL portions of Et₂O, and the combined organic phases were washed with 30 mL of brine, dried over Na₂SO₄, filtered, and concentrated to give 1.62 g of an orange oil which was immediately dissolved in 12 mL of CH₃CN and transferred to a 100-mL pear flask equipped with a rubber septum and argon inlet needle. Water (0.08 mL, 4.43 mmol, 1.0 equiv) and Et₃N (0.95 mL, 6.82 mmol, 1.5 equiv) were added, and a solution of methanesulfonyl azide (0.880 g, 7.27 mmol, 1.6 equiv) in 14 mL of CH₃CN was added dropwise via cannula over 30 min (1 mL CH₃CN wash). The resulting solution was stirred at room temperature for 3 h and then diluted with 50 mL of Et₂O and washed with three 20-mL portions of 10% aq NaOH solution and 50 mL of brine, dried over Na₂SO₄, filtered, and concentrated to afford 1.199 g of a yellow oil. Column chromatography on 53 g of silica gel (gradient elution with 10-20% EtOAc-hexanes) furnished 1.004 g (94%) of diazo ketone **30** as a yellow solid: mp 58-59 °C; IR (neat) 3124, 2982, 2103, 1752, 1617, 1496, 1396, 1336, 1255, 1154 cm⁻¹; UV (CH₃CN) λ_{max}, nm (ε) 294 (20642), 245 (19339), 229 (22817); ¹H NMR (500 MHz, CDCl₃) δ 7.70-7.72 (m, 1H), 7.23-7.25 (m, 1H), 6.51-6.52 (m, 1H), 5.63 (s, 1H), 1.62 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ 181.6, 148.3, 126.9, 122.0, 121.4, 110.3, 85.4, 54.2, 28.1; HRMS (ESI) (*m/z*) [M + Na]⁺ calculated for C₁₁H₁₃N₃NaO₃: 258.0849, found: 258.0848.

Ynamide Reactivity Comparison Study

2-Hexyl-3-(*N*-benzyl-*N*-(methoxycarbonyl))-2-cyclobuten-1-one (23**)**—Ketene was generated by pyrolysis of acetone over an electrically heated metal filament using the apparatus described by Williams and Hurd.⁴¹ A 12x75 mm test tube fitted with a rubber septum was charged with ynamide **21**^{13a} (0.098 g, 0.358 mmol, 1.0 equiv) and 3.0 mL of toluene. The septum was fitted with an inlet needle connected to the ketene generator and an outlet needle connected via tubing to a column of calcium sulfate leading into a trap of water. Ketene was bubbled into the solution at rt over a period of 6 h. The reaction mixture was then concentrated to afford 0.127 g of an orange oil. Column chromatography on 12 g of silica gel (elution with 15% EtOAc-hexane) yielded 0.051 g of unreacted ynamide **21** as a colorless oil and 0.045 g (40%) of cyclobutenone **23**⁴² as a yellow oil: IR (thin film) 2956, 2929, 2857, 1737, 1611, 1453, 1393, 1364, 1233 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.37 (t, *J* = 7.0 Hz, 2H), 7.30 (t, *J* = 7.0 Hz, 1H), 7.19 (d, *J* = 7.5 Hz, 2H), 4.99 (s, 2H), 3.84 (s, 3H), 3.54 (t, *J* = 1.5 Hz, 2H), 1.97 (t, *J* = 7.5 Hz, 2H), 1.30-1.38 (m, 2H), 1.20-1.30 (m, 2H), 1.10-1.20 (m, 4H), 0.84 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 188.1, 159.6, 153.7, 136.5, 129.1, 127.9, 127.7, 125.9, 54.4, 51.2, 50.9, 31.6, 29.4, 28.5, 24.0, 22.7, 14.2; HRMS (ESI) (*m/z*) [M + H]⁺ calcd for C₁₉H₂₆NO₃, 316.1907, found 316.1911.

Ketene [2 + 2] Cycloaddition Competition Experiment: *N*-Carbomethoxy vs. *N*-Tosyl Ynamides—Ketene was generated by pyrolysis of acetone over an electrically heated metal filament using the apparatus described by Williams and Hurd. A 5-mL conical vial fitted with a rubber septum was charged with ynamide **21** (0.103 g, 0.377 mmol), ynamide **22**^{14b} (0.140 g, 0.379 mmol) and 1.5 mL of CDCl₃ containing 1,3,5-trimethoxybenzene (0.063 g, 0.375 mmol) as an internal standard. The septum was fitted with an inlet needle connected to the ketene generator and an outlet needle connected via tubing to a column of calcium sulfate leading into a trap of water. Ketene was bubbled into the solution at rt over a period of 4 h. Aliquots (ca. 0.1 mL) of the reaction mixture were

taken at intervals, diluted with CDCl_3 , and examined by ^1H NMR (500 MHz) with a relaxation time $d1 = 20$ s to ensure accurate integration and auto phasing or manual phasing to ensure a level baseline. For cyclobutenone **23** the resonance at 4.96 ppm was integrated; for cyclobutenone **24**⁴² the resonance at 5.04 ppm was integrated. For ynamide **21** the resonance at 4.61 ppm was integrated; for ynamide **22** the resonance at 4.45 ppm was integrated.

2-Hexyl-3-(*N*-benzyl-*N*-(diethylphosphoryl))-2-cyclobuten-1-one (26)—Ketene was generated by pyrolysis of acetone over an electrically heated metal filament using the apparatus described by Williams and Hurd. A 10x75 mm test tube fitted with a rubber septum was charged with ynamide **25**^{14e} (0.061 g, 0.174 mmol, 1.0 equiv) and 1.5 mL of toluene. The septum was fitted with an inlet needle connected to the ketene generator and an outlet needle connected via tubing to a column of calcium sulfate leading into a trap of water. Ketene was bubbled into the solution at rt over a period of 40 min. The reaction mixture was then concentrated to afford 0.082 g of an orange oil. Column chromatography on 7 g of silica gel (elution with 30% EtOAc-hexane) yielded 0.062 g (90%) of cyclobutenone **26** as a yellow oil: IR (thin film) 2929, 2857, 1751, 1602, 1455, 1384, 1351, 1270, 1020 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.37 (t, $J = 7.5$ Hz, 2H), 7.25-7.34 (m, 3H), 4.90 (d, $J = 11.5$ Hz, 2H), 3.90-4.20 (m, 4H), 3.37 (t, $J = 1.5$ Hz, 2H), 1.90 (t, $J = 7.5$ Hz, 2H), 1.29-1.39 (m, 8H), 1.18-1.29 (m, 2H), 1.04-1.18 (m, 4H), 0.84 (t, $J = 7.3$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 187.1, 161.6 (d, $J = 5.1$ Hz), 137.4, 129.0, 127.9, 126.3, 125.6 (d, $J = 9.8$ Hz), 64.2 (d, $J = 5.8$ Hz), 52.2 (d, $J = 5.0$ Hz), 49.7, 31.7, 29.5, 28.6, 23.9, 22.7, 16.2 (d, $J = 6.9$ Hz) 14.2; ^{31}P NMR (121 MHz, CDCl_3) δ 2.17; HRMS (ESI) (m/z) [$M + H$]⁺ calcd for $\text{C}_{21}\text{H}_{33}\text{NO}_4\text{P}$, 394.2142, found 394.2141.

Ketene [2 + 2] Cycloaddition Competition Experiment: *N*-Diethylphosphoryl vs. *N*-Tosyl Ynamides—Ketene was generated by pyrolysis of acetone over an electrically heated metal filament using the apparatus described by Williams and Hurd. A 13x100 mm test tube equipped with a rubber septum was charged with 3.5 mL CDCl_3 and then fitted with an inlet needle connected to the ketene generator and an outlet needle connected via tubing to a column of calcium sulfate leading into a trap of water. Ketene was bubbled into the solution at rt for 5 min, then a solution of ynamide **25** (0.124 g, 0.353 mmol), ynamide **22** (0.130 g, 0.352 mmol), and 1,3,5-trimethoxybenzene (0.059 g, 0.351 mmol) as an internal standard in 0.5 mL of CDCl_3 was added via syringe, and the addition of ketene continued over a period of 2 h. Aliquots (ca. 0.1 mL) of the reaction mixture were taken at intervals, diluted with CDCl_3 , and examined by ^1H NMR (500 MHz) with a relaxation time $d1 = 20$ s to ensure accurate integration and auto phasing or manual phasing to ensure a level baseline. For cyclobutenone **26** the resonance at 4.90 ppm was integrated; for cyclobutenone **24** the resonance at 5.04 ppm was integrated. For ynamide **25** the resonance at 4.40 ppm was integrated; for ynamide **22** the resonance at 4.45 ppm was integrated.

Ketene [2 + 2] Cycloaddition Competition Experiment: *N*-Carbomethoxy vs. *N*-Diethylphosphoryl Ynamides—Ketene was generated by pyrolysis of acetone over an electrically heated metal filament using the apparatus described by Williams and Hurd. A 12x75 mm test tube fitted with a rubber septum was charged with 3.5 mL CDCl_3 then fitted with an inlet needle connected to the ketene generator and an outlet needle connected via tubing to a column of calcium sulfate leading into a trap of water. Ketene was bubbled into the solution at rt for 10 min, then a solution of ynamide **21** (0.110 g, 0.404 mmol), ynamide **25** (0.142 g, 0.404 mmol), and 1,3,5-trimethoxybenzene (0.068 g, 0.404 mmol) as an internal standard in 0.5 mL of CDCl_3 was added via syringe, and the addition of ketene continued over a period of 2 h. Aliquots (ca. 0.1 mL) of the reaction mixture were taken at

intervals, diluted with CDCl_3 , and examined by ^1H NMR (500 MHz) with a relaxation time $d1 = 20$ s to ensure accurate integration and auto phasing or manual phasing to ensure a level baseline. For cyclobutenone **23** the resonance at 4.96 ppm was integrated; for cyclobutenone **26** the resonance at 4.90 ppm was integrated. For ynamide **21** the resonance at 4.61 ppm was integrated; for ynamide **25** the resonance at 4.40 ppm was integrated.

Scope of Benzannulation Reactions

3-[N-(Carbomethoxy)-N-(methyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (18) by Batch Photochemical Benzannulation—A 20-cm quartz tube (7 mm i.d., 10 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with ynamide **17**⁴² (0.106 g, 0.537 mmol, 1.0 equiv) and 1.7 mL of CH_2Cl_2 . A 10-mL Pyrex tube fitted with a rubber septum and argon inlet needle was charged with diazo ketone **16**^{18a} (0.202 g, 1.34 mmol, 2.5 equiv) and 3.3 mL of CH_2Cl_2 . Both solutions were degassed via three cycles of freeze-pump-thaw (-196 °C, 0.05 mmHg). The solution of diazo ketone was taken up in a 5-mL glass syringe wrapped in aluminum foil and fitted with a 20-gauge, 20 cm long steel needle. The upper portion (ca. 5-7 cm length) of the quartz tube was wrapped in aluminum foil. The quartz tube was positioned ca. 20 cm proximity to the Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water). The diazo ketone solution was added to the reaction tube via syringe pump during irradiation under argon at 28 °C. After the addition was completed (ca. 6 h), the Pyrex tube was rinsed with 0.5 mL CH_2Cl_2 and added with the same syringe in one portion to the reaction mixture. Irradiation was continued for 4 h. The resulting reaction mixture was concentrated to 0.320 g of orange oil and transferred to a 10-ml Pyrex tube with the aid of 3.5 mL of toluene. The solution was heated at reflux for 1.5 h and then cooled to rt and concentrated to afford 0.347 g of dark orange oil. Purification by column chromatography on 13 g of silica gel (elution with 10% EtOAc-hexanes) afforded 0.146 g (85%) of the aniline **18** as a viscous yellow oil: IR (thin film) 3408, 2960, 2929, 1688, 1455, 1261, and 1101 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 6.53 (s, minor rotamer), 6.49 (s, 1H), 4.95 (s, 1H), 3.79 (s, minor rotamer), 3.64 (s, 3H), 3.19 (s, 3H), 2.73-2.67 (m, 2H), 2.60 (t, $J = 6.3$ Hz, 2H), 2.49-2.45 (m, 2H), 1.88-1.83 (m, 2H), 1.79-1.72 (m, 2H), 1.53-1.47 (m, 2H), 1.41-1.35 (m, 2H), 1.33-1.27 (m, 4H), and 0.89 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) 156.9, 152.4, 139.6, 136.6, 123.4, 122.4, 120.1, 53.1, 38.9, 31.9, 30.1, 29.4, 29.3, 25.9, 23.1, 22.8, 22.7, and 14.3; HRMS (ESI) (m/z) $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{30}\text{NO}_3$; 320.2220, found 320.2229.

3-[N-(Carbomethoxy)-N-(methyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (18) by Thermal Benzannulation—A 10-mL round-bottomed flask equipped with a reflux condenser, rubber septum, and argon inlet needle was charged with ynamide **17** (0.060 g, 0.306 mmol, 1.0 equiv) and 1.5 mL of toluene. A 10-mL pear flask fitted with a rubber septum and argon inlet needle was charged with diazo ketone **16** (0.092 g, 0.612 mmol, 2.0 equiv) and 3 mL of toluene. Both solutions were degassed with a stream of argon for 15 min. The solution of diazo ketone was taken up in a 5-mL glass syringe wrapped in aluminum foil and fitted with a 20-gauge, 20-cm long steel needle. The reaction mixture was heated at reflux and the diazo ketone solution was added through the condenser via syringe pump. After the addition was completed (ca. 8 h), the pear flask was rinsed with 0.5 mL of toluene and added with the same syringe in one portion to the reaction mixture. Heating was continued for 4 h. After cooling to rt, the reaction mixture was concentrated to afford 0.148 g of orange oil. Purification by column chromatography on 20 g of silica gel (elution with 5% EtOAc-10% benzene-85% hexanes) afforded 0.074 g of a yellow oil that was further purified by column chromatography on 16 g of silica gel (elution with 5% EtOAc-10% benzene-85% hexanes) to afford 0.055 g (56%) of phenol **18** as a

yellow oil with spectral data consistent with that reported in the procedure for the batch process given above.

3-[*N*-(Diethylphosphoryl)-*N*-(benzyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (29) – Batch Reactor

—A 20-cm quartz tube (7 mm i.d., 10 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **16** (0.073 g, 0.486 mmol, 1.1 equiv), ynamide **25**^{14c} (0.151 g, 0.430 mmol, 1.0 equiv), and 2.0 mL of CH₂Cl₂. The yellow solution was degassed for 10 min with a stream of argon. The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 6 h. The reaction mixture was transferred to a 25-mL pear flask with the aid of two 4-mL portions of CH₂Cl₂ and concentrated to afford 0.222 g of orange oil. This material was dissolved in 4.4 mL of toluene, the flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 2 h. The resulting mixture was concentrated to afford 0.234 g of orange oil. Column chromatography on 30 g of silica gel (gradient elution with 20-40% EtOAc-hexanes) furnished 0.149 g of an yellow oil which was further purified by column chromatography on 25 g of silica gel (elution with 1% MeOH-CH₂Cl₂) to afford 0.135 g (66%) of phenol **29** as a yellow oil: IR (neat) 3296 (broad), 2929, 2858, 1572, 1439, 1421, 1211, 1125, 1097, 1055, 1029, 966 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.20-7.26 (m, 3H), 7.14-7.20 (m, 2H), 6.32 (s, 1H), 4.80 (s, 1H), 4.50-4.66 (m, 1H), 4.21-4.35 (m, 1H), 3.96-4.16 (m, 4H), 2.37-2.67 (m, 6H), 1.75-1.84 (m, 2H), 1.65-1.75 (m, 2H), 1.54-1.65 (m, 1H), 1.22-1.38 (m, 12H), 1.06-1.21 (m, 1H), 0.90 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 152.7, 138.4 (d, *J* = 2.5 Hz), 138.1 (d, *J* = 3.8 Hz), 135.7 (d, *J* = 1.8 Hz), 129.9, 128.2, 127.6, 125.3 (d, *J* = 4.0 Hz), 123.5, 122.1 (d, *J* = 1.8 Hz), 62.9 (d, *J* = 5.6 Hz), 56.0 (d, *J* = 5.1 Hz), 32.0, 30.5, 29.5, 29.1, 26.4, 23.2, 23.0, 22.9, 16.4 (d, *J* = 6.9 Hz), 14.2; ³¹P NMR (121 MHz, CDCl₃) δ 6.65; HRMS (ESI) (*m/z*) [*M* + *H*]⁺ calculated for C₂₇H₄₁NO₄P, 474.2768; found 474.2780.

3-[*N*-(Diethylphosphoryl)-*N*-(benzyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (29) – Flow Reactor

—Reactor A with a Pyrex filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **16** (0.115 g, 0.766 mmol, 1.1 equiv), ynamide **25** (0.240 g, 0.683 mmol, 1.0 equiv), and 2.9 mL of DCE, and the yellow solution was degassed via three freeze-pump-thaw (−196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.057 mL/min by syringe pump. The solution of **16** and **25** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.057 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The 25-mL pear-shaped collection flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 40 h and then concentrated to afford 0.386 g of orange oil. This material was dissolved in 5 mL of CH₂Cl₂ and concentrated onto 5 g of silica gel. The free-flowing powder was deposited on a column of 55 g of silica gel and eluted with 35% EtOAc-hexanes to furnish 0.204 g of an orange oil which was further purified by column chromatography on 33 g of silica gel (gradient elution with 0-1% MeOH-CH₂Cl₂) to afford 0.176 g (53%) of phenol **29** as an orange oil with spectral data consistent with that reported in the procedure for the batch process given above.

3-[*N*-(Diethylphosphoryl)-*N*-(benzyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (29) – Thermal Benzannulation

—A 25-mL round-bottomed flask equipped with a rubber septum and argon inlet needle was charged with ynamide **25** (0.154 g, 0.438 mmol, 1.0 equiv), diazo ketone **16** (0.070 g, 0.466 mmol, 1.1

equiv), and 5.5 mL of toluene. The septum was replaced with a condenser fitted with an argon inlet adapter and the yellow reaction mixture was stirred at reflux for 7 h. The resulting orange mixture was concentrated to yield 0.220 g of orange oil. Column chromatography on 42 g of silica gel (elution with 20% EtOAc-hexanes + 5% Et₃N) provided 0.154 g of an orange oil that was further purified by column chromatography on 21 g of silica gel (elution with 20% EtOAc-hexanes) to afford 0.137 g (65%) of phenol **29** as a pale yellow oil with spectral data consistent with that reported in the procedure for the batch process given above.

3-[N-(Carbomethoxy)-N-(methyl)amino]-2-hexylnaphthalen-1-ol (20) – Batch Reactor—A 25-cm quartz tube (5 mm i.d., 7 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **19**^{18a} (0.081 g, 0.552 mmol, 1.1 equiv), ynamide **17** (0.099 g, 0.502 mmol, 1.0 equiv), and 1.7 mL of CH₂Cl₂. The yellow reaction mixture was degassed via three cycles of freeze-pump-thaw (−196 °C, 0.05 mmHg). The quartz tube was positioned ca. 20 cm proximity to the Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 2.5 h. The reaction mixture was concentrated to 0.244 g of yellow oil, which was transferred with 2 mL of toluene to a 10-mL reaction tube. The solution was heated at reflux for 2.5 h, cooled to rt, and concentrated to give 0.262 g orange oil, which was dissolved in 1 mL of CH₂Cl₂ and concentrated on 0.5 g of silica gel. The resulting free flowing powder was deposited onto a column of 12 g of silica gel and eluted with 20% EtOAc-hexanes to afford 0.102 g (64%) of **20** as an orange wax: IR (thin film) 3385, 2955, 2928, 2857, 1684, 1573, 1461, 1374, 1194, and 1164 cm^{−1}; ¹H NMR (500 MHz, CDCl₃) δ 8.12 (d, *J* = 9 Hz, 1 H), 7.95 (bs, minor rotamer), 7.74 (d, *J* = 9 Hz, 1H), 7.60 (bs, minor rotamer), 7.46 (app dt, *J* = 3.8, 7.0 Hz, 2H), 7.34 (bs, minor rotamer), 7.26 (s, 1H), 7.24 (s, minor rotamer), 6.19 (s, minor rotamer), 5.85 (s, 1H), 3.88 (s, minor rotamer), 3.64 (s, 3H), 3.30 (s, 3H), 3.28 (s, minor rotamer), 2.67 (t, *J* = 7.1 Hz, 2H), 1.61-1.55 (m, 2H), 1.43-1.39 (m, 2H), 1.33-1.27 (m, 4H), and 0.90 (t, *J* = 6.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 157.0, 150.1, 140.6, 132.8, 127.7, 126.3, 125.8, 124.0, 121.3, 121.0, 118.8, 53.1, 39.1, 31.9, 30.0, 29.5, 26.0, 22.8, and 14.3; HRMS (EI) (*m/z*) M⁺ calcd for C₁₉H₂₅NO₃: 315.1829, found 315.1823.

3-[N-(Carbomethoxy)-N-(methyl)amino]-2-hexylnaphthalen-1-ol (20) – Flow Reactor—Reactor A with a Pyrex filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **19** (0.088 g, 0.602 mmol, 1.1 equiv), ynamide **17** (0.108 g, 0.548 mmol, 1.0 equiv), and 2.2 mL of DCE and the yellow solution was degassed via three freeze-pump-thaw (−196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.057 mL/min by syringe pump. The solution of **19** and **17** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.057 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The 25-mL pear-shaped collection flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 48 h and then concentrated to afford 0.188 g of orange oil. This material was dissolved in 3 mL of CH₂Cl₂ and concentrated onto 2 g of silica gel. The free-flowing powder was deposited on a column of 25 g of silica gel and eluted with 10% EtOAc-10% benzene-80% hexanes to furnish 0.114 g (66%) of phenol **20** as a yellow oil with spectral data consistent with that reported in the procedure for the batch process given above.

3-[N-(Carbomethoxy)-N-(methyl)amino]-2-hexylnaphthalen-1-ol (20) – Large Scale Flow Reactor—Reactor B with a Pyrex filter was employed for this reaction. A 25-

mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **19** (0.184 g, 1.26 mmol, 1.1 equiv), ynamide **17** (0.223 g, 1.13 mmol, 1.0 equiv), and 4.8 mL of DCE. and the yellow solution was degassed via three freeze-pump-thaw (−196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.32 mL/min by syringe pump. The solution of **19** and **17** was transferred to a 10-mL glass syringe and pumped through the tubing at a rate of 0.32 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.6, 0.6, and then 9.0 mL). The 50-mL pear-shaped collection flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 40 h and then concentrated to afford 0.500 g of orange oil. This material was dissolved in 5 mL of CH₂Cl₂ and concentrated onto 5 g of silica gel. The free-flowing powder was deposited on a column of 50 g of silica gel and eluted with 10% EtOAc-10% benzene-80% hexanes to furnish 0.220 g (63%) of phenol **20** as a yellow oil with spectral data consistent with that reported in the procedure for the batch process given above.

3-[Bis(trimethylsilyl)amino]-2-phenylnaphthalen-1-ol (28)—A 20-cm quartz tube (7 mm i.d., 10 mm o.d.) equipped with a stir bar, rubber septum, and argon inlet needle was charged with diazo ketone **19** (0.048 g, 0.325 mmol, 1.0 equiv), ynamine **27**²² (0.65 mL, 0.5 M in benzene, 0.325 mmol, 1.0 equiv), and 0.7 mL of toluene. The yellow solution was degassed (three freeze-pump-thaw cycles at −196 °C, 0.05 mmHg) and backfilled with argon. The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp equipped with a uranium filter (quartz immersion well, cooled by recirculating tap water) and irradiated under argon with stirring at 25 °C for 45 h. The reaction mixture was transferred to a 15-mL threaded Pyrex tube (20 mm o.d.) with the aid of two 0.8-mL portions of toluene. The reaction tube was sealed with a threaded Teflon cap and the reaction mixture was heated at 110 °C (bath temperature) for 2.5 h. The resulting mixture was concentrated to afford 0.116 g of a dark yellow oil. Column chromatography on 20 g of silica gel (elution with 94:5:1 hexanes-EtOAc-Et₃N) furnished 0.075 g (61%) of phenol **28** as a yellow solid: mp 79–81 °C; IR (neat) 3540, 3054, 2954, 1630, 1590, 1494, 1435, 1385, 1251, 1127, 1075, 942, 876, 839, and 822 cm^{−1}; ¹H NMR (500 MHz, CDCl₃) δ 8.18 (app d, *J* = 8.4 Hz, 1H), 7.70 (app d, *J* = 8.2 Hz, 1H), 7.52–7.56 (m, 2H), 7.38–7.49 (m, 5H), 7.10 (s, 1H), 5.47 (s, 1H), and −0.06 (s, 18H); ¹³C NMR (125 MHz, CDCl₃) δ 149.1, 145.9, 135.2, 134.0, 132.1, 129.5, 128.4, 126.8, 126.7, 124.2, 122.9, 122.6, 121.8, 119.2, and 2.3; HRMS (ESI) (*m/z*) [M + H]⁺ calculated for C₂₂H₃₀NOSi₂, 380.1860; found 380.1889.

1-(tert-Butoxycarbonyl)-6-hexyl-7-hydroxy-5-[N-(methoxycarbonyl)-N-(methyl)amino]indole (31) – Batch Reactor—A 20-cm quartz tube (7 mm i.d., 10 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **30** (0.135 g, 0.574 mmol, 1.1 equiv), ynamide **17** (0.103 g, 0.522 mmol, 1.0 equiv), and 2.2 mL of CH₂Cl₂. The solution was degassed for 10 min with a stream of argon. The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 8 h. The reaction mixture was transferred to a 25-mL pear flask with the aid of two 4-mL portions of CH₂Cl₂ and concentrated to afford 0.222 g of an orange oil. This material was dissolved in 5.2 mL of toluene, the flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 2 h. The resulting mixture was concentrated to afford 0.234 g of an orange oil. Column chromatography on 25 g of silica gel (gradient elution with 10-20% EtOAc-hexanes) furnished 0.084 g (40%) of phenol **31** as a yellow oil: IR (neat) 3162 (broad), 2930, 2857, 1708, 1688, 1584, 1452, 1380, 1353, 1242, 1156, 1120 cm^{−1}; ¹H NMR (500 MHz, CDCl₃) δ 11.13 (s, 1H), 11.09 (s, minor rotamer), 7.36-7.42 (m, 1H), 6.88 (s, minor rotamer), 6.82 (s, 1H), 6.43-6.49 (m, 1H), 3.80 (s, minor

rotamer), 3.62 (s, 3H), 3.24 (s, 3H), 3.22 (s, minor rotamers), 2.53-2.68 (m, 2H), 1.65 (s, 9H), 1.50-1.60 (m, 2H), 1.36-1.45 (m, 2H), 1.27-1.36 (m, 4H), 0.90 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 157.0, 152.7, 143.6, 139.5, 131.1, 126.7, 124.8, 123.3, 110.7, 109.4, 86.3, 53.0, 38.9, 32.0, 30.2, 29.4, 28.2, 26.6, 22.8, 14.3; HRMS (ESI) (m/z) [$\text{M} + \text{Na}$] $^+$ calculated for $\text{C}_{22}\text{H}_{32}\text{N}_2\text{NaO}_5$, 427.2203; found 427.2208.

1-(tert-Butoxycarbonyl)-6-hexyl-7-hydroxy-5-[N-(methoxycarbonyl)-N-(methyl)amino]indole (31) – Flow Reactor—Reactor A with a Pyrex filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **30** (0.134 g, 0.570 mmol, 1.1 equiv), ynamide **17** (0.100 g, 0.507 mmol, 1.0 equiv), and 2.2 mL of DCE and the yellow solution was degassed via three freeze-pump-thaw (-196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.057 mL/min by syringe pump. The solution of **30** and **17** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.057 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The reaction mixture was then concentrated to afford 0.209 g of orange oil. This material was dissolved in 5.0 mL of toluene in the same 25-mL pear-shaped collection flask. The flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 2 h. The resulting mixture was concentrated to afford 0.189 g of a dark orange oil. Column chromatography on 25 g of silica gel (elution with 12% EtOAc-hexanes) furnished 0.075 g (37%) of phenol **31** as a yellow oil with spectral data consistent with that reported in the procedure for the batch process given above.

5-Hexyl-4-(trifluoromethylsulfonyloxy)-6-[N-(diethylphosphoryl)-N-(benzyl)amino]indole (33) – Batch Reactor—A 20-cm quartz tube (7 mm i.d., 10 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **32**^{2b} (0.137 g, 0.582 mmol, 1.1 equiv), ynamide **17** (0.103 g, 0.522 mmol, 1.0 equiv), and 2.2 mL of CH_2Cl_2 . The yellow reaction mixture was degassed via three freeze-pump-thaw (-196 °C, 0.05 mmHg) cycles. The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 5.5 h. The reaction mixture was transferred to a 25-mL pear flask with the aid of two 4-mL portions of CH_2Cl_2 and concentrated to afford 0.249 g of a dark orange oil. This material was dissolved in 5.2 mL of toluene, the flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 2 h. The resulting mixture was concentrated to afford 0.262 g of a dark orange oil. Column chromatography on 30 g of silica gel (elution with 15% EtOAc-hexanes) furnished 0.152 g of crude benzannulation product as an orange oil.

A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with the crude benzannulation product (0.152 g, 1.0 equiv), 4-DMAP (0.092 g, 0.753 mmol, 2.0 equiv), and 2.7 mL of CH_2Cl_2 . The orange solution was cooled to 0 °C and triflic anhydride (0.07 mL, 0.416 mmol, 1.1 equiv) was added dropwise by syringe over ca. 2 min. The reaction mixture was allowed to warm to rt, stirred for 2 h, and then diluted with 20 mL of CH_2Cl_2 and washed with two 10-mL portions of 1 M HCl solution. The combined aqueous layers were extracted with 10 mL of CH_2Cl_2 , and the combined organic layers were washed with 20 mL of satd aq NaHCO_3 solution and 20 mL of brine, dried over MgSO_4 , filtered, and concentrated to afford 0.187 g of orange oil. Column chromatography on 28 g of silica gel (elution with 10% EtOAc-hexanes) afforded 0.160 g (57% over two steps from the ynamide) of triflate **33** as an orange oil: IR (neat) 2958, 2932, 2860, 1744, 1716, 1529, 1449, 1409, 1373, 1347, 1304, 1249, 1215, 1154 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 8.12 (minor

rotamer), 8.06 (s, 1H), 7.57-7.67 (m, 1H), 6.65-6.72 (m, 1H), 3.81 (minor rotamer), 3.63 (s, 3H), 3.29 (s, 3H), 2.61-2.76 (m, 2H), 1.66 (s, 9H), 1.46-1.57 (m, 2H), 1.32-1.40 (m, 2H), 1.25-1.32 (m, 4H), 0.89 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 156.7, 149.2, 139.7, 139.5, 135.0, 129.0, 128.1, 124.0, 118.9 (q, $J = 318$ Hz), 115.6, 103.9, 85.0, 53.2, 39.1, 31.8, 29.8, 29.8, 28.3, 26.6, 22.7, 14.2; HRMS (ESI) (m/z) [$M + \text{Na}$] $^+$ calculated for $\text{C}_{23}\text{H}_{31}\text{F}_3\text{N}_2\text{NaO}_7\text{S}$, 559.1696; found 559.1703.

5-Hexyl-4-(trifluoromethylsulfonyloxy)-6-[*N*-(diethylphosphoryl)-*N*-(benzyl)amino]indole (33) – Flow Reactor

—Reactor A with a Pyrex filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **32** (0.570 g, 0.134 mmol, 1.1 equiv), ynamide **17** (0.100 g, 0.507 mmol, 1.0 equiv), and 2.2 mL of DCE and the yellow solution was degassed via three freeze-pump-thaw (-196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.057 mL/min by syringe pump. The solution of **32** and **17** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.057 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The reaction mixture was concentrated to afford 0.202 g of orange oil. This material was dissolved in 5.0 mL of toluene in the same 25-mL pear-shaped collection flask. The flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 2 h. The resulting mixture was concentrated to afford 0.232 g of a dark orange oil. Column chromatography on 25 g of silica gel (elution with 12% EtOAc-hexanes) furnished 0.113 g of ca. 85-90% pure benzannulation product as an orange oil.

A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with the impure benzannulation product (0.113 g, 1.0 equiv), 4-DMAP (0.070 g, 0.573 mmol, 2.0 equiv), and 3.0 mL of CH_2Cl_2 . The orange solution was cooled to 0 °C and triflic anhydride (0.06 mL, 0.357 mmol, 1.3 equiv) was added dropwise by syringe over ca. 2 min. The reaction mixture was allowed to warm to rt, stirred for 2 h, and then diluted with 20 mL of CH_2Cl_2 and washed with two 10-mL portions of 1 M HCl solution. The combined aqueous layers were extracted with 10 mL of CH_2Cl_2 , and the combined organic layers were washed with 20 mL of satd aq NaHCO_3 solution and 20 mL of brine, dried over MgSO_4 , filtered, and concentrated to afford 0.151 g of orange oil. Column chromatography on 18 g of silica gel (elution with 10% EtOAc-hexanes) afforded 0.127 g (47% over two steps from the ynamide) of triflate **33** as an orange oil with spectral data consistent with that reported in the procedure for the batch process given above.

5-Hexyl-4-(trifluoromethylsulfonyloxy)-6-[*N*-(*p*-tolylsulfonyl)-*N*-(benzyl)amino]benzo[*b*]thiophene (35) – Flow Reactor

—Reactor A with a uranium filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **34** (0.062 g, 0.407 mmol, 1.1 equiv), ynamide **22** (0.134 g, 0.363 mmol, 1.0 equiv), and 1.6 mL of DCE and the yellow solution was degassed via three freeze-pump-thaw (-196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.036 mL/min by syringe pump. The solution of **34** and **22** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.036 mL/min. Once the addition was complete, the tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The 25-mL pear-shaped collection flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 42 h and then concentrated to afford 0.195 g of

red oil. Column chromatography on 20 g of silica gel (elution with 10% EtOAc-hexanes) furnished 0.102 g of ca. 85-90% pure benzannulation product as an orange oil.

A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with the impure benzannulation product (0.102 g, 1.0 equiv), 4-DMAP (0.050 g, 0.409 mmol, 2.0 equiv), and 1.6 mL of CH₂Cl₂. The orange solution was cooled to 0 °C and triflic anhydride (0.04 mL, 0.238 mmol, 1.2 equiv) was added dropwise by syringe over ca. 2 min. The reaction mixture was allowed to warm to rt, stirred for 4 h, and then diluted with 10 mL of CH₂Cl₂ and washed with two 5-mL portions of 1 M HCl solution. The combined aqueous layers were extracted with 5 mL of CH₂Cl₂, and the combined organic layers were washed with 15 mL of satd aq NaHCO₃ solution and 15 mL of brine, dried over MgSO₄, filtered, and concentrated to afford 0.125 g of a red oil. Column chromatography on 13 g of silica gel (elution with 7% EtOAc-hexanes) afforded 0.106 g (47% over two steps from the ynamide) of triflate **35** as an off-white solid with spectral data consistent with that reported in the procedure for the batch process given above.

5-Hexyl-4-hydroxy-6-[N-(diethylphosphoryl)-N-(benzyl)amino]benzo[b]thiophene (36) – Batch Reactor

—A 20-cm quartz tube (5 mm i.d., 7 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **34** (0.049 g, 0.322 mmol, 1.1 equiv), ynamide **25** (0.099 g, 0.282 mmol, 1.0 equiv), and 1.3 mL of CH₂Cl₂. The yellow reaction mixture was degassed via three freeze-pump-thaw (−196 °C, 0.05 mmHg) cycles. The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 5.5 h. The reaction mixture was transferred to a 25-mL pear flask with the aid of two 4-mL portions of CH₂Cl₂ and concentrated to afford 0.143 g of an orange oil. This material was dissolved in 3 mL of toluene, the flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 1.5 h. The resulting mixture was concentrated to afford 0.159 g of a dark orange oil. Column chromatography on 15 g of silica gel (elution with 30% EtOAc-hexanes) furnished 0.115 g of an orange oil which was further purified by column chromatography on 15 g of silica gel (elution with 0.5% MeOH-CH₂Cl₂) to afford 0.105 g (78%) of phenol **36** as an orange oil: IR (neat) 3225 (broad), 2924, 2856, 1605, 1552, 1495, 1424, 1392, 1330, 1215, 1126, 1051, 970 cm^{−1}; ¹H NMR (500 MHz, CDCl₃) δ 7.32 (dd, *J* = 0.8 Hz, *J* = 5.5 Hz, 1H) 7.18-7.24 (m, 3H), 7.10-7.14 (m, 2H), 7.09 (d, *J* = 5.5 Hz), 6.95-6.97 (m, 1H), 6.74 (s, 1H), 4.59-4.67 (m, 1H), 4.02-4.32 (m, 5H), 2.58-2.68 (m, 2H), 2.45-2.55 (m, 2H), 1.57-1.70 (m, 1H), 1.21-1.42 (m, 12H), 0.99-1.11 (m, 1H), 0.90 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 150.4, 137.9 (d, *J* = 1.4 Hz), 137.7 (d, *J* = 3.4), 137.6 (d, *J* = 2.9 Hz), 129.9, 129.1 (d, *J* = 1.1 Hz), 128.3, 127.7, 124.2, 123.4 (d, *J* = 4.0 Hz), 120.3, 116.3, 63.2 (m), 55.9 (d, *J* = 4.5 Hz), 31.9, 30.5, 29.2, 26.2, 23.0, 16.4 (m), 14.4; ³¹P NMR (121 MHz, CDCl₃) δ 6.18; HRMS (ESI) (*m/z*) [M + H]⁺ calculated for C₂₅H₃₅NO₄PS, 476.2019; found 476.2034.

5-Hexyl-4-hydroxy-6-[N-(diethylphosphoryl)-N-(benzyl)amino]benzo[b]thiophene (36) – Flow Reactor

—Reactor A with a Pyrex filter was employed for this reaction. A 25-mL pear flask equipped with a rubber septum and argon inlet needle was charged with diazo ketone **34** (0.069 g, 0.448 mmol, 1.1 equiv), ynamide **25** (0.143 g, 0.407 mmol, 1.0 equiv), and 1.7 mL of DCE and the yellow solution was degassed via three freeze-pump-thaw (−196 °C, 0.05 mmHg) cycles. The FEP tubing of the reactor was flushed with 3 mL of degassed DCE, the mercury lamp was turned on, and degassed DCE was pumped through the tubing for 5 min at a rate of 0.057 mL/min by syringe pump. The solution of **34** and **25** was transferred to a 5-mL glass syringe and pumped through the tubing at a rate of 0.057 mL/min. Once the addition was complete, the

tubing was flushed with three portions of degassed DCE (0.4, 0.4, and then 4.0 mL). The 25-mL pear-shaped collection flask was equipped with a stir bar and cold-finger condenser, and the reaction mixture was heated at reflux for 42 h and then concentrated to afford 0.199 g of red oil. This material was dissolved in 2 mL of CH₂Cl₂ and concentrated onto 2 g of silica gel. The free-flowing powder was deposited on a column of 20 g of silica gel and eluted with 30% EtOAc-hexanes to furnish 0.159 g of an orange oil which was further purified by column chromatography on 20 g of silica gel (elution with 1% MeOH-CH₂Cl₂) to afford 0.154 g (79%) of phenol **36** as an orange oil with spectral data consistent with that reported in the procedure for the batch process given above.

Reductive Cleavage of *N*-Phosphorylaniline 29. 3-[*N*-(Benzyl)amino]-2-hexyl-5,6,7,8-tetrahydronaphthalen-1-ol (37**)**—A 25-mL, pear flask equipped with a rubber septum and argon inlet needle was charged with phosphoramidate **29** (0.132 g, 2.79 mmol, 1.0 equiv) and 4 mL of toluene. Red-Al (65 wt% in toluene, 0.65 mL, 2.13 mmol, 8 equiv) was added and the resulting solution was heated at 85 °C for 2 h. The reaction mixture was quenched at room temperature by the addition of 10 mL of satd aq K₂CO₃ solution, diluted with 10 mL of ether, and the organic layer was washed with 10 mL of satd aq K₂CO₃ solution. The combined aqueous layers were extracted with three 10-mL portions of Et₂O. The combined organic layers were washed with 20 mL of brine, dried over Na₂SO₄, filtered, and concentrated to give 0.122 g of red oil. Column chromatography on 11 g of silica gel (elution with 2% EtOAc-hexanes) provided 0.077 g (84%) of amine **37** as a red solid: mp 58–60 °C; IR (neat) 3556 (broad), 3452 (broad) 2926, 2856, 1621, 1582, 154, 1438, 1352, 1317, 1198, 1143 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.36–7.45 (m, 4H), 7.31 (tt, *J* = 7.0 Hz, *J* = 1.5 Hz, 1H), 6.09 (s, 1H), 4.62 (s, 1H), 4.36 (s, 2H), 3.87 (s, 1H), 2.68 (t, *J* = 6.3 Hz, 2H), 2.52–2.60 (m, 4H), 1.83–1.89 (m, 2H), 1.73–1.79 (m, 2H), 1.52–1.60 (m, 2H), 1.38–1.47 (m, 2H), 1.29–1.38 (m, 4H), 0.91 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 151.8, 144.5, 140.1, 136.2, 128.8, 127.7, 127.3, 112.1, 110.7, 104.1, 48.9, 32.0, 30.0, 29.9, 28.8, 24.1, 23.3, 23.2, 22.9, 22.6, 14.3; HRMS (ESI) (*m/z*) [M + H]⁺ calculated for C₂₃H₃₂NO, 338.2478; found 338.2470.

Tandem Benzannulation/Ring Closing Metathesis

3-[*N*-(Carbomethoxy)-*N*-(allyl)amino]-2-[but-3-en-2-(*tert*-butyldimethylsilyloxy)yl]naphthalen-1-ol (38**)**—A 25-cm quartz tube (5 mm i.d., 7 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with diazo ketone **19** (0.075 g, 0.510 mmol, 1.1 equiv), ynamide **15** (0.150 g, 0.469 mmol, 1.0 equiv), and 1.8 mL of CH₂Cl₂. The yellow reaction mixture was degassed via three cycles of freeze-pump-thaw (–196 °C, 0.05 mmHg). The quartz tube was positioned ca. 20 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by recirculating tap water) and irradiated under argon at 25 °C for 2.5 h. The reaction mixture was concentrated to afford 0.260 g of yellow oil, which was transferred with 2 mL of toluene to a 10-mL reaction tube. The solution was heated at reflux for 2.5 h, allowed to cool to rt, and then concentrated to give 0.291 g orange oil. Purification by column chromatography on 16 g of silica gel (elution with 10% EtOAc-hexanes) afforded 0.036 g (24%) of unreacted ynamide **15** and 0.129 g (62%) of **38** as a viscous yellow oil: IR (thin film) 3238, 2955, 2859, 1716, 1636, 1597, 1574, 1448, 1386, and 1256 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.16 (s, minor rotamer), 9.01 (s, 1H), 8.28–8.30 (m, 1H), 7.71–7.74 (m, 1H), 7.45–7.47 (m, 2H), 7.20–7.23 (m, 1H), 5.84–6.08 (m, 2H), 5.11–5.30 (m, 4H), 4.47–4.54 (m, 2H), 3.81–3.95 (m, 2H), 3.66 (s, 3H), 3.60 (s, minor rotamer), 3.03–3.11 (m, 1H), 2.82–2.89 (m, 1H), 0.92 (s, 9H), 0.08 (s, minor rotamer), 0.05 (s, 3H), –0.05 (s, minor rotamer), and –0.15 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 156.5, 152.9, 139.8, 139.1, 139.0, 133.5, 133.1, 127.4, 126.6, 124.6, 125.3, 122.8, 119.5, 118.7, 117.3, 115.7, 54.4, 53.0, 35.5, 35.0, 25.9, 18.2, –4.3, and –5.0, (Peaks corresponding to the minor rotamer: δ 153.2, 133.2, 127.3,

122.9, 119.2, 118.5, 117.0, 54.4, 53.1, 26.0, 18.3, -4.2, and -4.8); HRMS (ESI) (m/z) [$M + Na$]⁺ calcd for C₂₅H₃₅NNaO₄Si: 464.2228, found: 464.2217.

Methyl 5-(*tert*-Butyldimethylsilyloxy)-7-hydroxy-5,6-dihydro-2*H*-naphtheno[*b*]azocine-1-carboxylate (40)—A 100-mL, round-bottomed flask fitted with a rubber septum and argon inlet needle was charged with a solution of the Grubbs catalyst **39** (0.012 g, 0.014 mmol, 0.05 equiv) in 20 mL of CH₂Cl₂. Naphthol **38** (0.124 g, 0.281 mmol, 1.0 equiv) in 8 mL of CH₂Cl₂ was added to the reaction flask via cannula in one portion. The rubber septum was replaced with a reflux condenser fitted with an argon inlet adapter. The pale pink reaction mixture was heated at reflux for 6 h, allowed to cool to rt, and then concentrated to afford ca. 0.134 g of yellow-brown foam. Column chromatography on 12 g of silica gel (elution with 15% EtOAc-hexane) yielded 0.088 g (76%) of azocine **40** as an off-white solid: mp 74–76 °C; IR (thin film) 3351, 2955, 2857, 1709, 1684, 1572, 1450, 1388, 1250 and 1065 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.24–8.26 (m, 1H), 7.70 (s, 1H), 7.72–7.75 (m, 1H), 7.44 (app dd, $J = 3.1, 6.2$ Hz, 2H), 7.26 (bs, 1H), 5.57 (app d, $J = 11.6$ Hz, 1H), 5.34–5.41 (m, 1H), 5.14 (app s, 1H), 4.44 (bs, 1H), 4.20 (dd, $J = 7.9, 14.4$ Hz, 1H), 3.72 (bs, 3H), 3.24 (app d, $J = 2.7$ Hz, 2H), 1.01 (s, 9H), 0.26 (s, 3H), and 0.19 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 156.3, 150.6, 140.6, 137.7, 133.1, 127.5, 127.3, 126.3, 125.3, 124.7, 122.5, 118.9, 117.7, 72.6, 53.3, 48.0, 35.9, 26.0, 18.4, -4.5, and -4.8; HRMS (EI) (m/z) [$M + Na$]⁺ calcd for C₂₃H₃₁NNaO₄Si, 436.1915; found: 436.1922.

***N*-(Methoxycarbonyl)-*N*-allyl-(2-allyl-4,5-dimethyl-3-hydroxyphenyl)amine (43)**—A 25-mL recovery flask equipped with a condenser, rubber septum and argon inlet needle was charged with ynamide **41**⁷ (0.146 g, 0.815 mmol, 1.0 equiv) and 6.0 mL of toluene. A 25-mL pear flask fitted with a rubber septum and argon inlet needle was charged with diazo ketone **42**^{18a} (0.244 g, 1.96 mmol, 2.4 equiv) and 8 mL of toluene. Both solutions were degassed with a stream of argon for 15 min. The solution of diazo ketone was taken up in a 10-mL glass syringe wrapped in aluminum foil and fitted with a 20-gauge, 20-cm long steel needle. The reaction mixture was heated at reflux and the diazo ketone solution was added through the condenser via syringe pump. After the addition was completed (ca. 13.5 h), the pear flask was rinsed with 0.5 mL of toluene and added with the same syringe in one portion to the reaction mixture. Heating was continued for 5 h. After cooling to rt, the reaction mixture was concentrated to afford 0.450 g of orange oil that was diluted with 5 mL of MeOH and 3 mL of 5 M KOH solution. After heating at 60–70 °C for 2 h, the resulting mixture was cooled to rt, quenched with 16 mL of 1 M HCl solution, and diluted with 20 mL of Et₂O. The aqueous layer was extracted with two 20-mL portions of Et₂O and the combined organic layers were washed with 40 mL of satd aq NaHCO₃ solution and 40 mL of brine, dried over MgSO₄, filtered, and concentrated to provide 0.308 g of orange oil. Purification by column chromatography on 39 g of silica gel (gradient elution with 15–20% EtOAc-hexanes) afforded 0.146 g of aniline **43** (65%) as a yellow solid: mp 70–72 °C; IR (neat) 3412, 3078, 2925, 2857, 1686, 1575, 1455, 1389, 1320, 1257, 1196 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 6.58 (s, 1 H), 5.85–6.02 (m, 2 H), 5.18–5.30 (m, 3 H), 5.07–5.15 (m, 2 H), 4.21–4.32 (m, 1 H), 3.85–4.03 (m, 1 H), 3.78 (s, minor rotamer), 3.62 (s, 3 H), 3.26–3.39 (m, 2 H), 2.24 (s, 3 H), 2.15 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 156.4, 153.5, 137.8, 136.8, 135.9, 133.4, 123.4, 122.1, 120.2, 118.4, 117.3, 54.1, 53.1, 30.9, 20.2, 12.0; HRMS (ESI) (m/z) [$M + H$]⁺ calculated for C₁₆H₂₂NO₃, 276.1594; found 276.1598.

Methyl 6-Hydroxy-7,8-dimethyl-2,5-dihydro-benzo[*b*]azepine-1-carboxylate (44)—A 100-mL, round-bottomed flask equipped with a rubber septum and argon inlet needle was charged with a solution of the Grubbs catalyst **39** (0.014 g, 0.016 mmol, 0.05 equiv) in 30 mL of CH₂Cl₂. The aniline **43** (0.094 g, 0.341 mmol, 1.0 equiv) in 6 mL of

CH₂Cl₂ was added to the reaction flask via cannula in one portion and the rubber septum was replaced with a reflux condenser fitted with an argon inlet adapter. The pale pink reaction mixture was heated at reflux for 40 min, allowed to cool to rt, and then concentrated onto 2 g of silica gel. The free-flowing powder was deposited on a column of 20 g of silica gel and eluted with 25% EtOAc-hexanes to afford 0.081 g (96%) of dihydrobenzoazepine **44** as a tan wax: IR (neat) 3399, 3024, 2955, 1685, 1615, 1579, 1456, 1395, 1333, 1262, 1193, 1088 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 6.70 (s, minor rotamer), 6.64 s, 1 H), 5.75-5.84 (m, 1 H), 5.43-5.52 (m, 1 H), 4.72-5.16 (m, 1 H), 4.69 (s, 1H), 4.22 (s, minor rotamer), 3.79 (s, minor rotamer) 3.67 (s, 3 H), 3.34-3.66 (m, 3 H), 2.26 (s, 3 H), 2.24 (s, minor rotamer), 2.15 (s, 3 H), 2.12 (s, minor rotamer); ¹³C NMR (125 MHz, CDCl₃ mixture of two rotamers) for major rotamer: δ 156.4, 150.4, 139.5, 135.8, 127.3, 126.0, 125.2, 123.8, 121.5, 53.4, 48.3, 23.3, 20.4, 12.3; additional resonances appeared for the minor rotamer at δ: 126.9, 121.4, 48.1, 23.1; HRMS (ESI) (*m/z*) [M + H]⁺ calculated for C₁₄H₁₈NO₃, 248.1281; found 248.1272.

Synthesis of Indoles via Tandem Ynamide Benzannulation/Cyclization Strategies

2-(Trimethylsilyl)ethyl Benzyl[2-(2,2-dimethoxyethyl)-3-hydroxy-4,5-dimethylphenyl]carbamate (46)—A solution of ynamide **45**⁸ (0.340 g, 0.936 mmol, 1.0 equiv) in 3.5 mL of CH₂Cl₂ was distributed evenly between two 25-cm quartz reaction tubes (13 mm i.d., 15 mm o.d.) fitted with rubber septa and argon inlet needles. A 15-mL Pyrex tube fitted with a rubber septum and argon inlet needle was charged with a solution of diazo ketone **42** (0.310 g, 2.07 mmol, 2.4 equiv) in 7 mL of CH₂Cl₂. All solutions were purged with argon for 10 min. The upper portions (ca. 10 cm) of the quartz tubes were wrapped in aluminum foil and the tubes were positioned ca. 4 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by tap water). The quartz reaction tubes and the immersion well were partially submerged in a room-temperature water bath contained in a 2-L beaker wrapped in aluminum foil. The diazo ketone solution was taken up in two 5-mL glass syringes fitted with 20 gauge, 20-cm steel needles and wrapped in aluminum foil, and the solution was added to the reaction mixture using a syringe pump over 5 h (0.5 mL CH₂Cl₂ rinse). Irradiation was continued for 1 h, and then the reaction mixtures were combined in a 25-mL round-bottomed flask, concentrated, and the residue was diluted with 5 mL of toluene. The flask was fitted with a cold-finger condenser with an argon inlet sidearm, and the solution was heated at reflux for 1 h, allowed to cool to rt, and then concentrated to give 0.625 g of an orange oil. Column chromatography on 20 g of grade II neutral alumina (gradient elution with 0–5% EtOAc–hexanes) gave 0.292 g of a mixture of unreacted ynamide **45** and phenol **46**. Mixed fractions (0.057 g) were purified by chromatography on 5 g of neutral alumina (gradient elution with 5–10% EtOAc–hexanes) to give additional 0.034 g of the mixture of **45** and **46**. The combined mixtures were then further purified on 25 g of silica gel (elution with 10:10:80 Et₂O:benzene:CH₂Cl₂) to give 0.255 g (59%) of phenol **46** as a yellow oil: IR (film) 3314, 2952, 1698, 1619, 1573, 1454, 1407, 1311, 1251, 1116, 1093, 1074, 1041 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 8.03 (br s, 1H), 7.20–7.28 (m, 5H), 6.43 (br s, minor rotamer), 6.35 (br s, 1H), 4.68 (s, 2H), 4.00–4.36 (br m, 3H), 3.32 (s, 3H), 3.23 (s, 3H), 2.52 (d, *J* = 5.6 Hz, 2H), 2.13 (s, 3H), 2.12 (s, 3H), 1.08 (br m, minor rotamer), 0.86 (br m, 2H), 0.04 (s, minor rotamer), -0.10 (s, 9H); ¹³C NMR (100 MHz, CDCl₃), (mixture of two rotamers) for major rotamer: δ 156.6, 154.5, 138.3, 137.7, 137.1, 129.5, 128.5, 127.7 124.6, 121.8, 118.7 105.7, 64.2, 55.1, 54.8, 53.6, 30.8, 20.1, 18.1, 12.3, -1.5; additional resonances appeared for the minor rotamer at δ 129.0; HRMS (ESI) [M + Na]⁺ calcd for C₂₅H₃₇NNaO₅Si: 482.2333, found 482.2336.

***N*-Benzyl-*N*-2-(trimethylsilyl)ethoxycarbonyl-[1-hydroxy-2-(2,2-dimethoxyethyl)-5,6,7,8-tetrahydronaphthalen-3-yl]amine (48)**—A solution of ynamide **45** (0.313 g, 0.86 mmol, 1.0 equiv) in 2.8 mL of CH₂Cl₂ was distributed equally

between two 25-cm quartz reaction tubes (13 mm i.d., 15 mm o.d.) fitted with rubber septa and argon inlet needles. A 15-mL Pyrex tube fitted with a rubber septum and argon inlet needle was charged with diazo ketone **16** (0.310 g, 2.07 mmol, 2.4 equiv) and 5.8 mL of CH₂Cl₂. Both solutions were degassed by three freeze-pump-thaw cycles at -196 °C, 0.05 mmHg. The solution of diazo ketone was taken up in two 5-mL glass syringes wrapped in aluminum foil and fitted with 20-gauge, 20-cm steel needles. The upper portions (ca. 18-cm length) of the quartz tubes were wrapped in aluminum foil and the tubes were positioned ca. 3 cm from a Hanovia 450W medium pressure mercury lamp cooled in a quartz immersion well. The quartz tubes and the immersion well were partially submerged in a room temperature water bath contained in a 2-L beaker wrapped in aluminum foil. The diazo ketone solution was added to the reaction tubes via syringe pump over ca. 5 h (0.5 mL CH₂Cl₂ wash) while the reaction mixture was irradiated and then irradiation was continued for 90 min. The combined reaction mixture was concentrated to provide 0.583 g of an orange oil which was transferred with the aid of 5 mL of toluene to a 25-mL, one-necked, round-bottomed flask equipped with a cold-finger condenser with argon inlet side arm. The solution was heated at reflux for 1 h, allowed to cool to room temperature, and then concentrated to give 0.550 g of an orange oil. This material was dissolved in ca. 10 mL of CH₂Cl₂ and concentrated onto 3.5 g of silica gel. The free-flowing powder was added to the top of a column of 55 g of silica gel and eluted with 15% EtOAc-hexanes to afford 0.224 g of acetal **48** as a viscous yellow oil and 0.059 g of a mixture of phenol **48** and unreacted ynamide **45**. This mixture was dissolved in ca. 5 mL of CH₂Cl₂ and concentrated onto 0.5 g of silica gel. The free flowing powder was added to the top of a column of 6 g of silica gel and eluted with 15% EtOAc-hexanes to afford 0.027 g of phenol **48** as a viscous yellow oil. The combined yield of phenol **48** was 0.251 g (60%): IR (film) 3315, 2935, 2858, 2838, 1698, 1617, 1572, 1496, 1436, 1406, 1333, 1299, 1249, 1116, and 1072 cm⁻¹; ¹H NMR (500 MHz, CDCl₃, ca. 68:32 mixture of rotamers) for major rotamer: δ 8.13 (s, minor rotamer), 8.09 (s, 1H), 7.18-7.35 (m, 5H), 6.38 (s, minor rotamer), 6.28 (s, 1H), 4.75 (d, *J* = 14.4 Hz, 1H), 4.66 (d, *J* = 14.4 Hz, 1H), 4.05-4.47 (m, 3H), 3.40 (s, minor rotamer), 3.37 (s, 3H), 3.31 (s, minor rotamer), 3.28 (s, 3H), 2.51-2.71 (m, 6H), 1.68-1.84 (m, 4H), 1.05-1.19 (m, minor rotamer), 0.83-1.01 (m, 2H), 0.07 (s, minor rotamer), -0.07 (s, 9H); ¹³C NMR (125 MHz, CDCl₃, mixture of two rotamers) for major rotamer: δ 156.5, 154.1, 138.3, 137.7, 137.6, 129.4, 128.4, 127.5, 125.3, 120.8, 117.6, 105.6, 64.1, 55.0, 54.7, 53.5, 30.9, 30.6, 29.4, 23.5, 22.8, 17.9, and -1.5; additional resonances appeared for the minor rotamer at: δ 156.0, 139.0, 128.8, 120.4, 117.8, 55.2, 54.9, and -1.4; HRMS (ESI) [M+Na]⁺ calcd for C₂₇H₃₉NNaO₅Si: 508.2490, found 508.2497.

2-(Trimethylsilyl)ethyl Benzyl[5-(2,2-dimethoxyethyl)-6-hydroxy-2,3-dihydro-1H-phenalen-4-yl]carbamate (54)—

Two 25-cm quartz tubes (13 mm i.d., 15 mm o.d.) were each charged with ynamide **45** (0.300 g, 0.825 mmol, 1.0 equiv) and diazo ketone **53**^{2b} (0.231 g, 1.24 mmol, 1.5 equiv). A stir bar was added to each reaction tube, the tubes were fitted with rubber septa and argon inlet needles, and purged with argon. Dichloromethane (8.5 mL) was then added to each tube and the reaction tubes were positioned ca. 3 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by tap water) in a room-temperature water bath contained in a 2-L beaker wrapped in aluminum foil. The reaction solutions were irradiated for 2.5 h and then combined and concentrated to give 1.129 g of an orange oil. A solution of this material in 20 mL of toluene was transferred to a flask fitted with a cold-finger condenser. The solution was heated at reflux for 2 h, allowed to cool to rt, and concentrated to give 1.15 g of an orange oil. Column chromatography on 50 g of grade II neutral alumina (elution with 10% EtOAc-hexanes) gave 0.680 g (79%) of phenol **54** as a yellow oil: IR (film) 3291, 2949, 2835, 1698, 1620, 1583, 1496, 1398, 1357, 1318, 1250, 1208, 1184, 1141, 1115 cm⁻¹; ¹H NMR (400 MHz, CDCl₃, ca. 80:20 mixture of rotamers) for major rotamer: δ 8.50 (s, 1H),

8.48 (s, minor rotamer), 8.14 (d, $J = 8.4$ Hz, 1H), 8.13 (d, $J = 8.3$ Hz, minor rotamer), 7.41–7.36 (m, 1H), 7.31–7.10 (m, 6H), 4.84 (d, $J = 13.9$ Hz, 1H), 4.58 (d, $J = 13.9$ Hz, 1H), 4.48 (d, $J = 14.2$ Hz, minor rotamer), 4.44–4.04 (m, 3H), 3.39 (s, minor rotamer), 3.37 (s, 3H), 3.28 (s, minor rotamer), 3.27 (s, 3H), 3.01 (t, $J = 5.9$ Hz, 2H), 2.73 (d, $J = 6.9$ Hz, minor rotamer), 2.69 (d, $J = 6.9$ Hz, 2H), 2.84–2.42 (m, 2H), 1.89 (pq, $J = 6.1$ Hz, minor rotamer), 1.81 (pq, $J = 6.0$ Hz, 2H), 1.26–1.18 (m, minor rotamer), 0.95–0.75 (m, 2H), 0.12 (s, minor rotamer), and 0.10 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3 , mixture of two rotamers) for major rotamer: δ 156.7, 150.8, 137.0, 136.3, 135.0, 130.1, 129.9, 128.5, 128.0, 125.7, 125.5, 125.2, 124.8, 120.6, 114.9, 106.0, 64.2, 55.1, 54.4, 53.0, 31.36, 31.34, 26.8, 22.8, 18.1, –1.5; additional resonances appeared for the minor rotamer at δ : 155.7, 136.9, 135.5, 130.4, 130.2, 128.0, 125.1, 124.6, 115.4, 106.2, 64.3, 55.6, 54.3, 52.9, 31.5, 22.9, 18.2, –1.2; HRMS (ESI) $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{30}\text{H}_{39}\text{NNaO}_5\text{Si}$: 544.2490, found 544.2502.

4-[Benzyl([2-(trimethylsilyl)ethoxy]carbonyl)amino]-5-(2,2-dimethoxyethyl)-2,3-dihydro-1H-phenalen-6-yl Trifluoromethanesulfonate (55)—A 25-mL recovery

flask was charged with phenol **54** (0.233 g, 0.447 mmol, 1 equiv) and 4 mL of THF and cooled to 0 °C. Sodium hydride (60% wt oil disp., 0.025 g, 0.63 mmol, 1.4 equiv) was then added and the reaction mixture was stirred for ca. 5 min until gas evolution ceased. A solution of *N*-phenyltriflimide (0.224 g, 0.627 mmol, 1.4 equiv) in 2 mL of THF was added, and the resulting brown mixture was allowed to warm to rt over 15 min. The reaction mixture was cooled to 0 °C, and 3 mL of satd aq NaHCO_3 solution was added. The resulting mixture was diluted with 15 mL of Et_2O , and the aqueous layer was separated and extracted with two 15-mL portions of Et_2O . The combined organic phases were washed with two 20-mL portions of 2.5 M NaOH solution and 20 mL of brine, dried over MgSO_4 , filtered and concentrated to give 0.329 g of a yellow oil. Column chromatography on 15 g of silica gel (elution with 10% EtOAc –hexanes) gave 0.219 g (75%) of trifluoromethanesulfonate **55** as a viscous colorless oil: IR (film) 2953, 2835, 1703, 1597, 1403, 1318, 1248, 1215, 1138, 1121, 1074, 943 and 729 cm^{-1} ; ^1H NMR (600 MHz, CDCl_3) δ 8.00 (d, $J = 8.5$ Hz, 1H), 7.59–7.60 (m, 1H), 7.38–7.14 (m, 6H), 5.35 (d, $J = 14.3$ Hz, 1H), 5.19 (d, $J = 14$ Hz, minor rotamer), 4.76–4.78 (m, 1H), 4.70–4.72 (m, minor rotamer), 4.16–4.56 (m, 3H), 3.35–3.46 (m, 3H), 3.29–3.35 (m, 3H), 2.96–3.13 (m, 3H), 2.54–2.60 (m, 1H), 2.14–2.27 (m, 1H), 1.27–1.96 (m, 4H), 0.87–1.00 (m, 2H), 0.20 (s, minor rotamer), 0.00 (s, 9H); ^{13}C NMR (150 MHz, CDCl_3 , mixture of two rotamers) for major rotamer: δ 156.4, 142.4, 137.3, 137.1, 136.6, 134.7, 130.4, 130.3, 128.6, 127.6, 127.2, 126.87, 125.7, 119.6, 103.6, 64.5, 54.5, 54.1, 53.2, 34.8, 32.1, 31.1, 27.5, 22.1, 18.2, –1.5; additional resonances appeared for the minor rotamer at δ : 155.2, 142.5, 137.6, 137.5, 136.4, 134.9, 130.0, 128.7, 127.4, 126.93, 125.6, 104.3, 64.7, 55.2, 54.4, 52.2, 32.6, 31.7, 27.3, 22.8, 18.3, –1.3; HRMS (ESI) $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{31}\text{H}_{38}\text{F}_3\text{NNaO}_7\text{SSi}$: 676.1983, found 676.1986.

Methyl Allyl(2-[2-(*tert*-butyldimethylsilyloxy)ethyl]-3-hydroxy-4,5-dimethylphenyl)carbamate (61)—Two 25-cm quartz tubes (13 mm i.d., 15 mm o.d.)

fitted with rubber septa and argon inlet needles were charged with solutions of ynamide **58**⁸ (flask A: 0.168 g, 0.565 mmol, 1.0 equiv; flask B: 0.151 g, 0.508 mmol, 1.0 equiv, each in 2 mL of CH_2Cl_2). A 25-mL recovery flask was charged with a solution of diazo ketone **42** (0.340 g, 2.74 mmol, 2.6 equiv) in 7 mL of CH_2Cl_2 . All solutions were purged with argon for 15 min. The upper portions (ca. 10 cm) of the quartz reaction tubes were wrapped in aluminum foil and the tubes were positioned ca. 5 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by tap water). The reaction tubes and the immersion well were partially submerged in a room-temperature water bath contained in a 2-L beaker wrapped in aluminum foil.⁴⁴ The diazo ketone solution was taken up into two 5-mL gastight syringes wrapped in aluminum foil and fitted with 20 gauge, 20-cm steel needles and added to the reaction tubes using a syringe pump over 7 h (1 mL CH_2Cl_2

rinse).⁴⁵ Irradiation was continued for 20 min, and then the reaction mixtures were combined and concentrated in a 25-mL round-bottomed flask. The residue was diluted with 7 mL of toluene, the flask was fitted with a cold-finger condenser with an argon inlet sidearm, and the solution was heated at reflux for 3 h. The reaction mixture was allowed to cool to rt and concentrated to give 0.7 g of a dark orange oil. Column chromatography on 50 g of silica gel (elution with 10% EtOAc–hexanes) gave 0.328 g (78%) of phenol **61** as a yellow oil which crystallized upon refrigeration: mp 48–49 °C; IR (neat) 3273, 2859, 2930, 1709, 1620, 1574, 1449, 1388, 1313, 1259, 1214, 1192, 1147, 1092, 1039 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.59 (s, 1H), 6.51 (s, 1H), 5.85–5.95 (m, 1H), 5.12 (m, 2H), 3.70–4.30 (m, 4H), 3.62 (s, 3H), 2.78 (m, 2H), 2.23 (s, 3H), 2.17 (s, 3H), 0.92 (s, 9H), 0.08 (d, *J* = 5.6 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 156.5, 154.7, 137.6, 136.7, 133.3, 124.7, 122.4, 121.2, 118.3, 65.3, 54.3, 53.0, 29.5, 25.9, 20.2, 18.4, 12.3, -5.5; HRMS (ESI) [M + Na]⁺ calcd for C₂₁H₃₅NNaO₄Si: 416.2228, found 416.2218.

Methyl Allyl[4-(2-[(*tert*-butyldimethylsilyloxy)ethyl]-3-hydroxybenzocyclobutene]-5-carbamate (64**)**—

A 25-cm quartz tube (13 mm i.d., 15 mm o.d.) equipped with two stir bars, a rubber septum, and argon inlet needle was charged with ynamide **58** (0.310 g, 1.04 mmol, 1.0 equiv) and 3 mL of CH₂Cl₂. An 18x160 mm test tube was charged with a solution of diazo ketone **50** (0.328 g, 2.69 mmol, 2.6 equiv) in 8 mL of CH₂Cl₂. Both solutions were purged with argon for 15 min. The upper portion (ca. 10 cm) of the quartz reaction tube was wrapped in aluminum foil and the tube was positioned ca. 15 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by tap water). The diazo ketone solution was taken up into a 10-mL gastight syringe wrapped in aluminum foil and fitted with a 20 gauge, 20-cm steel needle and added to the reaction tubes using a syringe pump over 5 h (1 mL CH₂Cl₂ rinse). Irradiation was continued for 5 h and the reaction mixture was concentrated in a 50-mL round-bottomed flask. The residue was diluted with 15 mL of toluene, the flask was fitted with a reflux condenser with an argon inlet, and the solution was heated at reflux for 2.5 h. The resulting red-brown solution was allowed to cool to rt and then concentrated to give 0.74 g of a red-brown oil. Column chromatography on 15 g of silica gel (elution with 10% EtOAc–hexanes) yielded 0.099 g (32%) of unreacted ynamide **58** as a colorless oil and 0.251 g (62%; 91% based on recovered ynamide) of carbamate **64** as a white solid: mp 129–130 °C; IR (KBr pellet) 3317, 2928, 1679, 1593, 1457, 1432, 1310, 1258, 1148, 1089, 982, 921, 859, 834 and 773 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.64 (s, 1H), 6.45 (s, 1H), 5.87–5.93 (m, 1H), 5.11–5.14 (m, 2H), 4.31 (dd, *J* = 19.1, 10.1 Hz, 1H), 3.59–3.91 (m, 6H), 3.09–3.15 (m, 4H), 2.74–2.85 (m, 2H), 0.92 (s, 9H), 0.10 (d, *J* = 4.1 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 156.6, 151.0, 145.7, 133.2, 130.9, 124.2, 118.5, 118.4, 115.5, 65.3, 54.4, 53.0, 29.7, 29.2, 27.4, 26.0, 18.5, -5.4; Anal. calcd for C₂₁H₃₃NO₄Si: C, 64.41; H, 8.49; N, 3.58. Found: C, 64.50; H, 8.36; N, 3.53.

1-Benzyl-5,6-dimethyl-4-hydroxyindole (47**)**—A 25-mL recovery flask fitted with a rubber septum and argon inlet needle was charged with a solution of phenol **46** (0.325 g, 0.712 mmol, 1.0 equiv) in 3.5 mL of THF. The solution was cooled at 0 °C while TBAF solution (1 M in THF, 3.5 mL, 3.5 mmol, 5 equiv) was added via syringe over 3 min. The resulting brown solution was stirred at rt for 18 h, and then cooled at 0 °C while 6 M aqueous HCl solution (3.5 mL, 21.0 mmol, 30 equiv) was added dropwise over ca. 2 min. The resulting light yellow solution was stirred at 0 °C for 5 min, and then heated at reflux for 40 min. The resulting blue-brown mixture was allowed to cool to rt, diluted with 80 mL of Et₂O, and washed with two 30 mL portions of satd aq NaHCO₃ solution. The combined aqueous layers were extracted with three 15 mL portions of Et₂O, and the combined organic phases were washed with 50 mL of brine, dried over Na₂SO₄, filtered, and concentrated to give 0.184 g of a brown oil. Column chromatography on 20 g of silica gel (elution with 10%

EtOAc–hexanes) gave 0.146 g (82%) of indole **47** as a tan solid: IR (KBr pellet) 3592, 3054, 2986, 2924, 1570, 1491, 1454, 1442, 1297, 1265, 1236, 909, 705 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.25–7.32 (m, 3H), 7.09–7.11 (m, 2H), 6.98 (d, $J = 3.2$ Hz, 1H), 6.75 (s, 1H), 6.47 (dd, $J = 3.2, 0.4$ Hz, 1H), 5.27 (s, 2H), 4.85 (s, 1H), 2.35 (s, 3H), 2.25 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 146.2, 137.9, 136.2, 132.5, 128.9, 127.6, 126.8, 126.6, 116.6, 111.6, 103.5, 97.1, 50.1, 21.4, 11.3; HRMS (ESI) $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{18}\text{NO}$: 252.1383, found 252.1371.

1-Benzyl-5,6,7,8-tetrahydro-1H-benz[*f*]indol-4-yl Acetate (49).⁴⁶—A 25-mL, two-necked, round-bottomed flask fitted with a rubber septum and argon inlet adapter was charged with a solution of carbamate **48** (0.305 g, 0.63 mmol, 1.0 equiv) in 3.2 mL of THF. The solution was cooled at 0 °C while TBAF solution (1 M in THF, 3.2 mL, 3.2 mmol, 5 equiv) was added via syringe over 3 min. The resulting orange solution was stirred at rt for 8 h and then cooled at 0 °C while 6 M aq HCl solution (3.2 mL, 19.2 mmol, 30 equiv) was added dropwise over 3 min. The resulting dark orange solution was stirred at rt for 41 h. The resulting dark green mixture was diluted with 50 mL of Et_2O and washed with two 25 mL portions of satd aq NaHCO_3 solution. The combined aqueous layers were extracted with two 20 mL portions of Et_2O , and the combined organic phases were washed with 40 mL of brine, dried over Na_2SO_4 , filtered, and concentrated to give 0.207 g of a dark brown oil. This indole was immediately protected as the acetate derivative due to its propensity to oxidize. The unpurified indole was transferred to a 25-mL, one-necked, round-bottomed flask equipped with a rubber septum and argon inlet needle, and 6.3 mL of CH_2Cl_2 and Et_3N (0.138 g, 0.190 mL, 1.36 mmol, 2.2 equiv) were added. The resulting brown solution was cooled at 0 °C while acetyl chloride (0.061 g, 0.055 mL, 0.77 mmol, 1.2 equiv) was added in one portion. The resulting solution was stirred at room temperature for 2.5 h and then diluted with 10 mL of CH_2Cl_2 and washed with 10 mL of 1 M HCl solution. The aqueous layer was extracted with 10 mL of CH_2Cl_2 and the combined organic phases were washed with 10 mL of satd aq NaHCO_3 solution, 10 mL of H_2O , and 10 mL of brine, dried over MgSO_4 , filtered, and concentrated to provide 0.240 g of a brown foam. This material was dissolved in ca. 10 mL of CH_2Cl_2 and concentrated onto 2.4 g of silica gel. The free flowing powder was added to the top of a column of 35 g of silica gel and eluted with 15% EtOAc–hexanes to give 0.115 g (57% over two steps) of acetate **49** as a pale yellow solid: mp 112–115 °C; IR (KBr pellet) 3348, 3102, 3031, 2935, 2855, 1752, 1636, 1563, 1509, 1478, 1453, 1437, 1367, and 1203 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.24–7.35 (m, 3H), 7.10–7.15 (m, 2H), 7.01 (d, $J = 3.2$ Hz, 1H), 6.93 (s, 1H), 6.32 (dd, $J = 3.2, 0.7$ Hz, 1H), 5.26 (s, 2H), 2.89 (t, $J = 5.6$ Hz, 2H), 2.70 (t, $J = 6.0$ Hz, 2H), 2.42 (s, 3H), 1.75–1.86 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.2, 141.1, 137.6, 136.5, 132.6, 128.8, 128.2, 127.6, 126.9, 120.5, 120.0, 107.6, 97.8, 50.1, 30.5, 23.3, 23.3, 23.1, and 20.9; Anal. Calcd for $\text{C}_{21}\text{H}_{21}\text{NO}_2$: C, 78.97; H, 6.63; N, 4.39. Found: C, 78.75; H, 6.46; N, 4.34.

In a separate experiment, an impure sample of hydroxy indole **S1** was isolated and observed to have the following spectroscopic properties: IR (film) 3425, 2926, 2855, 1633, 1568, 1510, 1495, 1453, 1360, 1296, 1240, and 1218 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.23–7.34 (m, 3H), 7.12 (app d, $J = 6.8$ Hz, 2H), 6.98 (d, $J = 3.2$ Hz, 1H), 6.66 (s, 1H), 6.50 (d, $J = 3.2$ Hz, 1H), 5.26 (s, 2H), 4.91 (s, 1H), 2.87 (t, $J = 6.2$ Hz, 2H), 2.78 (t, $J = 6.5$ Hz, 2H), 1.86–1.91 (m, 2H), and 1.77–1.82 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 146.0, 137.9, 136.7, 133.1, 128.8, 127.6, 126.9, 126.8, 116.4, 112.2, 102.2, 97.2, 50.1, 30.8, 23.5, and 22.7.

N-Benzyl-5,6-dihydro-1H-cyclobuta[*f*]indol-4-yl Trifluoromethanesulfonate (52)—A 20-cm quartz reaction tube (13 mm i.d., 15 mm o.d.) fitted with a rubber septum and an argon inlet needle was charged with ynamide **45** (0.351 g, 0.966 mmol, 1.0 equiv) and

flushed with argon. Dichloromethane (3 mL) was then added and the reaction tube was positioned ca. 15 cm from a Hanovia 450W medium pressure mercury lamp (quartz immersion well, cooled by tap water). A solution of the diazo ketone **50**^{2b} (0.287 g, 2.35 mmol, 2.4 equiv) in 7 mL of CH₂Cl₂ was added to the vigorously stirred reaction mixture via a syringe wrapped in aluminum foil over 3.5 h. Irradiation was continued for 6.5 h and then the reaction mixture was concentrated to give 0.7 g of a dark red-orange oil. A solution of this oil in 15 mL of toluene was transferred to a flask fitted with a cold-finger condenser with argon inlet side arm and heated at reflux for 2 h. Concentration gave 0.68 g of a red-brown oil. Column chromatography on 15 g of silica gel (elution with 15% EtOAc–hexanes) afforded 0.262 g (59%) of phenol **51** as a yellow oil: IR (film) 3583, 3297, 2952, 2832, 1698, 1671, 1594, 1431, 1251, 1116, 1073, 859, 838 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): 8.08 (s, 1H), 6.37 (s, minor rotamer), 6.28 (s, 1H), 4.14–4.84 (m, 5H), 3.29–3.37 (m, 6H), 3.03–3.11 (m, 4H), 2.60 (d, 2H), 0.89–1.60 (m, 2H), 0.08 (s, minor rotamer), –0.06 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 156.5, 150.6, 145.8, 140.9, 137.5, 130.7, 129.4, 128.9, 128.4, 127.6, 120.4, 116.1, 105.7, 64.2, 55.1, 54.8, 53.5, 34.7, 30.8, 29.0, 27.2, 18.0, –1.56.

A 25-mL recovery flask fitted with a rubber septum and an argon inlet needle was charged with a solution of phenol **51** (0.170 g, 0.372 mmol, 1.0 equiv) in 2 mL of THF. The solution was cooled at 0 °C while TBAF solution (1 M in THF, 2 mL, 2 mmol, 5 equiv) was added via syringe over 1 min. The resulting brown solution was stirred at rt for 24 h, and then cooled to 0 °C while 6 M HCl solution (2 mL, 12 mmol, 32 equiv)⁴⁷ was added dropwise over ca. 2 min. The resulting solution was stirred at 0 °C for 5 min, and then heated at reflux for 1 h. The resulting blue-brown mixture was allowed to cool to rt, diluted with 20 mL of Et₂O, and washed with 20 mL of satd aq NaHCO₃ solution. The combined aqueous layers were extracted with Et₂O, and the combined organic phases were washed with brine, dried over MgSO₄ containing a small amount of decolorizing carbon, filtered, and concentrated to give 0.191 g of hydroxy indole **52** as a brown oil used in the next step without purification: ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.24 (m, 3H), 7.09–7.11 (m, 2H), 6.98 (d, *J* = 3.2 Hz, 1H), 6.64 (m, 1H), 6.61 (s, 1H), 5.26 (s, 2H), 3.54 (s, 1H) 3.11–3.15 (m, 4H).

The hydroxy indole prepared in the previous step was transferred into a 25-mL recovery flask equipped with a rubber septum and an argon inlet needle, dissolved in 3 mL of THF, and cooled to 0 °C. Sodium hydride (60% wt oil disp., 0.025 g, 0.63 mmol, 1.7 equiv) was added and the reaction mixture was stirred for 2 min until no more gas evolved. A solution of *N*-phenyltriflimide (0.210 g, 0.588 mmol, 1.6 equiv) in 2 mL of THF was added, and the resulting mixture was stirred at rt for 1 h. Satd aq NaHCO₃ (3 mL), was added, and the resulting mixture was extracted with three 20 mL-portions of Et₂O. The combined organic phases were washed with two 20-mL portions of 10% NaOH solution and 20 mL of brine, dried over MgSO₄, filtered, and concentrated to give 0.225 g of a brown oil. Column chromatography on 5 g of silica gel (elution with 10% EtOAc–hexanes) gave 0.130 g of an oily white solid. Column chromatography on 5 g of silica gel (elution with 5% EtOAc–hexanes) gave 0.093 g of colorless oil. The oil was taken up in 4 mL of pentane, and the solution was cooled to –20 °C. The resulting white crystals were washed with cold pentane and dried under vacuum. A second crop was collected by concentrating the mother liquor and the washes to a volume of ca. 0.5 mL and cooling to –20 °C. Crystallization provided 0.089 g (63% from **45**) of indole **52** as a white solid: mp 77–78 °C; IR (KBr pellet) 2926, 1497, 1421, 1250, 1200, 1187, 998, 916, 839, 727, 617 and 603 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.34 (m, 3H), 7.14 (d, *J* = 3.2 Hz, 1H), 7.07–7.13 (m, 2 H), 7.00 (s, 1H), 6.62 (d, *J* = 3.2 Hz, 1H), 5.31 (s, 2H), 3.18–3.36 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 140.5, 138.6, 137.0, 135.4, 129.1, 128.3, 128.0, 126.9, 126.0, 120.6, 119.0 (q, *J* = 320 Hz, CF₃), 105.3, 98.4, 50.9, 28.5, 27.1; Anal. calcd for C₂₁H₃₃NO₄Si: C, 64.41; H, 8.49; N, 3.58. Found: C, 64.50; H, 8.36; N, 3.53.

N-Benzyl-1,2,3,10-tetrahydronaphtho[1,8-fg]indol-7-yl

Trifluoromethanesulfonate (56)—A 25-mL recovery flask was charged with carbamate **55** (0.189 g, 0.289 mmol, 1 equiv) and 5 mL of CH₂Cl₂. Trifluoromethanesulfonic acid (0.225 mL, 2.88 mmol, 10 equiv) was added via syringe in one portion producing a dark red-brown solution. The reaction mixture was stirred vigorously at rt for 5 min and then 2 mL of water was added. This biphasic reaction mixture turned light yellow within 30 sec. Satd aq NaHCO₃ (5 mL) was then added,⁴⁸ and the reaction mixture was diluted with 20 mL CH₂Cl₂ and separated. The organic phase was washed with 10 mL of water and brine, dried over MgSO₄, filtered, and concentrated to give 0.117 g of a tan solid. Column chromatography on 10 g of silica gel (elution with 20% benzene–hexanes) followed by a column chromatography on 6 g of silica gel (elution with 20% benzene–hexanes) furnished 0.098 g (76%) of indole **56** as a white powder: mp 112–113 °C (dec); IR (film) 3055, 2987, 1422, 1266, 737 cm⁻¹; ¹H NMR (600 MHz, CDCl₃) δ 7.97 (d, *J* = 8.7 Hz, 1H), 7.36–7.40 (m, 1H), 7.25–7.35 (m, 4H), 7.15 (dd, *J* = 10.0, 3.9 Hz, 1H), 6.96 (d, *J* = 7.3 Hz, 2H), 6.82 (d, *J* = 3.4 Hz, 1H), 5.69 (s, 2H), 3.41 (t, *J* = 6.2 Hz, 2H), 3.06 (t, *J* = 6.2 Hz, 2H), 2.02–1.99 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 139.0, 136.6, 136.1, 135.0, 134.4, 129.2, 127.8, 127.2, 125.6, 124.5, 123.9, 122.6, 121.7, 119.1 (q, *J* = 310 Hz), 118.6, 118.3, 98.1, 53.2, 31.4, 26.9, 22.7; HRMS (ESI) [*M* + *H*]⁺ calcd for C₂₃H₁₉F₃NO₃S: 446.1032, found 446.1020.

N-Benzyl-7-ethyl-1,2,3,10-tetrahydronaphtho[1,8-fg]indole (57)—A 25-mL pear flask was charged with Pd(OAc)₂ (2.5 mg, 0.011 mmol, 5 mol %), SPhos (6.6 mg, 0.016 mmol, 7.5 mol %), and Cs₂CO₃ (0.208 g, 0.638 mmol, 3 equiv). The flask was equipped with a rubber septum fitted with an argon inlet needle and flushed with argon. A solution of triflate **56** (0.095 g, 0.213 mmol, 1 equiv) in 4.5 mL of THF and a solution of Et₃B (1 M in THF, 0.533 mL, 0.533 mmol, 2.5 equiv) were added. The orange-yellow reaction mixture was stirred at rt for 1 h, diluted with 15 mL of Et₂O, and filtered through 5 g of silica gel with the aid of two 20-mL portions of Et₂O. The resulting yellow solution was concentrated to give 0.061 g of an off-white solid. This material was dissolved in 50 mL of hot pentane and crystallized at –78 °C to give 0.055 g (79%) of indole **57** as an off-white powder: mp 114–115 °C; IR (film) 3054, 2936, 1453, 1386, 1355, 1265, 739 cm⁻¹; ¹H NMR (600 MHz, CDCl₃) δ 8.07 (d, *J* = 9 Hz, 1H), 7.27–7.34 (m, 5H), 7.13 (d, *J* = 3.6 Hz, 1H), 7.01 (d, *J* = 7.2 Hz, 2H), 6.81 (d, *J* = 3.0 Hz, 1H), 5.70 (s, 2H), 3.45–3.48 (m, 4H), 3.11 (t, *J* = 6 Hz, 2H), 2.02–2.06 (m, 2H), 1.49 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 140.0, 136.3, 134.5, 134.1, 129.9, 129.3, 129.0, 127.5, 127.4, 126.2, 125.8, 122.2, 114.8, 100.1, 53.2, 32.1, 27.2, 23.1, 22.9, 15.5; HRMS (ESI) [*M* + *H*]⁺ calcd for C₂₄H₂₄N: 326.1903, found 326.1900.

Methyl 2-(2-[(*tert*-Butyldimethylsilyloxy)ethyl]-3-hydroxy-4,5-dimethylphenyl(2-oxoethyl)carbamate (62)

—A 50-mL recovery flask fitted with a rubber septum and argon inlet needle was charged with alkene **61** (0.325 g, 0.826 mmol, 1.0 equiv), 6 mL of THF, 2 mL of H₂O, and OsO₄ (4 wt% in H₂O, 0.11 mL, 0.11 g, 0.018 mmol, 0.02 equiv). The solution was stirred at rt for 10 min and then NMO (0.130 g, 1.11 mmol, 1.3 equiv) was added. The resulting cloudy tan mixture was stirred at rt for 24 h. Aqueous NaHSO₃ solution (1.0 M, 9.0 mL, 9.0 mmol, 11 equiv) was added, and the reaction mixture was stirred at rt for 10 min, diluted with 15 mL of brine, and extracted with three 20-mL portions of EtOAc. The combined organic phases were dried over MgSO₄, filtered, and concentrated to give 0.401 g of diol **S3** as a pale yellow oil used in the next step without purification: IR (neat) 3281, 2955, 2931, 2884, 2860, 1685, 1619, 1573, 1455, 1391, 1321, 1260, 1197, 1164, 1092, 1039, 900, 839, 781, 732 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) (ca. 50:50 mixture of rotamers) δ 8.69 and 8.60 (s, 1H), 6.60 and 6.42 (s, 1H), 3.81–3.73 (m,

2H), 3.73–3.59 (m, 6H), 2.94–2.70 (m, 3H), 2.22 (d, $J = 8.9$ Hz, 3H), 2.17 (s, 3H), 0.92 (s, 9H), 0.10–0.08 (m, 6H).

A 25-mL recovery flask was charged with NaIO₄ supported on silica gel (0.68 mmol/g, 2.5 g, 1.7 mmol, 2.0 equiv) and 5 mL of CH₂Cl₂. A solution of the diol prepared above in 5 mL of CH₂Cl₂ was added to the stirring suspension via pipette (0.5 mL CH₂Cl₂ rinse). The reaction mixture was stirred at rt for 0.5 h and then filtered through a sintered glass funnel with the aid of three 3-mL portions of CH₂Cl₂. Concentration of the filtrate gave 0.316 g (97%) of aldehyde **62** as a pale yellow oil: IR (neat) 3259, 2955, 2931, 2885, 2859, 1706, 1619, 1574, 1451, 1387, 1331, 1260, 1136, 1093, 1053, 1038, 899, 839, 782 cm⁻¹; ¹H NMR (600 MHz, CDCl₃) δ 9.71 (s, 1H), 8.65 (s, minor rotamer), 8.59 (s, 1H), 6.61 (s, 1H), 4.43 (d, $J = 18$ Hz, 1H), 4.34 (d, $J = 18$, minor rotamer), 4.03 (d, $J = 18$ Hz, 1H), 3.87 (m, 2H), 3.75 (s, minor rotamer), 3.68 (s, 3H), 2.88 (m, 2H), 2.23 (s, 3H), 2.17 (s, 3H), 0.92 (s, 9H), 0.09 (d, $J = 6$ Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 197.3, 156.9, 154.9, 137.8, 137.2, 125.3, 122.2, 120.6, 65.2, 61.1, 53.5, 29.3, 25.9, 20.1, 18.4, 12.2, -5.5; additional resonance appeared for the minor rotamer at δ 120.4; HRMS (ESI) [M + H]⁺ calcd for C₂₀H₃₄NO₅Si: 396.2201, found 396.2216.

4,5-Dimethyl-6-hydroxy-7-(2-hydroxyethyl)-1-methoxycarbonylindole (63)—A 50-mL recovery flask equipped with a cold finger condenser with an argon inlet sidearm was charged with aldehyde **62** (0.316 g, 0.800 mmol, 1.0 equiv), 40 mL of isopropanol, and K₂CO₃ (0.11 g, 0.80 mmol, 1.0 equiv). The pale yellow reaction mixture was heated at 75 °C for 2.5 h. The oil bath was removed, and the warm mixture was diluted with 15 mL of H₂O and then treated with 5 mL of aq 1 M HCl to adjust the pH to 1. The resulting mixture was allowed to stir at rt for 0.5 h and then concentrated to a volume of ca. 10 mL. The resulting heterogeneous mixture was diluted with 200 mL of Et₂O and washed with two 40-mL portions of H₂O and 40 mL of brine. The aqueous layers were extracted with two 20-mL portions of Et₂O, and the combined organic phases were dried over MgSO₄, filtered, and concentrated to afford 0.209 g of a brown oil. Column chromatography on 7 g of silica gel (elution with 50% EtOAc–hexanes) afforded 0.130 g (62%) of indole **63** as a white solid: mp 116–117 °C; IR (KBr pellet) 3472, 3311, 3153, 3113, 2994, 2952, 2891, 1719, 1560, 1450, 1351, 1294, 1252, 1156, 1041, 930 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.79 (s, 1H), 7.42 (d, $J = 4.0$ Hz, 1H), 6.56 (d, $J = 4.0$ Hz, 1H), 4.36 (m, 2H), 3.93 (s, 3H), 3.13 (t, $J = 4.8$ Hz, 2H), 2.43 (br s, 1H), 2.41 (s, 3H), 2.30 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.0, 152.4, 133.7, 127.8, 125.6, 125.4, 121.5, 111.9, 107.4, 66.3, 54.0, 31.1, 15.8, 12.6; HRMS (ESI) [M + H]⁺ calcd for C₁₄H₁₈NO₄: 264.1230, found 264.1231.

Methyl 2-Oxoethyl[4-(2-[(*tert*-butyldimethylsilyl)oxy]ethyl)-3-hydroxybenzocyclobutene]-5-carbamate (65)—A 100-mL recovery flask fitted with a rubber septum and argon inlet needle was charged with alkene **64** (1.22 g, 3.12 mmol, 1.0 equiv), 22.5 mL of THF, 7.5 mL of H₂O, OsO₄ (4 wt% in H₂O, 0.5 mL, 0.08 mmol, 0.03 equiv), and NMO (0.552 g, 4.71 mmol, 1.5 equiv). The resulting yellow mixture was stirred at rt for 26 h. Aqueous NaHSO₃ solution (2.0 M, 9.0 mL, 9.0 mmol, 9 equiv) was added, and the reaction mixture was stirred at rt for 10 min. The resulting mixture was transferred to a separatory funnel containing 60 mL of brine and extracted with three 40-mL portions of Et₂O. The combined organic phases were dried over MgSO₄, filtered, and concentrated to give 1.4 g of diol **S4** as an off-white foam used in the next step without purification: IR (neat) 3345, 2939, 2858, 1680, 1595, 1455, 1391, 1254, 1084, 914, 836 and 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), (ca. 50:50 mixture of rotamers): δ 8.75 (s, 1H), 8.63 (s, minor rotamer), 6.55 (s, 1H), 6.38 (s, minor rotamer), 3.55–3.91 (m, 9H), 3.35–3.50 (m, 3H), 3.05–3.15 (m, 4H), 2.61–2.85 (m, 2H), 0.90 (s, 9H), 0.09 (m, 6H).

A 100-mL recovery flask was charged with NaIO₄ supported on silica gel (0.68 mmol/g, 10.6 g, 7.2 mmol, 2.3 equiv) and 20 mL of CH₂Cl₂. A solution of the diol prepared above in 15 mL of CH₂Cl₂ was added to the stirring suspension via pipette (two 5 mL CH₂Cl₂ rinses). The reaction mixture was stirred at rt for 20 min and then filtered through 1 g of silica gel with the aid of CH₂Cl₂. Concentration of the filtrate gave 1.08 g (88%) of aldehyde **65** as a white powder: mp 134–135 °C; IR (KBr pellet) 3275, 3055, 2932, 1737, 1703, 1592, 1451, 1382, 1265, 1057, 914, 839 and 738 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.69 (s, 1H), 8.71 (s, minor rotamer), 8.65 (s, 1H), 6.58 (s, 1H), 3.75–4.51 (m, 4H), 3.68 (s, 3H), 3.35 (s, minor rotamer), 3.05–3.14 (m, 4H), 2.87–2.90 (m, 2H), 0.92 (s, 9H), 0.10 (d, *J* = 2.8 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 197.1, 156.9, 151.2, 146.2, 140.5, 131.4, 124.0, 115.1, 65.2, 61.2, 53.5, 29.6, 29.1, 27.4, 26.0, 18.5, -5.45; Anal. calcd for C₂₀H₃₁NO₅Si: C, 61.04; H, 7.94; N, 3.56. Found: C, 60.77; H, 7.96; N, 3.52.

***N*-Carbomethoxy-4-(2-[(*tert*-butyldimethylsilyloxy)ethyl]-5-hydroxy-6,7-dihydro-3*H*-cyclobuta[e]indole (**66**))**—A 300-mL round-bottom flask was charged with aldehyde **65** (0.735 g, 1.87 mmol, 1.0 equiv), 100 mL of isopropanol, and K₂CO₃ (0.260 g, 1.89 mmol, 1.0 equiv). The reaction flask was equipped with a reflux condenser fitted with an argon inlet adapter, and the mixture was heated at 80 °C for 10 h. The resulting black mixture was allowed to cool to rt, neutralized with 3.3 mL of aq 0.6 M HCl, and concentrated to a volume of ca. 20 mL. This mixture was extracted with three 50 mL- portions of Et₂O, and the combined organic phases were washed with brine, dried over MgSO₄, and concentrated to afford 0.82 g of a brown oil. This oil was taken up in 200 mL of hexanes and filtered, and the filtrate was concentrated to give 0.577 g of a light brown oil. Column chromatography on 25 g of silica gel (elution with 5% EtOAc–hexanes) afforded 0.337 g (48%) of indole **66** as a white powder: mp 90–91 °C; IR (KBr pellet) 3248, 2928, 2857, 1751, 1608, 1548, 1465, 1440, 1371, 1336, 1273, 1256, 1174, 1145, 1041, 901, 786 and 719 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.14 (s, 1H), 7.45 (d, *J* = 4.0 Hz, 1H), 6.40 (d, *J* = 4.0 Hz, 1H), 4.30 (m, 2H), 3.94 (s, 3H), 3.24–3.30 (m, 4H), 3.08 (t, *J* = 4.4 Hz, 2H), 0.98 (s, 9H), 0.18 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 152.4, 149.2, 136.9, 136.7, 128.0, 127.1, 121.4, 114.2, 105.6, 67.2, 54.0, 32.3, 28.6, 28.0, 26.1, 18.5, -5.31; HRMS (ESI) [*M* + *H*]⁺ calcd for C₁₄H₁₈NO₄: 264.1230, found 264.1231; Anal. calcd for C₂₀H₂₉NO₄Si: C, 63.97; H, 7.78; N, 3.73. Found: C, 64.20; H, 7.50; N, 3.46.

Methyl 4-(2-[(*tert*-Butyldimethylsilyloxy)ethyl]-5-hydroxy-7,9-dioxo-8-phenyl-6a,7,8,9,9a,10-hexahydroisoidolo[5,6-*e*]indole-3(6*H*)-carboxylate (67**))**—A 10-cm threaded Pyrex tube (14 mm i.d., 25 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with *N*-phenylmaleimide (0.183 g, 1.06 mmol, 3.6 equiv) and flushed with argon. A solution of indole **66** (0.110 g, 0.293 mmol, 1 equiv) in 12 mL of *p*-xylene was added, the septum was replaced with a threaded Teflon cap, and reaction mixture was heated at 180 °C (bath temperature) for 40 h. The reaction mixture was allowed to cool to rt and concentrated to give 0.376 g of a dark orange oil. Column chromatography on 15 g of silica gel (elution with 20% EtOAc–hexanes) gave 0.070 g (44%) of indole **67** as a white powder: mp 172–174 °C; IR (KBr pellet) 3449, 2926, 1625, 1569, 1497, 1455, 1438, 1380, 1320, 1287, 1250, 1220, 1140, 998, 916, 840 and 727 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.24 (s, 1H), 7.19–7.50 (m, 4H), 6.97 (d, *J* = 7.2 Hz, 2H), 6.59 (d, *J* = 3.6 Hz, 1H), 4.227–4.28 (m, 2H), 3.94 (s, 3H), 2.85–3.80 (m, 8H), 0.92 (s, 9H), 0.11 (s, 3H), 0.09 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 179.0, 178.8, 152.22, 152.17, 134.6, 132.1, 129.4, 129.1, 128.5, 126.7, 126.5, 123.8, 119.8, 113.6, 106.2, 67.2, 54.1, 40.29, 40.26, 31.8, 26.2, 26.0, 22.7, 18.4, -5.42, -5.46; HRMS (ESI) [*M* + *H*]⁺ calcd for C₃₀H₃₇N₂O₆Si: 549.2415, found 549.2419.

7-Ethyl 3-Methyl 4-(2-[(*tert*-Butyldimethylsilyloxy)ethyl]-5-hydroxy-3*H*-benzo[*e*]indole-3,7(6*H*,9*H*)-dicarboxylate (69) and 8-Ethyl 3-Methyl 4-(2-[(*tert*-Butyldimethylsilyloxy)ethyl]-5-hydroxy-3*H*-benzo[*e*]indole-3,8(6*H*,9*H*)-dicarboxylate (70)—A 10-cm threaded Pyrex tube (14 mm i.d., 25 mm o.d.) equipped with a rubber septum and argon inlet needle was charged with a solution of indole **66** (0.109 g, 0.290 mmol, 1 equiv) in 10 mL of *p*-xylene followed by ethyl propiolate (0.108 mL, 1.07 mmol, 3.7 equiv). The septum was replaced with a threaded Teflon cap and the reaction mixture was heated at 180 °C (bath temperature) for 40 h. The reaction mixture was allowed to cool to rt and concentrated to give 0.159 g of an orange solid. Column chromatography on 15 g of silica gel (elution with 10% EtOAc–hexanes) gave 0.085 g (62%) of an inseparable ca. 60:40 mixture of isomers **69** and **70** as a white powder: IR (KBr pellet) 3250, 2956, 2931, 2884, 2858, 1750, 1714, 1667, 1440, 1337, 1292, 1257, 1230, 1144, 1038, 918, 838, 781 and 719 cm⁻¹; ¹H NMR (600 MHz, CDCl₃), (ca. 60:40 mixture of regioisomers) for major isomer: δ 9.24 (s, 1H), 7.53 (d, *J* = 2.8 Hz, 1H), 7.34–7.35 (m, 1H), 6.67 (d, *J* = 2.4 Hz, 1H), 4.35–4.39 (m, 4H), 4.02 (s, 3H), 3.72–3.82 (m, 2H), 3.67–3.71 (m, 2H), 3.22–3.24 (m, 2H), 1.43–1.45 (m, 3H), 1.03 (s, 9H), 0.21 (s, 6H); for the minor isomer: δ 9.23 (s, 1H), 7.52 (d, *J* = 2.4 Hz, 1H), 7.24–7.35 (m, 1H), 6.56 (d, *J* = 2.4 Hz, 1H), 4.35–4.39 (m, 4H), 4.02 (s, 3H), 3.72–3.82 (m, 2H), 3.67–3.71 (m, 2H), 3.22–3.24 (m, 2H), 1.43–1.45 (m, 3H), 1.0 (s, 9H), 0.21 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 153.0, 152.6, 152.2, 137.2, 134.9, 134.1, 129.0, 127.0, 125.8, 124.7, 123.8, 123.3, 123.1, 119.1, 117.6, 112.5, 112.4, 106.5, 106.3, 67.3, 60.7, 60.6, 54.1, 31.7, 31.7, 28.4, 26.7, 26.3, 26.1, 26.0, 24.4, 18.5, 18.4, 14.6, –5.3, –5.4; HRMS (ESI) [M + H]⁺ calcd for C₂₅H₃₆NO₆Si: 474.2306, found 474.2312.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Science Foundation (CHE-1111567), the National Institutes of Health (GM 28273), Merck Research Laboratories, and Boehringer Ingelheim Pharmaceuticals for generous financial support. T. P. W. was supported in part by an AstraZeneca Graduate Fellowship and David A. Johnson, Merck, and George Büchi Summer Fellowships. T. Y. L. was supported in part by a fellowship from the C. P. Chu and Y. Lai Fund. Y.-P. Wang was supported in part by an Eli Lilly Graduate Fellowship.


Supporting Information Available: ¹H and ¹³C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References


- (a) Kotha S, Misra S, Halder S. *Tetrahedron*. 2008; 64:10775–10790. Reviews: (b) Saito S, Yamamoto Y. *Chem. Rev.* 2000; 100:2901–2915. [PubMed: 11749309]
- (a) Danheiser RL, Gee SK. *J. Org. Chem.* 1984; 49:1672–1674. (b) Danheiser RL, Brisbois RG, Kowalczyk JJ, Miller RF. *J. Am. Chem. Soc.* 1990; 112:3093–3100.
- (a) Danheiser, RL.; Dudley, GB.; Austin, WF. Alkenylketenes. In: Danheiser, RL., editor. *Science of Synthesis*. Vol. 23. Thieme; Stuttgart: 2006. p. 493–568. For reviews of the chemistry of vinylketenes, see (b) Tidwell, TT. *Ketenes*. Second. John Wiley & Sons; Hoboken, NJ: 2006. p. 206–214.
- (a) Liebeskind LS, Iyer S, Jewell CF. *J. Org. Chem.* 1986; 51:3065–3067. A related, complementary strategy has been reported by Liebeskind and Moore. See ref. 3a and. (b) Perri ST, Foland LD, Decker OHH, Moore HW. *J. Org. Chem.* 1986; 51:3067–3068.
- Dudley GB, Takaki KS, Cha DD, Danheiser RL. *Org. Lett.* 2000; 2:3407–3410. For examples, see, and references cited therein. [PubMed: 11029223]
- (a) Kowalski CJ, Lal GS. *J. Am. Chem. Soc.* 1988; 110:3693–3695. For additional examples of the application of this benzannulation strategy in natural product synthesis, see. (b) Smith AB III,

- Adams CM, Kozmin SA, Paone DV. *J. Am. Chem. Soc.* 2001; 123:5925–5937. [PubMed: 11414825]
7. Mak XY, Crombie AL, Danheiser RL. *J. Org. Chem.* 2011; 76:1852–1873. [PubMed: 21322545]
 8. Lam TY, Wang Y-P, Danheiser RL. *J. Org. Chem.* 2013; 78:9396–9414. (DOI: 10.1021/jo401635c). [PubMed: 23952525]
 9. (a) Wakamatsu T, Miyachi N, Ozaki F, Ban Y. *Heterocycles.* 1987; 26:1445–1448. For example, see. (b) Curini M, Epifano F, Marcotullio MC, Rosati O, Guo M, Guan Y, Wenkert E. *Helv. Chim. Acta.* 2005; 88:330–338.
 10. (a) Kirmse W. *Eur. J. Org. Chem.* 2002:2193–2256. For reviews on the Wolff rearrangement, see: (b) Meier H, Zeller K-P. *Angew. Chem. Int. Ed.* 1975; 14:32–43.
 11. For the purpose of this paper, we refer to *N*-sulfonyl and *N*-phosphoryl ynamines as ynamides
 12. (a) Evano G, Jouvin K, Coste A. *Synthesis.* 2013; 45:17–26. For recent reviews on the synthesis and chemistry of ynamides, see: (b) DeKorver KA, Li H, Lohse AG, Hayashi R, Lu Z, Zhang Y, Hsung RP. *Chem. Rev.* , 110. 2010:5064–5106. [PubMed: 20429503] (c) Evano G, Coste A, Jouvin K. *Angew. Chem. Int. Ed.* 2010; 49:2840–2859.
 13. (a) Dunetz JR, Danheiser RL. *Org. Lett.* 2003; 5:4011–4014. [PubMed: 14535766] (b) Kohnen AL, Dunetz JR, Danheiser RL. *Org. Synth.* 2007; 84:88–101. [PubMed: 20657723]
 14. (a) Frederick MO, Mulder JA, Tracey MR, Hsung RP, Huang J, Kurtz KCM, Shen L, Douglas . J. *Am. Chem. Soc.* 2003; 125:2368–2369. [PubMed: 12603105] (b) Zhang Y, Hsung RP, Tracey MR, Kurtz KCM, Vera EL. *Org. Lett.* 2004; 6:1151–1154. [PubMed: 15040745] (c) Zhang X, Zhang Y, Huang J, Hsung RP, Kurtz KCM, Oppenheimer J, Petersen ME, Sagamanova IK, Shen L, Tracey MR. *J. Org. Chem.* 2006; 71:4170–4177. [PubMed: 16709057] (d) Sagamanova IK, Kurtz KCM, Hsung RP. *Org. Synth.* 2007; 84:359–367.(e) DeKorver KA, Walton MC, North TD, Hsung RP. *Org. Lett.* 2011; 13:4862–4865. [PubMed: 21848304]
 15. Hofmeister H, Annen K, Laurent H, Wiechert R. *Angew. Chem. Int. Ed.* 1984; 23:727–729.
 16. (a) Hamada T, Ye X, Stahl SS. *J. Am. Chem. Soc.* 2008; 130:833–835. For other useful coupling routes to ynamides, see. [PubMed: 18166058] (b) Jouvin K, Couty F, Evano G. *Org. Lett.* 2010; 12:3272–3275. [PubMed: 20565123] (c) Jouvin K, Coste A, Bayle A, Legrand F, Karthikeyan G, Tadiparthi K, Evano G. *Organometallics.* 2012; 31:7933–7947.(d) Jouvin K, Heimburger J, Evano G. *Chem. Sci.* 2012; 3:756–760.
 17. (a) Maas G. *Angew. Chem. Int. Ed.* 2009; 48:8186–8195. For reviews on the synthesis and chemistry of diazo ketones, see: (b) Zhang Y, Wang J. *Chem. Commun.* 2009:5350–5361.(c) Zhang Y, Wang J. *Tetrahedron.* 2008; 64:6577–6605.(d) Doyle, MP.; McKervey, MA.; Ye, T. *Modern Catalytic Methods for Organic Synthesis of Diazo Compounds: from Cyclopropanes to Ylides.* Wiley & Sons; New York; 1998. *Synthesis of α - Diazocarbonyl Compounds*; p. 1-60.(e) Regitz, M.; Maas, G. *Diazo Compounds: Properties and Synthesis.*; Academic Press; Orlando, FL; 1986.
 18. (a) Danheiser RL, Miller RF, Brisbois RG, Park SZ. *J. Org. Chem.* 1990; 55:1959–1964.(b) Danheiser RL, Miller RF, Brisbois RG. *Org. Synth.* 1996; 73:134–140.
 19. Chataigner I, Panel C, Gérard H, Pietre SR. *Chem. Commun.* 2007:3288–3290.
 20. (a) Witulski B, Schweikert T, Schollmeyer D, Nemkovich NA. *Chem. Commun.* 2010; 46:2953–2955. For examples involving the reactivity of ynamide derivatives, see. (b) Li H, Hsung RP, DeKorver KA, Wei Y. *Org. Lett.* 2010; 12:3780–3783. [PubMed: 20690601]
 21. For details, see Experimental Section
 22. (a) Weigmann RH, Würthwein E-U. *Tetrahedron Lett.* 1989; 30:6147–6150.(b) Schulte N, Möller MH, Rodewald U, Würthwein E-U. *Chem. Ber.* 1994; 127:1287–1293.
 23. The moisture sensitivity of *N,N*-bis(trimethylsilyl) ynamines also limits their utility as benzannulation partners. To date our attempts to prepare more stable bis-silyl ynamines have not been fruitful
 24. (a) Knowles JP, Elliott LD, Booker-Milburn KI. *Beilstein J. Org. Chem.* 2012; 8:2025–2052. For reviews of continuous flow photochemical reactions, see. [PubMed: 23209538] (b) Oelgemöller M, Shvydkiv O. *Molecules.* 2011; 16:7522–7550. [PubMed: 21894087]
 25. (a) Nguyen JD, Reiss B, Dai C, Stephenson CRJ. *Chem. Commun.* 2013; 49:4352–4354. For recent examples of continuous flow photochemical reactions, see. (b) Zhang Y, Blackman ML,


- Leduc AB, Jamison TF. *Angew. Chem. Int. Ed.* 2013; 52:4251–4255.(c) Maskill KG, Knowles JP, Elliott LD, Alder RW, Booker-Milburn KI. *Angew. Chem. Int. Ed.* 2013; 52:1499–1502.(d) Šterk D, Juki M, asar Z. *Org. Process. Res. Dev.* 2013; 17:145–151.(e) Junkers T, Conradi M. *J. Photochem. Photobiol., A.* 2013; 259:41–46.(f) Harrowven DC, Mohamed M, Gonçalves TP, Whitby RJ, Bolien D, Sneddon HF. *Angew. Chem. Int. Ed.* 2012; 51:4405–4408.(g) Lévesque F, Seeberger PH. *Angew. Chem. Int. Ed.* 2012; 51:1706–1709.(h) Anderson BG, Bauta WE, Cantrell WR Jr. *Org. Process. Res. Dev.* 2012; 16:967–975.(i) Terao K, Nishiyama Y, Tanimoto H, Morimoto T, Oelgemöller M, Kakiuchi K. *J. Flow Chem.* 2012; 2:73–76.
26. Hook BDA, Dohle W, Hirst PR, Pickworth M, Berry MB, Booker-Milburn KI. *J. Org. Chem.* 2005; 70:7558–7564. [PubMed: 16149784]
 27. For additional details regarding the construction of the reactor and its operation see the experimental section
 28. Shintani R, Murakami M, Hayashi T. *Org. Lett.* 2009; 11:457–459. For example, see: [PubMed: 19072063]
 29. Park KK, Lee JL, Ryu J. *Tetrahedron.* 2003; 59:7651–7659. For the photo-Fries rearrangement of sulfonamides, see. and references cited therein.
 30. School M, Ding S, Lee CW, Grubbs RH. *Org. Lett.* 1999; 1:953–956. [PubMed: 10823227]
 31. Jiménez MC, Márquez F, Miranda MA, Tormos R. *J. Org. Chem.* 1994; 59:197. For a discussion on the photocyclization of allylphenols and related compounds, see: and references therein.
 32. Improved yields in the benzannulation were obtained by treating the crude product with KOH in methanol. This serves to hydrolyze any esters that were formed by trapping of the hydroxyl group in the product phenol with ketene intermediates during the reaction
 33. In this case reaction with TBAF led to cleavage of the phenolic triflate
 34. (a) Mehta G, Kotha S. *Tetrahedron.* 2001; 57:625–659. For reviews on cyclobutane chemistry, see. (b) Sadana AK, Saini RK, Billups WE. *Chem. Rev.* 2003; 103:1539–1602. [PubMed: 12683790]
 35. Barder TE, Walker SD, Martinelli JR, Buchwald SL. *J. Am. Chem. Soc.* 2005; 127:4685–4696. [PubMed: 15796535]
 36. (a) Charlton JL, Alauddin MM. *Tetrahedron.* 1987; 43:2873–2889. Reviewed in. (b) Segura JL, Martin N. *Chem. Rev.* 1999; 99:3199–3246. [PubMed: 11749515]
 37. Zhong Y-L, Shing Y-L, Zhong TKM. *J. Org. Chem.* 1997; 62:2622–2624. [PubMed: 11671605]
 38. (a) Watson SC, Eastham JF. *J. Organomet. Chem.* 1967; 9:165–167. For the titration of *n*-butyllithium see: (b) Ellison RA, Griffin R, Kotsonis FN. *J. Organomet. Chem.* 1972; 36:209–213. (c) Lin H-S, Paquette L. *Synth. Commun.* 1994; 24:2503–2505. For the titration of *i*-PrMgBr see:
 39. Roulland E, Monneret C, Florent J-C. *J. Org. Chem.* 2002; 67:4399–4406. [PubMed: 12076134]
 40. He Y, Lin M, Li Z, Liang X, Li G, Antilla JC. *Org. Lett.* 2011; 13:4490–4493. [PubMed: 21834569]
 41. Williams JW, Hurd CD, Hanford WE, Sauer JC. *J. Org. Chem. Org. React.* 1940; 1958; 53:122, 108–140. See also.
 42. Kohnen AL, Mak XY, Lam TY, Dunetz JR, Danheiser RL. *Tetrahedron.* 2006; 62:3815–3822. [PubMed: 21394235]
 43. Fu N, Allen AD, Chan W, Kobayashi S, Tidwell TT, Tahmassebi D, Aguilar A, Cabrera EP, Godoy J. *Can. J. Chem.* 2008; 86:333–341.
 44. The reaction tubes were positioned such as to allow efficient stirring with a magnetic stir motor. This required that the bottoms of the reaction tubes were lower than the lamp's anode, as opposed to having the solution aligned with the center of the arc
 45. The bath temperature rose to 29 °C during the irradiation
 46. All solvents and solutions used in this reaction were degassed prior to use by bubbling argon through the solution for 10 min
 47. Because of the facile air oxidation of the product indole to the corresponding *p*-quinone, HCl and all of the workup and chromatography solutions were purged with argon prior to use
 48. Quick neutralization is essential once the product indole forms to avoid acid-promoted dimerization/decomposition




Scheme 1.
Tandem Benzannulation-Cyclization Strategies Based on Cyclobutenones as Vinylketene Precursors




Scheme 2.
Benzannulation Disconnection for Polycyclic Compounds




Scheme 3.
Pericyclic Cascade Mechanism of Benzannulations Based on Diazo Ketones



Scheme 4.
Tandem Photochemical Benzannulation-RCM




Scheme 5.
Tandem Thermal Benzannulation-RCM



Scheme 6.
Tricyclic Indoles via Diels-Alder Reactions of *o*-Quinodimethanes

Table 1

Optimization of the Photochemical Benzannulation



entry	diazo ketone (equiv)	solvent	irradiation ^a time (h)	yield (%) ^b
1	1.0	CH ₂ Cl ₂	8	48
2	2.5	CH ₂ Cl ₂	5.5	67
3	1.5	CH ₂ Cl ₂	4.5	65 ^c
4	2.5	CH ₂ Cl ₂	10	85 ^c
5	2.5	ClCH ₂ CH ₂ Cl	9	73 ^{cd}

^aIrradiation with a 450 W medium-pressure mercury lamp (220-600 nm).


^bIsolated yields of products purified by column chromatography.

^cA solution of diazo ketone (0.4 M) was added to the ynamide solution via syringe pump (ca. 0.009 mL/min).


^dHeated in DCE (reflux, 24 h) after irradiation.

Table 2

Photochemical Benzannulations Performed in Batch and Flow



entry	diazo ketone	ynamide	product	% yield ^a (method) ^{b,c}	entry	diazo ketone	ynamide	product	% yield ^a (method) ^{b,c}
1		17		85 ^d (batch)	4		17		40 (batch) 37 (flow)
2		25		66 (batch) 53 (flow)	5		17		57 ^e (batch) 47-50 ^e (flow)
3		17		64-76 (batch) 66 (flow)	6		22		52 ^e (batch) 47 ^e (flow)



entry	diazo ketone	ynamide	product	% yield ^a (method) ^{b, c}	entry	diazo ketone	ynamide	product	% yield ^a (method) ^{b, c}
									78 (batch) 79 (flow)
					7				

^a Isolated yield of products purified by column chromatography

^b Batch reactions: 1.1 equiv of diazo ketone, irradiation (220-600 nm) for 3-8 h in CH_2Cl_2 ; then reflux, toluene, 1.5-2.5 h. For entry 6, irradiation at >340 nm (uranium filter) for 50 h.


^c Continuous-flow reactions: 1.1 equiv of diazo ketone, irradiation at >280 nm (Pyrex filter) in DCE, residence time = ca. 21 min; then reflux, DCE, 40-48 h (entries 2, 3, 6, and 7) or reflux, toluene, 2 h (entries 4 and 5). For entry 6, irradiation at >340 nm (uranium filter), residence time = ca. 33 min.

^d Slow addition of 2.5 equiv of diazo ketone.

^e Overall yield for 2 steps. Benzannulation product was treated with 1.2 equiv TiO_2 , 2 equiv 4-DMAP, CH_2Cl_2 , 0 °C to rt, 2-4 h.

Table 3

Tandem Benzannulation/Cyclization Strategy for the Synthesis of Indoles



entry	diazo ketone	benzannulation product	yield (%) ^b	cyclization method	indole	yield (%) ^b
1			59	A		82
2			60	A		57 ^c
3			59	A		63 ^d
4			79 75 ^e	B		76

^aIrradiation (220-600 nm) of **45** and 2.4 equiv of diazo ketone (1.5 equiv in entry 4). In entries 1-3 diazo ketone was added over 3.5-5 h followed by 1.0-6.5 h of further irradiation. In entry 4 a solution of **45** and **53** was irradiated for 2.5 h.

^bIsolated yield of products purified by column chromatography.









^cOverall yield from **48** for cyclization and acetylation (1.2 equiv AcCl, 2.2 equiv Et₃N, CH₂Cl₂, 0 °C to rt, 2.5 h). In this case reaction with HCl was conducted at rt for 41 h.

^dOverall yield from **51** for cyclization and trifluoromethylsulfonylation (1.7 equiv NaH, 1.6 equiv PhNTf₂, THF, 0 °C to rt, 1 h).

^eTrifluoromethylsulfonylation of **54** by reaction with 1.4 equiv NaH, 1.4 equiv PhNTf₂, THF (0 °C to rt, 15 min).

Table 4

Tandem Benzannulation-Aromatic Substitution Strategy for the Synthesis of Indoles

entry	diazo ketone	benzannulation product	yield (%) ^d	aldehyde	yield (%) ^d	indole	yield (%) ^d
1			76-78		89-97		59-62
2			62-75		88-92		48-62

^a Irradiation (220-600 nm) of **58** and 2.6 equiv of diazo ketone. Diazo ketone was added over 5-7 h followed by 0.3-5 h of further irradiation.^b Oxidation conditions: 2-3 mol % OsO₄, 1.3-1.5 equiv NMO, THF/H₂O (rt, 24-26 h); then 2.0-2.3 equiv NaIO₄-SiO₂, CH₂Cl₂ (rt, 10-30 min).^c Cyclization conditions: 1.0 equiv K₂CO₃, *t*-PrOH, 75-80 °C, 2.5 h (entry 1) or 10 h (entry 2), then 1 M HCl to pH 1 (entry 1) or neutralize with 0.6 M HCl (entry 2).^d Isolated yield of products purified by column chromatography.