

Thermoregulation and Stress Hormone Recovery After Exercise Dehydration: Comparison of Rehydration Methods

Brendon P. McDermott, PhD, ATC*; Douglas J. Casa, PhD, ATC, FNATA, FACSM†; Elaine Lee, PhD†; Linda Yamamoto, PhD†; Kathleen Beasley, MS†; Holly Emmanuel, MS, ATC†; Jeffrey Anderson, MD†; Linda Pescatello, PhD, FACSM†; Lawrence E. Armstrong, PhD, FACSM†; Carl Maresh, PhD, FACSM†

*Athletic Training Education Program, Department of Health, Human Performance and Recreation, University of Arkansas, Fayetteville; †Human Performance Laboratory, Department of Kinesiology, University of Connecticut, Storrs

Context: Athletic trainers recommend and use a multitude of rehydration (REHY) methods with their patients. The REHY modality that most effectively facilitates recovery is unknown.

Objective: To compare 5 common REHY methods for thermoregulatory and stress hormone recovery after exercise dehydration (EXDE) in trained participants.

Design: Randomized, cross-over, controlled study.

Patients or Other Participants: Twelve physically active, non-heat-acclimatized men (age = 23 ± 4 years, height = 180 ± 6 cm, mass = 81.3 ± 3.7 kg, $\text{Vo}_2\text{max} = 56.9 \pm 4.4$ mL·min⁻¹·kg⁻¹, body fat = $7.9\% \pm 3\%$) participated.

Intervention(s): Participants completed 20-hour fluid restriction and 2-hour EXDE; they then received no fluid (NF) or REHY (half-normal saline) via ad libitum (AL), oral (OR), intravenous (IV), or combination IV and OR (IV + OR) routes for 30 minutes; and then were observed for another 30 minutes.

Main Outcome Measure(s): Body mass, rectal temperature, 4-site mean weighted skin temperature, plasma stress hormone concentrations, and environmental symptoms questionnaire (ESQ) score.

Results: Participants were hypohydrated (body mass $-4.23\% \pm 0.22\%$) post-EXDE. Rectal temperature for the NF group was significantly greater than for the IV group ($P = .023$) at 30 minutes after beginning REHY (REHY30) and greater than OR, IV, and IV + OR ($P \leq .009$) but not AL ($P = .068$) at REHY60. Mean weighted skin temperature during AL was less than during IV + OR at REHY5 ($P = .019$). The AL participants demonstrated increased plasma cortisol concentrations compared with IV + OR, independent of time ($P = .015$). No differences existed between catecholamine concentrations across treatments ($P > .05$). The ESQ score was increased at REHY60 for NF, AL, OR, and IV ($P < .05$) but not for IV + OR ($P = .217$). The NF ESQ score was greater than that of IV + OR at REHY60 ($P = .012$).

Conclusions: Combination IV + OR REHY reduced body temperature to a greater degree than OR and AL REHY when compared with NF. Future studies addressing clinical implications are needed.

Key Words: heat stress, heat illness, hypohydration

Key Points

- A combination of intravenous and oral fluid replacement after exercise dehydration reduced heat-related symptoms more than a single mode of rehydration.
- If intravenous fluid replacement is warranted, the addition of simultaneous oral rehydration seems to add benefit to an efficient recovery.

The importance of rehydration (REHY) after exercise dehydration (EXDE) is emphasized by the many physiologic systems affected by hypohydration. Many of the adverse effects of hypohydration can impair recovery or hamper future exercise performance.^{1,2} The popular mantra surrounding REHY is that intravenous (IV) fluid is more rapid and advantageous to the athlete. Whether IV fluid administration is necessary in many instances can be questioned.³ Furthermore, it may not be ethical or legal in athletes without a significant medical reason.⁴

Previous researchers^{5–14} have identified that cardiovascular and thermoregulatory strain corresponds with greater

body fluid losses. When blood volume is limited, evaporative sweat losses are impeded and overall heat dissipation decreases.^{5,9,14–17} After EXDE, if dehydration is lasting, body temperature decreases at a slower rate, and this can delay recovery.⁴ Cellular metabolism remains elevated in conjunction with high temperatures, further complicating recovery.¹⁵ A delay in exercise recovery could have deleterious implications, hindering the benefits of exercise sessions or increasing the amount and duration of cellular stress.¹⁵ Finish-line medical tent staffs treat many patients after endurance races (triathlons, marathons, ultramarathons) for EXDE.³ The recovery time for these patients is the limiting factor in the number of patients that




Figure 1. Experimental flow chart. Urine values consist of urine color, specific gravity, and osmolality. Abbreviations: CATS, plasma concentration of norepinephrine, epinephrine, and dopamine; CORT, plasma cortisol concentration; ESQ, Environmental Symptoms Questionnaire; T_{msk}, mean weighted skin temperature.

a tent staff can treat at once and should be expedited when possible.³

Decreased water content and heat stress negatively affect cell function and integrity. Circulating catecholamine concentrations are typically used to quantify sympathetic stress.^{18–22} Previous researchers who compared modes of REHY offer a glimpse of cellular recovery. These studies^{18,19,21} demonstrate no appreciable differences between modes of REHY on cellular stress recovery. However, no previous authors have included trials during which participants consumed fluids as thirst dictated or combined oral (OR) and IV fluids.

Comparisons of modes of REHY demonstrate the importance of fluid replacement after EXDE.^{18,19,21,23,24} A lack of REHY delays recovery and profoundly decreases subsequent exercise performance.⁴ When equal volumes of fluid are provided, IV and OR fluid administration are essentially the same.^{19,21,23} No previous researchers have tested post-EXDE recovery facilitated by ad libitum (AL) consumption or a combination of IV and OR fluids. Therefore, the purpose of our study was to compare modes of REHY after heat stress and EXDE in recovery of body temperature and sympathetic stress hormone concentrations. We hypothesized that a combination trial, including simultaneous IV and OR fluids, would maximize the rate at which body temperature and hormone levels return to baseline.

METHODS

Research Design

We used a randomized, crossover, controlled comparison. All participants completed 5 trials consisting of baseline, EXDE, and REHY components. The only difference between trials was the mode of REHY, which occurred in random order. To establish scientific control, 1 trial included no fluid (NF) but mirrored data collection for all other trials. Other REHY trials consisted of (1) fluid REHY via AL consumption, in which participants were instructed to drink “as little or as much as they desired, according to thirst”; (2) metered OR REHY, which was quantified and separated into equal boluses provided at REHY0 and every 5 minutes throughout REHY (at 0, 5, 10, 15, 20, and 25 minutes); (3) IV REHY, or (4) a combination of IV and OR REHY, in which half of the fluid received was via IV and the other half was OR (IV + OR). Metered fluids were based on a goal of REHY back to -2% body mass.

Dependent recovery variables included rectal temperature (T_{rec}), mean weighted skin temperature (T_{msk}), stress

hormone levels, and other blood and perceptual measures of stress and thermoregulatory strain. Measurements were recorded according to the experimental flow chart in Figure 1.

Participants. Twelve well-trained, non-heat-acclimatized men (age = 18–39 years) from the college community were recruited. The men were participating in a moderate fitness program involving a minimum of 8 hours of moderately intense exercise training per week. Screening information was obtained to ensure participants did not have (1) chronic health problems; (2) a previous history of heat illness within the past 3 years; (3) a history of cardiovascular, metabolic, or respiratory disease; (4) a history of consuming any muscle-building supplements within the past year; or (5) a history of smoking within the past 2 years. The participants were instructed to continue regular involvement in their training regimes in order to maintain cardiovascular fitness throughout the study. Physical characteristics of our participants were age = 23 ± 4 years, height = 180 ± 6 cm, mass = 81.3 ± 3.7 kg, $\dot{V}O_{2\max} = 56.9 \pm 4.4$ mL·min⁻¹·kg⁻¹, and body fat = $7.9\% \pm 3\%$. This study was approved by the University of Connecticut institutional review board, and all participants attended a prestudy briefing during which they had ample time to ask questions. This study was approved by the college institutional review board, and all participants signed a human subjects informed consent form. Participants were financially compensated.

Baseline Testing. Participants initially attended baseline testing and familiarization visits for anthropometric measurements and to ensure an understanding of the experimental protocol. Baseline skin fold calculated body fat percentage (3 site, Jackson and Pollock²⁵) was collected by the same trained clinician. Maximal oxygen uptake ($\dot{V}O_{2\max}$) testing was completed via an incremental run to exhaustion on a treadmill (ParvoMedics, Sandy, UT). The $\dot{V}O_{2\max}$ testing occurred in the heat (35°C , 40% relative humidity) to match experimental conditions. Participants also practiced exercising on the treadmill and cycle ergometer to verify exercise intensity for experimental trials as a percentage of $\dot{V}O_{2\max}$ and to familiarize themselves with the testing protocol. We chose to use multiple modalities (walking and cycling) for EXDE to prevent boredom and specific muscular fatigue. Euhydrated baseline body mass was collected by having participants report to the laboratory on 3 consecutive days after a 12-hour fast and consuming extra water the night before and morning of baseline measures.²⁶ Euhydration was verified by urine specific gravity (USG) ≤ 1.020 .^{1,2}

Experimental Protocol. The fluid provided in the IV and IV + OR trial via IV was half-normal saline (0.45% NaCl), and the OR fluid for AL, OR, and IV + OR trials was the same with a noncaloric lemon flavoring (Supervalu Inc, Eden Prairie, MN) added. All REHY fluid was stored in the same insulated cooler in the laboratory (ambient temperature $\sim 22^{\circ}\text{C}$) outside the environmental chamber until REHY began to assure no difference in temperature among fluids. The EXDE/REHY trials were separated by at least 6 days to minimize fatigue, allow normal training to continue, and eliminate acclimation (average separation = 10 ± 5 days).

Experimental trials began with a baseline blood draw. Participants reported to the laboratory after a 12-hour overnight fast (no food or fluid, except water) and 24 hours without exercising, drinking alcohol, or consuming stimulants (including caffeine). To ensure euhydration, participants were asked to drink an extra 36 fluid ounces (1.06 L) of water the night before each baseline visit and 36 fluid ounces of water the morning of each visit before they arrived at the laboratory. Participants provided a small urine sample and voided their bladders, and researchers recorded body mass. Urine was assessed for color, USG, and osmolality via freezing-point depression (model 3250; Advanced Instruments, Norwood, MA). Participants then completed a baseline environmental symptoms questionnaire (ESQ) and thermal sensation scale. Beginning at 1 PM, participants did not consume fluids and were asked to eat low-moisture foods and refrain from ingesting stimulants (including caffeine). After the blood draw, participants left the laboratory and reported back the following day.

Upon arrival, participants were weighed and provided a urine sample to verify hydration state (with a goal of between -1% and -2% hypohydration). They ate a standard meal (bagel, banana, 3 g cream cheese; 490 kcal, 11 g fat, 86 g carbohydrate, 12 g protein, 356 mg Na^+) provided by the researchers. We held meal consistent between trials by weighing the volitional amount consumed for trial 1 and weighing subsequent meals to assure the same amount of all food was consumed for the 5 trials. After breakfast, participants had 20-gauge Teflon cannulas inserted into both arms at the antecubital veins that were kept patent with normal saline and heparin (9:1).

Participants then entered the environmental chamber (model 2000; Minus Eleven, Inc, Malden, MA) for the remainder of the testing protocol. Environmental conditions for all trials averaged temperature of $35.5 \pm 1.5^{\circ}\text{C}$, relative humidity of $33 \pm 8\%$, and barometric pressure of 748 ± 6 mm Hg. After a 15-minute seated equilibration, pre-EXDE blood was drawn. For EXDE, participants alternated between walking on a treadmill at 5.6 to 6.8 $\text{km}\cdot\text{h}^{-1}$ with 5% to 10% grade (40% – 60% $\dot{V}\text{O}_2\text{max}$), and cycling on an ergometer at a power output preset to elicit an output of 40% to 60% $\dot{V}\text{O}_2\text{max}$ for 30-minute increments. Body mass was monitored at 30-minute increments. Participants were provided a small snack at EXDE90 (230 or 240 kilocalories with equal amounts of macronutrients, Na^+ , K^+ ; Powerbar, Glendale, CA). Each participant was provided with the same weight and snack bar flavor in each trial after eating volitionally (up to the provided nutrition information) during the first trial.

Participants walked or cycled for 120 minutes with the goal of decreasing total body mass by -4% .

Post-EXDE, participants remained seated and received 1 of 5 REHY methods (except in the NF trial) with a goal of -2% by reducing baseline body mass within 30 minutes. For 30 minutes post-REHY, the participant was observed (Figure 1). Sweat rates were calculated for the EXDE and REHY periods using standard calculations.¹

Nutrition Analysis. Dietary records for food intake were completed by the participants for the 3 days preceding the EXDE experimental protocol. Participants were asked to match their diet for these days leading into each trial, and copies of previous records were provided for each trial. Dietary records were entered into nutrition analysis software (Nutritionist Pro, version 4.2.0; Axxya Systems, Stafford, TX) to analyze total caloric, carbohydrate, protein, and fat intake.

Body Temperature Measures. Participants' T_{rec} was measured via flexible thermistors inserted 10 cm beyond the anal sphincter (Yellow Springs Instruments, Yellow Springs, OH). Temperature measures were recorded every 15 minutes during EXDE and every minute during REHY. Skin temperature measurements were recorded (Ototemp 3000; Exergen, Watertown, MA) in succession at the chest, forearm, thigh, and calf. Participants' T_{msk} was then calculated using these data according to Ramanathan et al.²⁷

Physiologic Measures. Blood was collected using tubes containing EDTA (Vacutainer; Becton Dickinson and Company, Franklin Lakes, NJ) and then centrifuged at 1500 rpm for 15 minutes for component separation. Plasma was then aliquoted into sealed plasma storage containers and frozen at -80°C until later analysis. Plasma cortisol concentrations ([CORT]) were quantified using a standard, commercially available enzyme-linked immunosorbent assay (ELISA) kit (Assay Designs, Inc, Ann Arbor, MI). Coefficient of variation for [CORT] analysis was 6.4%.

Standard sample extraction via commercially available methods (ESA, Inc, Chelmsford, MA), followed by high-performance liquid chromatography (HPLC) was used to quantify plasma concentrations of norepinephrine (NE), epinephrine (EPI), and dopamine (DOP). Data analysis after HPLC revealed coefficients of variation of 3.3% for NE, 4.6% for EPI, and 5.0% for DOP.

Perceptual Measures. A previously validated brief environmental symptoms questionnaire (ESQ) was used for perceptual heat stress and hypohydration symptom reporting. The ESQ was administered via laptop computer at the specified time points labeled in Figure 1. The thermal sensation scale was used to quantify perception of the surrounding environment. This scale includes whole and half numbers and verbal cues ranging from 0.0 (unbearably cold) to 9.0 (unbearably hot). The ESQ and thermal sensation scales were used to quantify subjective thermal strain throughout.

Statistical Analysis. We completed an a priori power analysis to establish an estimate of power and to project adequate sample size. Using data from previous studies,^{18,19,21,22} common dependent variables were considered: T_{rec} and plasma [NE] during EXDE and REHY. A 2-sided test with $\alpha = .05$ and desired power level of .80 was completed for both variables and the

Table 1. Body Mass Measurements During Experimental Protocol (Mean ± SD)

Trial	Pretrial Body Mass, kg	Post-EXDE Body Mass, kg	REHY Fluid Consumed, L	Post-REHY Body Mass, kg	Post-REHY Body Mass Loss, %
No fluid	81.2 ± 3.9	77.8 ± 3.9	0 ^a	77.6 ± 3.9 ^a	4.5 ± 1.2 ^b
Ad libitum	81.4 ± 3.8	77.9 ± 4.2	1.8 ± 0.5	79.5 ± 3.9	2.3 ± 1.3
Oral	81.8 ± 3.7	78.0 ± 4.0	1.8 ± 0.4	79.6 ± 3.9	2.7 ± 0.7
Intravenous	81.3 ± 3.9	77.9 ± 4.1	2.1 ± 0.5	79.8 ± 4.1	1.9 ± 1.0
Intravenous + oral	80.9 ± 4.0	77.5 ± 4.3	2.1 ± 0.4	79.5 ± 4.3	1.8 ± 0.9

Abbreviations: EXDE, exercise dehydration; REHY, rehydration.

^a No fluid significantly less than all other trials ($P < .001$).

^b No fluid significantly greater than all other trials ($P < .001$).

estimated adequate sample size was 9 or 10 participants. Given the findings of these 2 estimates, we recruited 12 participants to adequately power this study.

We used 5 × 5 (trial × time) repeated-measures analysis of variance to compare differences among trials and over time. A *t* test post hoc comparison with appropriate Bonferroni correction was conducted to detect differences within or among trials when trial × time interactions were present. Select variables were compared using the Pearson *r* correlation as well. The .05 level of significance was selected for identified differences. Data are presented as mean ± SD, unless otherwise noted.

RESULTS

Environmental conditions were not different among trials ($P > .05$). Nutritional analysis revealed no differences among trials in kilocalorie, fat, carbohydrate, protein, Na⁺, or K⁺ intake ($P > .05$). Exercise $\dot{V}O_2$ percentages for EXDE

treadmill walking and ergometer cycling were not different among trials ($P > .05$).

Trial body mass measurements are presented in Table 1. Participants lost 4.32% ± 0.22% of their body mass through fluid restriction and EXDE and were rehydrated to -2.13% ± 0.47% body mass in the controlled OR, IV, and IV + OR trials. After the 60-minute REHY period, the NF trial hypohydration level remained constant (4.5% ± 1.2%). The NF group consumed less fluid than in all other trials ($P > .05$), but there were no statistical differences among other REHY trials ($P < .05$). Sweat rates were not different among trials for EXDE and averaged 0.96 ± 0.18 L·h⁻¹ ($P = .699$). Rehydration sweat rates averaged 0.27 ± 0.12 L·h⁻¹ and were not different among trials ($P = .675$). Urine color (3 ± 1 to 5 ± 1), USG (1.012 ± .006 to 1.024 ± .004), and urine osmolality (405 ± 200 to 889 ± 116 mOsm·L⁻¹) increased from baseline to pre-EXDE ($P < .001$).




Figure 2. Rectal temperature responses during rehydration. ^a No fluid was greater than intravenous ($P = .02$). ^b No fluid was greater than oral, intravenous, and intravenous + oral fluids ($P \leq .009$) but not ad libitum ($P = .07$). Overall, independent of trial, rectal temperature was less than after exercise dehydration at all subsequent time points ($P < .001$).




Figure 3. Mean weighted skin temperature responses during rehydration. ^a Ad libitum was less than intravenous + oral fluids ($P = .02$).

Body Temperature Measures

Participants' T_{rec} was not different among trials during EXDE ($P > .05$). Participants' T_{rec} responses during REHY are shown in Figure 2. Data for T_{msk} during REHY are shown in Figure 3. Correlative data for T_{msk} and T_{rec} throughout REHY are presented in Table 2.

Physiologic Measures

Plasma [CORT] responses during experimental trials are shown in Figure 4. Plasma [NE] demonstrated an increase over time, with post-EXDE greater than at all other time points ($P \leq .037$). All REHY time points demonstrated elevated plasma [NE] compared with baseline and pre-EXDE for all trials ($P \leq .018$). Plasma [EPI] and [DOP] did not show time or trial differences during our protocol ($P \geq .083$).

Perceptual Measures

The ESQ score responses during the experimental protocol are shown in Figure 5. Thermal sensation

demonstrated an increase, independent of trial, from pre- to post-EXDE ($P = .011$). During REHY, thermal sensation decreased over time, independent of trial ($P < .05$). Thermal sensation at 5 minutes after beginning REHY (REHY5) was greater than at REHY30 through REHY60, and at REHY15 was greater than at REHY60 ($P < .05$). No significant between-trials differences of thermal sensation during REHY were seen ($P \geq .324$).

DISCUSSION

We are the first to compare common REHY modes (AL, OR, IV, and IV + OR) and the first to focus on how these modes might be applied to acute medical recovery from EXDE and heat stress. Our participants lost more than 4% body mass and were partially rehydrated to -2% body mass loss within 30 minutes; both values represent common scenarios in competitive athletic settings. The most important finding from our study is that supplementing IV with OR REHY during recovery from thermal strain and EXDE gave a slight recovery advantage compared with IV or OR fluid alone.

A recent examination of previous IV versus OR studies concluded that neither mode offered greater benefits compared with the other.⁴ Oral REHY provides oropharyngeal stimulation, important for perceptual and some physiologic responses.²⁸⁻³⁰ Bypassing this reflex with IV REHY, however, eliminates the potential disadvantage of delayed gastric emptying or intestinal fluid absorption or discomfort.³¹⁻³³ Direct comparisons are obviously contingent upon OR fluid tolerance, which can be impaired during significant dehydration.³ These instances require the use of IV fluids, which rapidly increase plasma volume without gastric filling.^{21,23} Although the suggestion had been

Table 2. Relationship Between Mean Weighted Skin Temperature and Rectal Temperature According to Rehydration Method

Trial	<i>r</i>	<i>P</i>
No fluid	-0.968	.007
Ad libitum	-0.886	.046
Oral	-0.955	.011
Intravenous	-0.974	.005
Oral + intravenous	-0.945	.015
Overall	-0.870	<.001




Figure 4. Plasma cortisol concentration responses during experimental protocol. Ad libitum demonstrated increased plasma cortisol concentration compared with intravenous + oral fluids, independent of time ($P = .02$). ^a After-exercise dehydration was greater than baseline ($P = .001$), before-exercise dehydration ($P = .001$), and 60-min rehydration ($P = .03$). ^b The 30-min rehydration was greater than 60-min rehydration ($P = .048$).

presented, REHY approaches including a combination trial of IV + OR fluid had not been directly compared. We are the first to directly research this comparison and provide evidence for any potential, added benefit of IV + OR hydration.

Our study also helped evaluate the efficacy of AL in a comprehensive comparison among different REHY methods. Recently, due to an increase in exertional hyponatremia incidence, some experts³⁴ recommended that athletes consume fluids according to their level of thirst, or AL




Figure 5. Environmental symptoms questionnaire responses during experimental protocol. ^a All trials except intravenous + oral fluids ($P = .22$) demonstrated greater scores at 60-min rehydration compared with before-exercise dehydration ($P < .05$). ^b Intravenous + oral fluids were less than no fluid ($P = .01$).

instead of attempting to replace estimated sweat losses. Ad libitum fluid consumption most often prevents the overdrinking or fluid intoxication that athletes sometimes experience based on abnormal hydration strategies. However, athletes attempting to maximize performance during exercise should not consider dehydration advantageous.^{1,2,5,10,12} Previous research^{28,31} comparing metered and AL fluid consumption identified the popular phenomenon of “voluntary dehydration.” However, no investigators have compared physiologic recovery from EXDE with other REHY modes. Our results suggest that AL REHY may not effectively facilitate recovery to homeostasis after EXDE.

Body Temperature Measures

As expected, NF during recovery from EXDE demonstrated a physiologic thermoregulatory disadvantage. This finding provides further evidence to refute arguments that hypohydration does not affect body temperature.^{35,36} A novel finding was that AL was not statistically better at reducing thermoregulatory strain than NF, although clinical relevance could be questioned here. This is remarkable, however, considering that the amount of fluid consumed during AL was not different from that consumed during OR, which did reduce T_{rec} more than NF. Consistent with previous researchers, we found minimal differences in T_{rec} among OR, IV, and IV + OR throughout REHY. Previous authors^{18,19,21,23} identified similar marginal and relatively short-term differences among REHY methods. It seems that OR, AL, IV, and IV + OR REHY offer clinically equivalent advantages in facilitating thermoregulatory recovery after EXDE compared with NF.

The most efficient way for humans to dissipate heat is through sweat evaporation. This mechanism is maximized with increased skin blood flow, which transfers internal heat to the periphery for removal. Our results suggest a threshold-initiated oropharyngeal reflex inhibiting skin temperature increase. That is, we identified a graded response in decreased skin temperature with increased OR fluid consumption in acute REHY. At REHY5, IV + OR T_{msk} increased in comparison with post-EXDE, whereas OR and AL decreased T_{msk} temporarily. More OR fluid consumed initially during REHY was matched with a decreased skin temperature (Figure 3). Participants consumed the most fluid in the first 10 minutes of REHY during AL and exhibited an obvious, but slight, decrease in skin temperature. The OR trial showed a skin temperature decrease but to a lesser extent than AL, and this trial included less acute oropharyngeal stimulation. Our IV + OR trial showed a surprising increase in T_{msk} despite OR fluid consumption. This demonstrates a potential threshold for oropharyngeal blunting of the skin temperature increase during recovery. A potential explanation for this response is a limited blood volume after EXDE and an immediate perfusion of gastric blood vessels in order to facilitate fluid absorption.^{9,13,17} The IV and IV + OR trials demonstrate a mechanism of direct increases in total blood volume, bypassing and potentially overcoming the oropharyngeal reflex.

The T_{msk} responses demonstrated an inverse relationship with T_{rec} . This correlation held constant regardless of the

acute response of skin temperature, increase or decrease. This finding is consistent with previous results^{9,13,17} on thermoregulation and heat dissipation in similar environments. Heat dissipation is effectively facilitated via blood flow directed peripherally, increased peripheral temperatures, and sweat evaporation. Increased gastric filling during recovery seems to blunt this response when overall blood volume is limited.

Physiologic Measures

As a means of quantifying the amount of stress our participants underwent with our EXDE protocol, we measured stress hormone responses. We also used stress hormones to quantify the effect of REHY mode on stress recovery. Previous authors^{18,19} have found that NE and EPI show similar increases in response to sympathetic stress. Varying responses were identified with previous IV versus OR REHY comparisons.⁴ Catecholamine responses demonstrated similarities among trials, regardless of REHY method. This observation is similar to previous reports and summaries on IV versus OR REHY after EXDE.^{4,18,19,23} Significant stress was induced by our protocol; however, recovery was equivalent among modes of REHY.

Plasma [CORT] was increased post-EXDE, indicating that participants were under stress from the heat and EXDE. During REHY, IV + OR reduced plasma [CORT] by a greater amount than AL. This indicates a potential advantage in overall stress reduction for IV + OR above and beyond that of AL fluid consumption. The differences we identified were slight but may indicate future areas for study. A potential explanation for our results is that participants during AL consumed more fluid than in other trials within the first 10 minutes of REHY. Our data suggest this may actually increase sympathetic stress for a brief period of time before the return to homeostasis. Oral REHY did not demonstrate this increase, but the amount of fluid provided was metered and equal every 5 minutes. Previous researchers^{18,19,21} comparing IV versus OR REHY after EXDE showed no differences in plasma [CORT].

Perceptual Measures

Our participants demonstrated an increased ESQ score during EXDE. We found a difference, with IV + OR being less than NF at the end of REHY. This demonstrates accentuated perceptual recovery from EXDE with multiple REHY methods. No other trial offered a decreased ESQ score compared with NF. Concomitant REHY with IV + OR limited heat stress and dehydration symptoms better than other means of REHY.

Thermal sensation during EXDE increased and then decreased for all trials during REHY. Participants were no longer creating metabolic heat with muscular contractions during REHY and were rehydrated during 4 of the 5 trials, contributing to decreased thermal sensation. Previous authors²⁴ did identify minor thermal sensation differences, with IV being greater than OR during subsequent exercise. However, others²¹ identified no differences between REHY modes. Interestingly, our lowest value was seen with IV + OR, followed by IV, although this was not significant. This suggests potential clinical relevance, as participants felt better with rapid IV + OR versus IV or OR alone.

Practical Implications

Body temperature recovery after EXDE is important for restoring homeostasis so that body systems can function normally. Rapid REHY using IV offers slight advantages over other modes, but these are short-term benefits. The benefits of REHY on the thermoregulatory system are seen with AL, OR, IV, and IV + OR REHY after EXDE, allowing the body to restore body temperature within minutes. The amount of stress on the body after EXDE seems to be minimized favorably by IV + OR beyond that of other methods. Interestingly, environmental symptoms are also attenuated with this method of REHY. Clinically, this study demonstrated a potential interplay of mechanisms contributing to recovery from EXDE, facilitated by oropharyngeal reflexes, coupled with rapid plasma volume expansion above and beyond that of a solitary mode of REHY.

CONCLUSIONS

The importance of REHY in acute recovery from EXDE should not be downplayed, as the potential benefits are profound and numerous. The mode of REHY during acute recovery had a limited effect. The few differences among methods of REHY were short lived and inadequate to alter recommended care after EXDE for athletes. However, if IV fluids are used for EXDE, OR REHY should be used to potentially accentuate benefits and more rapidly replace fluid deficits. Future research should be conducted to determine if the recovery benefits of IV and IV + OR offer ergogenic advantages compared with OR REHY.

REFERENCES

1. Casa DJ, Armstrong LE, Hillman SK, et al. National Athletic Trainers' Association position statement: fluid replacement for athletes. *J Athl Train*. 2000;35(2):212–224.
2. American College of Sports Medicine, Sawka MN, Burke LM, et al. American College of Sports Medicine position stand: exercise and fluid replacement. *Med Sci Sports Exerc*. 2007;39(2):377–390.
3. Pyne S. Intravenous fluids post marathon: when and why? *Sports Med*. 2007;37(4–5):434–436.
4. Casa DJ, Ganio MS, Lopez RM, McDermott BP, Armstrong LE, Maresh CM. Intravenous versus oral rehydration: physiological, performance and legal considerations. *Curr Sports Med Rep*. 2008; 7(4):S41–S49.
5. Armstrong LE, Maresh CM, Gabaree CV, et al. Thermal and circulatory responses during exercise: effects of hypohydration, dehydration, and water intake. *J Appl Physiol*. 1997;82(6):2028–2035.
6. Candas V, Libert JP, Brandenberger G, Sagot JC, Amoros C, Kahn JM. Hydration during exercise: effects on thermal and cardiovascular adjustment. *Eur J Appl Physiol*. 1986;55(2):113–122.
7. Claremont AD, Costill DL, Fink W, Van Handel P. Heat tolerance following diuretic induced dehydration. *Med Sci Sports Exerc*. 1976; 8(4):239–243.
8. Ebert TR, Martin DT, Bullock N, et al. Influence of hydration status on thermoregulation and cycling hill climbing. *Med Sci Sports Exerc*. 2007;39(2):323–329.
9. Fortney SM, Wenger CB, Bove JR, Nadel ER. Effect of blood volume on forearm venous volume and cardiac stroke volume during exercise. *J Appl Physiol*. 1983;55(3):884–890.
10. Montain SJ, Coyle EF. Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *J Appl Physiol*. 1992;73(4):1340–1350.

11. Nadel ER, Fortney SM, Wenger CB. Effect of hydration state on circulatory and thermal regulations. *J Appl Physiol*. 1980;49(4):715–721.
12. Sawka MN, Coyle EF. Influence of body water and blood volume on thermoregulation and exercise performance in the heat. *Exerc Sport Sci Rev*. 1999;27:167–218.
13. Sawka MN, Gonzalez RR, Young AJ, Dennis RC, Valeri CR, Pandolf KB. Control of thermoregulatory sweating during exercise in the heat. *Am J Physiol*. 1989;257(2 pt 2):R311–R316.
14. Sawka MN, Young AJ, Francesconi RP, Muza SR, Pandolf KB. Thermoregulatory and blood responses during exercise at graded hypohydration levels. *J Appl Physiol*. 1985;59(5):1394–1401.
15. American College of Sports Medicine, Armstrong LE, Casa DJ, et al. American College of Sports Medicine position stand: exertional heat illness during training and competition. *Med Sci Sports Exerc*. 2007; 39(3):556–572.
16. Fortney SM, Nadel ER, Wenger CB, Bove JR. Effect of blood volume on sweating rate and body fluids in exercising humans. *J Appl Physiol*. 1981;51(6):1594–1600.
17. Nose H, Takamata A. Integrative regulations of body temperature and body fluid in humans exercising in a hot environment. *Int J Biometeorol*. 1997;40(1):42–49.
18. Casa DJ, Maresh CM, Armstrong LE, et al. Intravenous versus oral rehydration during a brief period: stress hormone responses to subsequent exhaustive exercise in the heat. *Int J Sport Nutr Exerc Metab*. 2000;10(4):361–374.
19. Castellani JW, Maresh CM, Armstrong LE, et al. Endocrine responses during exercise-heat stress: effects of prior isotonic and hypotonic intravenous rehydration. *Eur J Appl Physiol*. 1998;77(3): 242–248.
20. Costill DL, Sparks KE. Rapid fluid replacement following thermal dehydration. *J Appl Physiol*. 1973;34(3):299–303.
21. Kenefick RW, O'Moore KM, Mahood NV, Castellani JW. Rapid IV versus oral rehydration: responses to subsequent exercise heat stress. *Med Sci Sports Exerc*. 2006;38(12):2125–2131.
22. Maresh CM, Whittlesey MJ, Armstrong LE, et al. Effect of hydration state on testosterone and cortisol responses to training-intensity exercise in collegiate runners. *Int J Sports Med*. 2006;27(10):765–770.
23. Casa DJ, Maresh CM, Armstrong LE, et al. Intravenous versus oral rehydration during a brief period: responses to subsequent exercise in the heat. *Med Sci Sports Exerc*. 2000;32(1):124–133.
24. Maresh CM, Gabaree-Boulant CL, Armstrong LE, et al. Effect of hydration status on thirst, drinking, and related hormonal responses during low-intensity exercise in the heat. *J Appl Physiol*. 2004;97(1):39–44.
25. Jackson AS, Pollock ML. Generalized equations for predicting body density of men. *Br J Nutr*. 1978;40(3):497–504.
26. Baker LB, Lang JA, Kenney WL. Change in body mass accurately and reliably predicts change in body water after endurance exercise. *Eur J Appl Physiol*. 2009;105(6):959–967.
27. Ramanathan NL. A new weighting system for mean surface temperature of the human body. *J Appl Physiol*. 1964;19(3):531–533.
28. Greenleaf JE, Sargent F II. Voluntary dehydration in man. *J Appl Physiol*. 1971;20(4):719–724.
29. Maresh CM, Herrera-Soto A, Armstrong LE, et al. Perceptual responses in the heat after brief intravenous versus oral rehydration. *Med Sci Sports Exerc*. 2001;33(6):1039–1045.
30. Maughan RJ, Leiper JB, Shirreffs SM. Factors influencing the restoration of fluid and electrolyte balance after exercise in the heat. *Br J Sports Med*. 1997;31(3):175–182.
31. Hubbard RW, Sandick BL, Matthew WT, et al. Voluntary dehydration and allesthesia for water. *J Appl Physiol*. 1984;57(3): 868–873.
32. Mitchell JB, Grandjean PW, Pizza FX, Starling RD, Holtz RW. The effect of volume on rehydration and gastric emptying following exercise-induced dehydration. *Med Sci Sports Exerc*. 1994;26(9): 1135–1143.

33. Shirreffs SM, Taylor AJ, Leiper JB, Maughan RJ. Post-exercise rehydration in man: effects of volume consumed and drink sodium content. *Med Sci Sports Exerc.* 1996;28(10):1260–1271.
34. Noakes TD. Hydration in the marathon: using thirst to gauge safe fluid replacement. *Sports Med.* 2007;37(4–5):463–466.
35. Godek SF, Bartolozzi AR, Godek JJ. Sweat rate and fluid turnover in American football players compared with runners in a hot and humid environment. *Br J Sports Med.* 2005;39(4):205–211.
36. Laursen PB, Suriano R, Quod MJ, et al. Core temperature and hydration status during an Ironman triathlon. *Br J Sports Med.* 2006;40(4):320–325.

Address correspondence to Brendon P. McDermott, PhD, ATC, University of Arkansas, 155 Stadium Drive, HPER 326B, Fayetteville, AR 72703. Address e-mail to brendonpmcdermott7@gmail.com.