

Published in final edited form as:

Infect Genet Evol. 2013 January ; 13: . doi:10.1016/j.meegid.2012.08.024.

Analysis of salivary gland transcripts of the sand fly *Lutzomyia ayacuchensis*, a vector of Andean-type cutaneous leishmaniasis

Hiroto Kato^{a,*}, Ryan C. Jochim^b, Eduardo A. Gomez^c, Hiroshi Uezato^d, Tatsuyuki Mimori^e, Masataka Korenaga^f, Tatsuya Sakurai^a, Ken Katakura^a, Jesus G. Valenzuela^b, and Yoshihisa Hashiguchi^{f,g,h}

^aLaboratory of Parasitology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Japan

^bVector Molecular Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, NIH, USA

^cPrograma de Control de Leishmaniasis, Servicio Nacional de Erradicacion de la Malaria (SNEM), Ministerio de Salud Publica, Ecuador

^dDepartment of Dermatology, Faculty of Medicine, University of the Ryukyus, Okinawa, Japan

^eDepartment of Microbiology, School of Health Sciences, Kumamoto University, Kumamoto, Japan

^fDepartment of Parasitology, Kochi Medical School, Kochi University, Kochi, Japan

^gPrometeo, Secretaria Nacional de Educacion Superior, Ciencia, Tecnologia e Innovacion (SENESCYT), Ecuador

^hUniversidad Central del Ecuador Centro de Biomedicina, Quito, Ecuador

Abstract

The saliva of blood sucking insects contains potent pharmacologically active components that assist them in counteracting the host hemostatic and inflammatory systems during blood feeding. In addition, sand fly salivary proteins affect host immunity and have the potential to be a vaccine against *Leishmania* infection. In the present study, the salivary gland transcripts of *Lutzomyia ayacuchensis*, a vector of cutaneous leishmaniasis in Ecuadorian and Peruvian Andes, were analyzed by sequencing randomly selected clones of the salivary gland cDNA library of this sand fly. This resulted in the identification of the most abundant transcripts coding for secreted proteins. These proteins were homologous to the salivary molecules present in other sand flies including the RGD-containing peptide, PpSP15/SL1 family protein, yellow-related protein, putative apyrase, antigen 5-related protein, D7 family protein, and 27 kDa salivary protein. Of note, homologues of maxadilan, an active vasodilator abundantly present in saliva of *Lutzomyia longipalpis*, were not identified. This analysis is the first description of salivary proteins from a sand fly of the subgenus *Helcoerytomyia* and from vector of cutaneous leishmaniasis in the New World. The present analysis will provide further insights into the evolution of salivary components in blood sucking arthropods.

© 2012 Elsevier B.V. All rights reserved.

* Corresponding author. Address: Laboratory of Parasitology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, North 18 West 9, Kita-ku, Sapporo, Hokkaido 060-0818, Japan. Tel./fax: +81 11 706 5196. hkato@vetmed.hokudai.ac.jp (H. Kato).

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version, at <http://dx.doi.org/10.1016/j.meegid.2012.08.024>

Keywords

Lutzomyia ayacuchensis; Salivary gland; Transcript; Bioinformatics; cDNA library

1. Introduction

Phlebotomine sand flies are hematophagous insects of the family Psychodidae in the order Diptera, and approximately 80 species can transmit *Leishmania* protozoa, the causative agents of leishmaniasis (Munstermann, 2005; Kato et al., 2010a). During blood-feeding, sand flies inject saliva containing various physiologically active substances including anticoagulants, vasodilators, and inhibitors of platelet aggregation to facilitate successful acquisition of a blood meal (Ribeiro, 1995; Ribeiro and Francischetti, 2003; Valenzuela, 2005). The infection of mammalian hosts with *Leishmania* protozoa occurs during the blood feeding of an infected female sand fly where the parasites are co-inoculated into the hosts together with sand fly saliva (Rohoušová and Volf, 2006; Oliveira et al., 2009). Sand fly saliva was shown to facilitate the transmission of *Leishmania* to mammalian hosts and to exacerbate the disease (Titus and Ribeiro, 1988; Theodos et al., 1991; Lima and Titus, 1996). Alternatively, cellular immune response to the salivary components protects the host from *Leishmania* infection (Belkaid et al., 1998, 2000; Kamhawi et al., 2000; Valenzuela et al., 2001; Gomes et al., 2008; Oliveira et al., 2008; Tavares et al., 2011). Humoral immunity to sand fly salivary components was shown in the case of *Leishmania (Viannia) braziliensis* to exacerbate infection (Oliveira et al., 2008). In addition, recent studies are showing that sand fly salivary proteins can be used as a specific marker of sand fly exposure and to use them to evaluate potential risks of *Leishmania* transmission (Teixeira et al., 2010; Vlkova et al., 2011). The profile of salivary components has been defined in 8 Old World *Phlebotomus* species; *Phlebotomus (P.) papatasi* [a vector of *Leishmania (Leishmania) major*] (Valenzuela et al., 2001), *P. ariasi* [a vector of *L. (L.) infantum*] (Oliveira et al., 2006), *P. perniciosus* [a vector of *L. (L.) infantum*] (Anderson et al., 2006), *P. argentipes* [a vector of *L. (L.) donovani*] (Anderson et al., 2006), *P. duboscqi* [a vector of *L. (L.) major*] (Kato et al., 2006), *P. arabicus* [a vector of *L. (L.) tropica*] (Hostomská et al., 2009), *P. tobbi* [a vector of *L. (L.) infantum*] (Rohoušová et al., 2012), and *P. sergenti* [a vector of *L. (L.) tropica*] (Rohoušová et al., 2012). On the other hand, the salivary components of the New World *Lutzomyia* species have been analyzed only in *Lutzomyia (Lu.) longipalpis*, a vector of *L. (L.) infantum* [a synonym of *L. (L.) chagasi*] causing visceral leishmaniasis (Charlab et al., 1999; Valenzuela et al., 2004). To date, no information is available on the repertoire of salivary transcripts from vectors of the New World cutaneous leishmaniasis. In the present study, to obtain further insight into the salivary components of *Lutzomyia* species, the salivary gland transcriptome analysis was performed on *Lu. ayacuchensis*, which is a proven vector of *L. (L.) mexicana* (Takaoka et al., 1990; Kato et al., 2005, 2008) and *L. (V.) peruviana* (Caceres et al., 2004; Kato et al., 2008), causative agents of cutaneous leishmaniasis in Andean areas of Ecuador and Peru, respectively. This is the first report of salivary proteins from a sand fly of the subgenus *Helcocyrtomyia* and from vector of cutaneous leishmaniasis in the New World.

2. Materials and methods

2.1. *Lutzomyia ayacuchensis* salivary glands

Sand flies were captured using protected human bait at Huigra (2°20'S, 78°58'W; 1200 m above sea level) in the Department of Chimborazo, Ecuador, where Andean-type cutaneous leishmaniasis caused by *L. (L.) mexicana* is endemic. The sand flies were dissected and identified at the species level based mainly on the morphology of their spermathecae (Young

and Duncan, 1994). Salivary glands of *Lu. ayacuchensis* were dissected and transferred to tubes containing RNAlater (Ambion, Austin, TX) and kept at -20°C .

2.2. Construction of salivary gland cDNA library

Lu. ayacuchensis salivary gland mRNA was isolated from 38 pairs of salivary glands using the Micro FastTrack mRNA isolation kit (Invitrogen, San Diego, CA). The PCR-based cDNA library was prepared following the instructions for the SMART cDNA library construction kit (BD-Clontech, Palo Alto, CA) with some modifications (Valenzuela et al., 2004). The quality of the cDNA was checked by agarose gel electrophoresis and the absence of smaller fragments derived from degraded mRNA was confirmed. The obtained cDNA library was fractionated using a Chromaspin 1000 column (BD-Clontech) into small (approximately 400–800 bp), medium (approximately 800–1200 bp) and large (>1200 bp) transcripts based on their electrophoresis profiles on a 1.1% agarose gel. Pooled fractions were ligated into Lambda TriplEx2 vector (BD-Clontech) and packaged into lambda phage (Stratagene, La Jolla, CA).

2.3. Sequence analysis of cDNA library

Single isolated plaques were picked from the plate using sterile wooden sticks and placed into 50 μl of water. Amplification of cDNA was performed in a volume of 15 μl using a pair of primers, PT2F1 (5'-AAG TAC TCT AGC AAT TGT GAG C-3') and PT2R1 (5'-CTC TTC GCT ATT ACG CCA GCT G-3'), Premix Taq (Takara Bio, Shiga, Japan) and 3 μl of template DNA. After an initial denaturation at 75°C for 3 min and following 95°C for 4 min, PCR amplification was performed with 35 cycles of denaturation (95°C , 1 min), annealing (50°C , 1 min) and polymerization (72°C , 2 min). PCR products were cleaned using Multiscreen PCR cleaning plates (Millipore Corporation, Bedford, MA) and used as templates for cycle-sequencing kit (Applied Biosystems, Foster City, CA) with PT2F3 primer (5'-TCT CGG GAA GCG CGC CAT TGT-3'). Cycle-sequencing products were cleaned using sephadex and MultiScreen HV plates (Millipore Corporation), dried and stored at -20°C . Sequencing was performed on an ABI 3730xl DNA sequencer (Applied Biosystems).

2.4. Bioinformatics

Expressed sequence tags (ESTs) were trimmed of primer and vector sequences and clustered. The ESTs were grouped based on nucleotide homology of 95% identity over 100 residues using the BLASTn algorithm (Altschul et al., 1997). The assembly of the ESTs into transcript contigs was done using the CAP3 algorithm, generating a consensus sequence (Huang, 1992). Contigs and singletons (contig containing only one sequence) were compared using BLASTx or BLASTn (Altschul et al., 1997) of the non-redundant (NR) protein database of the National Center of Biological Information (NCBI), the gene ontology database (GO) (Ashburner et al., 2000), and the Conserved Domains Database (CDD) that includes all Pfam (Bateman and Birney, 2000), SMART (Schultz et al., 1998) and COG protein domains in the NCBI (Marchler-Bauer et al., 2002). Additionally, contigs were compared using BLASTn (Altschul et al., 1997) to custom databases of mitochondrial (mit-pla) and rRNA (rrna) nucleotide sequences. Identification of putative secreted proteins was conducted using SignalP server (Bendtsen et al., 2004).

2.5. T cell epitope prediction

For the T cell epitope prediction, ProPred MHC Class-II Binding Peptide Prediction Server (Singh and Raghava, 2001) was utilized. HLA class II-binding peptides were searched on the 51 different HLA-DR alleles, and the promiscuous epitopes were selected from the *Lu. ayacuchensis* salivary protein sequences tested that were predicted to bind at least 20 alleles.

2.6. Phylogenetic analysis

The sequences that had homologies with secreted proteins by BLASTx analyses were aligned with CLUSTAL W software (Thompson et al., 1994) and examined using the program MEGA (Molecular Evolutionary Genetics Analysis) version 4.0 (Tamura et al., 2007). Phylogenetic trees by the neighbor-joining method were constructed with the distance algorithms available in the MEGA package. Bootstrap values were determined on 1000 replicates of the data sets.

3. Results and discussion

3.1. Sequencing of *Lutzomyia ayacuchensis* salivary gland cDNA library

Lu. ayacuchensis salivary gland cDNA library was constructed, and sequencing was performed on randomly selected 1152 clones. As a result, 768 high-quality sequences were obtained. Three categories of expressed genes were derived from the manual annotation of the contigs: secreted, housekeeping and unknown. The putative secreted category comprised 45.7% of the clusters and 74.6% of the total sequences. The high ratio of transcripts encoding secreted proteins were also reported in other phlebotomine sand flies (Valenzuela et al., 2001; Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009; Rohoušová et al., 2012). The housekeeping category had 35.0% of the clusters and 16.3% of the total sequences. Finally, the category of “unknowns” comprised 19.3% of the cluster and 9.1% of the sequences.

3.2. Housekeeping genes

The clusters of sequences attributed to housekeeping genes (94 clusters with 125 sequences in total) were further divided into 18 subgroups, according to their possible function (Table 1). The two largest subgroups were associated with “Energy production and conversion” (24 sequences in 20 clusters) and “Translation, ribosomal structure and biogenesis” (23 sequences in 19 clusters). Six sequences in 6 clusters, which represent conserved proteins with unknown function, were classified as “unknown conserved”. Other sequences were identified with homology to housekeeping genes and were associated with transport, metabolism, signal transduction, and cell structure, among other potential activities.

3.3. Putative secreted proteins

The transcripts coding for secreted proteins were further analyzed using the BLASTx program for comparison to the NCBI non-redundant protein database. Table 2 lists clusters for the most abundant transcripts coding for the putative secreted salivary gland proteins from *Lu. ayacuchensis*. The table was arranged from the most abundant to the least abundant transcripts found in the cDNA library. The nomenclature for the transcripts on the cDNA library is the following: Lay = *Lu. ayacuchensis*, S = salivary glands, and the number (i.e. 45) denotes the contig number on the cDNA library where a contig is a cluster of identical transcripts. Many of the isolated transcripts code for proteins previously identified from the saliva of sand flies, *Lu. longipalpis* and *P. perniciosus*, including RGD-containing peptide, 14 kDa salivary protein, antigen 5-related protein, yellow-related protein and SL1-like protein (Table 2). Table 3 shows the classification of transcripts coding for putative secreted proteins in *Lu. ayacuchensis* salivary glands. Of the 573 transcripts associated with putative secreted proteins, the most abundant transcript (18.7%) had homology with that of RGD-containing peptide from salivary glands of *Lu. longipalpis*. The following abundant transcripts were SL1-like protein (14.3%) and 14 kDa salivary protein (12.4%), both of which belong to a family of PpSP15/SL1 originally identified as 15 kDa protein from *P. papatasi* saliva. Additional transcripts coding for secreted proteins include homologs to yellow-related protein (11.1%), putative apyrase (7.9%), D7-related salivary protein (7.3%),

27 kDa salivary protein (6.6%), antigen 5-related protein (6.5%), 16.4 kDa salivary protein (5.6%), and others (Table 3).

3.3.1. RGD-containing peptide—The most abundant sequences found in the *Lu. ayacuchensis* salivary gland transcriptome contain C-terminal RGD (Arg-Gly-Asp) sequence; a motif that has been shown to bind integrins such as $\alpha v\beta 3$ and $\alpha v\beta 5$ (Hynes, 1992). Ten transcripts of RGD-containing peptide (LayS38–LayS47) were identified in the cDNA library of *Lu. ayacuchensis* salivary glands. These *Lu. ayacuchensis* RGD-containing molecules have a predicted molecular mass of 5.3 kDa in the mature form and a calculated isoelectric point of 3.44 (Table 2). The only homologous protein has been identified in the salivary gland transcriptome of *Lu. longipalpis* (LuloRGD), and speculated to act as an inhibitor of platelet aggregation via its RGD motif; although, the function has not yet been characterized (Charlab et al., 1999). Alignment of the RGD-containing peptide showed a high level of homology among contigs, and an RGD-motif in the C-terminal was strictly conserved in all molecules (Fig. S1).

3.3.2. PpSP15/SL1 family—The PpSP15/SL1 family of proteins has been reported only in sand flies and the biological function during blood-feeding remains unknown. Interestingly, immunization of mice with PpSP15 protected against *Leishmania* major infection co-injected with *P. papatasi* saliva, suggesting that this family of protein is a vaccine candidate for *Leishmania* infection (Valenzuela et al., 2001). Two groups of molecules were identified from the *Lu. ayacuchensis* salivary gland cDNA library, SL-1-like proteins and 14 kDa salivary proteins (PpSP14-like protein), as members of the PpSP15/SL1 family. Twenty-five contigs (LayS48–LayS72) showed the highest homology with SL1 protein from *Lu. longipalpis*, and six contigs (LayS32–LayS37) coding for a 14 kDa salivary protein were more homologous to the 14.6 kDa salivary protein from *P. perniciosus* (PpeSP09) (Table 2). Alignment of PpSP15/SL1 family showed a high level of homology among SL-1-like proteins and among 14 kDa salivary proteins (Figs. S2 and S3). When both groups of proteins from *Lu. ayacuchensis* were aligned with PpSP15/SL1 family proteins from other sand flies, only six cysteine residues and three other amino acids (P, F, and A) were conserved in the amino acid sequence of mature proteins (Fig. 1A), reflecting the divergence among PpSP15/SL1 family proteins in sand flies (Anderson et al., 2006). Since the proteins of this family are immunogenic and potential vaccines for *Leishmania* infection (Valenzuela et al., 2001), T cell epitopes were analyzed using ProPred MHC Class-II Binding Peptide Prediction Server. We identified two potential T cell epitopes, YRITKKHIE and IIHYRRCVV, in *Lu. ayacuchensis* SL-1-like proteins (Fig. S2), and two potential epitopes, IKNKAVDGS and FKYESAINSY, in 14 kDa salivary proteins (Fig. S3). A phylogenetic analysis of sand fly PpSP15/SL1 family proteins revealed that SL-1-like proteins and 14 kDa salivary proteins from *Lu. ayacuchensis* constructed separate clades (Fig. 1B).

3.3.3. Yellow-related protein—Yellow-related proteins are abundantly expressed in salivary glands of sand flies (Valenzuela et al., 2001; Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009; Rohoušová et al., 2012). The proteins of this family are immunogenic and host antibody responses to this protein can be a potential marker for sand fly exposure (Teixeira et al., 2010; Vlkova et al., 2011). Interestingly, prior DNA vaccination of mice with PpSP44 coding for a 44 kDa yellow-related protein from *P. papatasi* saliva primed strong humoral immunity and exacerbated subsequent *L. major* infection in the presence of sand fly saliva (Oliveira et al., 2008). In contrast, a yellow-related protein from *Lu. longipalpis* saliva, LJM11, conferred protective cellular immunity in mice against *L. major* infection plus sand fly saliva. Other *Lu. longipalpis* yellow-related proteins, LJM111 and LJM17, were not protective, suggesting that structural features are a

determinant for the host immunity against yellow-related proteins (Xu et al., 2011). Regarding the biological function, yellow-related proteins, LJM11, LJM111 and LJM17, were shown to act as high affinity binders of proinflammatory biogenic amines such as serotonin, catecholamines and histamine, suggesting that the proteins play a role for the reduction of inflammation during sand fly blood-feeding (Xu et al., 2011). In the *Lu. ayacuchensis* cDNA library, five contigs (LayS22–LayS24, LayS117 and LayS118) are putative yellow-related proteins, of which LayS117 was truncated in the 5' region. When these proteins were aligned with salivary yellow-related proteins from other sand fly species, four cysteine residues were conserved in the amino acid sequence of mature proteins (Fig. 2A and S4). In addition, an amino acid motif, T-x(52,63)-Y-Q-x(85,90)-[FY]-x(44,46)-F-x(54)-[IVL]-x(45,46)-[FY]-x-[TS]-D-x(13)-[NT]-x-[QHFL], recently identified in the ligand binding pocket of yellow-related proteins (Xu et al., 2011) were conserved in those from *Lu. ayacuchensis* (Fig. 2A), suggesting that these proteins work as anti-inflammatory agents in *Lu. ayacuchensis* saliva by binding biogenic amines. Two potential N-glycosylation sites were present in yellow-related proteins from *Lu. ayacuchensis*, but the positions of LayS118 were different from those of others (LayS22–LayS24) (Fig. 2A). T cell epitopes were analyzed on the immunogenic proteins of this family, and we identified seven potential epitopes in LayS22–LayS24 (Fig. S4A) and 5 potential epitopes in LayS118 (Fig. S4B). A phylogenetic analysis of sand fly yellow-related proteins revealed that LayS22, LayS23 and LayS24 were closer related to LJM11 whereas LayS118 were closer related to LJM17 from *Lu. longipalpis* saliva (Fig. 2B). The result suggested that effect on the host immunity of LayS118 is different from those of LayS22, LayS23 and LayS24.

3.3.4. Apyrase—Apyrases are nucleoside triphosphate-diphosphohydrolases present in a variety of organisms. In the saliva of blood sucking arthropods, apyrases function to hydrolyze ADP in a Ca²⁺-dependent manner and inhibit ADP-induced platelet aggregation to facilitate blood feeding (Ribeiro and Francischetti, 2003; Faudry et al., 2004). Apyrases identified in sand fly saliva belong to the *Cimex* apyrase family (Valenzuela et al., 2001; Hamasaki et al., 2009). Thirteen contigs (LayS8–LayS14 and LayS16–LayS21) were homologous to putative apyrase from *Lu. longipalpis* (LuloAPY). A multiple sequence alignment analysis together with LuloAPY confirmed the close relationship among putative apyrases of *Lu. ayacuchensis* (Fig. S5).

3.3.5. D7 family—D7 family proteins are abundantly expressed in salivary glands of blood-feeding Diptera such as mosquitoes, sand flies, black flies, and biting midges (James et al., 1991; Valenzuela et al., 2002; Campbell et al., 2005; Andersen et al., 2009). To date, the function of sand fly D7 proteins remains to be elucidated; however, several D7 family proteins have been functionally characterized in mosquitoes. Hamadarin from *Anopheles stephensi* saliva was identified as a blood coagulation inhibitor affecting the activation of the plasma contact system (Isawa et al., 2002). Salivary D7 proteins from *Anopheles gambiae* and *Aedes aegypti* were characterized as binders of biogenic amines such as serotonin, histamine and norepinephrine (Calvo et al., 2006). Nine contigs (LayS95–LayS103) encoding a 26.6 kDa protein were homologous to the 27 kDa D7-related salivary protein from *P. argentipes* (PagSP10) and *P. perniciosus* (PpeSP04b). In a multiple sequence alignment analysis of D7 family proteins from sand flies, 10 cysteine residues were strictly conserved in the amino acid sequence of mature proteins (Fig. 3A and Fig. S6). Alignment and phylogenetic analysis together with PpeSP04b showed that D7 family proteins of *Lu. ayacuchensis* had close relationships (Fig. 3B).

3.3.6. 27 kDa Salivary protein—The protein family is also known as PpSP32-like family. Eleven contigs (LayS83–LayS93) encoding a 27 kDa protein were homologous to the 29.2 kDa salivary protein from *Lu. longipalpis* (Table 2). These transcripts are also

present in the salivary glands from Old World sand flies such as *P. ariasi*, *P. duboscqi*, *P. arabicus*, *P. argentipes*, *P. papatasi*, *P. perniciosus*, *P. tobbi*, and *P. sergenti* (Valenzuela et al., 2001; Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009; Rohoušová et al., 2012). To date, no homologues of these proteins have been reported in organisms other than sand flies and the biological function remains to be clarified. Although no conserved domains were found in the amino acid sequences, these proteins contain glycine-rich sequences in the middle region (amino acids 66–179 in LayS91) (Fig. S7).

3.3.7. Antigen 5-related protein—This family of proteins belongs to the cysteine-rich secretory proteins (CRISPs) and is related to venom allergens in social wasps and ants (Lu et al., 1993; Hoffman, 1993; King and Spangfort, 2000). Transcripts coding for the members of this protein family have been identified in the salivary glands of blood sucking insects such as mosquitoes (Francischetti et al., 2002; Arcà et al., 2007), sand flies (Valenzuela et al., 2001; Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009; Rohoušová et al., 2012) and triatomine bugs (Ribeiro et al., 2004; Santos et al., 2007; Assumpção et al., 2008; Kato et al., 2010b). Although the antigen 5 family of protein has been widely identified in blood-sucking arthropods, the function in their saliva has yet to be determined. Nine contigs (LayS73–LayS81) from *Lu. ayacuchensis* salivary glands were included in this family. A multiple sequence alignment analysis of antigen 5-related proteins with that of *Lu. longipalpis* revealed the close relationship among the proteins of *Lu. ayacuchensis* and strict conservation of 12 cysteine residues in mature forms (Fig. S8).

3.3.8. 16.4 kDa Salivary protein—Thirteen contigs (LayS120–LayS132) coded for 16.4 kDa protein containing a C-type lectin/C-type lectin-like domain (Table 2). This putative domain may function as a Ca²⁺-dependent carbohydrate-binding pocket involved in extracellular matrix organization, pathogen recognition and cell-to-cell interactions (Weis et al., 1998). Homologous proteins with molecular weight of 16.2–16.5 kDa have been identified from *Lu. longipalpis* saliva (Charlab et al., 1999; Valenzuela et al., 2004), and they were similar to the previously described anticoagulant from *Lu. longipalpis* saliva (Charlab et al., 1999). A recombinant protein of mature LayS127 protein was expressed in *Escherichia coli*, and its activity on coagulation pathway was assessed; however, the protein inhibited neither intrinsic nor extrinsic coagulation pathways (data not shown). At present, the biological function of these proteins from *Lu. ayacuchensis* remains to be clarified. In a multiple sequence alignment analysis of *Lu. ayacuchensis* 16.4 kDa proteins together with *Lu. longipalpis* 16.5 kDa protein, three cysteine residues were strictly conserved in the amino acid sequence of mature proteins (Fig. S9).

3.3.9. 11.5 kDa Salivary protein—Four contigs encoding 10.7–11.9 kDa proteins (LayS4–LayS7) were identified in the cDNA library of *Lu. ayacuchensis* salivary glands. The different size of these proteins was mainly resulted from the number of “SSDGSSG” repeat sequences with unknown function (Fig. S10). The only homologous protein has been identified in salivary glands of *Lu. longipalpis* (9 kDa salivary protein) with unknown function. The amino acid sequences of these proteins are rich in serine from the middle to the C-terminal regions (Fig. S10).

3.3.10. Other putative secreted proteins—Three contigs (LayS25–LayS27) coding for a 33.6 kDa protein had homology with a 34 kDa salivary protein from *P. ariasi*. Three contigs (LayS141–LayS143) coded for a 9.3 kDa protein. Three singletons (LayS109–LayS111) coding for a 15.9 kDa protein had homology with a 15.5 kDa salivary protein from *P. argentipes* with unknown function. Two contigs (LayS2 and LayS3) coded for a 25.9 kDa protein homologous to allergen from *Culex quinquefasciatus* belonging to the

sperm-coating protein (SCP) superfamily. Two singletons (LayS167 and LayS168) and one singleton (LayS153) coded for homologous proteins with a 32.4 kDa salivary protein and a 71 kDa salivary protein from *Lu. longipalpis*, respectively, although the sequence of LayS168 was truncated in the 5' region. A singleton (LayS147) coded for homologous protein with the 43.7 kDa salivary protein from *Lu. longipalpis*, a putative endonuclease, but the sequence was truncated in the 5' region. Singletons, LayS106 and LayS215, coded for homologous protein with the 27 kDa salivary protein from *P. ariasi* and trypsin inhibitor like cysteine rich domain containing protein from *Drosophila*, respectively.

4. Conclusions

The present study identified abundant salivary gland transcripts of *Lu. ayacuchensis*, a proven vector of *L. (L.) mexicana* (Takaoka et al., 1990; Kato et al., 2005, 2008) and *L. (V.) peruviana* (Caceres et al., 2004; Kato et al., 2008) causative agents of cutaneous leishmaniasis in Andean areas of Ecuador and Peru, respectively. The relatively divergent transcripts resulting in the generation of a variety of contigs noted for each component probably reflects population of the sand flies used in this study; that is, our cDNA library was constructed from field-captured *Lu. ayacuchensis* while most studies prepared the cDNA libraries from colonized insects. To date, the salivary components of the New World *Lutzomyia* species have been analyzed only in *Lu. longipalpis* (Charlab et al., 1999; Valenzuela et al., 2004), a subgenus *Lutzomyia* species, transmitting *L. (L.) infantum*, a causative agent of visceral leishmaniasis (Young and Duncan, 1994). On the other hand, *Lu. ayacuchensis* is a vector species of cutaneous leishmaniasis and belongs to the subgenus *Helcocyrtomyia* (Young and Duncan, 1994). Therefore, the present analysis is the first description of salivary proteins of a sand fly in the subgenus *Helcocyrtomyia* and a vector of cutaneous leishmaniasis in the New World.

In the present analysis, transcripts homologous to maxadilan, an active vasodilator abundantly present in saliva of *Lu. longipalpis*, were not identified. Maxadilan was originally isolated from saliva of *Lu. longipalpis* (Lerner et al., 1991), and its homologue was reported from *Lu. neivai* (Aires et al., 2005). Conversely, maxadilan-related proteins have never been identified in Old World *Phlebotomus* species (Valenzuela et al., 2001; Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009). *Phlebotomus* sand flies except for *P. duboscqi* lack adenosine deaminase (ADA) that hydrolyze adenosine and adenosine monophosphate (AMP) (Oliveira et al., 2006; Anderson et al., 2006; Kato et al., 2006; Hostomská et al., 2009), but contain large amounts of adenosine and AMP, very active vasodilators, in their saliva (Ribeiro et al., 1999; Katz et al., 2000; Ribeiro and Modi, 2001). On the other hand, *Lu. longipalpis* has ADA (Charlab et al., 1999, 2000; Valenzuela et al., 2004) and lacks adenosine and AMP in the saliva (Ribeiro et al., 1989). Therefore, with an exception, salivary vasodilators of sand flies are considered to be adenosine and AMP in *Phlebotomus* species and maxadilan in *Lutzomyia* species. Unexpectedly, neither ADA nor maxadilan were identified in *Lu. ayacuchensis*, which is similar to *Phlebotomus* species, suggesting that salivary vasodilators of this species may be adenosine and AMP. Further analysis of small molecules in the saliva of *Lu. ayacuchensis* will elucidate this issue.

In conclusion, the most abundant proteins of *Lu. ayacuchensis* saliva were identified in this study. These results will provide further insights into the evolution of salivary components in blood sucking arthropods. In addition, the cDNAs and future recombinant proteins prepared from these transcripts will result in the discovery of vaccine candidates and markers of sand fly exposure as well as novel pharmacologically active compounds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to Dr. José M. C. Ribeiro (Vector Biology Section, Laboratory of Malaria and Vector Research, NIAID, NIH, USA) for the development and training of all custom bioinformatics programs used in this research. This study was supported by the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (Grant Nos. 18256004, 10037385 and 23580424).

References

- Aires JM, Chociay MF, Nascimento MM, Figueiredo JF, Roselino AM. Maxadilan (MAX) - Salivary protein of *Lutzomyia longipalpis*: detection of antibodies anti-MAX in American tegumentar leishmaniasis (ATL), and genetic and protein expression of MAX in *Lutzomyia neivai*. *An. Bras. Dermatol.* 2005; 80:S333–S338.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 1997; 25:3389–4002. [PubMed: 9254694]
- Andersen JF, Pham VM, Meng Z, Champagne DE, Ribeiro JMC. Insight into the sialome of the Black Fly, *Simulium vittatum*. *J. Proteome Res.* 2009; 8:1474–1488. [PubMed: 19166301]
- Anderson JM, Oliveira F, Kamhawi S, Mans BJ, Reynoso D, Seitz AE, Lawyer P, Garfield M, Pham M, Valenzuela JG. Comparative salivary gland transcriptomics of sandfly vectors of visceral leishmaniasis. *BMC Genomics.* 2006; 7:52. [PubMed: 16539713]
- Arcà B, Lombardo F, Francischetti IMB, Pham VM, Mestres-Simon M, Andersen JF, Ribeiro JMC. An insight into the sialome of the adult female mosquito *Aedes albopictus*. *Insect Biochem. Mol Biol.* 2007; 37:107–127. [PubMed: 17244540]
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* 2000; 25:25–29.
- Assumpção TC, Francischetti IMB, Andersen JF, Schwarz A, Santana JM, Ribeiro JMC. An insight into the sialome of the blood-sucking bug *Triatoma infestans*, a vector of Chagas' disease. *Insect Biochem. Mol Biol.* 2008; 38:213–232. [PubMed: 18207082]
- Bateman A, Birney E. Searching databases to find protein domain organization. *Adv. Protein Chem.* 2000; 54:137–157. [PubMed: 10829227]
- Belkaid Y, Kamhawi S, Modi G, Valenzuela J, Noben-Trauth N, Rowton E, Ribeiro J, Sacks DL. Development of a natural model of cutaneous leishmaniasis: powerful effects of vector saliva and saliva preexposure on the long-term outcome of *Leishmania major* infection in the mouse ear dermis. *J. Exp. Med.* 1998; 188:1941–1953. [PubMed: 9815271]
- Belkaid Y, Valenzuela JG, Kamhawi S, Rowton E, Sacks DL, Ribeiro JMC. Delayed-type hypersensitivity to *Phlebotomus papatasi* sand fly bite: an adaptive response induced by the fly? *Proc. Natl. Acad. Sci USA.* 2000; 97:6704–6709. [PubMed: 10841567]
- Bendtsen JD, Nielsen H, von Heijne G, Brunak S. Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* 2004; 340:783–795. [PubMed: 15223320]
- Caceres AG, Villaseca P, Dujardin JC, Bañuls AL, Inga R, Lopez M, Arana M, Le Ray D, Arevalo J. Epidemiology of Andean cutaneous leishmaniasis: incrimination of *Lutzomyia ayacuchensis* (Diptera: psychodidae) as a vector of *Leishmania* in geographically isolated, upland valleys of Peru. *Am. J. Trop. Med. Hyg.* 2004; 70:607–612. [PubMed: 15211000]
- Calvo E, Mans BJ, Andersen JF, Ribeiro JMC. Function and evolution of a mosquito salivary protein family. *J. Biol. Chem.* 2006; 281:1935–1942. [PubMed: 16301315]
- Campbell CL, Vandyke KA, Letchworth GJ, Drolet BS, Hanekamp T, Wilson WC. Midgut and salivary gland transcriptomes of the arbovirus vector *Culicoides sonorensis* (Diptera: Ceratopogonidae). *Insect Mol. Biol.* 2005; 14:121–136. [PubMed: 15796745]

- Charlab R, Valenzuela JG, Rowton ED, Ribeiro JMC. Toward an understanding of the biochemical and pharmacological complexity of the saliva of a hematophagous sand fly *Lutzomyia longipalpis*. Proc. Natl. Acad. Sci. USA. 1999; 96:15155–15160. [PubMed: 10611354]
- Charlab R, Rowton ED, Ribeiro JMC. The salivary adenosine deaminase from the sand fly *Lutzomyia longipalpis*. Exp. Parasitol. 2000; 95:45–53. [PubMed: 10864517]
- Faudry E, Rocha PS, Vernet T, Lozzi SP, Teixeira AR. Kinetics of expression of the salivary apyrases in *Triatoma infestans*. Insect Biochem. Mol. Biol. 2004; 34:1051–1058. [PubMed: 15475299]
- Francischetti IMB, Valenzuela JG, Pham VM, Garfield MK, Ribeiro JMC. Toward a catalog for the transcripts and proteins (sialome) from the salivary gland of the malaria vector *Anopheles gambiae*. J. Exp. Biol. 2002; 205:2429–2451. [PubMed: 12124367]
- Gomes R, Teixeira C, Teixeira MJ, Oliveira F, Menezes MJ, Silva C, de Oliveira CI, Miranda JC, Elnaïem DE, Kamhawi S, Valenzuela JG, Brodskyn CI. Immunity to a salivary protein of a sand fly vector protects against the fatal outcome of visceral leishmaniasis in a hamster model. Proc. Natl. Acad. Sci. USA. 2008; 105:7845–7850. [PubMed: 18509051]
- Hamasaki R, Kato H, Terayama Y, Iwata H, Valenzuela JG. Functional characterization of a salivary apyrase from the sand fly, *Phlebotomus duboscqi*, a vector of *Leishmania major*. J Insect Physiol. 2009; 55:1044–1049. [PubMed: 19651132]
- Hoffman DR. Allergens in Hymenoptera venom. XXV: The amino acid sequences of antigen 5 molecules and the structural basis of antigenic cross-reactivity. J. Allergy Clin. Immunol. 1993; 92:707–716. [PubMed: 8227862]
- Hostomská J, Volfová V, Mu J, Garfield M, Rohousová I, Volf P, Valenzuela JG, Jochim RC. Analysis of salivary transcripts and antigens of the sand fly *Phlebotomus arabicus*. BMC Genomics. 2009; 10:282. [PubMed: 19555500]
- Huang X. A contig assembly program based on sensitive detection of fragment overlaps. Genomics. 1992; 14:18–25. [PubMed: 1427824]
- Hynes RO. Integrins: versatility, modulation, and signaling in cell adhesion. Cell. 1992; 69:11–25. [PubMed: 1555235]
- Isawa H, Yuda M, Orito Y, Chinzei Y. A mosquito salivary protein inhibits activation of the plasma contact system by binding to factor XII and high molecular weight kininogen. J. Biol. Chem. 2002; 277:27651–27658. [PubMed: 12011093]
- James AA, Blackmer K, Marinotti O, Ghosn CR, Racioppi JV. Isolation and characterization of the gene expressing the major salivary gland protein of the female mosquito, *Aedes aegypti*. Mol. Biochem. Parasitol. 1991; 44:245–253. [PubMed: 2052024]
- Kamhawi S, Belkaid Y, Modi G, Rowton E, Sacks D. Protection against cutaneous leishmaniasis resulting from bites of uninfected sand flies. Science. 2000; 290:1351–1354. [PubMed: 11082061]
- Kato H, Uezato H, Katakura K, Calvopiña M, Marco JD, Barroso PA, Gomez EA, Mimori T, Korenaga M, Iwata H, Nonaka S, Hashiguchi Y. Detection and identification of *Leishmania* species within naturally infected sand flies in the andean areas of Ecuador by a polymerase chain reaction. Am. J. Trop. Med. Hyg. 2005; 72:87–93. [PubMed: 15728872]
- Kato H, Anderson JM, Kamhawi S, Oliveira F, Lawyer PG, Pham VM, Sangare CS, Samake S, Sissoko I, Garfield M, Sigutova L, Volf P, Doumbia S, Valenzuela JG. High degree of conservancy among secreted salivary gland proteins from two geographically distant *Phlebotomus duboscqi* sandflies populations (Mali and Kenya). BMC Genomics. 2006; 7:226. [PubMed: 16952314]
- Kato H, Cáceres AG, Gomez EA, Mimori T, Uezato H, Marco JD, Barroso PA, Iwata H, Hashiguchi Y. Molecular mass screening to incriminate sand fly vectors of Andean-type cutaneous leishmaniasis in Ecuador and Peru. Am. J. Trop. Med. Hyg. 2008; 79:719–721. [PubMed: 18981511]
- Kato H, Gomez EA, Cáceres AG, Uezato H, Mimori T, Hashiguchi Y. Molecular epidemiology for vector research on leishmaniasis. Int. J. Environ. Res Public Health. 2010a; 7:814–826. [PubMed: 20617005]
- Kato H, Jochim RC, Gomez EA, Sakoda R, Iwata H, Valenzuela JG, Hashiguchi Y. A repertoire of the dominant transcripts from the salivary glands of the blood-sucking bug, *Triatoma dimidiata*, a vector of Chagas disease. Infect. Genet. Evol. 2010b; 10:184–191. [PubMed: 19900580]

- Katz O, Waitumbi JN, Zer R, Warburg A. Adenosine, AMP, and protein phosphatase activity in sandfly saliva. *Am. J. Trop. Med. Hyg.* 2000; 62:145–150. [PubMed: 10761741]
- King TP, Spangfort MD. Structure and biology of stinging insect venom allergens. *Int. Arch. Allergy Immunol.* 2000; 123:99–106. [PubMed: 11060481]
- Lerner EA, Ribeiro JMC, Nelson RJ, Lerner MR. Isolation of maxadilan, a potent vasodilatory peptide from the salivary glands of the sand fly *Lutzomyia longipalpis*. *J. Biol. Chem.* 1991; 266:11234–11236. [PubMed: 2040631]
- Lima HC, Titus RG. Effects of sand fly vector saliva on development of cutaneous lesions and the immune response to *Leishmania braziliensis* in BALB/c mice. *Infect Immun.* 1996; 64:5442–5445. [PubMed: 8945606]
- Lu X, Deadman JJ, Williams JA, Kakkar VV, Rahman S. Synthetic RGD peptides derived from the adhesive domains of snake-venom proteins: evaluation as inhibitors of platelet aggregation. *Biochem J.* 1993; 296:21–24. [PubMed: 8250845]
- Marchler-Bauer A, Panchenko AR, Ariel N, Bryant SH. Comparison of sequence and structure alignments for protein domains. *Proteins.* 2002; 48:439–446. [PubMed: 12112669]
- Munstermann, LE. Phlebotomine sand flies, the Psychodidae. In: Marquardt, WC.; Black, WC.; Freier, JE.; Hagedorn, HH.; Hemingway, J.; Higgs, S.; James, AA.; Kondratieff, B.; Moore, CG., editors. *Biology of Disease Vectors.* second ed. San Diego, CA: Elsevier; 2005. p. 41-151.
- Oliveira F, Kamhawi S, Seitz AE, Pham VM, Guigal PM, Fischer L, Ward J, Valenzuela JG. From transcriptome to immunome: identification of DTH inducing proteins from a *Phlebotomus ariasi* salivary gland cDNA library. *Vaccine.* 2006; 24:374–390. [PubMed: 16154670]
- Oliveira F, Lawyer PG, Kamhawi S, Valenzuela JG. Immunity to distinct sand fly salivary proteins primes the anti-*Leishmania* immune response towards protection or exacerbation of disease. *PLoS Negl. Trop. Dis.* 2008; 2:e226. [PubMed: 18414648]
- Oliveira F, Jochim RC, Valenzuela JG, Kamhawi S. Sand flies, *Leishmania*, and transcriptome-borne solutions. *Parasitol. Int.* 2009; 58:1–5. [PubMed: 18768167]
- Ribeiro JMC. Blood-feeding arthropods: live syringes or invertebrate pharmacologists? *Infect. Agents Dis.* 1995; 4:143–152. [PubMed: 8548192]
- Ribeiro JMC, Francischetti IMB. Role of arthropod saliva in blood feeding: sialome and post-sialome perspectives. *Annu. Rev. Entomol.* 2003; 48:73–88. [PubMed: 12194906]
- Ribeiro JMC, Modi G. The salivary adenosine/AMP content of *Phlebotomus argentipes* Annandale and Brunetti, the main vector of human kala-aza. *J Parasitol.* 2001; 87:915–917. [PubMed: 11534660]
- Ribeiro JMC, Vachereau A, Modi GB, Tesh RB. A novel vasodilatory peptide from the salivary glands of the sand fly *Lutzomyia longipalpis*. *Science.* 1989; 243:212–214. [PubMed: 2783496]
- Ribeiro JMC, Katz O, Pannell LK, Waitumbi J, Warburg A. Salivary glands of the sand fly *Phlebotomus papatasi* contain pharmacologically active amounts of adenosine and 5'-AMP. *J. Exp. Biol.* 1999; 202:1551–1559. [PubMed: 10229701]
- Ribeiro JMC, Andersen J, Silva-Neto MA, Pham VM, Garfield MK, Valenzuela JG. Exploring the sialome of the blood-sucking bug *Rhodnius prolixus*. *Insect Biochem. Mol. Biol.* 2004; 34:61–79. [PubMed: 14976983]
- Rohoušová I, Volf P. Sand fly saliva: effects on host immune response and *Leishmania* transmission. *Folia Parasitol.* 2006; 53:161–171. [PubMed: 17120496]
- Rohoušová I, Subrahmanyam S, Volfová V, Mu J, Volf P, Valenzuela JG, Jochim RC. Salivary gland transcriptomes and proteomes of *Phlebotomus tobbi* and *Phlebotomus sergenti*, vectors of leishmaniasis. *PLoS Negl. TropDis.* 2012; 6:e1660.
- Santos A, Ribeiro JMC, Lehane MJ, Gontijo NF, Veloso AB, Sant'Anna MR, Nascimento Araujo R, Grisard EC, Pereira MH. The sialotranscriptome of the blood-sucking bug *Triatoma brasiliensis* (Hemiptera, Triatominae). *Insect Biochem. Mol. Biol.* 2007; 37:702–712. [PubMed: 17550826]
- Schultz J, Milpetz F, Bork P, Ponting CP. SMART, a simple modular architecture research tool: identification of signaling domains. *Proc. Natl. Acad. Sci. USA.* 1998; 95:5857–5864. [PubMed: 9600884]
- Singh H, Raghava GP. ProPred: prediction of HLA-DR binding sites. *Bioinformatics.* 2001; 17:1236–1237. [PubMed: 11751237]

- Takaoka H, Gomez EA, Alexander JB, Hashiguchi Y. Natural infections with *Leishmania* promastigotes in *Lutzomyia ayacuchensis* (Diptera: Psychodidae) in an Andean focus of Ecuador. *J. Med. Entomol.* 1990; 27:701–702. [PubMed: 2388245]
- Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 2007; 24:1596–1599. [PubMed: 17488738]
- Tavares NM, Silva RA, Costa DJ, Pitombo MA, Fukutani KF, Miranda JC, Valenzuela JG, Barral A, de Oliveira CI, Barral-Netto M, Brodskyn C. *Lutzomyia longipalpis* saliva or salivary protein LJM19 protects against *Leishmania braziliensis* and the saliva of its vector, *Lutzomyia intermedia*. *PLoS Negl. Trop. Dis.* 2011; 5:e1169. [PubMed: 21655303]
- Teixeira C, Gomes R, Collin N, Reynoso D, Jochim R, Oliveira F, Seitz A, Elnaïem DE, Caldas A, de Souza AP, Brodskyn CI, de Oliveira CI, Mendonca I, Costa CH, Volf P, Barral A, Kamhawi S, Valenzuela JG. Discovery of markers of exposure specific to bites of *Lutzomyia longipalpis*, the vector of *Leishmania infantum chagasi* in Latin America. *PLoS Negl. Trop. Dis.* 2010; 4:e638. [PubMed: 20351786]
- Theodos CM, Ribeiro JMC, Titus RG. Analysis of enhancing effect of sand fly saliva on *Leishmania* infection in mice. *Infect. Immun.* 1991; 59:1592–1598. [PubMed: 2019430]
- Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994; 22:4673–4680. [PubMed: 7984417]
- Titus RG, Ribeiro JMC. Salivary gland lysates from the sand fly *Lutzomyia longipalpis* enhance *Leishmania* infectivity. *Science.* 1988; 239:1306–1308. [PubMed: 3344436]
- Valenzuela, JG. Blood-feeding arthropod salivary glands and saliva. In: Marquardt, WC.; Black, WC.; Freier, JE.; Hagedorn, HH.; Hemingway, J.; Higgs, S.; James, AA.; Kondratieff, B.; Moore, CG., editors. *Biology of Disease Vectors*. second ed. San Diego, CA: Elsevier; 2005. p. 377-386.
- Valenzuela JG, Belkaid Y, Garfield MK, Mendez S, Kamhawi S, Rowton ED, Sacks DL, Ribeiro JMC. Toward a defined anti-*Leishmania* vaccine targeting vector antigens: characterization of a protective salivary protein. *J. Exp. Med.* 2001; 194:331–342. [PubMed: 11489952]
- Valenzuela JG, Charlab R, Gonzalez EC, de Miranda-Santos IK, Marinotti O, Francischetti IMB, Ribeiro JMC. The D7 family of salivary proteins in blood sucking diptera. *Insect Mol. Biol.* 2002; 11:149–155. [PubMed: 11966880]
- Valenzuela JG, Garfield M, Rowton ED, Pham VM. Identification of the most abundant secreted proteins from the salivary glands of the sand fly *Lutzomyia longipalpis*, vector of *Leishmania chagasi*. *J. Exp. Biol.* 2004; 207:3717–3729. [PubMed: 15371479]
- Vlkova M, Rohousova I, Drahota J, Stanneck D, Kruehwagen EM, Mencke N, Otranto D, Volf P. Canine antibody response to *Phlebotomus perniciosus* bites negatively correlates with the risk of *Leishmania infantum* transmission. *PLoS Negl. Trop. Dis.* 2011; 5:e1344. [PubMed: 22022626]
- Weis WI, Taylor ME, Drickamer K. The C-type lectin superfamily in the immune system. *Immunol. Rev.* 1998; 163:19–34. [PubMed: 9700499]
- Xu X, Oliveira F, Chang BW, Collin N, Gomes R, Teixeira C, Reynoso D, My Pham V, Elnaïem DE, Kamhawi S, Ribeiro JMC, Valenzuela JG, Andersen JF. Structure and function of a “yellow” protein from saliva of the sand fly *Lutzomyia longipalpis* that confers protective immunity against *Leishmania major* infection. *J. Biol. Chem.* 2011; 286:32383–32393. [PubMed: 21795673]
- Young, DG.; Duncan, MA. *Memoirs of the American Entomological Institute*. Vol. 54. Gainesville, FL: Associated Publishers—American Entomological Institute; 1994. Guide to the Identification and Geographic Distribution of *Lutzomyia* Sand Flies in Mexico, the West Indies, Central and South America (Diptera: Psychodidae).





Fig. 1. (A) Sequence alignment of PpSP15/SL1 family proteins from *Lutzomyia (Lu.) ayacuchensis* (LayS36, LayS37, LayS58, LayS59, LayS66, LayS68, LayS71, and LayS72) together with those from *Phlebotomus (P.) arabicus* (Pab), *P. argentipes* (Pag), *P. ariasi* (Par), *P. duboscqi* (Pdu), *P. papatasi* (Pp), *P. perniciosus* (Ppe), and *Lu. longipalpis* (Lulo). Black-shaded amino acids represent identical amino acids and gray-shaded amino acids represent conserved amino acids. Dashes indicate gaps introduced for maximal alignment. Asterisks at the top of the amino acids denote conserved cysteine residues, and an arrowhead indicates the predicted signal peptide cleavage site. (B) Phylogenetic tree analysis of amino acid

sequences of PpSP15/SL1 family proteins from sand flies. Accession numbers are in parentheses and node values indicate branch support.

A	LayS22	M	LFFLLVFTGILFOGILGVEITQG	KWRO	LYGDVTPGT	NPDTN	PFAFAHADGKH	FLT	PRKFKPIPHLA	VDETEKNPGVSGKRS	90										
	LayS23	M	LFFLLVFTGILFOGILGVEITQG	KWRO	LYGDVTPGT	NPDTN	PFAFAHADGKH	FLT	PRKFKPIPYLA	VDETEKNPGVSGKRS	90										
	LayS24	M	LFFLLVFTGILFOGILGVEITQG	KWRO	LYGDVTPGT	NPDTN	PFAFAHADGKH	FLT	PRKFKPIPYLA	VDETEKNPGVSGKRS	90										
	LayS118	M	FFLCFFVFLSQGIGHAYVEITG	SWSN	TFEGLDTRK	KPRNN	PFAFAHDPEGYK	FIS	PRRLQPVYVMA	LNTVMHFGYFVERA	90										
	LJM17	M	FFVFLAIVLFGQIGHAYVEITG	SLRN	TFEGLDTRD	NPKFN	PGLAVDPEGYR	FIA	PRKPKPYVPIA	LNMVMHFGYFVERA	90										
	LJM111	M	LFVFLVFTFGLVQTIFVGEIKQG	KWKN	LYEAGDTSEN	NPDDN	LFAFVDPEEQK	FLT	PRKYPEITMYLA	VDETEKNSFESGDS	89										
	LJM11	M	VFFSIFLTVLFOQTILGAD-TQG	KWKO	LYNNVTPGS	NPDDN	SNAFVDAEAGEK	FIA	PRKLRPVYPIA	VDTKNSLGVMKGDHS	89										
	PabSP26	M	IFLCLFAVVSFHGAFAHYHVERE	AWKN	TYEGIDQAS	NIENT	PFAFAHDAISK	FIT	PRRYQOVPIILT	LDTSKHPE---RSP	87										
	PagSP04	M	WFFLLSTIVFQOGLGVHVERE	AWRN	TVEGVNPS	NVLHS	PAGFVDAETQK	FVA	PRRYQOVPIILT	IERKKHPE---RSP	87										
	ParSP04	M	IFLCLFAAVSIQAGALASQIERE	AWKN	IYEGIDQGS	NIENS	PFAFAHDAASK	FIT	PRIN-QVPIILT	FDISKYFG---GSP	86										
	ParSP04b	M	IFMGLIAVVSIGALASHYVERE	AWKN	TVEGIDQAS	NIENS	PFAFVDAALSK	IYA	PRLYPQVPIILT	LDTTKHPE---RSP	87										
	PduM10	M	ETLSVLAIASFORVFCDDVERA	AWRN	SFVDTRGGT	NPEDV	PAGVTHDAKTK	YFG	PRLYMNIPIYLA	IDTKVNSREIRSP	90										
	PduM35	M	LILTVLALFLSLQVALSDDVGR	EWSK	DIYGVREPV	DSSNI	PAGVYVADSKM	FFC	PRKYSKVPIYA	LSTRSYNSERRDD	90										
	PpsP42	M	LILCLVLSPLSLQVALSDDVGR	EWSE	KLVGVRPNA	DSGNI	PAGVYVDAASKM	FFC	PRLYSRVPIYA	LSTRSYNSREIPINP	90										
	PpsP44	M	FFLSVLAIASFQVVCDDVERP	AWRN	TFEVDKRG	KPGDV	PAGVTHDAKTK	YFG	PRRYSNIPIYLA	IDTRVNSREIRSP	90										
	FpsSP03	M	IFLCLIAVVSIGQVLAYDIERE	AWKN	SFEGIDPAS	SVKNS	VAGFAHADSKK	FIT	PRLN-PVPIILT	LDTTKHPE---RSP	86										
	FpsSP03b	M	IFLCLIAVVSIGQVGVFHYERE	AWKN	SYEGVDPAL	NIDNI	PAGVYVDAINK	FIA	PRRSQIPFPIILT	LDTTKHPE---RSP	87										
	LayS22	E	---LLN	ESGHKSGNELT	YQP	D	CRRLW	VQVGSV	YRSRGAKYPSHRPA	A	DIK	PNYPE	VRHFFPVRLVE	KPTYFGCG	176						
	LayS23	E	---LLN	ESGHKSGKELT	YQP	D	CRRLW	VQVGSV	YRSRGAKYPSHRPA	A	DIK	PNYPE	VRHFFPVRLVE	KPTYFGCG	176						
	LayS24	E	---LLN	ESGHKSGKELT	YQP	D	CRRLW	VQVGSV	YRSRGAKYPSHRPA	A	DIK	PNYPE	VRHFFPVRLVE	KPTYFGCG	176						
	LayS118	E	---KLS	FTGO-SSKDLV	YQP	D	CRRLW	VDTGAV	YSGDDAGYKTKQKFA	V	DIK	DHYPE	GRVELPDSVAS	KPTSFGCG	175						
	LJM17	E	---SFE	FFKPE	ENGE-GKDKLV	YQP	D	CRRLW	LDIGKV	YTGDDADQYKPKGT	A	DIK	DHTPE	BRFEPIDDLYS	SQVFEFGCG	178					
	LJM111	E	---LLG	ESGHETGKELT	YQP	D	CRRLW	VQVGSV	RNSDGTGGQPEHNP	A	DIK	ANYPE	TRTYFFPNSIE	KPYFTEFGCG	176						
	LJM11	E	---LLN	ESGHKTKELT	YQP	D	CRRLW	VQVGSV	YRSRGAKYPSHRPA	A	DIK	PNYPE	VRHYFFTRLVE	KPTYFGCG	175						
	PabSP26	E	---LS	ESGK-SDDLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	ENYPE	DREYLFABVVK	NPLGFEFGCG	168						
	PagSP04	E	---LS	ESGK-SSKDLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	ENYPE	DREYLFABVVK	NPLGFEFGCG	170						
	ParSP04	E	---LS	ESGK-SDDLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	DNYPE	DRVEIPIPIAG	NPLGFEFGCG	167						
	ParSP04b	E	---LE	ESGK-SDKLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	DNYPE	DRVEIPIPIAG	NPLGFEFGCG	168						
	PduM10	E	---FS	ESGK-SKKPLT	YQP	D	CRRLW	LDVCEA	YKK-NGNEYTPKNPE	A	DIT	ENYPE	DRYKLGDDVAK	NPLGFEFGCG	173						
	PduM35	E	---LD	ESGK-SKKPLT	YQP	D	CRRLW	LDVCEA	YKK-NGNEYTPKNPE	A	DIT	DNYPE	DRYKLGDDVAK	NPLGFEFGCG	173						
	PpsP42	E	---LD	ESGK-SKQPLT	YQP	D	CRRLW	LDVCEA	YKK-NGNEYTPKNPE	A	DIT	DNYPE	DRYKLGDDVAK	NPLGFEFGCG	173						
	FpsSP44	E	---FS	ESGK-SGKPLT	YQP	D	CRRLW	LDVCEA	YKK-NGNEYTPKNPE	A	DIT	ENYPE	DRYKLGDDVAK	NPLGFEFGCG	173						
	FpsSP03	E	---LS	ESGK-SDKLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	DNYPE	DRVEIPIPIAG	NPLGFEFGCG	167						
	FpsSP03b	E	---LS	ESGK-SDKLI	YQP	D	CRRLW	VQAV	YKE-DEQKPKKNA	A	DIT	DNYPE	DRVEIPIPIAG	NPLGFEFGCG	169						
	LayS22	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	260
	LayS23	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	260
	LayS24	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	260
	LayS118	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	260
	LJM17	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	263
	LJM111	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	263
	LJM11	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	259
	PabSP26	T	DVNNPFDGCS	---K	E	YI	TN	LENT	I	DQEKD	WKISHG	E	KP	-HDSVLTHNG	-KEHKYKVC	F	GHITLQ	RDEG	R	PAMY	252
	PagSP04	A	DVNNPFDGCS	---D	E	YI	TN	LENT	I	DQEKD	WKISHG	E	KP	-HDSVLTHNG	-KEHKYKVC	F	GHITLQ	RDEG	R	PAMY	255
	ParSP04	T	DVNNPFDGCS	---K	E	YI	TN	LENT	I	DQEKD	WKISHG	E	KP	-HDSVLTHNG	-KEHKYKVC	F	GHITLQ	RDEG	R	PAMY	251
	ParSP04b	T	DVNNPFDGCS	---K	E	YI	TN	LENT	I	DQEKD	WKISHG	E	KP	-HDSVLTHNG	-KEHKYKVC	F	GHITLQ	RDEG	R	PAMY	251
	PduM10	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	262
	PduM35	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	262
	PpsP42	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	260
	FpsSP44	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	261
	FpsSP03	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	251
	FpsSP03b	A	DVNNPFDGCS	---E	EVY	TN	LSNA	F	DHKNQQ	VNYTDA	E	K	KA	-RPSTFDHOG	-KQYTYKCA	F	GHITLQ	RDKD	R	PAMY	253
	LayS22	AGS	IK	YS	NKKEIKQ	NKKNPELLG	NRCKLN	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKMP	RMQHTDV	SNTFRV	G	D	S			350
	LayS23	AGS	IK	YS	NKKEIKQ	NKKNPELLG	NRCKLN	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKMP	RMQHTDV	SNTFRV	G	D	S			350
	LayS24	AGS	IK	YS	NKKEIKQ	NKKNPELLG	NRCKLN	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKMP	RMQHTDV	SNTFRV	G	D	S			350
	LayS118	AGS	IK	YS	NKKEIKQ	GGSN	NPDLG	DRCPHT	A	ALAVDPKTKV	PABSD	PROV	SCW	VQTE	-TPKNTDV	SYARFI	G	D	S		348
	LJM17	AGS	IK	YS	NKKEIKQ	NGQLNPOLG	DRCPYIT	A	ALAVDPKTKV	PABSD	PROV	SCW	VNME	-TPKNTDV	SSARFT	G	D	S			352
	LJM111	VAS	IK	HS	NKKEIKQ	GSKVNYLIG	DRCEST	A	ALAVDPKTKV	PABSD	PROV	SCW	TQET	-TPKNTDV	HNADFS	G	D	S			348
	LJM11	AGS	IK	YS	NKKEIKQ	GKGLNPELLG	NRCKYN	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKMP	RMKNTDV	TSSRFV	G	D	S			349
	PabSP26	GGS	IK	FE	SNKAKKK	GAKFDPVRLG	DRCPKT	A	ALAVDPKTKV	PABAD	PROV	SCW	TQKP	-LNHKTVDV	ASAKFI	G	D	S			341
	PagSP04	GGS	IK	YK	SNKAKKK	GAKFDPVRLG	DRCPYIT	A	ALAVDPKTKV	PABSI	PROV	SCW	TQTP	-LNHKTVDV	ASARFL	G	D	S			344
	ParSP04	GGS	IK	FE	SNKAKKK	GAKFDPVRLG	DRCPHT	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKP	-LNHKTVDV	ASPEFI	G	D	S			340
	ParSP04b	GGS	IK	FE	SNKAKKK	GAKFDPVRLG	DRCPHT	A	ALAVDPKTKV	PABSD	PROV	SCW	TQKP	-LNHKTVDV	ASSKFI	G	D	S			341
	PduM10	AGS	IK	YN	SNKAKKK	VKSILP	PTLGDRCYKT	A	ALAVDPKTKV	PABSD	PROV	SCW	IQKD	-TPKNTDV	TNAYFV	G	D	M			350
	PduM35	AGS	IK	YR	SNKAKKK	GSKLVPKLIG	DRCPKT	A	ALAVDPKTKV	PABTD	PROV	SCW	IKKE	-TPKNTDV	SSAKLN	A	D	M			349
	FpsP42	AGS	IK	YR	SNKAKKK	GSKLVPKLIG	DRCPKT	A	ALAVDPKTKV	PABAD	PROV	SCW	IKKE	-TPKNTDV	ANPFNN	G	D	M			349
	FpsP44	AGS	IK	YS	SNKAKKK	GASLPRLLG	DRCPHT	A	ALAVDPKTKV	PABSD	PROV	SCW	IQKE	-TPKNTDV	TNAYFV	G	D	M			350
	FpsSP03	GGS	IK	FR	NKAKKK	ACQIETPLG	DRCSHS	A	ALAVDPKTKV	PABSN	PROV	SCW	TQKS	-LNPDNDV	HSPDFI	G	D	S			340
	FpsSP03b	AGS	IK	FE	SNKAKKK	GAKFDPVRLG	DRCPHT	A	ALAVDPKTKV	PABSD	PROV	SCW	TQKP	-LNHKTVDV	ASAKFI	G	D	S			342
	LayS22	DSKGT	WFM	NGHPE	VKNKQKFK	----	FDYPRYR	LN-	DTQAAIAGTN	-CEIKPR	-----									400	
	LayS23	DSKGT	WFM	NGHPE	VKNKQKFK	----	FDYPRYR	LN-	DTQAAIAGTN	-CEIKPR	-----									400	
	LayS24	DSKGT	WFM	NGHPE	VKNKQKFK	----	FDYPRYR	LN-	DTQAAIAGTN	-CEIKPR	-----									400	
	LayS118	DKKGT	WFM	NGHPE	VIKDAE	---K-LKFYDRKR	MR-	NTYVLPYSK	-CNPYKPGQGI	PIV										405	
	LJM17	DSKGT	WFM	NGHPE	VQEKIKW	-MRFVNRKR	MK-	DTERVFKYSR	-CNPYKPKPKIE	IV										397	
	LJM111	DSQDN	WFL	NGHPE	LENSDKFV	----	FTKPRYQ	EK-	NIQEAIAGTK	-CEKNL	-----									412	
	LJM11	DSKGT	WFM	NGHPE	IRKSEKFK	----	YDFPRYR	MRI	DTQEAIAGTA	-CDMNA	-----									399	
	PabSP26	DSKGT	WFL	NGHPE	VENLKL	----	FDK														

**Fig. 2.**

(A) Sequence alignment of yellow-related proteins from *Lu. ayacuchensis* (LayS22, LayS23, LayS24, and LayS118) together with those from *P. arabicus* (Pab), *P. argentipes* (Pag), *P. ariasi* (Par), *P. duboscqi* (Pdu), *P. papatasi* (Pp), *P. perniciosus* (Ppe), and *Lu. longipalpis* (LJM). Black-shaded amino acids represent identical amino acids and gray-shaded amino acids represent conserved amino acids. Dashes indicate gaps introduced for maximal alignment. Asterisks at the top of the amino acids denote conserved cysteine residues, and an arrowhead indicates the predicted signal peptide cleavage site. Closed circles show conserved amino acids contained in the ligand binding pocket of yellow-related protein family. Potential N-glycosylation sites of *Lu. ayacuchensis* yellow-related proteins are boxed. (B) Phylogenetic tree analysis of amino acid sequences of yellow-related proteins from sand flies. Accession numbers are in parentheses and node values indicate branch support.




Fig. 3. (A) Sequence alignment of D7 family proteins from *Lu. ayacuchensis* (LayS101–LayS103) together with those from *P. arabicus* (Pab), *P. argentipes* (Pag), *P. ariasi* (Par), *P. duboscqi* (Pdu), *P. papatasi* (Pp), *P. perniciosus* (Ppe), and *Lu. longipalpis* (Lulo). Black-shaded amino acids represent identical amino acids and gray-shaded amino acids represent conserved amino acids. Dashes indicate gaps introduced for maximal alignment. Asterisks at the top of the amino acids denote conserved cysteine residues, and an arrowhead indicates the predicted signal peptide cleavage site. (B) Phylogenetic tree analysis of amino acid sequences of D7 family proteins from sand flies. Accession numbers are in parentheses and node values indicate branch support.

Table 1Functional classification of housekeeping genes expressed in *Lutzomyia ayacuchensis* salivary glands

Type of transcripts	Cluster	%	Sequence	%
Energy production and conversion	20	21.3	24	19.2
Translation, ribosomal structure and biogenesis	19	20.2	23	18.4
General function prediction only	9	9.6	10	8.0
Signal transduction mechanisms	8	8.5	8	6.4
Posttranslational modification, protein turnover, chaperones	7	7.4	14	11.2
Lipid transport and metabolism	5	5.3	5	4.0
Amino acid transport and metabolism	4	4.3	4	3.2
Chromatin structure and dynamics	2	2.1	2	1.6
Inorganic ion transport and metabolism	2	2.1	2	1.6
Intracellular trafficking, secretion, and vesicular transport	2	2.1	3	2.4
RNA processing and modification	2	2.1	2	1.6
Secondary metabolites biosynthesis, transport and catabolism	2	2.1	2	1.6
Transcription	2	2.1	14	11.2
Carbohydrate transport and metabolism	1	1.1	3	2.4
Cell cycle control, cell division, chromosome partitioning	1	1.1	1	0.8
Cell wall/membrane/envelope biogenesis	1	1.1	1	0.8
Defense mechanisms	1	1.1	1	0.8
Unknown conserved	6	6.4	6	4.8
Total	94	100.0	125	100.0

Table 2

Most abundant salivary gland transcripts from *Lutzomyia ayacuchensis*

Cluster	Sequence name	No. of seq in cluster	Mature Mw (kDa)	pI	Best match to NR protein database			Accession No.
					Accession No.	Organism	% Identity	
LayS45	RGD-containing peptide	90	5.3	3.44	AAD32196	<i>Lu. longipalpis</i>	AK416785	
LayS37	14 kDa Salivary protein	58	14.1	8.21	ABA43057	<i>P. perniciosus</i>	AK416778	
LayS81	Antigen 5-related protein	24	28.5	9.08	AAD32191	<i>Lu. longipalpis</i>	AK416820	
LayS24	Yellow-related protein	23	43.5	9.37	AAS05318	<i>Lu. longipalpis</i>	AK416770	
LayS59	SL1-like protein	19	13.8	8.22	AAD32197	<i>Lu. longipalpis</i>	AK416798	
LayS91	27 kDa Salivary protein	17	27.2	10.14	AAS16906	<i>Lu. longipalpis</i>	AK416828	
LayS117*	Yellow-related protein	16			AAD32198	<i>Lu. longipalpis</i>	AB744655	
LayS118	Yellow-related protein	14	43.8	8.05	AAD32198	<i>Lu. longipalpis</i>	AK416844	
LayS127	16.4 kDa Salivary protein	11	16.4	8.05	ABB00903	<i>Lu. longipalpis</i>	AK416852	
LayS21	Putative apyrase	11	35.7	9.19	AAD33513	<i>Lu. longipalpis</i>	AK416767	
LayS142	9.3 kDa Protein	10	9.3	4.22	XP 001238639	<i>Eimeria tenella</i>	AK416859	
LayS103	D7-related salivary protein	9	26.6	8.81	ABA43052	<i>P. perniciosus</i>	AK416840	
LayS36	14 kDa Salivary protein	9	14.1	8.21	ABA43057	<i>P. perniciosus</i>	AK416777	
LayS6	11.5 kDa Salivary protein	8	11.5	6.67	AAS16919	<i>Lu. longipalpis</i>	AK416753	
LayS102	D7-related salivary protein	8	26.6	8.81	ABA43052	<i>P. perniciosus</i>	AK416839	
LayS19	Putative apyrase	8	35.8	9.09	AAD33513	<i>Lu. longipalpis</i>	AK416765	
LayS58	SL1-like protein	8	13.8	8.22	AAD32197	<i>Lu. longipalpis</i>	AK416797	
LayS68	SL1-like protein	8	13.9	8.61	AAD32197	<i>Lu. longipalpis</i>	AK416807	
LayS90	27 kDa Salivary protein	8	26.6	10.08	AAS16906	<i>Lu. longipalpis</i>	AK416827	
LayS101	D7-related salivary protein	7	26.6	8.81	ABA43052	<i>P. perniciosus</i>	AK416838	
LayS66	SL1-like protein	6	13.9	8.61	AAD32197	<i>Lu. longipalpis</i>	AK416805	
LayS20	Putative apyrase	6	35.5	9.00	AAD33513	<i>Lu. longipalpis</i>	AK416766	
LayS22	Yellow-related protein	6	43.4	9.39	AAS05318	<i>Lu. longipalpis</i>	AK416768	
LayS17	Putative apyrase	5	35.5	9.11	AAD33513	<i>Lu. longipalpis</i>	AK416763	
LayS18	Putative apyrase	5	36.0	9.46	AAD33513	<i>Lu. longipalpis</i>	AK416764	
LayS23	Yellow-related protein	5	43.5	9.37	AAS05318	<i>Lu. longipalpis</i>	AK416769	
LayS26	33.6 kDa Salivary protein	5	33.7	8.95	AAX55751	<i>P. ariasi</i>	AK416772	

Cluster	Sequence name	No. of seq in cluster	Mature Mw (kDa)	pI	Best match to NR protein database		E value	% Identity	Accession No.
					Accession No.	Organism			
LayS44	RGD-containing peptide	5	5.3	3.47	AAD32196	<i>Lu. longipalpis</i>	8E-07	46	AK416784
LayS71	SL1-like protein	5	13.8	8.22	AAD32197	<i>Lu. longipalpis</i>	2E-50	64	AK416810
LayS72	SL1-like protein	5	13.8	8.43	AAD32197	<i>Lu. longipalpis</i>	3E-51	64	AK416811
LayS100	D7-related salivary protein	4	26.6	8.61	ABA43052	<i>P. perniciosus</i>	4E-86	59	AK416837
LayS131	16.4 kDa Salivary protein	4	16.4	8.32	ABB00903	<i>Lu. longipalpis</i>	7E-26	39	AK416856
LayS3	Allergen-related protein	4	26.0	5.68	XP 001865175	<i>Culex quinquefasciatus</i>	8E-31	35	AK416750
LayS64	SL1-like protein	4	13.9	8.61	AAD32197	<i>Lu. longipalpis</i>	5E-48	59	AK416803
LayS97	D7-related salivary protein	4	26.6	8.81	ABA43052	<i>P. perniciosus</i>	2E-86	59	AK416834
LayS128	16.4 kDa Salivary protein	3	16.4	8.05	ABB00903	<i>Lu. longipalpis</i>	1E-29	39	AK416853
LayS129	16.4 kDa Salivary protein	3	16.4	7.69	ABB00903	<i>Lu. longipalpis</i>	2E-29	40	AK416854
LayS16	Putative apyrase	3	36.6	9.34	AAD35513	<i>Lu. longipalpis</i>	1E-125	64	AK416762
LayS27	33.6 kDa Salivary protein	3	33.7	8.69	AAX55751	<i>P. ariasi</i>	1E-90	52	AK416773
LayS46	RGD-containing peptide	3	5.3	3.44	AAD32196	<i>Lu. longipalpis</i>	1E-07	49	AK416786
LayS47	RGD-containing peptide	3	5.3	3.44	AAD32196	<i>Lu. longipalpis</i>	2E-07	47	AK416787
LayS70	SL1-like protein	3	13.9	7.98	AAD32197	<i>Lu. longipalpis</i>	5E-51	64	AK416809
LayS77	Antigen 5-related protein	3	28.6	9.21	AAD32191	<i>Lu. longipalpis</i>	1E-135	80	AK416816
LayS89	27 kDa Salivary protein	3	26.6	10.08	AAS16906	<i>Lu. longipalpis</i>	5E-90	58	AK416826
LayS98	D7-related salivary protein	3	26.6	8.81	ABA43052	<i>P. perniciosus</i>	3E-86	59	AK416835
LayS130	16.4 kDa Salivary protein	2	16.4	8.05	ABB00903	<i>Lu. longipalpis</i>	2E-29	40	AK416855
LayS132	16.4 kDa Salivary protein	2	16.4	8.05	ABB00903	<i>Lu. longipalpis</i>	2E-29	40	AK416857
LayS2	Allergen-related protein	2	26.0	5.46	XP 001865175	<i>Culex quinquefasciatus</i>	5E-30	34	AK416749
LayS60	SL1-like protein	2	13.9	7.98	AAD32197	<i>Lu. longipalpis</i>	7E-51	64	AK416799
LayS61	SL1-like protein	2	13.8	8.22	AAD32197	<i>Lu. longipalpis</i>	1E-51	65	AK416800
LayS62	SL1-like protein	2	13.8	8.52	AAD32197	<i>Lu. longipalpis</i>	6E-51	64	AK416801
LayS63	SL1-like protein	2	13.8	8.23	AAD32197	<i>Lu. longipalpis</i>	1E-51	64	AK416802
LayS65	SL1-like protein	2	13.8	7.98	AAD32197	<i>Lu. longipalpis</i>	1E-51	64	AK416804
LayS67	SL1-like protein	2	13.8	8.35	AAD32197	<i>Lu. longipalpis</i>	1E-50	64	AK416806
LayS69	SL1-like protein	2	13.9	8.22	AAD32197	<i>Lu. longipalpis</i>	4E-51	64	AK416808
LayS7	11.5 kDa Salivary protein	2	10.8	8.36	AAS16919	<i>Lu. longipalpis</i>	2E-07	34	AK416754
LayS78	Antigen 5-related protein	2	28.4	9.16	AAD32191	<i>Lu. longipalpis</i>	1E-133	80	AK416817

Cluster	Sequence name	No. of seq in cluster	Mature Mw (kDa)	pI	Best match to NR protein database			E value	% Identity	Accession No.
					Accession No.	Organism	Accession No.			
LayS79	Antigen 5-related protein	2	28.4	9.08	AAD32191	<i>Lu. longipalpis</i>	1E-135	80	AK416818	
LayS80	Antigen 5-related protein	2	28.5	9.01	AAD32191	<i>Lu. longipalpis</i>	1E-135	80	AK416819	
LayS92	27 kDa Salivary protein	2	27.3	10.14	AAS16906	<i>Lu. longipalpis</i>	9E-91	58	AK416829	
LayS93	27 kDa Salivary protein	2	27.2	10.14	AAS16906	<i>Lu. longipalpis</i>	1E-91	58	AK416830	
LayS95	D7-related salivary protein	2	26.6	8.81	ABA43052	<i>P. perniciosus</i>	5E-86	59	AK416832	
LayS96	D7-related salivary protein	2	26.6	8.81	ABA43052	<i>P. perniciosus</i>	1E-85	58	AK416833	
LayS99	D7-related salivary protein	2	26.6	8.81	ABA43052	<i>P. perniciosus</i>	6E-86	59	AK416836	

* Truncated in the 5' region.

Table 3Classification of transcripts coding for putative secreted proteins in *Lutzomyia ayacuchensis* salivary glands

	Cluster	Sequence	% Sequence
RGD-containing peptide	10	107	18.7
SL1-like protein	25	82	14.3
14 kDa Salivary protein	6	71	12.4
Yellow-related protein	5	64	11.1
Putative apyrase	13	45	7.9
D7-related salivary protein	9	42	7.3
27 kDa Salivary protein	11	38	6.6
Antigen 5-related protein	9	37	6.5
16.4 kDa Salivary protein	13	32	5.6
9.3 kDa Protein	3	16	2.8
11.5 kDa Salivary protein	4	12	2.1
33.6 kDa Salivary protein	3	9	1.6
Allergen-related protein	2	6	1.0
Others	9	12	2.1
Total	122	573	100.0