

RESEARCH

Open Access

# Anti-inflammatory effects of fish oil in ovaries of laying hens target prostaglandin pathways

Erfan Eilati,Carolynn C Small, Stacey R McGee, Nawneet K Kurrey and Dale Buchanan Hales\*

## Abstract

**Background:** An effective way to control cancer is by prevention. Ovarian cancer is the most lethal gynecological malignancy. Progress in the treatment and prevention of ovarian cancer has been hampered due to the lack of an appropriate animal model and absence of effective chemo-prevention strategies. The domestic hens spontaneously develop ovarian adenocarcinomas that share similar histological appearance and symptoms such as ascites and metastasis with humans. There is a link between chronic inflammation and cancer. Prostaglandin  $E_2$  ( $PGE_2$ ) is the most pro-inflammatory eicosanoid and one of the downstream products of two isoforms of cyclooxygenase (COX) enzymes: COX-1 and COX-2.  $PGE_2$  exerts its effects on target cells by coupling to four subtypes of receptors which have been classified as EP1-4. Fish oil is a source of omega-3 fatty acids (OM-3FAs) which may be effective in prevention of ovarian cancer. Our objective was to assess the potential impact of fish oil on expression of COX enzymes,  $PGE_2$  concentration, apoptosis and proliferation in ovaries of laying hens.

**Methods:** 48 white Leghorn hens were fed 50, 100, 175, 375 and 700 mg/kg fish oil for 21 days. The OM3-FAs and omega-6 fatty acids contents of egg yolks were determined by Gas Chromatography. Proliferation, apoptosis, COX-1, COX-2 and prostaglandin receptor subtype 4 (EP4) protein and mRNA expression and  $PGE_2$  concentration in ovaries were measured by PCNA, TUNEL, Western blot, quantitative real-time qPCR and ELISA, respectively.

**Results:** Consumption of fish oil increased the incorporation of OM-3FAs into yolks and decreased both COX-1 and COX-2 protein and mRNA expression. In correlation with COXs down-regulation, fish oil significantly reduced the concentrations of  $PGE_2$  in ovaries. EP4 protein and mRNA expression in ovaries of hens was not affected by fish oil treatment. A lower dose of fish oil increased the egg laying frequency. 175 and 700 mg/kg fish oil reduced proliferation and 700 mg/kg increased apoptosis in hen ovaries.

**Conclusions:** Our findings suggest that the lower doses of fish oil reduce inflammatory PG and may be an effective approach in preventing ovarian carcinogenesis. These findings may provide the basis for clinical trials utilizing fish oil as a dietary intervention targeting prostaglandin biosynthesis for the prevention and treatment of ovarian cancer.

**Keywords:** Fish oil, Laying hen, Omega-3 fatty acids, Inflammation, Cyclooxygenase, Prostaglandin  $E_2$

## Introduction

The best approach for reducing the impact of ovarian cancer is by prevention. Fish oil supplementation as a source of omega-3 fatty acids (OM-3FAs) might be effective in prevention and amelioration of the severity of the ovarian cancer by targeting inflammatory prostaglandin pathways.

Cyclooxygenase (COX) is the rate limiting enzyme in catalyzing the conversion of arachidonic acid (AA) to prostaglandins (PGs). Two isoforms of cyclooxygenase have been identified, COX-1 and COX-2. Although both COX isoforms have similar structure and function, they are encoded by different genes (PTGS1 and PTGS2) and show distinct expression patterns. COX-1 is expressed in most cells and tissues and remains constant under most physiologic conditions to play a housekeeping role whereas the COX-2 form is inducible and usually only expressed in response to various inflammatory stimuli

\* Correspondence: dhales@siu.edu  
Department of Physiology, Southern Illinois University Carbondale, School of Medicine, Life Science II, Room 250 (M/C 6512), 1125 Lincoln Drive, Carbondale, IL 62901, USA

[1]. COX enzymes may be involved in both tumor establishment [2] and maintenance of existing tumors [3]. Up-regulation of COX-2 has been reported in many malignancies including breast [4], lung [4] and ovarian cancer [5]. However, we and others have shown that COX-1 is over-expressed in ovarian cancer [3,6,7].

PGs are biologically active lipids that are associated with inflammation, fever, pain and tissue injury [8]. High concentrations of PGE<sub>2</sub> are found in different human cancers including colon, lung, breast, and head and neck cancers [9]. Prostaglandins and their metabolites enhance the cell proliferation, inhibit apoptosis and stimulate cell adhesion. PGE<sub>2</sub> exerts its autocrine/paracrine effects on target cells by coupling to four subtypes of G-protein coupled receptors, which have been classified as EP1-4 [10]. EP4 is over-expressed in many cancers such as castration-resistant prostate cancer [11] and colorectal cancer [12] and EP4 antagonists inhibit breast cancer metastasis in mice [13].

Ovarian cancer is the fifth leading cause of cancer death among women and the most lethal gynecological malignancy. Research into the discovery of early detection markers and therapeutics continues, but there are still no effective treatments for ovarian cancer today [14]. It is clear that poor diet is a well established risk factor for cancer. In fact, the American Institute for Cancer Research estimates that 30–40 percent of all cancers can be prevented by appropriate diets, physical activity and maintenance of appropriate body weight [15]. Omega-6 fatty acids (OM-6FAs) and OM-3FAs are essential poly unsaturated fats that must be obtained from diet. OM-6FAs can be found in red meat, chicken and vegetable seed oil such as sunflower seed oil. OM-3FAs can be found in flaxseed and in oily cold water fish such as salmon and tuna. The main OM-6FA, Linoleic Acid (LA), is a direct precursor of the Arachidonic Acid (AA). The OM-3FAs include alpha-linolenic acid (ALA), eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA). In early human diets, the ratio of OM-6FAs to OM-3FAs was approximately equal; however, with the evolution of the modern western diets, the ratio favored the OM-6FAs that have been attributed to the increased risk of cardiovascular disease and cancer [16]. In contrast, changing the ratio in favor of OM-3FAs has been shown to have growth suppressive effects on cancerous cells in animal studies [17]. Epidemiological evidence and preclinical data suggest that consumption of OM-3FAs might be associated with prevention of many cancers in humans likely due to reduction in inflammatory prostaglandins. However, the mechanism(s) by which the OM-3FAs in dietary fish oil, EPA and DHA, have anti-neoplastic activity remains unclear [18].

Hens spontaneously develop ovarian adenocarcinomas that are similar to human disease by histopathology [19],

expression molecular markers [20-22] and presentation of symptoms such as peritoneal fluid and peritoneal metastatic dissemination [19,23].

Previously we have shown that increased expression of both COX enzymes and concentration of PGE<sub>2</sub> in hen ovaries correspond to the increased incidence of ovarian cancer indicating their importance in early stages of ovarian cancer [24]. Moreover, we demonstrated that the long-term consumption of flaxseed (richest plant source of OM-3FAs) reduced the incidence and severity of ovarian cancer in hens which correlated with reduced COX-2 and PGE<sub>2</sub> [25]. Thus, targeting COX expression and prostaglandin biosynthesis by dietary intervention using fish oil starting at young ages might be an effective approach to prevent or suppress ovarian cancer later in life. The objective of this study was to assess the effect of fish oil on COX-1 and COX-2 enzyme expression and PGE<sub>2</sub> concentrations in normal ovaries of laying hens.

## Material and methods

### Animal care

48 single comb white Leghorn hens, aged one year were used. Hens were maintained two per cage and were provided with feed and water ad libitum and exposed to a photoperiod of 17 h light: 7 h dark, with lights on at 05:00 h and lights off at 22:00 h. Animal management and procedures were reviewed and approved by the Institutional Animal Care and Use Committees at the University of Illinois at Urbana-Champaign and Southern Illinois University at Carbondale.

### Experimental design

This study was performed in two parts. For the first part, 24 hens were randomly divided into four groups (n=6). Three groups were fed 175, 375 and 700 mg/kg/day of fish oil filled into the gelatin capsules and one group was fed empty gelatin capsules. For the second part, 24 hens were randomly divided into four groups (n=6) and three groups were fed 50, 100 and 175 mg/kg/day of fish oil and the empty gelatin capsules were given to the control group. Eggs were collected daily to determine ovulation rate which is reported as number of laid eggs per hen per week. To ensure that the OM-3FAs from fish oil were incorporated into the ovaries, four eggs from all groups were collected on days 0, 7, 14 and 21 of the study and measured for total OM-3FAs and OM-6FAs contents by gas chromatography. If enough eggs were not laid on a given day, the last laid eggs from previous day were used. EPA and DHA contents of fish oil capsules (500 mg omega-3 Salmon oil, Puritan's Pride, Oakdale, NY, USA) were measured and the manufactures reported amounts confirmed. No overt toxicity was observed at any dose.

### Collection of tissue

After 21 days, all hens were euthanized using CO<sub>2</sub> asphyxiation and necropsied. Ovaries were removed from hens and small yellow follicles (6–8 mm) and pre-ovulatory follicles (9–35 mm) were removed. The ovaries were dissected into several pieces. The first portion was frozen in liquid nitrogen and later stored at –80°C; the second portion was put into RNAlater solution and stored at 4°C before processing; the third and fourth portions were used for histological analysis and fixed in NBF fixative solution.

### Gas chromatography

Measuring OM-3FA in egg yolks is a non-invasive approach in the poultry industry to measure total body OM-3FA incorporation [26,27]. The yolk was separated from the rest of the egg. A solution containing 12.5 ug/ml C17:0 standard in methanol was then added to the egg yolks. Lipid extraction was performed using HPLC grade chloroform, water and HPLC grade hexane. The lipids were then dried under a stream of nitrogen and methylated using the Instant Metanolic HCl kit (Alltech, Lexington, KY, USA). After methylation, the lipids were extracted twice with hexane. Once dried under nitrogen, the lipids were re-dissolved in hexane and injected into the Shimadzu 2010 gas chromatograph equipped with a flame ionization detector. Helium was used as the carrier gas and nitrogen as the make up gas with an Omega-wax column. The injection volume was 1.0 µL, helium was used as the carrier gas (30 cm/sec, 205°C), and Helium as the make-up gas with an Omegawax (250, 30 m × 0.25 mm I.D., 0.25 µm film) column. The injector temperature was 250°C. A split injection technique (100:1) was used, and the temperature program was as follows: 175°C held for 8 min, increased to 200°C at 4.5°C/min, and held at 200°C for 8 min, then increased to 230°C at 6°C/min, and held at 230°C for 14 min. Individual FAME were identified by reference to external standards (PUFA-3, and C17:0 Sigma-Adrich). Protein levels were determined by Pierce BCA Protein Assay. Fatty acids were quantified by integration of area under peak after normalization to C17:0 standards and to ug/ug protein.

### RNA extraction and analysis

Total RNA was extracted from ovarian tissue using Trizol reagent. Quantification was performed by determination of absorbance at A260, and qualified by Experion RNA StdSens Analysis. All RNA samples used in this study had a 260:280 ratio range between 2 and 2.10, and had three bands: 5S, 18S and 28S for electrophoresis results. RNA samples were then treated with RQ1 RNase-free DNase prior to reverse transcription reaction. cDNA was synthesized from DNase treated RNA with the high capacity cDNA archive kit.

### Quantitative real-time PCR

The chicken-specific primers and plasmid standards used for each gene were designed. Primer sequences are shown in Table 1. qRT-PCR was conducted by amplifying cDNA with Eva-Green on CFX384 Real-Time System and analyzed with Bio-Rad CFX Manager software. Control reactions lacking template were run for each gene. Reactions were 10 µl in total volume and 200 nM of each primer. The amplification conditions were as follows: 50°C 5 S, 95°C 10 min, 40 cycles for 95°C 15 S, 60°C 1 min.

### Western blot

The proteins were extracted from snap frozen ovarian tissue samples and the western blot was performed to detect COX-1 and COX-2 as described previously [24].

### Apoptosis assay

The presence of apoptotic cells were detected with the TUNEL Apoptosis Detection biotin-labeled POD Kit according to manufacturer's protocol (GenScript, Piscataway, NJ, USA). Briefly, after deparaffinization of 5 µm PPFE sections, they were washed with phosphate buffered saline (PBS) and incubated with 0.02 mg/ml proteinase K at 37 C for 20 min. Cells were blocked with 3% hydrogen peroxide in methanol for 10 min at RT. Sections were incubated with a TUNEL labeling mixture of terminal deoxynucleotidyl transferase (TdT) and biotinylated dUTP at 37 C for 1 hr, followed by two washes in PBS for 5 min each. Apoptotic cells were visualized with a streptavidin conjugated antibody (Life Technologies, Carlsbad, CA, USA; Alexa Fluor 488) at RT for 30 min. After being washed with PBS, the sections were coverslipped using Fluormount G with Dapi mounting media. For control staining, the positive control section was treated for 15 min at 37 C with DNase I (25 U/ul) before incubation with TUNEL labeling mix while the negative control section was incubated in the TUNEL labeling mix without TdT.

**Table 1 Primer sequences**

COX-1	Forward	5'TCAGGTGGTTCTGGGACATCA 3'
	Reverse	5' TGTAGCCGTA CTACTGGGAGTTGAA 3'
COX-2	Forward	5' CTGCTCCCTCCCATGTCAGA 3'
	Reverse	5' CACGTGAAGAATCCGGTGTT 3'
EP4	Forward	5' GGTGTTTCATAGACTGGCGA 3'
	Reverse	5' GCAGATCACCGTAACCATGA 3'
SDHA	Forward	5' CAGGGATGTAGTGTCTGCT 3'
	Reverse	5' GGGAAATAGGCTCCTTAGTG 3'
TBP	Forward	5' CGTCAGGAAATAGGCA 3'
	Reverse	5' GACTGGCAGCAAGGAAG 3'
RPL4	Forward	5' TTATGCCATCTGTCTGTC 3'
	Reverse	5' GCGATTCCCTCATCTTACCCT 3'

### Proliferation assay

Proliferating Cell Nuclear Antigen (PCNA) was used to assess cell proliferation. Normal and cancerous ovaries were fixed in 10% normal buffered formalin and paraffin embedded blocks (PPFE) were sectioned at 5  $\mu$ m. Sections were deparaffinized in xylene and rehydrated in reducing concentrations of ethanol. Antigen retrieval was performed in a pressure cooker using unmasking solution (Sigma) for 20 min, depressurized for 10 min, and then allowed to cool to RT for 20 min changing  $\frac{1}{4}$  of the water every 5 min. The sections were treated with 3% hydrogen peroxide in methanol for 30 min at room temperature (RT) to block endogenous peroxidase activity and rinsed in Tris Buffered Saline with 0.5% tween (TTBS). Non specific binding was blocked using 2.5% normal horse serum for 1 hr at RT then incubation with the mouse monoclonal anti-PCNA antibody (Abcam, Cambridge, MA, USA) for 16–18 hr at 4 C. Slides were washed with TTBS and incubated with donkey anti mouse Alexa Fluor 488 secondary antibody (Jackson Immuno Research, West Grove, PA, USA) for 1 hr at RT. Coverslips were applied using Fluoromount G with Dapi mounting media (Southern Biotech, Birmingham, Alabama, USA). Negative controls were incubated without the primary antibody and showed no staining.

### TUNEL and PCNA analysis

Sections stained with either terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) or the proliferating cell nuclear antigen (PCNA) assay were analyzed for positive stained cells in relation to the total number of cells present. At least 3 animals from each treatment group were serially sectioned. Sections from three areas of the ovary were selected with at least 100  $\mu$ m between sections. Pictures of four different areas of each section were taken at 40 $\times$ . Image J software was utilized to randomly mark 10 regions of interest (ROI) in each area of each section. The total number of cells was determined in these ROIs by nuclear Dapi staining. Percent of proliferation or apoptosis was determined by the average positive stained cells divided by the average total number of cells multiplied by 100.

### PGE<sub>2</sub> EIA

The amount of PGE<sub>2</sub> in ovarian tissues from hens in all groups was measured using a specific enzyme immunoassay as shown previously [24].

### Statistical analysis

All experiments were performed in duplicate and differences in data from groups were analyzed with GraphPad InStat by using One-way ANOVA with Student–Newman–Keuls. A value of  $P < 0.05$  was considered significant whereas a value of  $P < 0.01$  was considered highly significant.

## Results

### Amounts of OM-3FAs and OM-6FAs in egg yolks

The results indicated that there were significantly higher amounts of DHA in yolks collected from hens fed 100 mg/kg ( $P < 0.05$ ) and 175 mg/kg ( $P < 0.01$ ) fish oil compared to amounts of DHA in yolks collected from control hens (Figure 1A). EPA was not detectable in most groups or it was extremely low. The yolks collected from hens fed 100 mg/kg fish oil had significantly lower amounts of AA compared to yolks collected from control hens (Figure 1B;  $P < 0.05$ ).

### Effect of fish oil on COX-1 mRNA and protein expression in ovaries

Comparing the COX-1 protein expression in ovaries of hens revealed that the groups of hens that were fed 100 and 175 mg/kg fish oil had significantly lower COX-1 protein expression compared to ovaries of control hens ( $P < 0.01$ ; Figure 2A). Similar to the protein expression, a significant reduction in expression of COX-1 mRNA was observed in ovaries of the hens fed 100 and 175 mg/kg fish oil compared to control hens ( $P < 0.01$ ; Figure 2B).

### Effect of fish oil on mRNA and protein COX-2 expression in ovaries

A significant decrease in expression of COX-2 protein and mRNA was seen in ovaries of hens fed 100 mg/kg fish oil compared to control hens ( $P < 0.05$ ; Figure 2C & D). The ovaries of hens fed 175 mg/kg fish oil had the lowest COX-2 protein expression among groups which was significantly lower than control hens ( $P < 0.01$ ). There was a significant decrease in COX-2 mRNA expression in ovaries of chickens fed 175 mg/kg fish oil in comparison to control hens ( $P < 0.05$ ; Figure 2D).


### Fish oil consumption decreased PGE<sub>2</sub> concentrations in ovaries

In parallel with the results that we obtained for expression of COX enzymes, significantly lower concentrations of PGE<sub>2</sub> were detected in ovaries of hens fed 100 mg/kg fish oil compared to ovaries of control hens ( $P < 0.001$ ; Figure 3A). Furthermore, feeding the hens with 175 mg/kg fish oil significantly reduced the PGE<sub>2</sub> concentrations in ovaries compared to ovaries of control hens ( $P < 0.01$ ).

### Effect of fish oil on EP4 mRNA and protein expression in ovaries

There were no significant differences in expression of EP4 among groups indicating that fish oil does not affect the expression of EP4 protein and mRNA in ovaries of hens (Figure 3B & D).





**Figure 1** The amounts of OM-3FAs and OM-6FAs in egg yolks. Four egg yolks were collected from each group (n=4). **(A)** More DHA was detected in yolks collected from hens treated with 100 mg/kg (P<0.05) and 175 mg/kg (P<0.01) fish oil compared to control hens. EPA was not detectable in most of the treated groups or it was extremely low. **(B)** The yolks collected from hens treated with 100 mg/kg fish oil had significantly lower amounts of AA compared to yolks collected from control hens (P<0.05). \*, P<0.05; \*\*, P<0.01. Bars indicate standard error.

#### Effect of fish oil on Egg laying frequency

At the end of first week of the study, the ovulation rate (egg laying frequency) of all fish oil-fed hens was compared to control group, no significant differences were seen. Fish oil did not have any effect on ovulation rate of hens compared to control hens in second week (P>0.05; Figure 3C). After being on fish oil for 3 weeks, the egg laying frequency of hens that were fed 100 mg/kg fish oil was significantly increased compared to the ovulation rate of the same group on first week and ovulation rate of control group on third week (P<0.05). The ovulation rate in 700 mg/kg fish oil-fed hens was significantly decreased in weeks two and three compared to week one (P<0.05).


#### Effect of fish oil on apoptosis and proliferation

As shown in Figure 4, 700 mg/kg fish oil increased the rate of apoptosis in ovaries, whereas the other doses failed to show such an effect (P<0.001). The assessment of cell

proliferation by PCNA staining is an indicator of proliferative growth. We observed that the population of PCNA-positive cells in ovary sections was substantially lower than controls when the hen's diet were supplemented with 175 and 700 mg/kg fish oil (P<0.05; Figure 5).

#### Discussion


Here, we report that the consumption of fish oil significantly increases the DHA incorporation into egg yolks, decreases the AA contents of yolks, and reduces both COX-1 and COX-2 protein and mRNA expression in hen ovaries. Moreover, in correlation with down-regulation of COX enzymes, fish oil significantly reduces the concentrations of PGE<sub>2</sub> in ovaries. EP4 protein and mRNA expression in ovaries of hens was not affected by fish oil treatment. A lower dose of fish oil increased the egg laying frequency. 175 and 700 mg/kg fish oil reduced proliferation and 700 mg/kg increased apoptosis in hen



ovaries. The efficacy of fish oil in reducing COX expression and PGE<sub>2</sub> concentration was in “U-shaped curve” manner meaning that the increasing doses of fish oil correlate with reduction of COX and PGE<sub>2</sub> until an optimal dose is reached beyond which further dose increases result in less effect. This finding was correlated to an “inverted U-shaped curve” of DHA amounts found in egg yolks. The concept of “U-shaped curve” refers to a nonlinear relationship between two variables, in this case fish oil dose and prostaglandin production (or COX expression). The nonlinear response, typical of many biological systems, indicates that in the optimum dose-range (100–175 mg/kg) prostaglandin production was maximally inhibited. At higher doses of fish oil the hens react in an adverse way and do not tolerate the treatment.

Incorporation of OM-3FAs into egg yolks of fish oil-fed hens provides a non-invasive measure of the incorporation of OM-3FAs into ovaries of the hens. EPA is normally found in much lesser abundance than DHA in

tissues and no specific physiological response is directly attributable to EPA [4,28]. Equal amounts of DHA and EPA were detected in fish oil (data not shown) but our data showed that EPA was not detectable or was extremely low in egg yolks collected from all groups. It is likely due to its conversion to DHA by  $\Delta^6$ -desaturase. DHA is the most prominent OM-3FA in the egg yolk [29] and we detected higher amounts of it compared to ALA and EPA in yolks. DHA contents of egg yolks collected from hens fed 100 and 175 mg/kg fish oil were significantly higher compared to controls. We did not detect higher amounts of DHA in egg yolks of hens fed higher doses of fish oil. Broughton *et al.* has reported that low EPA/DHA ingestion resulted in nearly as much total OM-3FA incorporation into ovarian tissue as high EPA/DHA ingestion, indicating that ovarian tissue OM-3FA incorporation may be saturable [30]. This may explain why when the fish oil was fed in higher than a certain dose, the amounts of OM-3FAs in the egg was not altered to a




greater extent. DHA contents of yolks of the hens fed 700 mg/kg were lower than DHA content of yolks collected from lower doses. Feeding the hens with 700 mg/kg fish oil for one week resulted in malabsorption of lipids and a diarrhetic response which may explain the lower DHA content of yolks. In general, our data indicated that fish oil increases the incorporation of OM3-FAs into the eggs which was in agreement with previous studies [31,32].

The amounts of AA in yolks collected from 100 mg/kg fish oil-fed hens were lower than the amounts of AA in yolks collected from the control group. It might be due to replacement of AA by DHA in membranes of ovarian cells as suggested by Schreiner *et al.* who reported that after treating hens with OM-3FAs, there was an almost complete exchange of AA and DHA in the *sn*-2 position of phosphatidylcholine and phosphatidylethanolamine in the egg yolk [29]. Meluzzi *et al.* reported that the feeding with fish oil caused a dramatic increase in all OM-3FAs of the yolk and caused an appreciable decrease in AA [32]. They proposed that this phenomenon is probably due to the greater utilization of  $\Delta^6$ -desaturase in the OM-3FA pathway with respect to the OM-6FA

pathway, as this enzyme acts in both pathways [32]. Moreover, they noted that a lower dose of fish oil decreased AA in yolks more than a higher dose. High concentrations of dietary OM-3FAs reduce the activity of the enzyme in the OM-6FAs pathway and the conversion of linoleic acid into AA which could be important for human health, as this acid is a precursor of pro-inflammatory eicosanoids namely PGE<sub>2</sub> [33].

The molecular basis for the health benefits of dietary fish oil is likely due to incorporation of EPA and DHA into the membrane [4]. The fatty acid composition of phospholipids is a key determinant of the ability of COX enzymes to function, and this can be influenced by diet [34]. We measured the expression of COX enzymes in ovaries to see if it is influenced by dietary fish oil. The expression of COX-1 protein and mRNA was significantly decreased in ovaries of 100 and 175 mg/kg fish oil-fed hens. The COX and PGE<sub>2</sub> endpoints are tightly correlated with the incorporation of DHA into the yolks. This was in agreement with the studies that showed the DHA [35] and EPA [4] found in fish oil are effective inhibitors of COX-1. We and others have shown that COX-1 is over-




expressed in ovarian cancer [3,6,7]. The regulation of PGE<sub>2</sub> synthesis by COX-1 in ovarian cancer cell lines [36] has been previously shown. Targeting COX-1 in ovarian cancer is one of the beneficial actions of fish oil which may attenuate and even suppress this disease.

Several studies have shown that OM-3FA enriched diets inhibit COX-2 and prostaglandins in both plasma and experimentally induced tumors [37] but the effect of fish oil on COX-2 expression in ovaries of hens was unknown. Our data indicated that COX-2 expression at the transcriptional level was decreased in ovaries of hens fed 100 and 175 mg/kg of fish oil resulting in decreased COX-2 protein. This was in agreement with a study that reported a significant decrease in COX-2 expression both in the colonic mucosa and in colon tumors of rats fed fish oil [38]. Aronson *et al.* showed that fish oil decreases the COX-2 expression in prostatic tissue of men [39]. The mechanism through which fish oil reduces the

expression of COX-2 remains to be completely elucidated but the inhibitory effect of DHA on COX-2 promoter activity, mRNA levels and protein expression in human endothelial cells has been reported [40]. NF- $\kappa$ B is a transcription factor that plays an important role in various inflammatory signaling pathways and controls effectors enzymes such as COX-2 [41]. Previous studies have shown that fish oil supplementation inhibited NF- $\kappa$ B activation [42,43] which in turn leads to down-regulation of COX-2.

Previous findings have demonstrated that fish oil supplementation in human diets decreases production of PGE<sub>2</sub> [44-47], but the effect of fish oil on PGE<sub>2</sub> concentration in hen ovaries had not been investigated. Our results indicated that there was a statistically significant reduction in concentration of PGE<sub>2</sub> in ovaries of hens fed 100 and 175 mg/kg fish oil compared to controls. The mechanisms by which the OM-3FAs inhibit PGE<sub>2</sub>





**Figure 5 PCNA staining of the ovaries.** The number of proliferative cells in ovary sections in hens fed 175 and 700 mg/kg fish oil was substantially lower than controls ( $P < 0.05$ ). Bars indicate standard error.

remain to be fully elucidated. Dietary OM-3FAs may modulate substrate pools available to COXs and lipoxygenases (LOX), thereby controlling the downstream eicosanoid formation and subsequent receptor activation [48]. COX enzymes convert OM-6FAs to 2-series PG products such as  $PGE_2$  whereas the end products of COX enzymes activity on OM-3FAs are 3-series prostaglandins such as  $PGE_3$ . The 3-series PG products are generally less pro-inflammatory than the 2-series products [49-51]. Prostaglandins are potent mediators of intercellular communication, and high concentrations of  $PGE_2$  are believed to be immunosuppressive of T cell-mediated immunity [52], increase angiogenesis [53], stimulate cell proliferation and inhibit apoptosis in ovarian cancer cell lines [54]. Elevation of  $PGE_2$  in ovarian cancer has been previously reported [55]. We have shown that at the age when ovarian cancer was detected for the first time in hens,  $PGE_2$  concentration and COX expression were elevated in

ovaries [24] pointing to the important role of COX enzymes and  $PGE_2$  in the early stages of ovarian carcinogenesis. Recently, we reported that consumption of flaxseed enriched diet, the richest plant source of OM-3FAs, reduced the ovarian cancer severity and incidence in hens which was correlated to decreased concentration of  $PGE_2$  and expression of COX-2 in ovaries [25,56].

EP4 receptor promotes tumor progression by increasing pro-angiogenic factor and tumor cell invasiveness in ovarian carcinoma cell lines. Spinella *et al.* reported that EP2-EP4 signaling regulates vascular endothelial growth factor production and ovarian carcinoma cell invasion [57]. Our study demonstrated that consumption fish oil does not affect expression of EP4 at the level of mRNA or protein.

COX-2 governs innumerable physiological processes including ovulation.  $PGE_2$  action on granulosa cells is essential for successful ovulation [58]. There was a concern that reduced expression of COX-2 and concentration of

PGE<sub>2</sub> may lead to decreased ovulation rate. We noted a higher number of ovulations in hens fed 100 mg/kg fish oil at the end of third week compared to first week of the study. This finding was in agreement with Broughton *et al.* who has shown that consumption of fish oil resulted in an elevated ovulation rate in rats [30]. Incorporation of OM3-FA appears to enhance ovulation by reducing excess inflammatory PGE<sub>2</sub> possibly through reduced AA availability and decreased COX-2 [30].

One of the hallmarks of cancer is the net gain of cancerous cells, either through excessive proliferation or through failure of programmed cell death (apoptosis). We postulated that fish oil reduces the proliferation and increases the apoptosis of hen ovarian cells. Our data indicated an increase in the rate of apoptosis in ovaries of the hens fed the highest fish oil dose. Proliferation was lower in the group of hens fed lower doses of fish oil. Our findings were in agreement with previous studies which have shown that fish oil increases apoptosis [59] and decreases proliferation in rat mammary tumors [60]; fish oil prevents colon carcinogenesis by enhancing apoptosis in the colon [61,62]. Thus measurement of apoptosis vs proliferation provides an index of therapeutic efficacy. By targeting apoptosis and proliferation, fish oil at the optimum concentration reduces the pro-carcinogenic environment potentially resulting in prevention of cancer initiation.

In conclusion, our findings show that dietary intervention with fish oil decreases the expression of COX-1, COX-2 and the concentration of PGE<sub>2</sub> which likely reduces the carcinogenic microenvironment in ovaries of the laying hens. Fish oil has been reported to be immune-suppressive at high concentrations and immune-stimulatory at lower concentration in chickens [31,63,64]. The reduction in PGE<sub>2</sub> appears to be primarily due to inhibition of both COX enzymes, and it is likely that substrate modulation of both COX enzymes further contributes to the reduction in PGE<sub>2</sub> levels. It is generally agreed that an effective way to control cancer is to find better ways of preventing it. Chemopreventive approaches are recommended for women who have a family history of cancer or express other known risk factors for ovarian cancer [65]. To our knowledge the present study provides the first insight into the efficacy of fish oil on the reduction of COX enzyme expression and PGE<sub>2</sub> concentration in the normal pre-cancerous ovary and further demonstrates the utility of the hen model for ovarian cancer. Our study provides new insight into the potential mechanism of action of fish oil in the reduction of ovarian cancer and will establish the foundation for clinical trials to test the efficacy of dietary intervention with fish oil for the prevention and suppression of ovarian cancer in women.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

EE and DBH designed the experiment and protocols and wrote the manuscript. EE performed animal care and treatment, sample collection, western blot, PCR, PGE<sub>2</sub> assay, statistical analysis. CCS conducted the gas chromatography analysis. SRM performed TUNEL and PCNA. NKK participated in animal handling, treatment and tissue collection and analysis. DBH and EE interpreted the data. All authors read and approved the final manuscript.

#### Acknowledgments

This work was funded by NIH Grant (R03 CA162511). We are grateful for the poultry farm management by Chet Utterback, Pam Utterback and Douglas Hilgendorf. We also thank Dr. Karen Hales for her suggestions and review of this manuscript.

Received: 3 September 2013 Accepted: 18 October 2013

Published: 24 October 2013

#### References

1. Williams CS, Mann M, DuBois RN: The role of cyclooxygenases in inflammation, cancer, and development. *Oncogene* 1999, **18**(55):7908–7916.
2. Gupta RA, Tejada LV, Tong BJ, Das SK, Morrow JD, Dey SK, DuBois RN: Cyclooxygenase-1 is overexpressed and promotes angiogenic growth factor production in ovarian cancer. *Cancer Res* 2003, **63**(5):906–911.
3. Daikoku T, Wang D, Tranguch S, Morrow JD, Orsulic S, DuBois RN, Dey SK: Cyclooxygenase-1 Is a Potential Target for Prevention and Treatment of Ovarian Epithelial Cancer. *Cancer Res* 2005, **65**(9):3735–3744.
4. Smith WL: Cyclooxygenases, peroxide tone and the allure of fish oil. *Curr Opin Cell Biol* 2005, **17**(2):174–182.
5. Denkert C, Kobel M, Pest S, Koch I, Berger S, Schwabe M, Siegart A, Reles A, Klosterhalfen B, Hauptmann S: Expression of cyclooxygenase 2 is an independent prognostic factor in human ovarian carcinoma. *Am J Pathol* 2002, **160**(3):893–903.
6. Urick ME, Johnson PA: Cyclooxygenase 1 and 2 mRNA and protein expression in the Gallus domesticus model of ovarian cancer. *Gynecol Oncol* 2006, **103**(2):673–678.
7. Hales DB, Zhuge Y, Lagman JA, Ansenberger K, Mahon C, Barua A, Luborsky JL, Bahr JM: Cyclooxygenases expression and distribution in the normal ovary and their role in ovarian cancer in the domestic hen (Gallus domesticus). *Endocrine* 2008, **33**(3):235–244.
8. Reese J, Zhao X, Ma WG, Brown N, Maziasz TJ, Dey SK: Comparative analysis of pharmacologic and/or genetic disruption of cyclooxygenase-1 and cyclooxygenase-2 function in female reproduction in mice. *Endocrinology* 2001, **142**(7):3198–3206.
9. Greene ER, Huang S, Serhan CN, Panigrahy D: Regulation of inflammation in cancer by eicosanoids. *Prostaglandins Other Lipid Mediat* 2011, **96**(1–4):27–36.
10. Gupta RA, Dubois RN: Colorectal cancer prevention and treatment by inhibition of cyclooxygenase-2. *Nat Rev Cancer* 2001, **1**(1):11–21.
11. Terada N, Shimizu Y, Kamba T, Inoue T, Maeno A, Kobayashi T, Nakamura E, Kamoto T, Kanaji T, Maruyama T, et al: Identification of EP4 as a potential target for the treatment of castration-resistant prostate cancer using a novel xenograft model. *Cancer Res* 2010, **70**(4):1606–1615.
12. Chell SD, Witherden IR, Dobson RR, Moorghen M, Herman AA, Qualtrough D, Williams AC, Paraskeva C: Increased EP4 receptor expression in colorectal cancer progression promotes cell growth and anchorage independence. *Cancer Res* 2006, **66**(6):3106–3113.
13. Ma X, Kundu N, Rifat S, Walser T, Fulton AM: Prostaglandin E receptor EP4 antagonism inhibits breast cancer metastasis. *Cancer Res* 2006, **66**(6):2923–2927.
14. Ansenberger K, Zhuge Y, Richards C, Barua A, Luborsky JL, Bahr JM, Hales DB: Decreased severity of ovarian cancer and increased survival in hens fed a flaxseed enriched diet for one year. *Gynecol Oncol* 2010, **117**:341–347.
15. Glade MJ: Food, nutrition, and the prevention of cancer: a global perspective. American Institute for Cancer Research/World Cancer Research Fund, American Institute for Cancer Research, 1997. *Nutrition* 1999, **15**(6):523–526.
16. Sharma A, Belna J, Espot J, Rodriguez G, Cannon VT, Hurteau JA: Effects of omega-3 fatty acids on components of the transforming growth factor beta-1 pathway: implication for dietary modification and prevention in ovarian cancer. *Am J Obstet Gynecol* 2009, **200**(5):e511–516.

17. Simopoulos AP: **Omega-3 fatty acids in health and disease and in growth and development.** *Am J Clin Nutr* 1991, **54**(3):438–463.
18. Calviello G, Serini S, Piccioni E: **n-3 polyunsaturated fatty acids and the prevention of colorectal cancer: molecular mechanisms involved.** *Curr Med Chem* 2007, **14**(29):3059–3069.
19. Ansenberger K, Zhuge Y, Lagman JA, Richards C, Barua A, Bahr JM, Hales DB: **E-cadherin expression in ovarian cancer in the laying hen, Gallus domesticus, compared to human ovarian cancer.** *Gynecol Oncol* 2009, **113**(3):362–369.
20. Hales DB, Zhuge Y, Lagman JAJ, Ansenberger K, Mahon C, Narua A, Luborsky J, Bahr JM: **Cyclooxygenases expression and distribution in the normal ovary and their role in ovarian cancer in the domestic hen (Gallus Domesticus).** *Endocrine* 2008. in press.
21. Zhuge Y, Lagman JA, Ansenberger K, Mahon CJ, Daikoku T, Dey SK, Bahr JM, Hales DB: **CYP1B1 expression in ovarian cancer in the laying hen Gallus domesticus.** *Gynecologic oncology* 2009, **112**(1):171–178.
22. Rodriguez-Burford C, Barnes MN, Berry W, Partridge EE, Grizzle WE: **Immunohistochemical expression of molecular markers in an avian model: a potential model for preclinical evaluation of agents for ovarian cancer chemoprevention.** *Gynecol Oncol* 2001, **81**(3):373–379.
23. Barua A, Bitterman P, Abramowicz JS, Dirks AL, Bahr JM, Hales DB, Bradaric MJ, Edassery SL, Rotmensch J, Luborsky JL: **Histopathology of ovarian tumors in laying hens: a preclinical model of human ovarian cancer.** *Int J Gynecol Cancer* 2009, **19**(4):531–539.
24. Eilati E, Pan L, Bahr JM, Hales DB: **Age dependent increase in prostaglandin pathway coincides with onset of ovarian cancer in laying hens.** *Prostaglandins Leukot Essent Fatty Acids* 2012, **87**(6):177–184.
25. Eilati E, Bahr JM, Hales DB: **Long term consumption of flaxseed enriched diet decreased ovarian cancer incidence and prostaglandin E in hens.** *Gynecol Oncol* 2013. DOI:10.1016/j.ygyo.2013.05.018.
26. Trebunova A, Vasko L, Svedova M, Kastel R, Tuckova M, Mach P: **The influence of omega-3 polyunsaturated fatty acids feeding on composition of fatty acids in fatty tissues and eggs of laying hens.** *Dtsch Tierarztl Wochenschr* 2007, **114**(7):275–279.
27. Bautista-Ortega J, Goeger DE, Cherian G: **Egg yolk omega-6 and omega-3 fatty acids modify tissue lipid components, antioxidant status, and ex vivo eicosanoid production in chick cardiac tissue.** *Poult Sci* 2009, **88**(6):1167–1175.
28. Jump DB: **The biochemistry of n-3 polyunsaturated fatty acids.** *J Biol Chem* 2002, **277**(11):8755–8758.
29. Schreiner M, Hulan HW, Razzazi-Fazeli E, Bohm J, Iben C: **Feeding laying hens seal blubber oil: effects on egg yolk incorporation, stereospecific distribution of omega-3 fatty acids, and sensory aspects.** *Poult Sci* 2004, **83**(3):462–473.
30. Broughton KS, Bayes J, Culver B: **High alpha-linolenic acid and fish oil ingestion promotes ovulation to the same extent in rats.** *Nutr Res* 2010, **30**(10):731–738.
31. Ebeid T, Eid Y, Saleh A, Abd El-Hamid H: **Ovarian follicular development, lipid peroxidation, antioxidative status and immune response in laying hens fed fish oil-supplemented diets to produce n-3-enriched eggs.** *Animal* 2008, **2**(1):84–91.
32. Meluzzi A, Sirri F, Manfreda G, Tallarico N, Franchini A: **Effects of dietary vitamin E on the quality of table eggs enriched with n-3 long-chain fatty acids.** *Poult Sci* 2000, **79**(4):539–545.
33. Simopoulos AP: **Human requirement for N-3 polyunsaturated fatty acids.** *Poult Sci* 2000, **79**(7):961–970.
34. Smith WL: **Nutritionally essential fatty acids and biologically indispensable cyclooxygenases.** *Trends Biochem Sci* 2008, **33**(1):27–37.
35. Corey EJ, Shih C, Cashman JR: **Docosahexaenoic acid is a strong inhibitor of prostaglandin but not leukotriene biosynthesis.** *Proc Natl Acad Sci USA* 1983, **80**(12):3581–3584.
36. Kino Y, Kojima F, Kiguchi K, Igarashi R, Ishizuka B, Kawai S: **Prostaglandin E2 production in ovarian cancer cell lines is regulated by cyclooxygenase-1, not cyclooxygenase-2.** *Prostaglandins Leukot Essent Fatty Acids* 2005, **73**(2):103–111.
37. Spencer L, Mann C, Metcalfe M, Webb M, Pollard C, Spencer D, Berry D, Steward W, Dennison A: **The effect of omega-3 FAs on tumour angiogenesis and their therapeutic potential.** *Eur J Cancer* 2009, **45**(12):2077–2086.
38. Singh J, Hamid R, Reddy BS: **Dietary fat and colon cancer: modulation of cyclooxygenase-2 by types and amount of dietary fat during the postinitiation stage of colon carcinogenesis.** *Cancer Res* 1997, **57**(16):3465–3470.
39. Aronson WJ, Gaspy JA, Reddy ST, Reese D, Heber D, Bagga D: **Modulation of omega-3/omega-6 polyunsaturated ratios with dietary fish oils in men with prostate cancer.** *Urology* 2001, **58**(2):283–288.
40. Massaro M, Habib A, Lubrano L, Del Turco S, Lazzzerini G, Bourcier T, Weksler BB, De Caterina R: **The omega-3 fatty acid docosahexaenoate attenuates endothelial cyclooxygenase-2 induction through both NAD(P)H oxidase and PKC epsilon inhibition.** *Proc Natl Acad Sci USA* 2006, **103**(41):15184–15189.
41. Ghosh S, Karin M: **Missing pieces in the NF-kappaB puzzle.** *Cell* 2002, **109**(Suppl):S81–96.
42. Xi S, Cohen D, Chen LH: **Effects of fish oil on cytokines and immune functions of mice with murine AIDS.** *J Lipid Res* 1998, **39**(8):1677–1687.
43. Wang TM, Chen CJ, Lee TS, Chao HY, Wu WH, Hsieh SC, Sheu HH, Chiang AN: **Docosahexaenoic acid attenuates VCAM-1 expression and NF-kappaB activation in TNF-alpha-treated human aortic endothelial cells.** *J Nutr Biochem* 2011, **22**(2):187–194.
44. Endres S, Ghorbani R, Kelley VE, Georgilis K, Lonemann G, van der Meer JW, Cannon JG, Rogers TS, Klempner MS, Weber PC, et al: **The effect of dietary supplementation with n-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells.** *N Engl J Med* 1989, **320**(5):265–271.
45. Caughey GE, Mantzioris E, Gibson RA, Cleland LG, James MJ: **The effect on human tumor necrosis factor alpha and interleukin 1 beta production of diets enriched in n-3 fatty acids from vegetable oil or fish oil.** *Am J Clin Nutr* 1996, **63**(1):116–122.
46. Meydani SN, Endres S, Woods MM, Goldin BR, Soo C, Morrill-Labrode A, Dinarello CA, Gorbach SL: **Oral (n-3) fatty acid supplementation suppresses cytokine production and lymphocyte proliferation: comparison between young and older women.** *J Nutr* 1991, **121**(4):547–555.
47. Swails WS, Kenler AS, Driscoll DF, DeMichele SJ, Babineau TJ, Utsunomiya T, Chavali S, Forse RA, Bistrian BR: **Effect of a fish oil structured lipid-based diet on prostaglandin release from mononuclear cells in cancer patients after surgery.** *JPEN J Parenter Enteral Nutr* 1997, **21**(5):266–274.
48. Moonen HJ, Dommels YE, van Zwam M, van Herwijnen MH, Kleinjans JC, Alink GM, de Kok TM: **Effects of polyunsaturated fatty acids on prostaglandin synthesis and cyclooxygenase-mediated DNA adduct formation by heterocyclic aromatic amines in human adenocarcinoma colon cells.** *Molecular carcinogenesis* 2004, **40**(3):180–188.
49. Kobel M, Reuss A, Bois A, Kommos S, Kommos F, Gao D, Kalloger SE, Huntsman DG, Gilks CB: **The biological and clinical value of p53 expression in pelvic high-grade serous carcinomas.** *J Pathol* 2010, **222**(2):191–198.
50. Wada M, DeLong CJ, Hong YH, Rieke CJ, Song I, Sidhu RS, Yuan C, Warnock M, Schmaier AH, Yokoyama C, et al: **Enzymes and receptors of prostaglandin pathways with arachidonic acid-derived versus eicosapentaenoic acid-derived substrates and products.** *J Biol Chem* 2007, **282**(31):22254–22266.
51. Bagga D, Wang L, Farias-Eisner R, Gaspy JA, Reddy ST: **Differential effects of prostaglandin derived from omega-6 and omega-3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion.** *Proc Natl Acad Sci USA* 2003, **100**(4):1751–1756.
52. Gebauer SK, Psota TL, Harris WS, Kris-Etherton PM: **n-3 fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits.** *Am J Clin Nutr* 2006, **83**(6 Suppl):1526S–1535S.
53. Rodriguez-Leyva D, Dupasquier CM, McCullough R, Pierce GN: **The cardiovascular effects of flaxseed and its omega-3 fatty acid, alpha-linolenic acid.** *Can J Cardiol* 2010, **26**(9):489–496.
54. Munkarah AR, Morris R, Baumann P, Deppe G, Malone J, Diamond MP, Saed GM: **Effects of prostaglandin E(2) on proliferation and apoptosis of epithelial ovarian cancer cells.** *J Soc Gynecol Investig* 2002, **9**(3):168–173.
55. Kushlinskii NE, Podistov Iu I, Laktionov KP, Karseladze AI, Babkina IV, Kerimova GI: **Prostaglandins E in the primary tumor, metastases, and ascitic fluid of patients with ovarian cancer.** *Biull Eksp Biol Med* 1997, **123**(1):83–86.
56. Eilati E, Hales K, Zhuge Y, Ansenberger Fricano K, Yu R, van Breemen RB, Hales DB: **Flaxseed enriched diet-mediated reduction in ovarian cancer severity is correlated to the reduction of prostaglandin E2 in laying hen ovaries.** *Prostaglandins Leukot Essent Fatty Acids* 2013, **89**(4):179–187.
57. Spinella F, Rosano L, Di Castro V, Natali PG, Bagnato A: **Endothelin-1-induced prostaglandin E2-EP2, EP4 signaling regulates vascular endothelial growth factor production and ovarian carcinoma cell invasion.** *J Biol Chem* 2004, **279**(45):46700–46705.

58. Duffy DM, McGinnis LK, Vandervoort CA, Christenson LK: **Mammalian oocytes are targets for prostaglandin E2 (PGE2) action.** *Reprod Biol Endocrinol* 2010, **8**:131.
59. Manna S, Chakraborty T, Ghosh B, Chatterjee M, Panda A, Srivastava S, Rana A: **Dietary fish oil associated with increased apoptosis and modulated expression of Bax and Bcl-2 during 7,12-dimethylbenz(alpha)anthracene-induced mammary carcinogenesis in rats.** *Prostaglandins Leukot Essent Fatty Acids* 2008, **79**(1-2):5-14.
60. Manna S, Chakraborty T, Damodaran S, Samanta K, Rana B, Chatterjee M: **Protective role of fish oil (Maxepa) on early events of rat mammary carcinogenesis by modulation of DNA-protein crosslinks, cell proliferation and p53 expression.** *Cancer Cell Int* 2007, **7**:6.
61. Hong MY, Bancroft LK, Turner ND, Davidson LA, Murphy ME, Carroll RJ, Chapkin RS, Lupton JR: **Fish oil decreases oxidative DNA damage by enhancing apoptosis in rat colon.** *Nutr Cancer* 2005, **52**(2):166-175.
62. Hong MY, Chapkin RS, Davidson LA, Turner ND, Morris JS, Carroll RJ, Lupton JR: **Fish oil enhances targeted apoptosis during colon tumor initiation in part by downregulating Bcl-2.** *Nutr Cancer* 2003, **46**(1):44-51.
63. Wang YW, Field CJ, Sim JS: **Dietary polyunsaturated fatty acids alter lymphocyte subset proportion and proliferation, serum immunoglobulin G concentration, and immune tissue development in chicks.** *Poult Sci* 2000, **79**(12):1741-1748.
64. Fritsche KL, Cassity NA, Huang SC: **Effect of dietary fat source on antibody production and lymphocyte proliferation in chickens.** *Poult Sci* 1991, **70**(3):611-617.
65. Wang D, Dubois RN: **Prostaglandins and cancer.** *Gut* 2006, **55**(1):115-122.

doi:10.1186/1476-511X-12-152

**Cite this article as:** Eilati et al.: Anti-inflammatory effects of fish oil in ovaries of laying hens target prostaglandin pathways. *Lipids in Health and Disease* 2013 **12**:152.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

